

On the resummed gluon anomalous dimension and structure functions at small x

Johannes Blümlein

DESY-Zeuthen, Platanenallee 6, D-15735 Zeuthen, Germany

Andreas Vogt

Institut für Theoretische Physik, Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany

(Received 19 September 1997; published 9 December 1997)

The impact of the recently evaluated “irreducible” contributions to the resummed next-to-leading logarithmic small- x anomalous dimension γ_{gg} is evaluated for the unpolarized parton densities and structure functions of the nucleon. These new terms diminish the gluon distribution and are found to overcompensate the enhancement caused by the resummed leading logarithmic small- x anomalous dimension and the quarkonic contributions beyond next-to-leading order. [S0556-2821(98)50201-9]

PACS number(s): 12.38.Cy, 13.60.Hb

The structure functions of the proton in unpolarized deep-inelastic scattering (DIS) show a steep rise towards very small Bjorken- x values, which becomes stronger with increasing resolution Q^2 . This behavior is a consequence both of the shape of the non-perturbative quark singlet and gluon initial distributions— $x\Sigma$ and xg rise roughly like $x^{-0.2}$ for a starting scale $Q_0^2 \approx 4 \text{ GeV}^2$ —and of the form of the evolution kernels governing the renormalization group equations of the mass singularities. The anomalous dimensions γ_{ij} for the evolution of the parton densities, as well as the Wilson coefficients C_n of the structure functions, contain large logarithmic small- x contributions. In order to arrive at a reliable theoretical framework at very small x , the resummation of these terms may be necessary to all orders in the strong coupling $\alpha_s(Q^2)$. For the unpolarized singlet case considered here the dominant contributions take, in Mellin- N space, the form $\alpha_s^{l+k}/(N-1)^k$, $l \geq 0$.

The resummation of γ_{gg} and γ_{gq} in the leading small- x approximation (Lx), $l=0$, was performed long ago [1]. $\gamma_{gg}^{(0)}(N)$ is obtained as the solution of

$$1 = \frac{\bar{\alpha}_s}{N-1} \chi_0(\gamma_{gg}^{(0)}) \quad (1)$$

with $\bar{\alpha}_s = C_A \alpha_s(Q^2)/\pi$, $C_A = N_c = 3$, $C_F = 4/3$ and

$$\chi_0(\gamma) = 2\psi(1) - \psi(\gamma) - \psi(1-\gamma), \quad (2)$$

furthermore $\gamma_{gq}^{(0)} = (C_F/C_A)\gamma_{gg}^{(0)}$. The quark anomalous dimensions γ_{qq} and γ_{qg} , on the other hand, receive contributions for $l \geq 1$ only. The $l=1$ terms were derived in Ref. [2], together with the corresponding resummations for the coefficient functions C_2 and C_L . The large effects of these quantities on the small- x behavior of the DIS structure functions were subsequently studied in detail [3–7]. In those investigations the small- x resummation of the gluon anomalous dimension γ_{gg} to next-to-leading order small- x (NLx) accuracy could not be taken into account. This resummation has now been performed

for the quarkonic contributions [8–11] proportional to the number of quark flavors N_f . Recently also the “irreducible” gluonic terms $\propto C_A$ have been derived [12], i.e., those contributions which are energy-scale independent in the framework of Ref. [10] underlying that calculation. The corresponding terms of γ_{gq} , however, still remain to be determined.

In this note we investigate the impact of these new resummed contributions to γ_{gg} on the evolution of the parton densities and the proton structure functions $F_2(x, Q^2)$ and $F_L(x, Q^2)$, for the first time including calculated subleading terms into the renormalization group analysis. Hence the comparison of the results to the findings of previous studies [3–7] should allow for improved estimates of the convergence of the small- x resummation approximation, despite a fully quantitative, scheme-independent NLx analysis not being possible at present.

As will be demonstrated below, the effect of the new contributions to the resummed anomalous dimension γ_{gg} is very large and opposite to that of the previously known resummed terms. This implies, already at the present stage, considerable changes particularly for gluon-dominated quantities, which partly modify conclusions obtained in previous numerical investigations [3–6]. A detailed account of the solution of the evolution equations in the presence of all-order anomalous dimensions and coefficient functions will be given in a forthcoming publication [13].

As shown in Ref. [9] the larger eigenvalue of the singlet anomalous dimension matrix, $\gamma_+(N)$, may be obtained in the Q_0 scheme [14] as the solution of

$$1 = \frac{\bar{\alpha}_s}{N-1} [\chi_0(\gamma_+) + \alpha_s \chi_1(\gamma_+)], \quad (3)$$

where the second term is the sum of [9,12]

$$\alpha_s \chi_1^{q\bar{q}} = \frac{N_f \alpha_s}{6\pi} \left[\frac{1}{2} (\chi_0^2(\gamma) + \chi_0'(\gamma)) - \frac{5}{3} \chi_0(\gamma) - \frac{1}{2N_c^2} \left(\frac{\pi}{\sin(\pi\gamma)} \right)^2 \frac{3 \cos(\pi\gamma)}{1-2\gamma} \frac{2+3\gamma(1-\gamma)}{(1+2\gamma)(3-2\gamma)} \right] \quad (4)$$

$$\begin{aligned} \alpha_s \chi_1^{gg} = & \frac{C_A \alpha_s}{4\pi} \left[-\frac{11}{6} (\chi_0^2(\gamma) + \chi_0'(\gamma)) + \left(\frac{67}{9} - \frac{\pi^2}{3} \right) \chi_0(\gamma) + \left(6\zeta(3) + \frac{\pi^2}{3\gamma(1-\gamma)} + \tilde{h}(\gamma) \right) \right. \\ & \left. - \left(\frac{\pi}{\sin(\pi\gamma)} \right)^2 \frac{\cos(\pi\gamma)}{3(1-2\gamma)} \left(11 + \frac{\gamma(1-\gamma)}{(1+2\gamma)(3-2\gamma)} \right) \right]. \end{aligned} \quad (5)$$

The function $\tilde{h}(\gamma)$ in Eq. (5) is given by

$$\tilde{h}(\gamma) = \sum_{k=1}^3 a_k \left(\frac{1}{k+\gamma} + \frac{1}{1+k-\gamma} \right) \quad (6)$$

with $a_1=0.72$, $a_2=0.28$ and $a_3=0.16$ [12]. From these results the irreducible NL x -contribution to γ_{gg} is then inferred by [15]

$$\gamma_{gg}^{(1)} - \frac{\beta_0}{4\pi} \alpha_s^2 \frac{d}{d\alpha_s} \ln(\gamma_{gg}^{(0)} \sqrt{-\chi_0'(\gamma_{gg}^{(0)})}) = \gamma^{+(1)} - \frac{C_F}{C_A} \gamma_{qg}^{(1)} = -\frac{\alpha_s \chi_1(\gamma_{gg}^{(0)})}{\chi_0'(\gamma_{gg}^{(0)})} - \frac{C_F}{C_A} \gamma_{qg}^{(1)}. \quad (7)$$

Our subsequent numerical analysis will be performed in the DIS factorization scheme. Here $\gamma_{gg}^{(1)}$ is represented as

$$\gamma_{gg, \text{DIS}}^{(1)} = \gamma_{gg, \varrho_0}^{(1)} + \frac{\beta_0}{4\pi} \alpha_s^2 \frac{d \ln R(\alpha_s)}{d\alpha_s} + \frac{C_F}{C_A} [1 - R(\alpha_s)] \gamma_{qg, \varrho_0}^{(1)} \quad (8)$$

$$= \alpha_s \sum_{k=1}^{\infty} \left[\frac{N_f}{6\pi} \left(d_{gg,k}^{q\bar{q},(a)} + \frac{C_F}{C_A} d_{gg,k}^{q\bar{q},(b)} \right) + \frac{C_A}{6\pi} d_{gg,k}^{gg} + \frac{\beta_0}{4\pi} \hat{r}_k \right] \left(\frac{\bar{\alpha}_s}{N-1} \right)^{k-1} \equiv \bar{\alpha}_s \sum_{k=0}^{\infty} b_k^{g,(1)} \left(\frac{\bar{\alpha}_s}{N-1} \right)^{k-1}, \quad (9)$$

with $R(\alpha_s)$ defined in Ref. [2] and $\beta_0 = (11/3) C_A - (2/3) N_f$. Tables of the expansion coefficients $d_{gg,k}^{q\bar{q},(a,b)}$, $d_{gg,k}^{gg}$ and \hat{r}_k may be found in Ref. [13]. Here we list for brevity only the numerical values of the first 15 coefficients $b_k^{g,(0)}$ and $b_k^{g,(1)}$ for the L x and NL x series for $N_f=4$, see Table I. Note that the new terms $b_{0,1}^{g,(1)}$ agree with the corresponding results from fixed-order perturbation theory already taking into account the ‘irreducible’ part of $\gamma_{gg}^{(1)}$ only. Collecting all presently available information, the anomalous dimensions for the resummed unpolarized singlet evolution in the DIS scheme are given by

$$\gamma(N, \bar{\alpha}_s)_{\text{DIS}} = \bar{\alpha}_s \gamma_0(N) + \bar{\alpha}_s^2 \gamma_1(N)_{\text{DIS}} + \sum_{k=2}^{\infty} \left(\frac{\bar{\alpha}_s}{N-1} \right)^{k+1} \left[\begin{pmatrix} 0 & 0 \\ C_F/C_A & 1 \end{pmatrix} b_k^{g,(0)} + (N-1) \begin{pmatrix} C_F/C_A & b_k^{q,(1)} \\ 0 & b_k^{g,(1)} \end{pmatrix} \right]. \quad (10)$$

Here γ_0 and γ_1 denote the leading and next-to-leading order singlet anomalous dimension matrices.

The resummed terms beyond $O(\alpha_s^2)$ in Eq. (10) do not comply with the energy-momentum sum rule for the parton densities, which requires

$$\gamma_{qq}(N, \alpha_s) + \gamma_{gq}(N, \alpha_s) = 0, \quad \gamma_{qg}(N, \alpha_s) + \gamma_{gg}(N, \alpha_s) = 0. \quad (11)$$

This relation is satisfied by the fixed-order anomalous dimensions order by order in α_s . The method to restore the sum rule with the least impact on the small- x results is a

(diagonal) subtraction at $N=2$, i.e., the addition of appropriate $\delta(1-x)$ contributions to the higher-order quark-quark and gluon-gluon splitting functions. We will label this prescription as (A) below. Other possibilities are the inclusion of somewhat less singular $1/(N-1)$ terms, later to be superseded by explicit calculations. In this manner a rough estimate can be obtained of the possible effect of subleading small- x contributions to the higher-order anomalous dimensions, particularly in those cases where only the first term is known currently. We will illustrate this procedure by the prescription (D) given by

TABLE I. The expansion coefficients $b_k^{g,(0)}$ and $b_k^{g,(1)}$ for the small- x resummed anomalous dimension γ_{gg} . The latter quantities are given for four active flavors. For comparison to previously employed estimates [5] also the ratios of these coefficients are shown.

k	$b_k^{g,(0)}$	$b_k^{g,(1)}$	$b_k^{g,(1)}/b_k^{g,(0)}$
0	1.000×10^0	-1.139×10^0	-1.14
1	0.000×10^0	-8.519×10^{-1}	
2	0.000×10^0	3.167×10^{-1}	
3	2.404×10^0	-1.166×10^1	-4.85
4	0.000×10^0	-9.104×10^0	
5	2.074×10^0	-1.554×10^1	-7.49
6	1.734×10^1	-1.511×10^2	-8.71
7	2.017×10^0	-1.350×10^2	-66.95
8	3.989×10^1	-4.513×10^2	-11.31
9	1.687×10^2	-2.226×10^3	-13.19
10	6.999×10^1	-2.533×10^3	-36.19
11	6.613×10^2	-1.006×10^4	-15.21
12	1.945×10^3	-3.540×10^4	-18.20
13	1.718×10^3	-5.245×10^4	-30.54
14	1.064×10^4	-2.060×10^5	-19.35

$$\gamma_{ij}(N) \rightarrow \gamma_{ij}(N) (1 - 2[N-1] + [N-1]^3)$$

for $ij = qq, qg, gq$. (12)

The terms $\propto [N-1]$ in γ_{gg} are taken from Eq. (9), hence only the terms $\propto [N-1]^3$ in this quantity are adjusted according to Eq. (11).

We are now ready to discuss the numerical effects of the small- x resummations. For definiteness, we choose the initial distributions of the Martin-Roberts-Stirling [MRS(A')] global fit [16]. Both the gluon and the sea quark densities behave as $x^{-0.17}$ for $x \rightarrow 0$ at the starting scale of $Q_0^2 = 4 \text{ GeV}^2$. The evolution is performed for four massless flavors, and also $\Lambda^{(4)} = 231 \text{ MeV}$ is adopted from the MRS analysis. We stress that these results are mainly theoretical illustrations. Detailed data analyses would require some flexibility of the input gluon density at small x , which is only indirectly constrained by structure function data, as well as the inclusion of heavy-flavor mass effects.

Figure 1 displays the evolution of the proton singlet quark and gluon (momentum) distributions, $x\Sigma(x, Q^2)$ and $xg(x, Q^2)$, in the DIS scheme. Different resummation approximations are compared with the NLO results (full lines). The results with the new $b_k^{g,(1)}$ terms omitted are marked by NLx_q .

In the quark sector the resummation corrections using prescription (A) are very large, e.g., they exceed a factor of four at $x = 10^{-5}$ and $Q^2 = 100 \text{ GeV}^2$. This huge correction is entirely dominated by the quarkonic (upper row) anomalous dimensions. Omitting the $b_k^{g,(1)}$ contributions, and even ignoring the Lx terms (that case is not shown in the figure), has an impact of less than about 20% on the singlet distribution




FIG. 1. The small- x evolution of the singlet quark and gluon densities including the resummed NLx_q kernels [2] and the new terms of the gluon-gluon anomalous dimension [9,12] as compared to the NLO results. Two prescriptions for implementing the momentum sum rule have been applied.




FIG. 2. The small- x behavior of the longitudinal structure function for the parton distributions shown in Fig. 1. The resummed coefficient functions of Ref. [2] have been employed, in (a) without, in (b) with a moderate subleading contribution.

[3,5]. The quark evolution is, however, very much affected by possible subleading contributions to γ_{qg} and γ_{qq} , even by such terms which are smaller than those now found for γ_{gg} . This is illustrated by prescription (D), where the less singular pieces actually overcompensate the effect of the leading $l=1$ contribution.

The effect of the new contribution $\gamma_{gg}^{(1)}$ to the anomalous dimension is, on the other hand, very substantial for the gluon density: the results even fall noticeably below the NLO evolution, and also below our previous lower estimate (D) at NLx_q accuracy [5]. There is no convergence so far, but the analysis of the known LO and NLO results supports some hope that the inclusion also of two further subleading terms (series), $l=2$ and $l=3$, may lead to a sufficiently stable result [13]. At this point the question arises how large a correction the presently unknown resummed $\gamma_{gg}^{(1)}$ term may introduce. Experience in NLO and the Lx and NLx_q resummations suggests that the impact of this contribution is of the order of 10% or less, for the evolution of both $x\Sigma$ and xg .

We now turn to the proton structure functions. As we are working in the DIS scheme, $x\Sigma(x, Q^2)$ already reflects F_2 up to the non-singlet pieces which are not relevant at small x .¹ Hence we turn directly to $F_L(x, Q^2)$. The resummed results, employing the parton densities shown in Fig. 1 and the $l=1$ resummation of the coefficient functions [2],

$$C_L(N) = \bar{\alpha}_s C_L^0(N) + \bar{\alpha}_s^2 C_L^1(N)_{\text{DIS}} + \sum_{k=2}^{\infty} c_{L,k} \bar{\alpha}_s^k \left(\frac{\bar{\alpha}_s}{N-1} \right)^k, \quad (13)$$

are depicted in Fig. 2(a). Here C_L^0 and C_L^1 stand for the leading and next-to-leading order [18] coefficient functions. Note that a full NLx calculation of $F_L(x, Q^2)$ requires the knowledge of the presently unknown next-order resummed coefficient function even in the DIS scheme.

The resummation corrections are exceedingly large at the lower Q^2 values shown. Here they are entirely dominated by the resummed coefficient functions. At very small values of x the resummation does even violate the condition $F_L \leq F_2$ at $Q^2 \simeq 4 \text{ GeV}^2$, thus requiring more terms in the resummation or an adjustment of the input densities at Q_0^2 . At high $Q^2 \gtrsim 100 \text{ GeV}^2$, due to the decrease of α_s and the parton evolution, the effects of the quarkonic anomalous dimensions and the coefficient functions are of the same order. However, besides the anomalous dimensions also the coefficient function will receive subleading corrections, which are presently unknown. To estimate the possible consequences of these terms in C_L mentioned above, Fig. 2(b) repeats the calculation illustrated by Fig. 2(a), but with an estimate for those unknown contributions $C_L \rightarrow C_L(1 - 2[N-1])$ beyond next-to-leading order. As in the quark evolution, already a moderate correction can lead to an even drastic overcompensation of the $l=1$ effect, calling for the evaluation of the next resummation contributions to C_L .

Let us summarize: Recent results in Refs. [9,12] for the first time allow a determination of a subleading

¹The resummation of the small- x terms (Lx) for the non-singlet structure functions was performed in [17]. The corresponding corrections are smaller than 1% over the whole x range.

small- x resummed anomalous dimension, namely the “irreducible” part of γ_{gg} in the unpolarized case. We have extracted the corresponding expansion coefficients in the usual DIS scheme. The coefficients of the subleading small- x poles turn out to be mainly of opposite sign than the Lx pieces, and they are typically much larger in N -space. The numerical impact of these additional terms on the proton’s parton densities and structure functions has been studied. It is largest for the gluon evolution, where a substantial overcompensation of the positive leading resummation effect takes place, but less important for $x\Sigma$ and F_L , which are dominated by the quarkonic anomalous dimensions

and the coefficient functions. Note, however, that the “energy-dependent” contributions still need to be derived. All in all, more terms need to be calculated in the small- x expansions, both of the anomalous dimensions and the coefficient functions, in order to arrive at stable resummation predictions.

We would like to thank M. Ciafaloni, L. Lipatov, S. Catani, and G. Camici for useful discussions. This work was supported in part by the German Federal Ministry for Research and Technology (BMBF) under contract No. 05 7WZ91P (0).

-
- [1] Y. Balitsky and L. Lipatov, *Sov. J. Nucl. Phys.* **28**, 822 (1978).
 - [2] S. Catani and F. Hautmann, *Nucl. Phys.* **B427**, 475 (1994).
 - [3] R. K. Ellis, F. Hautmann, and B. Webber, *Phys. Lett. B* **348**, 582 (1995).
 - [4] R. Ball and S. Forte, *Phys. Lett. B* **351**, 313 (1995); **358**, 365 (1995); J. Forshaw, R. Roberts, and R. Thorne, *ibid.* **356**, 79 (1995).
 - [5] J. Blümlein, S. Riemersma, and A. Vogt, in *QCD and QED in Higher Orders*, Proceedings of the Zeuthen Workshop, Rheinsberg, Germany, 1996, edited by J. Blümlein *et al.* [*Nucl. Phys. B (Proc. Suppl.)* **51C**, 30 (1996)].
 - [6] R. Thorne, *Phys. Lett. B* **392**, 463 (1997); hep-ph/9701241; I. Bojak and M. Ernst, *Phys. Lett. B* **397**, 296 (1997); hep-ph/9702282.
 - [7] J. Blümlein and A. Vogt, hep-ph/9706371, DESY 97–106.
 - [8] S. Catani, M. Ciafaloni, and F. Hautmann, *Nucl. Phys.* **B366**, 135 (1991).
 - [9] G. Camici and M. Ciafaloni, *Phys. Lett. B* **386**, 341 (1996).
 - [10] V. Fadin and L. Lipatov, *Yad. Fiz.* **50**, 1141 (1989); *Nucl. Phys.* **B406**, 259 (1993); **B477**, 767 (1996); V. Fadin, R. Fiore, and A. Quartalo, *Phys. Rev. D* **50**, 2265 (1994); **50**, 5893 (1994); V. Fadin, R. Fiore, and M. Kotsky, *Phys. Lett. B* **359**, 181 (1995); **387**, 593 (1996).
 - [11] V. Del Duca, *Phys. Rev. D* **54**, 989 (1996); **54**, 4474 (1996).
 - [12] G. Camici and M. Ciafaloni, hep-ph/9707390.
 - [13] J. Blümlein and A. Vogt (to appear).
 - [14] M. Ciafaloni, *Phys. Lett. B* **356**, 74 (1995).
 - [15] G. Camici and M. Ciafaloni, *Nucl. Phys.* **B496**, 305 (1997).
 - [16] A. D. Martin, R. G. Roberts, and W. J. Stirling, *Phys. Lett. B* **354**, 155 (1995).
 - [17] J. Blümlein and A. Vogt, *Phys. Lett. B* **370**, 149 (1996); *Acta Phys. Pol. B* **27**, 1309 (1996).
 - [18] E. Zijlstra and W. van Neerven, *Nucl. Phys.* **B383**, 525 (1992); J. Sanchez Guillén *et al.*, *ibid.* **B353**, 337 (1991).