

Nonperturbative determination of anisotropy coefficients in lattice gauge theories

S. Ejiri, Y. Iwasaki, and K. Kanaya

Center for Computational Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

(Received 8 June 1998; published 6 October 1998)

We propose a new nonperturbative method to compute the derivatives of gauge coupling constants with respect to anisotropic lattice spacings (anisotropy coefficients), which are required in an evaluation of thermodynamic quantities from numerical simulations on the lattice. Our method is based on a precise measurement of the finite temperature deconfining transition curve in the lattice coupling parameter space extended to anisotropic lattices by applying the spectral density method. We test the method for the cases of SU(2) and SU(3) gauge theories at the deconfining transition point on lattices with the lattice size in the time direction $N_t=4-6$. In both cases, there is a clear discrepancy between our results and perturbative values. A long standing problem, when one uses the perturbative anisotropy coefficients, is a nonvanishing pressure gap at the deconfining transition point in the SU(3) gauge theory. Using our nonperturbative anisotropy coefficients, we find that this problem is completely resolved: we obtain $\Delta p/T^4=0.001(15)$ and $-0.003(17)$ on $N_t=4$ and 6 lattices, respectively. [S0556-2821(98)05521-0]

PACS number(s): 11.15.Ha, 05.70.Ce, 12.38.Gc, 12.38.Mh

I. INTRODUCTION

In order to study the nature of the quark-gluon plasma in heavy ion collisions and in the early Universe, it is important to evaluate the energy density ϵ and the pressure p near the transition temperature of the deconfining phase transition. These quantities are defined by derivatives of the partition function in terms of the temperature T and the physical volume V of the system:

$$\epsilon = -\frac{1}{V} \frac{\partial \ln Z}{\partial T^{-1}}, \quad p = T \frac{\partial \ln Z}{\partial V}. \quad (1)$$

The lattice formulation of QCD provides us with a nonperturbative way to compute these quantities by numerical simulations. On a lattice with a size $N_s^3 \times N_t$, V and T are given by $V=(N_s a_s)^3$ and $T=1/(N_t a_t)$, with a_s and a_t the lattice spacings in spatial and temporal directions. Because N_s and N_t are discrete parameters, the partial differentiations in Eq. (1) are performed by varying a_s and a_t independently on anisotropic lattices.

The anisotropy on a lattice is realized by introducing different coupling parameters in temporal and spatial directions. For an SU(N_c) gauge theory, the standard plaquette action on an anisotropic lattice is given by

$$S = -\beta_s \sum_{x,i < j \neq 4} P_{ij}(x) - \beta_t \sum_{x,i \neq 4} P_{i4}(x), \quad (2)$$

where $P_{\mu\nu}(x) = (1/N_c) \text{Re Tr} \{ U_\mu(x) U_\nu(x + \hat{\mu}) U_\mu^\dagger(x + \hat{\nu}) U_\nu^\dagger(x) \}$ is the plaquette in the (μ, ν) plane. With this action, the energy density and pressure are given by [1,2]

$$\epsilon = -\frac{3N_t^4 T^4}{\xi^3} \left\{ \left(a_t \frac{\partial \beta_s}{\partial a_t} - \xi \frac{\partial \beta_s}{\partial \xi} \right) (\langle P_s \rangle - \langle P \rangle_0) + \left(a_t \frac{\partial \beta_t}{\partial a_t} - \xi \frac{\partial \beta_t}{\partial \xi} \right) (\langle P_t \rangle - \langle P \rangle_0) \right\}, \quad (3)$$

$$p = \frac{N_t^4 T^4}{\xi^3} \left\{ \xi \frac{\partial \beta_s}{\partial \xi} (\langle P_s \rangle - \langle P \rangle_0) + \xi \frac{\partial \beta_t}{\partial \xi} (\langle P_t \rangle - \langle P \rangle_0) \right\}, \quad (4)$$

where $\langle P_{s(t)} \rangle$ is the space- (time-)like plaquette expectation value and $\langle P \rangle_0$ the plaquette expectation value on a zero-temperature lattice. Here, for later convenience, we have chosen a_t and $\xi \equiv a_s/a_t$ as independent variables to vary the lattice spacings, instead of a_s and ξ adopted in Ref. [2].

In order to compute ϵ and p from Eqs. (3) and (4) using numerical results from simulations, the values for the derivatives of gauge coupling constants with respect to the anisotropic lattice spacings,

$$a_t \frac{\partial \beta_s}{\partial a_t}, \quad a_t \frac{\partial \beta_t}{\partial a_t}, \quad \frac{\partial \beta_s}{\partial \xi}, \quad \frac{\partial \beta_t}{\partial \xi}, \quad (5)$$

which we call the anisotropy coefficients, are required. They can be computed from a requirement that, in the scaling region, the effects of anisotropy in the physical observables can be absorbed by a renormalization of the coupling parameters. Similar to the case of the renormalization group beta-function, the anisotropy coefficients do not depend on the temperature, because the renormalization is independent of the temperature.

The calculation of these anisotropy coefficients in the lowest order perturbation theory has been done by Karsch [2]. However, the perturbative coefficients are known to lead to pathological results such as a negative pressure and a nonvanishing pressure gap at the deconfining transition in SU(3) gauge theory. Therefore, nonperturbative values of the anisotropy coefficients are required in order to study the thermodynamic quantities near the phase transition when N_t is not sufficiently large.

We are interested in the values of the anisotropy coefficients for isotropic lattices ($\beta_s = \beta_t \equiv \beta$, i.e., $\xi = 1$) where most simulations are performed. In this case, we have

$$\left(a_t \frac{\partial \beta_s}{\partial a_t} \right)_{\xi=1} = \left(a_t \frac{\partial \beta_t}{\partial a_t} \right)_{\xi=1} = a \frac{d\beta}{da} = 2N_c a \frac{dg^{-2}}{da},$$

where $a(dg^{-2}/da)$ is the beta function at $\xi=1$, whose non-perturbative values are well studied both in SU(2) and SU(3) gauge theories [3–6]. Furthermore, a combination of the remaining two anisotropy coefficients is known to be related to the beta function [2] by¹

$$\left(\frac{\partial\beta_s}{\partial\xi} + \frac{\partial\beta_t}{\partial\xi}\right)_{a_t: \text{fixed}, \xi=1} = \frac{3}{2} a \frac{d\beta}{da}. \quad (6)$$

Therefore, only one additional input is required to determine the anisotropy coefficients for isotropic lattices.

A nonperturbative determination of the anisotropy coefficients was attempted in Refs. [7–10] using a method that we call ‘‘the matching method’’ in the following. One first determines ξ as a function of β_s and β_t by matching spacelike and timelike Wilson loops on anisotropic lattices, and then numerically determines $\partial\gamma/\partial\xi$ at $\xi=1$, where $\gamma=\sqrt{\beta_t/\beta_s}$. Interpolation of the Wilson loop data at different sizes or interpolation of ξ at different γ 's using an Ansatz is required to evaluate $\partial\gamma/\partial\xi$ at $\xi=1$.

Alternatively, we can evaluate a nonperturbative value of pressure directly from the Monte Carlo data by ‘‘the integral method’’ [11]: Assuming homogeneity expected when the spatial lattice size is sufficiently large, we obtain the relation $p=-f$, where $f=-(T/V)\ln Z$ is the free energy density, which can be evaluated by numerically integrating the plaquette difference $\langle P_s \rangle + \langle P_t \rangle - 2\langle P \rangle_0$ in terms of β on isotropic lattices. The resulting value of the pressure, in turn, provides us with a nonperturbative estimate of an anisotropy coefficient [4,5]. In actual numerical simulations, as the value of p in the confining phase and near the deconfining transition point is quite small compared with the magnitude of errors, it is difficult to determine the anisotropy coefficients

near the transition point [9].

In this paper, we propose a new method to directly compute the anisotropy coefficients at the deconfining transition point. Our method is described in Sec. II. We test the method in the cases of SU(2) gauge theory in Sec. III. The more realistic case of SU(3) gauge theory is studied in Sec. IV. As an application of our nonperturbative anisotropy coefficients, we study the gaps for ϵ and p at the SU(3) deconfining transition for $N_t=4$ and 6. A summary is given in Sec. V.

II. METHOD

Our method is based on an observation that, in the scaling region, the transition temperature $T_c=1/\{N_t a_t(\beta_s, \beta_t)\}$ must be independent of the anisotropy of the lattice. Therefore, when we change the coupling constants along the transition curve in the (β_s, β_t) plane as $(\beta_s, \beta_t) \rightarrow (\beta_s + d\beta_s, \beta_t + d\beta_t)$ on a lattice with fixed N_t , the lattice spacing in the time direction a_t does not change:

$$da_t = \frac{\partial a_t}{\partial \beta_s} d\beta_s + \frac{\partial a_t}{\partial \beta_t} d\beta_t = 0. \quad (7)$$

We denote the slope of the transition curve at $\xi=1$ by r_t ,

$$\begin{aligned} r_t &= \frac{d\beta_s}{d\beta_t} = - \left(\frac{\partial a_t}{\partial \beta_t} \right)_{\xi=1} / \left(\frac{\partial a_t}{\partial \beta_s} \right)_{\xi=1} \\ &= \left(\frac{\partial \beta_s}{\partial \xi} \right)_{\xi=1} / \left(\frac{\partial \beta_t}{\partial \xi} \right)_{\xi=1}, \end{aligned} \quad (8)$$

where we used an identity

$$\begin{pmatrix} \frac{\partial \beta_s}{\partial a_t} & \frac{\partial \beta_t}{\partial a_t} \\ \frac{\partial \beta_s}{\partial \xi} & \frac{\partial \beta_t}{\partial \xi} \end{pmatrix} = \frac{1}{(\partial \xi / \partial \beta_t)(\partial a_t / \partial \beta_s) - (\partial \xi / \partial \beta_s)(\partial a_t / \partial \beta_t)} \begin{pmatrix} \frac{\partial \xi}{\partial \beta_t} & -\frac{\partial \xi}{\partial \beta_s} \\ -\frac{\partial a_t}{\partial \beta_t} & \frac{\partial a_t}{\partial \beta_s} \end{pmatrix}. \quad (9)$$

Hence, the derivatives of β_s and β_t in terms of ξ are expressed as

$$\begin{aligned} \left(\frac{\partial \beta_s}{\partial \xi} \right)_{\xi=1} &= \frac{3r_t}{2(1+r_t)} a \frac{d\beta}{da}, \\ \left(\frac{\partial \beta_t}{\partial \xi} \right)_{\xi=1} &= \frac{3}{2(1+r_t)} a \frac{d\beta}{da}. \end{aligned} \quad (10)$$

Introducing the conventional notation $\gamma=\sqrt{\beta_t/\beta_s}$ and $\beta=\sqrt{\beta_s\beta_t}$, we obtain

$$\left(\frac{\partial \gamma}{\partial \xi} \right)_{a_t: \text{fixed}, \xi=1} = \left(\frac{\partial \gamma}{\partial \xi} \right)_{a_s: \text{fixed}, \xi=1} = \frac{3}{4\beta} \frac{1-r_t}{1+r_t} a \frac{d\beta}{da}. \quad (11)$$

Finally, the customarily used forms for the anisotropy coefficients (Karsch coefficients) [2] are given by

$$c_s = \left(\frac{\partial g_s^{-2}}{\partial \xi} \right)_{a_s: \text{fixed}, \xi=1} = \frac{1}{2N_c} \left\{ \beta + \frac{r_t-2}{2(1+r_t)} a \frac{d\beta}{da} \right\},$$

¹In Ref. [2], a corresponding equation is given for $(\partial\beta_{s(t)}/\partial\xi)_{a_s: \text{fixed}}$.

$$c_t = \left(\frac{\partial g_t^{-2}}{\partial \xi} \right)_{a_s: \text{fixed}, \xi=1} = \frac{1}{2N_c} \left\{ -\beta + \frac{1-2r_t}{2(1+r_t)} a \frac{d\beta}{da} \right\}, \quad (12)$$

where $\beta_s = 2N_c g_s^{-2} \xi^{-1}$ and $\beta_t = 2N_c g_t^{-2} \xi$. Therefore, when the value for the beta function is available, we can determine these anisotropy coefficients by measuring r_t from the finite temperature transition curve in the (β_s, β_t) plane.²

In order to determine the transition curve in the coupling parameter space, we compute the rotated Polyakov loop

$$L = z \frac{1}{N_s^3} \sum_x \frac{1}{N_c} \text{Tr} \prod_{t=1}^{N_t} U_4(\vec{x}, t) \quad (13)$$

as a function of (β_s, β_t) , where z is a $Z(N_c)$ phase factor ($z^{N_c}=1$) such that $\arg(L) \in (-\pi/N_c, \pi/N_c]$. We define the transition point as the peak position of the susceptibility $\chi = N_s^3 (\langle L^2 \rangle - \langle L \rangle^2)$ in β for each fixed γ .

We compute the coupling parameter dependence of χ in the (β_s, β_t) plane by applying the spectral density method [13] extended to anisotropic lattices. This enables us to compute the anisotropy coefficients directly from simulations at $\xi \approx 1$ without introducing an interpolation *Ansatz*. Another good feature of the spectral density method is that the method works well even with data obtained only on isotropic lattices. Therefore, we can use data from previous high statistical simulations performed on isotropic lattices, when the time histories of the Polyakov loop and spacelike and time-like plaquettes are available near the transition point.

Fitting the transition curve with a polynomial

$$\beta_c(\gamma) = \sum_{n=0}^{n_{\max}} f_n (\gamma - 1)^n, \quad (14)$$

with f_n the fitting parameters, the slope r_t is given by

$$r_t = \left(\frac{d(\beta_c/\gamma)}{d(\beta_c \gamma)} \right)_{\xi=1} = \frac{(d\beta_c/d\gamma)_{\xi=1} - \beta_c}{(d\beta_c/d\gamma)_{\xi=1} + \beta_c}, \quad (15)$$

where $(d\beta_c/d\gamma)_{\xi=1} = f_1$. The range of β and γ in which the spectral density method is reliable is estimated by the condition that the statistical error for the reweighting factor (which is $\langle e^{-\Delta S} \rangle$ when the number of simulation points is 1) is less than 0.5%. We confirm that the results are completely stable under a variation of n_{\max} when we restrict ourselves to the range discussed above. Choosing a range of γ around 1 in such a way that the transition curve is almost straight, we use $n_{\max}=3$ for the final results.

III. RESULTS FOR SU(2)

We first test the method for the case of SU(2) gauge theory at the transition point β_c for $N_t=4$ and 5. Although the method should work well with data only from isotropic lattices, in order to confirm it, we perform Monte Carlo




FIG. 1. Polyakov loop susceptibility in the SU(2) gauge theory on a $16^3 \times 4$ lattice at $\gamma=0.995, 1.0$, and 1.005 . Errors are estimated by a jackknife method.

simulations also on several anisotropic lattices for SU(2). On a $16^3 \times 4$ lattice, we perform simulations at $(\beta_s, \beta_t) = (2.300, 2.300), (2.302, 2.302), (2.296, 2.306),$ and $(2.307, 2.298)$. On a $20^3 \times 5$ lattice, we simulate at $(\beta_s, \beta_t) = (2.373, 2.373), (2.375, 2.375), (2.380, 2.370),$ and $(2.368, 2.378)$. At each (β_s, β_t) on the $N_t=4$ (5) lattice, we accumulate 500 000 (1 250 000) configurations, each separated by 10 heat-bath sweeps, after thermalization. The statistical errors are estimated using the jackknife method with the bin size of 1000 configurations. We confirm that the errors are stable under a wide variation of the bin size around this value.

Computing the susceptibility in the (β_s, β_t) plane using data at each simulation point, we check that the results agree well with each other, i.e., the results for the susceptibility from isotropic lattices coincide with the results from anisotropic lattices. For the rest of this section, we combine the results for all four (β_s, β_t) combinations to compute the susceptibility with the spectral density method. In Fig. 1, we plot the susceptibility for $N_t=4$ at $\gamma=0.995, 1.000,$ and 1.005 . The results for the peak position β_c of the susceptibility computed at various values of γ are summarized in Fig. 2 for $N_t=4$ and 5.

Fitting the results for the transition curve, we obtain the values for β_c and r_t at $\xi=1$, as summarized in Table I. Combining the values of r_t with a result of the SU(2) beta function [4] at $\beta_c(\xi=1)$, we obtain the anisotropy coefficients (11) and (12). The results are summarized in Table II. Because no errors for the beta function are given in Ref. [4], we disregard their contribution to the errors of the anisotropy coefficients.

In Fig. 3, we compare our results for the Karsch coefficients with the results of the perturbation theory (dot-dashed curves) [2] and the integral method (dotted curves) [4]. We find significant discrepancies between our results and the results of the perturbation theory. On the other hand, our results are consistent with the results from the integral method.

²A similar approach was proposed in Ref. [12].




FIG. 2. Polyakov loop susceptibility in the SU(2) gauge theory as a function of (β_s, β_t) obtained on (a) $16^3 \times 4$ and (b) $20^3 \times 5$ lattices. Simulation points are shown by filled circles. The bold lines represent the peak position of the susceptibility and the dashed lines their errors. The magnitude of the susceptibility is shown by tone for the range (a) $8.2 < \chi < 10.4$ and (b) $9.0 < \chi < 11.2$, respectively, where different tone corresponds to a difference $\Delta\chi = 0.2$.

TABLE I. Results for β_c and the slopes at $\xi=1$ in the SU(2) gauge theory. The column “ γ range” is for the range of γ used in the fit for the slope $d\beta_c/d\gamma$.

lattice	β_c	γ range	$d\beta_c/d\gamma$	r_t
$16^3 \times 4$	2.30177(9)	0.995–1.005	−0.370(12)	−1.384(14)
$20^3 \times 5$	2.37430(8)	0.995–1.005	−0.312(15)	−1.303(17)

TABLE II. SU(2) anisotropy coefficients at $\xi=1$ using the beta function adg^{-2}/da obtained by the Bielefeld group [4].

lattice	$\partial\gamma/\partial\xi$	c_s	c_t	adg^{-2}/da
$16^3 \times 4$	0.683(21)	0.203(12)	−0.161(12)	−0.08439
$20^3 \times 5$	0.725(35)	0.182(21)	−0.144(21)	−0.07544

IV. RESULTS FOR SU(3)

Let us now study the more realistic case of the SU(3) gauge theory. We analyze the high statistic data for the SU(3) gauge theory obtained by the QCDPAX Collaboration [14]. Simulations were performed at the deconfining transition point for $N_t=4$ and 6. For $N_t=4$, the lattice sizes are $24^2 \times 36 \times 4$ and $12^3 \times 24 \times 4$, with 712 000 and 910 000 pseudo-heat-bath iterations, respectively. For $N_t=6$, data on $36^2 \times 48 \times 6$, $24^3 \times 6$, and $20^3 \times 6$ lattices with 1 112 000, 480 000, and 376 000 iterations are available. The Polyakov loop and the plaquettes are measured every iteration. Details of the simulation parameters are given in Ref. [14]. For the bin size in the jack-knife analysis, we adopt the same values as in Ref. [14].

A. Anisotropy coefficients

The results for the susceptibility on the largest spatial lattices are given in Figs. 4 and 5. Because the transition is of first order for SU(3), the peak of the susceptibility is quite clear when the spatial lattice size is large enough, as shown in Figs. 4 and 5. (Note the difference in the vertical scales between Figs. 1 and 4.)




FIG. 3. Anisotropy coefficients c_s and c_t for the SU(2) gauge theory. Our nonperturbative results are given by filled circles. The dot-dashed curves are the results of the perturbation theory [2]. The dotted curves are the results from the integral method [4]. No errors are published for these curves.




FIG. 4. Polyakov loop susceptibility in the SU(3) gauge theory obtained (a) on the $24^2 \times 36 \times 4$ lattice at $\gamma=0.9975$, 1.0, and 1.0025, and (b) on the $36^2 \times 48 \times 6$ lattice at $\gamma=0.999$, 1.0, and 1.001.

Our results for the slope r_t are summarized in Table III. Except for the case of the $24^3 \times 6$ lattice where the simulation point is slightly off the transition point, the errors become larger with decreasing spatial volume, because the peak of the susceptibility becomes less clear on small lattices. From Table III, we find that the slopes at $N_t=4$ with different spatial lattice volumes completely agree with each other. As shown in Figs. 4 and 5, the peak of the susceptibility for $N_t=6$ is less sharp compared with that for $N_t=4$ with the same relative spatial volume $(N_s/N_t)^3$ due to the fact that the transition is weaker for $N_t=6$ [14]. Therefore, with comparable statistics, r_t has a larger statistical error for $N_t=6$. Unlike in the case of $N_t=4$, the central values for the slope for $N_t=6$ given in Table III vary with the spatial vol-




FIG. 5. The same as Fig. 2 for the SU(3) gauge theory on (a) $24^3 \times 36 \times 4$ and (b) $36^2 \times 48 \times 6$ lattices. The range of χ plotted and the width $\Delta\chi$ for a tone are (a) 0.0–45.0, 5.0 and (b) 2.5–22.5, 2.5, respectively.

ume by about one standard deviation. However, because the volume dependence is not uniform, we consider that it is caused by statistical fluctuations. We use the values obtained on the largest spatial lattices for our final results.

Our results for the anisotropy coefficients are summarized in Table IV. For our final results, we adopt the beta function computed from a recent string tension data by the SCRI group [6]. See a subsection below for a discussion about the influence on the results from the choice of the beta function.

B. Pressure gap and latent heat

As an application of our nonperturbative anisotropy coefficients, we reanalyze the thermodynamic quantities ϵ and p

TABLE III. The same as Table I for SU(3) using the data by the QCDPAX Collaboration [14].

lattice	β_c	γ range	$d\beta_c/d\gamma$	r_t
$24^2 \times 36 \times 4$	5.69245(23)	0.9975–1.0025	–0.5193(23)	–1.2008(10)
$12^2 \times 24 \times 4$	5.69149(42)	0.995–1.005	–0.5183(52)	–1.2004(22)
$36^2 \times 48 \times 6$	5.89379(34)	0.999–1.001	–0.5844(83)	–1.2201(35)
$24^3 \times 6$	5.89292(87)	0.999–1.001	–0.542(33)	–1.202(14)
$20^3 \times 6$	5.8924(14)	0.9975–1.0025	–0.622(34)	–1.236(14)

at the deconfining transition point using the plaquette data by the QCDPAX Collaboration [14]. In terms of the slope r_t and the beta function, the conventional combinations $\epsilon - 3p$ and $\epsilon + p$ are given by

$$(\epsilon - 3p)/T^4 = -3N_t^4 a \frac{d\beta}{da} \{ \langle P_s \rangle + \langle P_t \rangle - 2\langle P \rangle_0 \}, \quad (16)$$

$$(\epsilon + p)/T^4 = 3N_t^4 a \frac{d\beta}{da} \frac{r_t - 1}{r_t + 1} \{ \langle P_s \rangle - \langle P_t \rangle \}. \quad (17)$$

At a first order transition point, we have a finite gap for energy density, the latent heat, but expect no gap for pressure. It is known that the perturbative anisotropy coefficients have a difficulty which leads to a nonvanishing pressure gap at the deconfining transition point: $\Delta p/T^4 = -0.32(3)$ and $-0.14(2)$ at $N_t = 4$ and 6 [14].

New values for the gaps in ϵ and p using our nonperturbative anisotropy coefficients are summarized in Table V. For the pressure gap, we obtain

$$\Delta p/T^4 = \begin{cases} 0.001(15) & \text{for } N_t = 4, \\ -0.003(17) & \text{for } N_t = 6. \end{cases} \quad (18)$$

We find that the problem of nonzero pressure gap is completely resolved with our nonperturbative anisotropy coefficients.

C. Choice of the beta function

In Table IV, we study the influence of the choice of the beta function on the anisotropy coefficients. We compare (i) the beta function computed from a recent string tension data by the SCRI group [6], (ii) that from a Monte Carlo renormalization group (MCRG) study by the QCDTARO Collaboration [3], and (iii) that from a study of $\beta_c(N_t)$ by the Bielefeld group [5]. The SCRI beta function is computed using a fit of the string tension for $5.6 \leq \beta \leq 6.5$. We note that the QCDTARO beta function is based on a fit of mean-field improved gauge coupling constant using the results of plaquette at $\beta > 5.8$; i.e., $\beta_c(N_t = 4) \approx 5.69$ is slightly off the range of validity [3,15]. Also the beta function by the Bielefeld group seems to be problematic around $\beta_c(N_t = 4)$, because it is largely affected by the data of $\beta_c(N_t = 3)$ where we cannot expect universal scaling. Accordingly, the beta-function of the Bielefeld group shows a systematic deviation from the data of a MCRG study at $\beta \leq 6$ [5].

TABLE IV. SU(3) anisotropy coefficients at $\xi = 1$, using the values for the beta function adg^{-2}/da by the SCRI group [6], the QCDTARO Collaboration [3], and the Bielefeld group [5]. For our final results, we take the values obtained on the largest spatial lattices using the SCRI beta function. Because the errors for the beta function are not given in the papers, we disregard their contribution to the errors of the anisotropy coefficients in this table. See text for details.

lattice	$\partial\gamma/\partial\xi$	c_s	c_t	adg^{-2}/da
$24^2 \times 36 \times 4$	0.6159(27)	0.3822(26)	–0.3466(26)	–0.07108 SCRI
	0.5575(25)	0.4359(23)	–0.4037(23)	–0.06434 QCDTARO
	0.6728(30)	0.3299(28)	–0.2910(28)	–0.07764 Bielefeld
$12^2 \times 24 \times 4$	0.6161(62)	0.3819(59)	–0.3464(59)	–0.07097 SCRI
	0.5573(56)	0.4360(53)	–0.4039(53)	–0.06418 QCDTARO
	0.6738(68)	0.3288(64)	–0.2900(64)	–0.07761 Bielefeld
$36^2 \times 48 \times 6$	0.7068(100)	0.3109(98)	–0.2650(98)	–0.09179 SCRI
	0.6936(98)	0.3235(96)	–0.2784(96)	–0.09008 QCDTARO
	0.6826(96)	0.3340(95)	–0.2897(95)	–0.08864 Bielefeld
$24^3 \times 6$	0.762(47)	0.257(46)	–0.211(46)	–0.09172 SCRI
	0.747(46)	0.271(45)	–0.226(45)	–0.08999 QCDTARO
	0.736(45)	0.282(45)	–0.237(45)	–0.08857 Bielefeld
$20^3 \times 6$	0.663(36)	0.354(35)	–0.308(35)	–0.09167 SCRI
	0.651(35)	0.366(35)	–0.321(35)	–0.08994 QCDTARO
	0.640(35)	0.375(34)	–0.331(34)	–0.08853 Bielefeld

TABLE V. Gaps for thermodynamic quantities in the SU(3) gauge theory at the deconfining transition point using our nonperturbative anisotropy coefficients. Plaquette data are taken from Ref. [14]. The low-temperature hadronic phase and the high-temperature quark-gluon-plasma phase are separated as described in Ref. [14]. We reanalyze $(\epsilon - 3p)/T^4$ also, using the SCRI beta function.

lattice	$24^2 \times 36 \times 4$	$36^2 \times 48 \times 6$
β	5.6925	5.8936
$\Delta(\epsilon + p)/T^4$	2.075(42)	1.565(51)
$\Delta(\epsilon - 3p)/T^4$	2.072(43)	1.578(42)
$\Delta\epsilon/T^4$	2.074(34)	1.569(40)
$\Delta p/T^4$	0.001(15)	-0.003(17)

These beta functions are plotted in Fig. 6. At $\beta_c(N_t = 6)$, different beta functions coincide with each other within 5%, while, at $\beta_c(N_t = 4)$, they vary by about 20%. Because only the SCRI beta function is reliable at $\beta_c(N_t = 4)$ as discussed in the previous paragraph, we adopt the SCRI beta function for our final results.

In order to compare the anisotropy coefficients from different references, however, it is important to check the effect of the beta-function on the results. From Table IV, we see that the results for the anisotropy coefficients using different beta-functions agree well with each other at $N_t = 6$. At $N_t = 4$, however, the anisotropy coefficients depend very much on the choice of the beta function. Accordingly, we find that the results for the latent heat are consistent with each other at $N_t = 6$: $\Delta\epsilon/T^4 = 1.569(40)$, $1.539(39)$, and $1.515(38)$ with SCRI, QCDTARO, and Bielefeld beta functions, respectively. At $N_t = 4$, we find a sizable dependence on the choice of the beta function: $\Delta\epsilon/T^4 = 2.074(34)$, $1.877(30)$, and $2.265(37)$ using SCRI, QCDTARO, and Bielefeld beta functions. For the pressure gap, on the other hand, because the beta function appears only as a common overall factor in Eqs. (16) and (17), the conclusion that Δp vanishes with our anisotropy coefficients does not depend on the choice of the beta function.




FIG. 6. Nonperturbative beta functions in the SU(3) gauge theory.




FIG. 7. Anisotropy coefficients in the SU(3) gauge theory. Our nonperturbative results are given by filled circles. The dot-dashed curves are the results of the perturbation theory [2]. The open squares are those from a matching of Wilson loops [9]. Open triangles and thin lines are the results of a matching method [10] combined with the SCRI beta function [6]. The dotted curves are the results from the integral method [5]. No errors are published for the results from the integral method.

D. Comparison with other methods

In Fig. 7, we summarize our results for the Karsch coefficients together with previous values; the perturbative results [2], results from the integral method [5], and those from the matching of Wilson loops on anisotropic lattices [9,10]. No errors are published for the results from the integral method. We find that all nonperturbative methods give values which deviate from the results in the perturbation theory.

Comparing the results from different nonperturbative methods, we find that, although the deviations from the perturbation theory are roughly consistent with each other, the central values are different by more than three standard deviations, when we take the published errors.

We think that one origin of the variation among different methods at $\beta_c(N_t = 4)$ is the beta function. Note that the results from Refs. [9] (matching method) and [5] (integral method) are computed using the beta function of the Bielefeld group, while our results and the results from Ref. [10] (matching method) are using the SCRI beta function. From Table IV, we note that, if we adopt the beta-function of the Bielefeld group, our results are consistent with those of Ref. [9] at $\beta_c(N_t = 4)$.

At $\beta_c(N_t = 6)$, on the other hand, the difference in the results is not due to the beta function, because the systematic error due to the choice of the beta function is small as discussed in the previous subsection. In order to see this, we study $\partial\gamma/\partial\xi$, which can be computed without using the beta function in the matching method. The values of $\partial\gamma/\partial\xi$ ob-

tained in Ref. [10] are reported to be consistent with those from the integral method [5], but are different to another result from the matching method [9]. Performing a quadratic interpolation in β , we find $\partial\gamma/\partial\xi \approx 0.64(1)$ [10], $0.66(2)$ [5], and $0.74(2)$ [9] at $\beta_c(N_t=6)$. Our result $0.707(10)$ given in Table IV is around the center of these values. A careful study of systematic errors in each method is required to understand the variation between different methods.

V. CONCLUSIONS

We have computed the anisotropy coefficients for the SU(2) and SU(3) gauge theories by measuring the transition curve of the deconfining transition in the (β_s, β_t) plane. One of the essential ingredients of our approach is the application of the spectral density method, which enables us to determine the anisotropy coefficients directly from simulations at $\xi \approx 1$. We note that the spectral density method is useful to avoid interpolation *Ansätze* also in the matching method.

Our nonperturbative results for the anisotropy coefficients are summarized in Tables II and IV. Our results shown in

Fig. 7 suggest that the Karsch coefficients converge to the perturbative values slightly faster than that suggested by the central values from Refs. [5] and [10]. Applying the results for SU(3), we reanalyzed the thermodynamic quantities at the deconfining transition point on $N_t=4$ and 6 lattices. We obtain vanishing pressure gaps with our nonperturbative anisotropy coefficients, thereby solving a long standing problem of nonzero pressure gap with the perturbative coefficients.

ACKNOWLEDGMENTS

We are grateful to O. Miyamura, A. Nakamura and H. Matsufuru for useful discussions and sending us the data for the QCDTARO beta function. We also thank A. Ukawa, T. Yoshié, Y. Aoki, T. Kaneko, R. Burkhalter, and H. P. Shanahan for helpful suggestions and comments. This work is in part supported by the Grants-in-Aid of Ministry of Education, Science and Culture (Grant Nos. 08NP0101 and 09304029). SE is supported by the Japan Society for the Promotion of Science.

-
- [1] J. Engels, F. Karsch, H. Satz, and I. Montvay, Nucl. Phys. **B205**, 545 (1982).
 - [2] F. Karsch, Nucl. Phys. **B205**, 285 (1982).
 - [3] K. Akemi *et al.*, Phys. Rev. Lett. **71**, 3063 (1993).
 - [4] J. Engels, F. Karsch, and K. Redlich, Nucl. Phys. **B435**, 295 (1995).
 - [5] G. Boyd *et al.*, Nucl. Phys. **B469**, 419 (1996).
 - [6] R. G. Edwards, U. M. Heller, and T. R. Klassen, Nucl. Phys. **B517**, 377 (1998).
 - [7] G. Burgers, F. Karsch, A. Nakamura, and I. O. Stamatescu, Nucl. Phys. **B304**, 587 (1988).
 - [8] M. Fujisaki *et al.*, Nucl. Phys. B (Proc. Suppl.) **53**, 426 (1997).
 - [9] J. Engels, F. Karsch, and T. Scheideler, Nucl. Phys. B (Proc. Suppl.) **63**, 427 (1998).
 - [10] T. R. Klassen, hep-lat/9803010.
 - [11] J. Engels *et al.*, Phys. Lett. B **252**, 625 (1990).
 - [12] I. Montvay and E. Pietarinen, Phys. Lett. **110B**, 148 (1982).
 - [13] A. M. Ferrenberg and R. H. Swendsen, Phys. Rev. Lett. **61**, 2635 (1988); **63**, 1195 (1989).
 - [14] Y. Iwasaki, K. Kanaya, T. Yoshié, T. Hoshino, T. Shirakawa, Y. Oyanagi, S. Ichii, and T. Kawai, Phys. Rev. D **46**, 4657 (1992).
 - [15] O. Miyamura and A. Nakamura (private communication).