

Probing the nonperturbative dynamics of the SU(2) vacuum

Paolo Cea*

*Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy
and INFN - Sezione di Bari, I-70126 Bari, Italy*

Leonardo Cosmai†

INFN - Sezione di Bari, I-70126 Bari, Italy

(Received 3 March 1999; revised manuscript received 9 July 1999; published 7 October 1999)

The vacuum dynamics of SU(2) lattice gauge theory is studied by means of a gauge-invariant effective action defined using the lattice Schrödinger functional. Numerical simulations are performed at zero temperature. The vacuum is probed using an external constant Abelian chromomagnetic field. The results suggest that the external field is screened in the continuum limit. [S0556-2821(99)04021-7]

PACS number(s): 11.15.Ha

I. INTRODUCTION

It is widely recognized that the effective action is a useful tool to investigate the quantum properties of field theories. In the case of gauge theories when including the quantum fluctuations one faces the problem of retaining in the effective action the gauge invariance that is manifest at the classical level. In the perturbative approach, however, the problem of the gauge invariance of the effective action is not so compelling. Indeed in order to perform the perturbative calculations we need to fix the gauge so that the gauge invariance is lost anyway. Obviously the physical quantities turn out to be gauge invariant. In the case of the perturbative evaluation of the effective action the problem of the gauge invariance can be efficiently resolved by the so-called method of the background effective action [1–3]. In the background field approach one separates the quantum field into the fluctuations $\eta(x)$ and the background field $\bar{A}_\mu^a(x)$. In order to define the background field effective action we introduce the partition function by coupling the external current to the fluctuations. Using the background field gauge fixing it is easy to see that the partition function is invariant against gauge transformation of the background field. In this way, after performing the usual Legendre transformation, one obtains an effective action which is invariant for background field gauge transformations.

The lattice approach to gauge theories allows the nonperturbative study of gauge systems without losing the gauge invariance. Thus, it is natural to seek for a lattice definition of the effective action. Previous attempts (both in three [4,5] and four [6–8] dimensions) in this direction introduced the background field by means of an external current coupled to the lattice gauge field. It turns out, however, that the current term added to the lattice gauge action in general is not invariant under the local gauge transformations belonging to the gauge group. For instance, if one considers Abelian background fields then the action with the current term turns out

to be invariant only for an Abelian subgroup of the gauge group. So that only in the case of Abelian U(1) gauge theory the lattice background field action is gauge invariant.

The aim of the present paper is to discuss in detail a recently proposed method [9] to define on the lattice the gauge-invariant effective action by using the so-called Schrödinger functional [10–12].

Let us consider the continuum Euclidean Schrödinger functional in Yang-Mills theories without matter field:

$$\mathcal{Z}[A^{(f)}, A^{(i)}] = \langle A^{(f)} | e^{-HT} \mathcal{P} | A^{(i)} \rangle. \quad (1)$$

In Eq. (1) H is the pure gauge Yang-Mills Hamiltonian in the fixed-time temporal gauge, T is the Euclidean time extension, while \mathcal{P} projects onto the physical states. $A_k^{a(i)}(\vec{x})$ and $A_k^{a(f)}(\vec{x})$ are static classical gauge fields, and the state $|A\rangle$ is such that

$$\langle A | \Psi \rangle = \Psi[A]. \quad (2)$$

From Eq. (1), inserting an orthonormal basis $|\Psi_n\rangle$ of gauge-invariant energy eigenstates, it follows:

$$\mathcal{Z}[A^{(f)}, A^{(i)}] = \sum_n e^{-E_n T} \Psi_n[A^{(f)}] \Psi_n^*[A^{(i)}]. \quad (3)$$

Note that we are interested in the lattice version of the Schrödinger functional, so that it makes sense to perform a discrete sum in Eq. (3) for the spectrum is discrete in a finite volume. Equation (3) shows that the Schrödinger functional is invariant under arbitrary gauge transformations of the fields $A^{(f)}$ and $A^{(i)}$.

Using standard formal manipulations and the gauge invariance of the Schrödinger functional it is easy to rewrite $\mathcal{Z}[A^{(f)}, A^{(i)}]$ as a functional integral [10,11]

$$\mathcal{Z}[A^{(f)}, A^{(i)}] = \int \mathcal{D}A e^{-\int_0^T dx_4 \int d^3x \mathcal{L}_{YM}(x)} \quad (4)$$

with the constraints

$$A_\mu(x_0=0) = A_\mu^{(i)}, \quad (5)$$

*Electronic address: cea@bari.infn.it

†Electronic address: cosmai@bari.infn.it

$$A_\mu(x_0=T) = A_\mu^{(f)}.$$

Strictly speaking we should include in Eq. (4) the sum over topological inequivalent classes. However, it turns out that [12] on the lattice such an average is not needed because the functional integral Eq. (4) is already invariant under arbitrary gauge transformations of $A_\mu^{(i)}$ and $A_\mu^{(f)}$.

On the lattice the natural relation between the continuum gauge fields and the corresponding lattice links is given by

$$U_\mu = P \exp \left\{ iag \int_0^1 dt A_\mu(x + at\hat{\mu}) \right\}, \quad (6)$$

where P is the path-ordering operator, a is the lattice spacing, and g is the gauge coupling constant.

The lattice implementation of the Schrödinger functional, Eq. (4), is now straightforward:

$$\mathcal{Z}[U^{(f)}, U^{(i)}] = \int \mathcal{D}U e^{-S}. \quad (7)$$

In Eq. (7) the functional integration is done over the links $U_\mu(x)$ with the fixed boundary values:

$$U(x)|_{x_4=0} = U^{(i)}, \quad U(x)|_{x_4=T} = U^{(f)}. \quad (8)$$

Moreover the action S is the standard Wilson action modified to take into account the boundaries at $x_4=0, T$ [12]:

$$S = \frac{1}{g^2} \sum_{x, \mu > \nu} W_{\mu\nu}(x) \text{Tr}[1 - U_{\mu\nu}(x)], \quad (9)$$

where $U_{\mu\nu}(x)$ are the plaquettes in the (μ, ν) plane and

$$W_{\mu\nu}(x) = \begin{cases} 1/2 & \text{spatial plaquettes at } x_4=0, T, \\ 1 & \text{otherwise.} \end{cases} \quad (10)$$

Moreover, it is possible to improve the lattice action S by modifying the weights $W_{\mu\nu}$'s [12]. Note that, due to the fact that $U^{(i)} \neq U^{(f)}$, one cannot impose periodic boundary conditions in the Euclidean time direction. On the other hand, one can assume periodic boundary conditions in the spatial directions.

Let us consider, now, a static external background field $\vec{A}^{\text{ext}}(\vec{x}) = \vec{A}_a^{\text{ext}}(\vec{x}) \lambda_a/2$, where $\lambda_a/2$ are the generators of the $SU(N)$ Lie algebra. We introduce, now, a new functional

$$\Gamma[\vec{A}^{\text{ext}}] = -\frac{1}{T} \ln \left\{ \frac{\mathcal{Z}[U^{\text{ext}}]}{\mathcal{Z}(0)} \right\}, \quad (11)$$

where

$$\mathcal{Z}[U^{\text{ext}}] = \mathcal{Z}[U^{\text{ext}}, U^{\text{ext}}], \quad (12)$$

and $\mathcal{Z}[0]$ means the Schrödinger functional Eq. (12) without external background field ($U_\mu^{\text{ext}} = 1$). The lattice link U^{ext} is obtained from the continuum background field \vec{A}^{ext} through Eq. (6).

From the previous discussion it is clear that $\Gamma[\vec{A}^{\text{ext}}]$ is invariant for lattice gauge transformations of the external links U_μ^{ext} . Moreover, from Eq. (3) it follows that

$$\lim_{T \rightarrow \infty} \Gamma[\vec{A}^{\text{ext}}] = E_0[\vec{A}^{\text{ext}}] - E_0[0], \quad (13)$$

where $E_0[\vec{A}^{\text{ext}}]$ is the vacuum energy in presence of the external background field. In other words $\Gamma[\vec{A}^{\text{ext}}]$ is the lattice gauge-invariant effective action for the static background field \vec{A}^{ext} . In particular, if we consider background fields that give rise to constant field strength, then due to the gauge invariance it is easy to show that $\Gamma[\vec{A}^{\text{ext}}]$ is proportional to the spatial volume V . In this case one is interested in the density of the effective action:

$$\varepsilon[\vec{A}^{\text{ext}}] = -\frac{1}{\Omega} \ln \left[\frac{\mathcal{Z}[\vec{A}^{\text{ext}}]}{\mathcal{Z}[0]} \right], \quad (14)$$

where $\Omega = V \cdot T$. We stress that our definition of the lattice effective action uses the lattice Schrödinger functional with the same boundary fields at $x_4=0$ and $x_4=T$. So that we can glue the two hyperplanes $x_4=0$ and $x_4=T$ together. This way we end up in a lattice with periodic boundary conditions in the time direction too. Therefore, our lattice Schrödinger functional turns out to be

$$\mathcal{Z}[U^{\text{ext}}] = \int \mathcal{D}U e^{-S}, \quad (15)$$

where the functional integral is defined over a four-dimensional hypertorus with the ‘‘cold-wall’’

$$U_\mu(x)|_{x_4=0} = U_\mu^{\text{ext}}. \quad (16)$$

Moreover, due to the lacking of free boundaries, the lattice action in Eq. (15) is now the familiar Wilson action

$$S = S_W = \frac{1}{g^2} \sum_{x, \mu > \nu} \text{Tr}[1 - U_{\mu\nu}(x)]. \quad (17)$$

In this paper we study the properties of the gauge-invariant lattice effective action in pure gauge non-Abelian theories. In particular, we consider the $SU(2)$ gauge theory in the presence of constant Abelian chromomagnetic field at zero temperature. The plan of the paper is as follows. In Sec. II we consider the $SU(2)$ gauge theory on the lattice in presence of constant Abelian chromomagnetic background field. Section III is devoted to the discussion of the Nielsen-Olesen instability on the lattice. In Sec. IV we present the numerical results of the Monte Carlo simulations [13]. Finally our conclusions are drawn in Sec. V.

II. $SU(2)$ IN A CONSTANT ABELIAN CHROMOMAGNETIC FIELD

In this paper we are interested in the case of a constant Abelian chromomagnetic field. Let us consider the $SU(2)$ gauge theory. In the continuum we have

$$\vec{A}_a^{\text{ext}}(\vec{x}) = \vec{A}^{\text{ext}}(\vec{x}) \delta_{a,3}, \quad \vec{A}_k^{\text{ext}}(\vec{x}) = \delta_{k,2} x_1 H. \quad (18)$$

The external links corresponding to $\vec{A}_a^{\text{ext}}(\vec{x})$ are easily evaluated from Eq. (6):

$$U_1^{\text{ext}}(\vec{x}, 0) = U_3^{\text{ext}}(\vec{x}, 0) = U_4^{\text{ext}}(\vec{x}, 0) = \mathbf{1}, \quad (19)$$

$$U_2^{\text{ext}}(\vec{x}, 0) = \cos\left(\frac{agHx_1}{2}\right) + i\sigma^3 \sin\left(\frac{agHx_1}{2}\right).$$

Our Schrödinger functional $\mathcal{Z}[\vec{A}^{\text{ext}}]$ is defined on a lattice with periodic boundary conditions, so that we impose that

$$U_2(x_1, x_2, x_3, x_4) = U_2(x_1 + L_1, x_2, x_3, x_4), \quad (20)$$

where L_1 is the lattice extension in the x_1 direction (in lattice units). As a consequence the magnetic field H turns out to be quantized

$$\frac{a^2 g H}{2} = \frac{2\pi}{L_1} n_{\text{ext}}, \quad (21)$$

with n_{ext} integer.

According to our previous discussion in evaluating the lattice functional integral Eq. (15) we impose that the links belonging to the time slice $x_4 = 0$ are frozen to the configuration Eq. (19). Moreover we impose also that the links at the spatial boundaries are fixed according to Eq. (19). In the continuum this last condition amounts to the usual requirement that the fluctuations over the background fields vanish at infinity.

Note that the background links must be consistent with the lattice geometry, and this has been achieved by imposing Eq. (20). On the other hand, the fluctuations $\eta(x)$ by definition satisfy the boundary conditions $\eta(x_4 = 0) = \eta(x_4 = L_4) = 0$. Therefore, the geometry of the lattice is immaterial as far as the dynamics of the fluctuating fields is concerned.

An alternative possibility is given by constraining the links belonging to the time slice $x_4 = 0$ and those at the spatial boundaries to the condition

$$U_2(x) = U_2^{\text{ext}}(\vec{x}), \quad (22)$$

while the links $U_\mu(x)$ with $\mu \neq 1, 2$ are unconstrained. The main advantage of the condition (22) resides in the fact that the timelike plaquettes nearest the frozen hypersurface $x_4 = 0$ behave symmetrically in the update procedure. Obviously in the thermodynamic limit both conditions should agree as the effective action is concerned.

III. THE NIELSEN-OLESEN INSTABILITY ON THE LATTICE

As it is well known in the continuum the perturbative evaluation of the effective action for the constant Abelian chromomagnetic field faces with the problem of Nielsen-Olesen unstable modes [14]. Let us briefly discuss the origin of the unstable modes in the continuum.

In order to evaluate the effective action in the continuum one writes

$$A_\mu^a(x) = \bar{A}_\mu^a(x) + \eta_\mu^a(x), \quad (23)$$

where $\bar{A}_\mu^a(x) = \delta_{\mu 2} \delta^{a3} x_1 H$ and $\eta_\mu^a(x)$ is the quantum fluctuation over the background field. In the background gauge

$$[\delta^{ab} \partial_\mu - g^{abc} \bar{A}_\mu^c(x)] \eta_\mu^b(x) = 0. \quad (24)$$

We rewrite the pure gauge action in the one-loop approximation as

$$S = S_{\text{class}} + \frac{1}{2} \int d^4x \eta_\mu^a(x) \mathcal{O}_{\mu\nu}^{ab}(x). \quad (25)$$

The one-loop effective action can be obtained by performing the Gaussian integration over the quantum fluctuations and including the Faddeev-Popov determinant. However, if we solve the eigenvalue equation

$$\mathcal{O}_{\mu\nu}^{ab} \phi_\nu^b(x) = \lambda \phi_\mu^a(x), \quad (26)$$

then we find that there are negative eigenvalues:

$$\lambda_u = p_0^2 + p_3^2 - gH. \quad (27)$$

As a matter of fact $\lambda_u < 0$ when $gH > p_0^2 + p_3^2$. If we perform formally the Gaussian functional integration in the one-loop approximation then the effective action picks up an imaginary part. The point is that in the functional integration over the unstable modes one must include the positive quartic term. It turns out that the unstable modes behave like a two-dimensional tachyonic charged scalar field. Thus the dynamics of the unstable modes resemble the dynamical Higgs mechanism. As a consequence the response of the gauge system to the external field turns out to be strong even in the nominally perturbative regime [15].

In order to ascertain if the Nielsen-Olesen one-loop instability survives the lattice regularization one should evaluate the Schrödinger functional Eq. (12) in the weak-coupling region. To this end we write the lattice version of Eq. (23):

$$U_\mu(x) = \exp(iagq_\mu(x)) U_\mu^{\text{ext}}, \quad (28)$$

where the fluctuations $q_\mu(x) = q_\mu^a(x) \sigma^a / 2$ satisfy the boundary condition

$$q_\mu(x)|_{x_4=0} = 0. \quad (29)$$

Inserting Eq. (28) into the plaquette

$$U_{\mu\nu}(x) = U_\mu(x) U_\nu(x + \hat{\mu}a) U_\mu^\dagger(x + \hat{\nu}a) U_\nu^\dagger(x), \quad (30)$$

we rewrite the Wilson action

$$S_W = \frac{4}{g^2} \sum_x \sum_{\mu > \nu} \left[1 - \frac{1}{2} \text{tr} U_{\mu\nu}(x) \right] \quad (31)$$

in the quadratic approximation as

$$S_W = S^{\text{ext}} + S^{(2)}, \quad (32)$$

where

$$S^{\text{ext}} = \frac{4\Omega}{g^2} \left[1 - \cos\left(\frac{gHa^2}{2}\right) \right], \quad (33)$$

with $\Omega = L_1 \times L_2 \times L_3 \times L_4$ the lattice volume. Note that the

external action S^{ext} in the naive continuum limit reduces to the classical action

$$S^{\text{cl}} = VT \frac{H^2}{2}. \quad (34)$$

As concerns the quadratic action $S^{(2)}$, following the method of Ref. [16] a standard calculation gives

$$\begin{aligned} S^{(2)} = & a^4 \sum_{x, \mu > \nu} \text{Tr}\{[D_\mu q_\nu(x) - D_\nu q_\mu(x)]^2 U_{\mu\nu}^{\text{ext}}(x)\} - a^4 \sum_{x, \mu > \nu} \text{Tr}\{[D_\nu q_\mu(x), D_\mu q_\nu(x)] U_{\mu\nu}^{\text{ext}}(x)\} \\ & - 2a^2 \sum_{x, \mu > \nu} \text{Tr}\{[q_\mu(x), q_\nu(x)] U_{\mu\nu}^{\text{ext}}(x)\} - a^3 \sum_{x, \mu > \nu} \text{Tr}\{([D_\nu q_\mu(x), q_\mu(x)] - [D_\mu q_\nu(x), q_\nu(x)]) U_{\mu\nu}^{\text{ext}}(x)\}, \end{aligned} \quad (35)$$

where D_μ is the lattice covariant derivative in the external background $U_\mu^{\text{ext}}(x)$:

$$D_\mu f(x) = \frac{1}{a} [U_\mu^{\text{ext}}(x) f(x + \hat{\mu}a) U_\mu^{\text{ext}\dagger}(x) - f(x)]. \quad (36)$$

Observing that

$$U_{\mu\nu}^{\text{ext}} = G_{\mu\nu} + iH_{\mu\nu} \quad (37)$$

with

$$G_{\mu\nu} = \begin{cases} \mathbf{1} & (\mu, \nu) \neq (1, 2), \\ \cos\left(\frac{a^2 g B}{2}\right) & (\mu, \nu) = (1, 2), \end{cases} \quad (38)$$

$$H_{\mu\nu} = \begin{cases} 0 & (\mu, \nu) \neq (1, 2), \\ \sin\left(\frac{a^2 g B}{2}\right) & (\mu, \nu) = (1, 2), \end{cases} \quad (39)$$

and integrating by parts, we rewrite the quadratic action as

$$\begin{aligned} S^{(2)} = & 2a^4 \sum_{x, \mu < \nu} G_{\mu\nu} \text{Tr}[q_\mu(x) (D_\nu^* D_\mu - \delta_{\mu\nu} D_\sigma^* D_\sigma) q_\nu(x)] \\ & + 2ia^4 \sum_{x, \mu > \nu} \text{Tr}[q_\mu(x) D_\nu^* D_\mu q_\nu(x) H_{\mu\nu}(x)] \\ & + 2ia^3 \sum_{x, \mu > \nu} \text{Tr}[q_\mu(x) (D_\nu^* + D_\nu) q_\mu(x) H_{\mu\nu}(x)] \\ & - 4ia^2 \sum_{x, \mu > \nu} \text{Tr}[q_\nu(x) q_\mu(x) H_{\mu\nu}(x)], \end{aligned} \quad (40)$$

where

$$\begin{aligned} D_\nu^* f(x) = & \frac{1}{a} [f(x) - (U_\mu^{\text{ext}})^{-1}(x - a\hat{\mu}) f(x - a\hat{\mu}) \\ & \times U_\mu^{\text{ext}}(x - a\hat{\mu})]. \end{aligned} \quad (41)$$

Taking into account that

$$q_\mu(x) = q_\mu^a(x) \frac{\sigma^a}{2} \quad (42)$$

and using Eq. (19), we perform the trace over the color indexes. After a rather long but otherwise elementary calculation we get

$$S^{(2)} = S^{(2)}(q^3) + S^{(2)}(q^+, q^-), \quad (43)$$

where

$$q_\mu^\pm(x) = q_\mu^1(x) \pm q_\mu^2(x). \quad (44)$$

We have

$$\begin{aligned} S^{(2)}(q^3) = & \frac{a^4}{2} \sum_x \sum_{\mu\nu} G_{\mu\nu} [q_\mu^3(x) \Delta_\nu^* \Delta_\mu q_\nu^3(x) \\ & - q_\nu^3(x) \Delta_\mu^* \Delta_\nu q_\mu^3(x)], \end{aligned} \quad (45)$$

where

$$\Delta_\mu f(x) = \frac{1}{a} [f(x + \hat{\mu}a) - f(x)], \quad (46)$$

$$\Delta_\mu^* f(x) = \frac{1}{a} [f(x) - f(x - \hat{\mu}a)].$$

Moreover we have

$$\begin{aligned}
 S^{(2)}(q^+, q^-) = & \frac{a^4}{4} \sum_x \sum_{\mu\nu} G_{\mu\nu}(q_\mu^-(x) \bar{D}_\nu^+ D_\mu^+ q_\nu^+(x) + q_\mu^+(x) \bar{D}_\nu^- D_\mu^- q_\nu^-(x) - q_\nu^-(x) \bar{D}_\mu^+ D_\mu^+ q_\nu^+(x) - q_\nu^+(x) \bar{D}_\mu^- D_\mu^- q_\nu^-(x)) \\
 & + \frac{ia^4}{2} \sum_x (q_1^-(x) \bar{D}_2^+ D_1^+ q_2^+(x) - q_1^+(x) \bar{D}_2^- D_1^- q_2^-(x)) H_{12} + \frac{ia^3}{2} \sum_x (q_1^-(x) \bar{D}_2^+ D_1^+ q_2^+(x) \\
 & - q_1^+(x) \bar{D}_2^- D_1^- q_2^-(x)) H_{12} + ia^2 \sum_x (q_2^-(x) q_1^+(x) - q_1^-(x) q_2^+(x)) H_{12}, \tag{47}
 \end{aligned}$$

with

$$\mathcal{D}_\mu^\pm f(x) = \Delta_\mu f(x) \pm \frac{2i}{a} \delta_{\mu 2} \sin\left(\frac{agH}{2} x_1\right) f(x + \hat{\mu}a), \tag{48}$$

$$\bar{\mathcal{D}}_\mu^\pm f(x) = \Delta_\mu^* f(x) \pm \frac{2i}{a} \delta_{\mu 2} \sin\left(\frac{agH}{2} x_1\right) f(x - \hat{\mu}a).$$

Obviously we need, now, to fix the gauge. To this end we add a gauge fixing term to the action and the associated Faddeev-Popov ghost field action. We use the background gauge condition

$$\sum_\mu D_\mu q_\mu(x) = 0. \tag{49}$$

In the Landau gauge the gauge-fixing term in the one-loop approximation is given by

$$S_{gf}^{(2)} = a^4 \sum_x \text{Tr} \left[\sum_\mu D_\mu q_\mu(x) \right]^2. \tag{50}$$

Moreover, in the same approximation we get the following Faddeev-Popov contribution:

$$S_{F-P}^{(2)} = -\ln \text{Det} \left[-\sum_\mu D_\mu^* D_\mu \right]. \tag{51}$$

The lattice version of the continuum operator $\mathcal{O}_{\mu\nu}^{ab}$ in Eq. (25) can be extracted from Eqs. (44), (47), and (50). Unlike the continuum case it is not possible to solve in closed form the lattice version of the eigenvalue equation (26). However, if we neglect the irrelevant terms and keep only the contributions that survive in the naive continuum limit $a \rightarrow 0$, then we are able to solve the eigenvalue equations and obtain the spectrum. In this approximation we replace $G_{\mu\nu}$ with the identity, so that $S^{(2)}(q^3)$ does not depend on the background field and we can discard it. Moreover the sum of $S^{(2)}(q^+, q^-)$ and $S_{gf}^{(2)}$ simplifies considerably. We get

$$\begin{aligned}
 S^{(2)}(q^+, q^-) + S_{gf}^{(2)} = & \frac{a^4}{2} \sum_x \sum_\mu q_\mu^+(x) [-\mathcal{O}_1] q_\mu^-(x) + \text{H.c.} \\
 & + \frac{a^4}{2} \sum_x \sum_{\mu, \nu} (\delta_{\mu 1} \delta_{\nu 2} \\
 & - \delta_{\mu 2} \delta_{\nu 1}) q_\mu^-(x) \mathcal{O}_2 q_\nu^+(x) + \text{H.c.}, \tag{52}
 \end{aligned}$$

where we restricted the quantum fluctuations to the class of function

$$q(x_1, x_2, x_3, x_4) = \sin(p_4 x_4) e^{i(p_2 x_2 + p_3 x_3)} f(x_1). \tag{53}$$

Note that the class of functions Eq. (53) is relevant for the constraint Eq. (19). Similar results can be obtained with the constraint Eq. (22). The periodic boundary conditions imply that

$$f(x_1 + L_1) = f(x_1) \tag{54}$$

$$p_\mu = \frac{2\pi}{L_\mu} n_\mu \quad \mu = 2, 3, 4, \tag{55}$$

and n_μ integer. Within the class of functions Eq. (53) we have

$$\begin{aligned}
 -\mathcal{O}_1 = & -\Delta_1^* \Delta_1 + \frac{4}{a^2} \sin^2\left(\frac{agHx_1}{2}\right) + \frac{2}{a^2} \sum_{\mu=1}^4 (1 - \cos p_\mu a) \\
 & - \frac{2}{a^2} \sin(agHx_1) \sin(p_2 a) - \frac{4}{a^2} \sin^2\left(\frac{agHx_1}{2}\right) \\
 & \times (1 - \cos(p_2 a)), \tag{56}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{O}_2 = & \frac{2i}{a^2} \sin\left(\frac{a^2 gH}{2}\right) + \frac{i}{a^2} \sin(gHa^2) \cos(gHax_1) \\
 & - \frac{2}{a^2} \sin(agHx_1) \sin(p_2 a) \frac{\Delta_1 + \Delta_1^*}{2}. \tag{57}
 \end{aligned}$$

By keeping only the relevant terms, the operators \mathcal{O}_1 and \mathcal{O}_2 further simplify as

$$\begin{aligned}
-\mathcal{O}_1 &= -\Delta_1^* \Delta_1 + \frac{4}{a^2} \sin^2\left(\frac{agHx_1}{2}\right) \\
&\quad - \frac{2}{a^2} \sin(agHx_1) \sin(p_2 a) + \frac{2}{a^2} \sum_{\mu=1}^4 (1 - \cos(p_2 a)),
\end{aligned} \tag{58}$$

$$\mathcal{O}_2 = \frac{2i}{a^2} \sin\left(\frac{a^2 gH}{2}\right) + \frac{i}{a^2} \sin(gHa^2) \simeq \frac{2i}{a^2} \sin\left(\frac{a^2 gH}{2}\right). \tag{59}$$

Introducing the complex scalar fields

$$\begin{aligned}
\phi &= q_+^-(x), & \phi^* &= q_-^+(x), \\
\psi &= q_-^-(x), & \psi^* &= q_+^+(x),
\end{aligned} \tag{60}$$

where $q_{\pm} = 1/\sqrt{2}(q_1 \pm q_2)$, we get

$$\begin{aligned}
&S^{(2)}(q^+, q^-) + S_{g-f}^{(2)} \\
&= \frac{a^4}{2} \sum_x \sum_{\nu=3,4} q_{\nu}^+(x) [-\mathcal{O}_1] q_{\nu}^- + \text{H.c.} \\
&\quad + \frac{a^4}{2} \sum_x \psi^*(x) [-\mathcal{O}_1 + \mathcal{O}_2] \psi(x) + \text{H.c.} \\
&\quad + \frac{a^4}{2} \sum_x \phi^*(x) [-\mathcal{O}_1 - \mathcal{O}_2] \phi(x) + \text{H.c.}
\end{aligned} \tag{61}$$

It is easy to verify that the contribution to the one-loop effective action due to the fluctuating fields q_{μ}^{\pm} , $\mu=3,4$ cancels the one due to the Faddeev-Popov determinant. So that we are left with the following quadratic action:

$$\begin{aligned}
S^{(2)}(\psi, \phi) &= a^4 \sum_x \psi^*(x) [-\mathcal{O}_1(x) + m^2] \psi(x) \\
&\quad + a^4 \sum_x \phi^*(x) [-\mathcal{O}_1(x) - m^2] \phi(x),
\end{aligned} \tag{62}$$

where

$$m^2 = \frac{2}{a^2} \sin(a^2 gH). \tag{63}$$

Let us introduce the operators

$$C = \Delta_1 - i \left[\frac{e^{ip_2 a} - 1}{a} \right] + \frac{2}{a} \sin\left(\frac{gHax_1}{2}\right), \tag{64}$$

$$C^* = -\Delta_1^* + i \left[\frac{e^{-ip_2 a} - 1}{a} \right] + \frac{2}{a} \sin\left(\frac{gHax_1}{2}\right). \tag{65}$$

By keeping the leading terms in the continuum limit it is not too hard to see that

$$-\mathcal{O}_1 = \frac{1}{2} [C^* C + C C^*] + \sum_{\nu=3}^4 \frac{2}{a^2} (1 - \cos p_{\nu} a) \tag{66}$$

and

$$[C, C^*] = \frac{4}{a^2} \sin\left(\frac{gHa^2}{2}\right). \tag{67}$$

So that we find that the eigenvalue equation

$$-\mathcal{O}_1 f_{\lambda}(x_1) = \lambda f_{\lambda}(x_1) \tag{68}$$

admits the solutions

$$\begin{aligned}
\lambda_n &= \frac{4n}{a^4} \sin\left(\frac{gHa^2}{2}\right) \\
&\quad + \frac{2}{a^2} \sum_{\mu=3}^4 (1 - \cos p_{\mu} a), \quad n=0,1,2,\dots
\end{aligned} \tag{69}$$

Note that the eigenvalues are degenerate. As a matter of fact the order of the degeneracy of the Landau levels turns out to be

$$g = \left(\frac{L_1 a}{2\pi}\right) \left(\frac{L_2 a}{2\pi}\right) \frac{2}{a^2} \sin\left(\frac{gHa^2}{2}\right). \tag{70}$$

We have numerically checked that the approximate spectrum Eq. (69) agrees quite well with the exact one as long as $L_1 \geq 32$ and for weak magnetic fields.

From Eqs. (68) and (69) we find the following eigenvalues:

$$\lambda_n^{\psi} = \sum_{\nu=3}^4 \frac{2}{a^2} (1 - \cos p_{\nu} a) + (2n+3) \frac{2}{a^2} \sin^2 \frac{gHa^2}{2}, \tag{71}$$

$$n=0,1,2,\dots,$$

$$\lambda_n^{\phi} = \sum_{\nu=3}^4 \frac{2}{a^2} (1 - \cos p_{\nu} a) + (2n-1) \frac{2}{a^2} \sin^2 \frac{gHa^2}{2}, \tag{72}$$

$$n=0,1,2,\dots$$

It is now evident that the ϕ mode with $n=0$ is the Nielsen-Olesen mode with eigenvalues

$$\lambda_u = \frac{2}{a^2} (1 - \cos p_3 a) + \frac{2}{a^2} (1 - \cos p_4 a) - \frac{2}{a^2} \sin^2\left(\frac{gHa^2}{2}\right), \tag{73}$$

which is the discretized version of Eq. (27).

IV. MONTE CARLO SIMULATIONS

Our numerical simulations have been done on a lattice of size $L_1 L_2 L_3 L_4$ with periodic boundary conditions. In order to project onto the ground state according to Eq. (13) we

need $L_4 \gg 1$. Moreover, in order to be close to the continuum limit, Eqs. (33) and (21) imply also $L_1 \gg 1$. As a consequence we performed the numerical simulations on lattices with $L_1 = L_4 = 32$. The transverse size of the lattice $L_\perp = L_2 = L_3$ has been varied from $L_\perp = 6$ up to $L_\perp = 32$. We are interested in the density of the effective action Eq. (14). We face the problem of computing a partition function which is the exponential of an extensive quantity [17]. To avoid this problem we consider the derivative of $\varepsilon[\vec{A}^{\text{ext}}]$ with respect to β by taking n_{ext} (i.e., gH) fixed [see Eq. (21)]. From Eqs. (14), (15), and (17) it follows that

$$\begin{aligned} \varepsilon'[\vec{A}^{\text{ext}}] &= \frac{\partial \varepsilon[\vec{A}^{\text{ext}}]}{\partial \beta} = -\frac{1}{\Omega} \left[\frac{1}{\mathcal{Z}[U^{\text{ext}}]} \frac{\partial \mathcal{Z}[U^{\text{ext}}]}{\partial \beta} \right. \\ &\quad \left. - \frac{1}{\mathcal{Z}[0]} \frac{\partial \mathcal{Z}[0]}{\partial \beta} \right] = \left\langle \frac{1}{\Omega} \sum_{x, \mu > \nu} \frac{1}{2} \text{Tr} U_{\mu\nu}(x) \right\rangle_0 \\ &\quad - \left\langle \frac{1}{\Omega} \sum_{x, \mu > \nu} \frac{1}{2} \text{Tr} U_{\mu\nu}(x) \right\rangle_{\vec{A}^{\text{ext}}}, \end{aligned} \quad (74)$$

where the subscripts on the average indicate the value of the external links at the boundaries. Obviously $\varepsilon[\vec{A}^{\text{ext}}]$ can be obtained by a numerical integration in β

$$\varepsilon[\vec{A}^{\text{ext}}, \beta] = \int_0^\beta d\beta' \varepsilon'[\vec{A}^{\text{ext}}, \beta'], \quad (75)$$

where we have taken into account that $\lim_{\beta \rightarrow 0} \varepsilon[\vec{A}^{\text{ext}}, \beta] = 0$.

It is evident that the contributions to $\varepsilon'[\vec{A}^{\text{ext}}]$ due to the frozen time slice at $x_4 = 0$ and to the fixed links at the spatial boundaries must be subtracted. In other words, only the dynamical links must be taken into account in evaluating $\varepsilon'[\vec{A}^{\text{ext}}]$. We recall that $\Omega = L_1 L_2 L_3 L_4$ is the total number of lattice sites (i.e., the lattice volume) belonging to the lattice Λ . If we denote with Ω_{ext} the lattice sites whose links are fixed according to Eq. (19):

$$\begin{aligned} \Omega_{\text{ext}} &= L_1 L_2 L_3 + (L_4 - 1)[L_1 L_2 L_3 - (L_1 - 2)(L_2 - 2) \\ &\quad \times (L_3 - 2)], \end{aligned} \quad (76)$$

then the volume occupied by the ‘‘internal’’ lattice sites is given by

$$\Omega_{\text{int}} = \Omega - \Omega_{\text{ext}}. \quad (77)$$

Accordingly, we define the derivative of the internal energy density $\varepsilon'_{\text{int}}[\vec{A}^{\text{ext}}]$ as

$$\begin{aligned} \varepsilon'_{\text{int}}[\vec{A}^{\text{ext}}] &= \left\langle \frac{1}{\Omega_{\text{int}}} \sum_{x \in \bar{\Lambda}, \mu > \lambda} \frac{1}{2} \text{Tr} U_{\mu\nu}(x) \right\rangle_0 \\ &\quad - \left\langle \frac{1}{\Omega_{\text{int}}} \sum_{x \in \bar{\Lambda}, \mu > \lambda} \frac{1}{2} \text{Tr} U_{\mu\nu}(x) \right\rangle_{\vec{A}^{\text{ext}}}, \end{aligned} \quad (78)$$




FIG. 1. The β derivative of the internal energy density Eq. (78) versus β at different values of the transverse lattice size L_\perp .

where $\bar{\Lambda}$ is the ensemble of the internal lattice sites.

To implement the constraint at the boundaries in the numerical simulations we update only the internal links, i.e., the links $U_\mu(x)$ with $x \in \bar{\Lambda}$.

We use the over-relaxed heat-bath algorithm to update the gauge configurations. Simulations have been performed by means of the APE100/Quadrics computer. Since we are measuring a local quantity such as the plaquette, a low statistics (from 1000 up to 5000 configurations) is required in order to get a good estimation of $\varepsilon'_{\text{int}}$.

In Fig. 1 we display the derivative of the energy density normalized to the derivative of the external energy density:

$$\varepsilon'_{\text{ext}} = 1 - \cos\left(\frac{gH}{2}\right) = 1 - \cos\left(\frac{2\pi}{L_1} n_{\text{ext}}\right) \quad (79)$$

versus β for $L_1 = L_4 = 32$ and $6 \leq L_\perp \leq 32$. From Fig. 1 we see that in the strong-coupling region $\beta \lesssim 1$ the external background field is completely shielded. Moreover $\varepsilon'_{\text{int}}$ displays a peak at $\beta \approx 2.2$ resembling the behavior of the specific heat [18,19]. This is not surprising since our previous studies [20] in U(1) showed that $\varepsilon'_{\text{int}}$ behaves like a specific heat.

In the weak-coupling region $\beta \gtrsim 3$ Fig. 1 shows that the ratio $\varepsilon'_{\text{int}}/\varepsilon'_{\text{ext}}$ stays constant. Actually the constant does depend on L_\perp and n_{ext} . Indeed in Fig. 1 the dependence on L_\perp for fixed external magnetic field is evident. On the other hand, in Fig. 2 we keep $L_\perp = 32$ fixed and vary n_{ext} . We see clearly that the weak-coupling plateau decreases by increasing the external field. In order to extract $\varepsilon_{\text{int}}(\beta, n_{\text{ext}})$, we can numerically integrate the data for $\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})/\varepsilon'_{\text{ext}}$ using the trapezoidal rule




FIG. 2. The β derivative of the internal energy density Eq. (78) versus β for a transverse lattice size $L_{\perp}=32$ at different values of applied external field strength.

$$\varepsilon_{\text{int}}(\beta, n_{\text{ext}}) = \varepsilon'_{\text{ext}} \int_0^{\beta} \frac{\beta' \varepsilon'_{\text{int}}(\beta', n_{\text{ext}})}{\varepsilon'_{\text{ext}}} d\beta'. \quad (80)$$

In Fig. 3 we display $\varepsilon_{\text{int}}(\beta, n_{\text{ext}})$ obtained from Eq. (80). The plateau of the derivative of the internal energy density in the weak-coupling region results in a linear rising term in the energy density. For $\beta \gg 1$ we get

$$\varepsilon_{\text{int}}(\beta, n_{\text{ext}}) \approx \beta \varepsilon'_{\text{ext}} a(n_{\text{ext}}). \quad (81)$$

Moreover, for $\beta \gg 1$ we get also




FIG. 3. The energy density Eq. (80) versus β for transverse lattice size $L_{\perp}=32$ at different values of applied external field strength. The solid lines are the linear fits Eq. (81).




FIG. 4. The data displayed in Fig. 2 near the peaks of the β derivative of the internal energy density Eq. (78) in correspondence of each value of n_{ext} . The solid lines are the fits Eq. (88).

$$\beta \varepsilon'_{\text{int}} = \beta \left(1 - \cos \frac{2\pi}{L_{\perp}} n_{\text{ext}} \right) \approx \frac{1}{2} H^2. \quad (82)$$

So that in the weak-coupling region

$$\varepsilon_{\text{int}}(\beta, n_{\text{ext}}) \approx a(n_{\text{ext}}) \frac{1}{2} H^2, \quad \beta \gg 1. \quad (83)$$

Figure 1 shows that $a(n_{\text{ext}}) \approx 1$ for $L_{\perp} \approx 6-8$ and $n_{\text{ext}}=1$. On the other hand, $a(n_{\text{ext}})$ decreases by increasing L_{\perp} or the external background field. This peculiar behavior can be compared with the Abelian case where we found that $a(n_{\text{ext}}) \approx 1$ independently on L_{\perp} and n_{ext} [9,20]. Previous theoretical studies [15] suggested that due to the presence of the Nielsen-Olesen modes the gauge system reacts strongly to the external perturbation even in the nominally perturbative regime. It turns out that the Nielsen-Olesen modes behave like a (1+1)-dimensional tachionic charged scalar field. The condensation of these modes takes place only in the thermodynamic limit. As a consequence the applied external background magnetic field is almost completely screened and there is a dramatic reduction of the vacuum magnetic energy. Indeed it turns out that in the infinite volume limit the perturbative vacuum and the magnetic condensate vacuum are degenerate for vanishing gauge coupling.

On the lattice the Nielsen-Olesen modes display the one-loop instability when λ_{μ} given by Eq. (73) becomes negative. In the approximation adopted in Sec. III we find that λ_{μ} gets negative by increasing L_{\perp} for fixed external field. Thus we can switch on and off the one-loop instability by varying L_{\perp} . This has been also noticed by the authors of Ref. [21]. For instance, by using Eqs. (21), (55), and (73) with $L_{\perp} = L_4 = 32$ and $n_{\text{ext}}=1$ we find that $\lambda_{\mu} \leq 0$ for $L_{\perp} \geq 11$.

Our numerical results in Fig. 1 show that for $n_{\text{ext}}=1$ and $L_{\perp} \lesssim 10$ there is no the Nielsen-Olesen instability and the




FIG. 5. The lattice data for $\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})/\varepsilon'_{\text{ext}}$ at different values of the transverse lattice size L_{\perp} and external field strength n_{ext} for $\beta = 5$ (a) and $\beta = \beta_{\text{peak}}$ (b), versus the scaling variable defined in Eq. (84). The solid lines are the fits Eq. (87) with $\alpha = 1.5$.

gauge system responds weakly to the external perturbation in the weak-coupling region. On the other hand, by increasing L_{\perp} (see Fig. 1) or n_{ext} (see Fig. 2) we increment the lattice Nielsen-Olesen modes. As a consequence we find a clear reduction of the vacuum energy density for both the peak values of $\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})$ and the coefficient $a(n_{\text{ext}})$ in Eq. (83) decreases towards zero in the thermodynamic limit. To see this we need to perform the infinite volume extrapolation. We can extract more information from our numerical data by expressing them versus

$$x = \frac{a_H}{L_{\text{eff}}}, \quad (84)$$

where

$$a_H = \sqrt{\frac{2\pi}{gH}} = \sqrt{\frac{L_1}{2n_{\text{ext}}}} \quad (85)$$

is the magnetic length and

$$L_{\text{eff}} = \Omega_{\text{int}}^{1/4} \quad (86)$$

is the lattice effective linear size. Indeed, we find that the data for $\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})/\varepsilon'_{\text{ext}}$ at the perturbative tail and at the




FIG. 6. The scaling curve obtained by rescaling all lattice data for $\varepsilon'_{\text{int}}(\tilde{\beta}, n_{\text{ext}}, L_{\text{eff}})$ according to Eq. (89).

peak for various lattice sizes and values of n_{ext} can be expressed as a function of the scaling variable x [defined in Eq. (84)]:

$$\frac{\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})}{\varepsilon'_{\text{ext}}} = \kappa(\beta)x^{\alpha}. \quad (87)$$

For the perturbative tail of $\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})/\varepsilon'_{\text{ext}}$ we keep the value of the ratio at $\beta = 5$. On the other hand, the peak values have been extracted by fitting the values around the peak to (see Fig. 4)

$$\frac{\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})}{\varepsilon'_{\text{ext}}} = \frac{a_1}{a_2(\beta - \beta_{\text{peak}})^2 + 1}. \quad (88)$$

From Eq. (88) we extract the peak value, a_1 , and the peak position β_{peak} . It has been found that the data are compatible with the scaling law Eq. (87) with $\alpha = 1.5$ (see Fig. 5). It is remarkable that the same power law arises if we adopt the alternative boundary conditions given by Eq. (22). So we see that both boundary conditions Eq. (20) or Eq. (22) lead to the same thermodynamic limit.

If we, further, take into account the shifts $\Delta\beta$ of the peak values, which turn out to depend only on n_{ext} , we are led to the universal scaling law

$$x^{-\alpha} \frac{\varepsilon'_{\text{int}}(\tilde{\beta}, n_{\text{ext}}, L_{\text{eff}})}{\varepsilon'_{\text{ext}}} = \kappa(\tilde{\beta}), \quad (89)$$

where $\tilde{\beta} = \beta - \Delta\beta$. Indeed Fig. 6 shows that all our numerical data (for all the values of a_H and L_{eff}) can be approximately arranged on the scaling curve $\kappa(\tilde{\beta})$. Remarkably we find that the peak in $\kappa(\tilde{\beta})$ is located at $\beta_c = 2.2209(68)$ which agrees with the peak position of the specific heat extrapolated to the infinite volume limit $\beta_c = 2.23(2)$ [19]. By

using Eq. (89) we can determine the infinite volume limit of the vacuum energy density ε_{int} . We have

$$\begin{aligned} \lim_{L_{\text{eff}} \rightarrow \infty} \varepsilon_{\text{int}}(\beta, n_{\text{ext}}, L_{\text{eff}}) &= \varepsilon'_{\text{ext}} \int_0^{\bar{\beta}} d\bar{\beta}' \kappa(\bar{\beta}') \lim_{L_{\text{eff}} \rightarrow \infty} \left(\frac{a_H}{L_{\text{eff}}} \right)^\alpha \\ &\simeq \frac{H^2}{2\bar{\beta}} \left[\int_0^{\bar{\beta}} d\bar{\beta}' \kappa(\bar{\beta}') \right] \\ &\times \lim_{L_{\text{eff}} \rightarrow \infty} \left(\frac{a_H}{L_{\text{eff}}} \right)^\alpha = 0, \end{aligned} \quad (90)$$

in the whole range of β . This in turn implies that in the continuum limit ($L_{\text{eff}} \rightarrow \infty, \beta \rightarrow \infty$) the SU(2) vacuum completely screens the external chromomagnetic Abelian field. In other words, the continuum vacuum behaves as an Abelian magnetic condensate medium in accordance with the dual superconductivity scenario.

V. CONCLUSIONS

We have studied the nonperturbative dynamics of the vacuum of SU(2) lattice gauge theory by means of the gauge-invariant effective action defined using the lattice Schrödinger functional.

Our numerical results indicate that in the continuum limit $L_{\text{eff}} \rightarrow \infty, \beta \rightarrow \infty$ we have

$$\varepsilon[\vec{A}^{\text{ext}}] = 0, \quad (91)$$

so that the vacuum screens completely the external chromomagnetic Abelian field. In other words, the continuum vacuum behaves as an Abelian magnetic condensate medium in accordance with the dual superconductivity scenario. In particular we have

$$\varepsilon[\vec{A}^{\text{ext}}] \sim H_k^a B_k^a = \frac{1}{\mu} F_{ij}^a F_{ij}^a, \quad (92)$$

where μ is the vacuum color magnetic permeability. Thus Eq. (91) implies that $\mu \rightarrow \infty$ in the continuum limit. As a consequence by Lorentz invariance the vacuum color dielectric constant tends to zero. This in turns implies that the vacuum does not support an isolated color charge, i.e., the color confinement.

Let us conclude by stressing that our method can be easily extended to the SU(3) gauge theory. Moreover we also feel that the lattice gauge-invariant effective action could be also employed to study different background fields.

-
- [1] J. Schwinger, Phys. Rev. **82**, 664 (1951).
 - [2] J. Honerkamp, Nucl. Phys. **B36**, 130 (1971); **B48**, 269 (1972).
 - [3] G. 't Hooft, Nucl. Phys. **B62**, 444 (1973).
 - [4] P. Cea and L. Cosmai, Phys. Rev. D **48**, 3364 (1993); Nucl. Phys. B (Proc. Suppl.) **34**, 234 (1994); hep-lat/9306007.
 - [5] H. D. Trottier and R. M. Woloshyn, Phys. Rev. Lett. **70**, 2053 (1993); **72**, 4155 (1994).
 - [6] P. Cea and L. Cosmai, Phys. Rev. D **43**, 620 (1991); Phys. Lett. B **264**, 415 (1991).
 - [7] J. Ambjorn, V. K. Mitryushkin, and V. G. Bornyakov, Phys. Lett. B **225**, 153 (1989); **245**, 575 (1990).
 - [8] A. R. Levi, Nucl. Phys. B (Proc. Suppl.) **34**, 161 (1994).
 - [9] P. Cea, L. Cosmai, and A. D. Polosa, Phys. Lett. B **392**, 177 (1997).
 - [10] G. C. Rossi and M. Testa, Nucl. Phys. **B163**, 109 (1980); **B176**, 477 (1980). For a recent review see G. C. Rossi, hep-th/9810177.
 - [11] D. J. Gross, R. D. Pisarski, and L. G. Yaffe, Rev. Mod. Phys. **53**, 43 (1981).
 - [12] M. Lüscher, R. Narayanan, P. Weisz, and U. Wolff, Nucl. Phys. **B384**, 168 (1992); M. Lüscher and P. Weisz, *ibid.* **B452**, 213 (1995).
 - [13] A partial account of the results of the present paper appeared in P. Cea and L. Cosmai, Mod. Phys. Lett. A **13**, 861 (1998); Nucl. Phys. B (Proc. Suppl.) **73**, 641 (1999).
 - [14] N. K. Nielsen and P. Olesen, Nucl. Phys. **B144**, 376 (1978).
 - [15] P. Cea, Phys. Rev. D **37**, 1637 (1988).
 - [16] R. Dashen and D. J. Gross, Phys. Rev. D **23**, 2340 (1981).
 - [17] A. Hasenfratz, P. Hasenfratz, and F. Niedermeyer, Nucl. Phys. **B329**, 739 (1990).
 - [18] B. Lautrup and M. Nauenberg, Phys. Rev. Lett. **45**, 1755 (1980).
 - [19] J. Engels and T. Scheideler, Nucl. Phys. B (Proc. Suppl.) **53**, 423 (1997).
 - [20] P. Cea, L. Cosmai, and A. D. Polosa, Phys. Lett. B **397**, 229 (1997).
 - [21] A. R. Levi and J. Polonyi, Phys. Lett. B **357**, 186 (1995).