

Period-doubling bifurcation in strongly anisotropic Bianchi I quantum cosmology

Michael Bachmann* and Hans-Jürgen Schmidt†

Institut für Mathematik, Universität Potsdam, PF 601553, D-14415 Potsdam, Germany
and Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, D-14195 Berlin, Germany

(Received 15 December 1999; published 25 July 2000)

We solve the Wheeler-DeWitt equation for the minisuperspace of a cosmological Bianchi type-I model with a minimally coupled massive scalar field ϕ as the source by generalizing the calculation of Lukash and Schmidt [Astron. Nachr. **309**, 25 (1988)]. Contrary to other approaches we allow strong anisotropy. Combining analytical and numerical methods, we apply an adiabatic approximation for ϕ , and as a new feature we find a period-doubling bifurcation. This bifurcation takes place near the cosmological quantum boundary, i.e., the boundary of the quasiclassical region with an oscillating ψ function where the WKB approximation is good. The numerical calculations suggest that such a notion of a “cosmological quantum boundary” is well defined, because, sharply beyond that boundary, the WKB approximation is no longer applicable at all. This result confirms the adequateness of the introduction of a cosmological quantum boundary in quantum cosmology.

PACS number(s): 98.80.Hw, 98.70.Vc

I. INTRODUCTION

The idea to consider the whole Universe as one quantum system has already attracted many researchers. One of the approaches to give this idea a physical meaning is to consider the superspace \mathcal{S} of all possible spatial three-geometries together with the matter field degrees of freedom as supercoordinates, and the supermomentum space \mathcal{S}^* is formed from the corresponding second fundamental forms together with the momenta of the matter fields. The dynamics will then be defined by the Einstein field equation accompanied by the matter field equations. In the last step, this system will be quantized.

Of course, one has to deal with an infinite number of degrees of freedom already from the geometric part. The only way to tackle such a system up to now is to linearize in almost all degrees of freedom and to take the nonlinearities only from the remaining finitely many ones. But even this system is almost untractable. So, the idea of the minisuperspace arose: One restricts to a finite-dimensional space of spatial geometries and to matter fields with finitely many components, disregarding or even ignoring all other degrees of freedom.

From the first glance one could believe that such a big simplification would lead to a picture which has nothing to do with the real Universe’s evolution. However, the results given in the last years are encouraging: The dynamics of the minisuperspace can be written as an equivalent mechanical system, and then the zero-energy-Schrödinger equation for this system carries the name “Wheeler-DeWitt equation,”¹ which can be solved for simple systems and gives already a surprisingly good picture of the evolution, even for the case

that the geometric part is restricted to only one degree of freedom, the “radius” of the Universe. This corresponds to the isotropic Friedmann Universe model. For this case, closed-form solutions for the Wheeler-DeWitt equation exist.

In the ordering with respect to simplicity, the next possible geometry is the Bianchi type-I model, where the spatial inner geometry is flat, but the expansion is allowed to be anisotropic. For this case, we have three degrees of freedom for the geometry: the expansion rates into the three spatial directions.² For this case, the Wheeler-DeWitt equation is already quite complicated, so one uses an approximation to solve it. One of the most powerful of these approximation schemes is the Wentzel-Kramers-Brillouin (WKB) approximation. In Ref. [1], the first-order WKB approximation for the massive scalar field in the Bianchi type-I model has been deduced, but—as far as we are aware—the higher-order WKB approximations have not yet been calculated up to now for this model. The analogous calculations for the Bianchi type-IX model have been done by Amsterdamski [2]. In both cases, the anisotropy degrees of freedom had been assumed to be small. In the present paper we allow also large anisotropies. Reference [3] deals with a similar model allowing large anisotropies, too; however, the authors of [3] use a simpler minisuperspace model, so the results are not directly comparable to ours.

The text is organized as follows: Sec. II presents the geometry of the Bianchi type-I cosmological model, Sec. III is the corresponding Wheeler-DeWitt equation, and Sec. IV its solutions both analytically and numerically. Section V shortly reviews Refs. [6–48], i.e., the earlier developments of the topic, and discusses the results.

In the Appendix, we show how the different versions to solve the factor-ordering problem influence the solutions of the Wheeler-DeWitt equation.

*Email address: mbach@physik.fu-berlin.de

†Email address: hjschmi@rz.uni-potsdam.de

¹Though this identification represents partially a misnomer, it is quite usual in the literature, see the more detailed explanation in Sec. V B.

²Strictly speaking, the axially symmetric Bianchi type-I model, where two of the three scale factors coincide, is even simpler, but as one will see from the calculations: restricting to axial symmetry does not really simplify the procedure.

II. THE BIANCHI TYPE-I MODEL

The metric of a spatially flat cosmological model is deduced as follows: One requires that an Abelian three-dimensional isometry group acts transitively on three-dimensional spatial hypersurfaces which consequently have to be flat three spaces. This restricts the possible metrics to

$$ds^2 = N^2 dt^2 - g_{ij} dx^i dx^j, \quad (2.1)$$

where $N = N(t)$ is the lapse function which can be put to $N = 1$ by a time reparametrization, $i, j = 1, 2, 3$, and g_{ij} represents a symmetric positive definite matrix whose components depend on t only. So we have six free components from the first consideration. However, restricting to metrics being a solution of Einstein's field equation we can simplify as follows: At any initial time, say $t = 0$, we can choose the initial condition

$$g_{ij} = \delta_{ij} = \text{diag}(1, 1, 1) \text{ at } t = 0 \quad (2.2)$$

without loss of generality, and then by a spatial rotation not changing Eq. (2.2), the second fundamental form, i.e., dg_{ij}/dt , can be brought into diagonal form. By use of the Einstein equation one can show that under these circumstances, g_{ij} will keep its diagonal form for all times. Thus: The Bianchi type-I model containing nondiagonal terms does *not* represent a generalization, and we can say that without loss of generality let

$$g_{ij} = \text{diag}(A^2(t), B^2(t), C^2(t)) \quad (2.3)$$

with certain positive functions A, B, C .

After an obvious rearrangement of the terms we get now from Eqs. (2.1) and (2.3)

$$ds^2 = N^2 dt^2 - e^{2\alpha} (e^{2s+2\sqrt{3}r} dx^2 + e^{2s-2\sqrt{3}r} dy^2 + e^{-4s} dz^2), \quad (2.4)$$

where α, r, s represent three arbitrary real functions depending on t . The Hubble parameter is the mean expansion, i.e., with a dot denoting d/dt we get

$$H = \frac{1}{3} \left(\frac{\dot{A}}{A} + \frac{\dot{B}}{B} + \frac{\dot{C}}{C} \right).$$

In the notation of Eq. (2.4), $H = d\alpha/dt$ is the Hubble parameter, and we restrict our considerations to the expanding Universe, i.e., to $H > 0$. The model goes over to the spatially flat isotropic Friedmann model if the dimensionless anisotropy parameter

$$\eta = \frac{1}{H} \sqrt{\dot{r}^2 + \dot{s}^2} \quad (2.5)$$

tends to zero.

It is essential to observe that this approach leading to the Wheeler-DeWitt equation breaks the Lorentz invariance of the system already at this level of geometry: The 3+1 decomposition of space time is already made from the beginning. The remaining degrees of freedom within the geometry

are the following: $\tilde{t}(t)$ as arbitrary function as long as $N(t)$ remains unspecified, and $\tilde{t}(t) = \pm t$ after having fixed $N = 1$. Permutations of the three spatial coordinates, and for the axially symmetric case also spatial rotations, such as if $r = 0$, then a rotation in the xy plane is an additional symmetry. We presented this geometric part so explicitly, because contradicting statements exist about this behavior in the literature; for example: In our interpretation, $ds^2 = dt^2 - dx^2 - dy^2 - dz^2$ and $ds^2 = dt^2 - t^2 dx^2 - dy^2 - dz^2$ represent different geometries in spite of the fact that they are locally isometric space times.

We use a one-component real massive scalar field ϕ as a source, and we interpret it as follows: Either it can really be such a scalar field (i.e., a spin zero field), e.g., a Higgs field which dominated the early Universe's evolution but disappeared after a symmetry-breaking effect. Or, it mimics any realistic matter field in a region where spin is negligible. Let m denote the mass of the scalar field. For the grand unified theory (GUT) mediated inflationary model, one considers the following order of magnitude: $m \approx 10^{-5} m_{\text{Pl}}$, where m_{Pl} is the Planck mass.³

Up to now, we did not restrict the spatial coordinates x, y , and z . At the classical level (i.e., for locally solving the Einstein field equation), it makes no difference, whether they cover all the reals or whether they are cyclic ones. But already for global classical considerations it *makes* a difference: Suppose, x, y , and z are cyclic, then even for the case $r = s = 0$ the metric is not spatially isotropic, because the natural length of a closed spatial geodesic depends on the chosen direction. However, we get the essential difference only after quantization. If nothing different is said, we assume x, y , and z to be cyclic with modulus 1, i.e., x and $x + 1$ represent the same point, etc.

III. THE WHEELER-DEWITT EQUATION

The Wheeler-DeWitt equation is always a zero-energy-Schrödinger equation⁴ due to time-reparametrization invariance of the gravitational action. However, this is not really a problem, because the Schrödinger equation with nonvanishing energy can be brought to the zero-energy form by a suitable shift of the potential, see, e.g., Ref. [4] for a class of transformations of this type.

In the present paper, we do not consider the entropy of the system; we only want to mention the recently published result of Kleinert [5] on how the incorporation of entropy can change the model.

To formulate the equations that lead to the Wheeler-DeWitt equation [6] as easily as possible, it proves useful to apply units such that $\hbar = c = 4\pi G/3 = 1$. Then the Lagrangian of the minimally coupled scalar field in Einstein's theory of gravity (see the first of Refs. [1] for details) is given by

$$L = L_g + L_m, \quad (3.1)$$

³In units with $c = 1$, the Planck mass is about 10^{19} GeV.

⁴But, see also, the discussion in Sec. V B. for this notation.

where the gravitational part is

$$L_g = \frac{1}{2N} e^{3\alpha} (\dot{r}^2 + \dot{s}^2 - \dot{\alpha}^2) \quad (3.2)$$

and the matter Lagrangian is

$$L_m = \frac{1}{2N} e^{3\alpha} (\dot{\phi}^2 - m^2 \phi^2 N^2). \quad (3.3)$$

Here we supposed the lapse function N to be constant in time. The minisuperspace is four-dimensional, its coordinates are $q^\mu = (q^0, q^1, q^2, q^3)$ with $q^0 = \alpha$, $q^1 = \phi$, $q^2 = r$, and $q^3 = s$. The usefulness of the Misner parametrization Eq. (2.4) of the anisotropy is obvious: The kinetic part of the Lagrangian is already in diagonal form. By defining the minisuperspace metric $f_{\mu\nu} = e^{3\alpha} \text{diag}(-1, 1, 1, 1)$ the Lagrangian (3.1) is compactly rewritten as

$$L = \frac{1}{2N} [f_{\mu\nu} \dot{q}^\mu \dot{q}^\nu - m^2 (q^1)^2 N^2 e^{3q^0}]. \quad (3.4)$$

Thus the introduction of canonical momenta

$$p_\mu = \frac{\partial L}{\partial \dot{q}^\mu} = \frac{1}{N} f_{\mu\nu} \dot{q}^\nu \quad (3.5)$$

allows us to write down the classical Hamiltonian of the minisuperspace

$$\mathcal{H} = \frac{N}{2} [p_\mu p^\mu + m^2 (q^1)^2 e^{3q^0}] = 0. \quad (3.6)$$

Time-reparametrization invariance implies $\partial \mathcal{H} / \partial N = 0$, i.e., $\mathcal{H} = 0$. The quantization in coordinate space is done as usual by going over to the appropriate operators $p_\mu = -i\hbar \partial / \partial q^\mu$ with the partial derivative ∂ .

From now on, we will choose two different versions for the operator ordering in $p_\mu p^\mu$: First, variant A, which is the most simple one, following the interpretation in Ref. [1] that the solutions of the forthcoming Wheeler-DeWitt equation will not essentially depend on this choice.

Second, variant B, solves the factor ordering problem by applying the supercovariance principle: It says that the essential equations have to be covariant ones with respect to the (mini-)superspace metric. Taking variant B, we have to replace p_μ by $p_\mu = -i\hbar \nabla / \nabla q^\mu$, where ∇ denotes the supercovariant derivative⁵ defined by $f_{\mu\nu}$.

To cover both variants simultaneously, we introduce a parameter ϵ as follows: $\epsilon = 0$ for variant A and $\epsilon = 1$ for variant B. Accordingly, we denote the two variants of the D'Alembert operator⁶ \square_ϵ by

$$\square_0 = -f^{\mu\nu} \frac{\partial^2}{\partial q^\mu \partial q^\nu}, \quad (3.7)$$

and

$$\square_1 = -f^{\mu\nu} \frac{\nabla^2}{\nabla q^\mu \nabla q^\nu}. \quad (3.8)$$

For the world function $\psi(q^\mu)$ we get the relations

$$\square_1 \psi = \square_0 \psi + 3 \psi_{,0} e^{-3\alpha} \quad (3.9)$$

where⁷

$$\square_0 \psi = e^{-3\alpha} [\psi_{,00} - \psi_{,11} - \psi_{,22} - \psi_{,33}]. \quad (3.10)$$

The stationary Schrödinger equation with zero energy in minisuperspace $\hat{\mathcal{H}}\psi(q^\mu) = 0$ with the wave function $\psi(q^\mu)$ and the Hamilton operator $\hat{\mathcal{H}}$ is called the *Wheeler-DeWitt equation* and possesses the form

$$\hat{\mathcal{H}}\psi(q^\mu) = \frac{N}{2} [\hbar^2 \square_\epsilon + m^2 (q^1)^2 e^{3q^0}] \psi(q^\mu) = 0 \quad (3.11)$$

for the Bianchi type-I Universe (2.4). In detail, we get after dividing by $N e^{3\alpha/2}$

$$0 = \left(\hbar^2 \left[\frac{\partial^2}{\partial \alpha^2} + 3\epsilon \frac{\partial}{\partial \alpha} - \frac{\partial^2}{\partial \phi^2} - \frac{\partial^2}{\partial r^2} - \frac{\partial^2}{\partial s^2} \right] + m^2 \phi^2 e^{6\alpha} \right) \psi(\alpha, \phi, r, s). \quad (3.12)$$

In what follows we derive an approximation scheme to solve the Wheeler-DeWitt equation.

IV. SOLUTIONS OF THE WHEELER-DeWITT EQUATION

Before we go over to solve Eq. (3.12) we want to clarify in subsection A what happens if we neglect the anisotropy from the beginning. In subsections B–D we analytically obtain solutions in a well-defined approximation, and in subsection E we present the results of our numerical calculation.

A. Isotropic models

For variant A, i.e., $\epsilon = 0$ in Eq. (3.11) we get

$$0 = \left(\hbar^2 \left[\frac{\partial^2}{\partial \alpha^2} - \frac{\partial^2}{\partial \phi^2} \right] + m^2 \phi^2 e^{6\alpha} \right) \psi(\alpha, \phi), \quad (4.1)$$

i.e., we simply remove all r and s dependences. For variant B, however, this recipe does not work because the covariant

⁵Here we apply the fact that the metric is always covariantly constant, so different orderings give the same result.

⁶The minus sign in Eqs. (3.7) and (3.8) is inserted to compensate for the factor i^2 in front of $p_\mu p^\mu$.

⁷If the dimension of the minisuperspace is D instead of 4, then the factor 3 in front of $\psi_{,0}$ in Eq. (3.9) will be replaced by $3(D-2)/2$, so \square_0 and \square_1 coincide for $D=2$.

derivatives mix the coordinates. After a short calculation one finds out that here also Eq. (4.1) is the correct Wheeler-DeWitt equation. The reason for this coincidence is as follows: The two-dimensional superspace is conformally flat, and the D'Alembertian is conformally invariant, cf. the second paper in Ref. [1] for more details.

Defining $a = e^{\alpha} > 0$ as new variable we have to replace $\partial/\partial\alpha$ by $a(\partial/\partial a)$, and we get from Eq. (4.1) now

$$0 = \left(\hbar^2 \left[a \frac{\partial}{\partial a} a \frac{\partial}{\partial a} - \frac{\partial^2}{\partial \phi^2} \right] + m^2 \phi^2 a^6 \right) \tilde{\psi}(a, \phi), \quad (4.2)$$

a version of the Wheeler-DeWitt equation favored in Ref. [7].

B. Separation ansatz

Now we look for the solutions of Eq. (3.12). First, we make a separation ansatz

$$\psi(\alpha, \phi, r, s) = \chi(\alpha, \phi) \cdot \rho(r) \cdot \sigma(s). \quad (4.3)$$

It turns out that $\rho_{,22}/\rho = c_2$ and $\sigma_{,33}/\sigma = c_3$ represent constants, i.e., we get

$$\rho(r) = \rho_1 \exp(\sqrt{c_2} r) + \rho_2 \exp(-\sqrt{c_2} r)$$

and

$$\sigma(s) = \sigma_1 \exp(\sqrt{c_3} s) + \sigma_2 \exp(-\sqrt{c_3} s)$$

with arbitrary constants ρ_i and σ_i . For negative values c_2 , the function $\rho(r)$ has a sinus-function behavior.

One can interpret these cases as follows: If both c_2 and c_3 are negative, then we have plane waves in the r and s direction. This is in agreement with the fact that a translation into the r or s direction can be compensated by a coordinate transformation in metric (2.4) (i.e., by multiplying x , y , and z with suitable constants), so that all r and s values should be equally probable. However, this interpretation works only in the case that we allow x , y , and z to cover all the reals. If we, however, restrict x , y , and z to be cyclic coordinates, then this argument in favor of equal distribution of the r and s values is no more valid.

If, on the contrary, one of the constants c_2 or c_3 is non-negative, then (up to singular exceptions) the product $\rho(r) \cdot \sigma(s)$ tends to $\pm\infty$ as r or s does. This allows the following interpretation: The probability to have small anisotropy is exponentially small.

For the moment we keep the further interpretation open and continue the calculation. We define the constant c_1 via $c_1 + c_2 + c_3 = 0$ and insert Eq. (4.3) into Eq. (3.12); we get

$$0 = \left(\hbar^2 \left[\frac{\partial^2}{\partial \alpha^2} + 3\epsilon \frac{\partial}{\partial \alpha} - \frac{\partial^2}{\partial \phi^2} \right] + c_1 \hbar^2 + m^2 \phi^2 e^{6\alpha} \right) \chi(\alpha, \phi). \quad (4.4)$$

This equation is for variant A just the Wheeler-DeWitt equation (4.1), however, now not the zero-energy equation but

the equation with energy proportional to the constant c_1 representing the anisotropic degrees of freedom.

We redefine the coordinate α to $v = a^3 = e^{3\alpha}$. The coordinate v is proportional to the spatial volume. Then Eq. (4.4) goes over to (after dividing by 9)

$$0 = \left(\hbar^2 \left[v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} + \epsilon v \frac{\partial}{\partial v} - \frac{\partial^2}{\partial \phi^2} \right] + c_1 \hbar^2 / 9 + m^2 \phi^2 v^2 / 9 \right) \tilde{\chi}(v, \phi) \quad (4.5)$$

As one can see in comparison with Eq. (4.2). Instead of the a^6 term we now have a v^2 potential.

C. Adiabatic scalar field

In this subsection, we assume the scalar field ϕ to be almost constant. Then γ defined by $\gamma^2 = m^2 \phi^2 / (9 \hbar^2)$ is an adiabatic constant. In Eq. (4.5) we omit the ϕ derivative corresponding to the adiabatic approach. To this end we consider the following equation for the exact dependence on v of a new function $\hat{\chi}$:

$$\left(\hbar^2 \left[v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} + \epsilon v \frac{\partial}{\partial v} \right] + c_1 \hbar^2 / 9 + \gamma^2 v^2 \hbar^2 \right) \hat{\chi}(v, \phi) = E(\phi) \cdot \hat{\chi}(v, \phi), \quad (4.6)$$

where $E(\phi)$ is an yet undetermined eigenvalue; here, ϕ plays the role of a parameter only. We define a new adiabatic constant Λ by $\Lambda^2 = E(\phi) \hbar^{-2} - c_1 / 9$. Then Eq. (4.6) becomes for $\epsilon = 0$

$$\left(v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} + \gamma^2 v^2 - \Lambda^2 \right) \hat{\chi} = 0. \quad (4.7)$$

Equation (4.7) is a Bessel-type ordinary differential equation. If we replace in Eq. (4.7) γv by x and $\hat{\chi}(v, \phi)$ by $y(x, \phi)$ we get exactly Bessel's form

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - \Lambda^2) y = 0 \quad (4.8)$$

whose solutions are (the constants C_i depend on Λ)

$$y = C_1 J_\Lambda(x) + C_2 J_{-\Lambda}(x). \quad (4.9)$$

The J_Λ are called cylinder or Bessel functions and have the development

$$J_\Lambda(x) = \sum_{k=0}^{\infty} (-1)^k \left(\frac{x}{2} \right)^{\Lambda+2k} (k!)^{-1} [\Gamma(\Lambda+k+1)]^{-1}. \quad (4.10)$$

Now we make the ansatz⁸

⁸If only a discrete set of values Λ appear, Eq. (4.11) will be replaced by an analogous sum.

$$\tilde{\chi}(v, \phi) = \int_{\Lambda} b_{\Lambda}(\phi) \hat{\chi}_{\Lambda}(v, \phi) d\Lambda, \quad (4.11)$$

where we assigned to $\hat{\chi}$ the subscript Λ to emphasize $\hat{\chi}_{\Lambda}$ being an eigenfunction with eigenvalue Λ . Thus the general solution of Eq. (4.7) is written as

$$\hat{\chi}_{\Lambda}(v, \phi) = C_1 J_{\Lambda}(\gamma v) + C_2 J_{-\Lambda}(\gamma v). \quad (4.12)$$

Inserting ansatz (4.11) into Eq. (4.5) we get

$$\begin{aligned} 0 = & \int_{\Lambda} -\hbar^2 \frac{\partial^2}{\partial \phi^2} [b_{\Lambda}(\phi) \hat{\chi}(v, \phi)] + b_{\Lambda}(\phi) \\ & \times \left[\hbar^2 \left(v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} + \frac{c_1}{9} \right) + \frac{m^2}{9} v^2 \phi^2 \right] \hat{\chi}(v, \phi) d\Lambda. \end{aligned} \quad (4.13)$$

In the second term of Eq. (4.13) we replace the term in the brackets [] by the eigenvalue $E(\phi)$ according to Eq. (4.6).

$E(\phi)$ depends on c_2 , c_3 , and Λ only, i.e., there is no ϕ dependence, and we write E instead of $E(\phi)$:

$$0 = \int_{\Lambda} -\hbar^2 \frac{\partial^2}{\partial \phi^2} [b_{\Lambda}(\phi) \hat{\chi}(v, \phi)] + b_{\Lambda}(\phi) E \hat{\chi}(v, \phi) d\Lambda. \quad (4.14)$$

In Eq. (4.14) we first write out the $\partial^2/\partial\phi^2$ applied to the product, but we assume that only $(\partial^2/\partial\phi^2)b_{\Lambda}(\phi)$ is essential, and disregarding the other terms⁹ and assuming that the $\hat{\chi}$'s are all independent we get

$$\left[-\hbar^2 \frac{\partial^2}{\partial \phi^2} + E \right] b_{\Lambda}(\phi) = 0, \quad (4.15)$$

which possesses solutions

$$b_{\Lambda}(\phi) = B_1 e^{\sqrt{E}\phi/\hbar} + B_2 e^{-\sqrt{E}\phi/\hbar}. \quad (4.16)$$

Finally, we get the general wave function

$$\begin{aligned} \psi_{c_2, c_3}^{(\phi)}(\alpha, \phi, r, s) = & (\rho_1 \exp\{\sqrt{c_2}r\} + \rho_2 \exp\{-\sqrt{c_2}r\}) (\sigma_1 \exp\{\sqrt{c_3}s\} + \sigma_2 \exp\{-\sqrt{c_3}s\}) \int_{\Lambda} [B_1 \exp\{\sqrt{\Lambda - (c_2 + c_3)/9}\phi\} \\ & + B_2 \exp\{-\sqrt{\Lambda - (c_2 + c_3)/9}\phi\}] [C_1 J_{\Lambda}(m\phi e^{3\alpha/3\hbar}) + C_2 J_{-\Lambda}(m\phi e^{3\alpha/3\hbar})] d\Lambda \end{aligned} \quad (4.17)$$

with real and for the moment continuous eigenvalues c_2, c_3 . The superscript (ϕ) at ψ indicates that we have treated the scalar field adiabatically. This solution has to be specified by appropriate choices for the constants ρ_1 , ρ_2 , σ_1 , σ_2 , B_1 , B_2 , C_1 , and C_2 from boundary conditions. Before discussing the result (4.17), however, we calculate the wave function by considering the scale factor α as the adiabatic variable.

D. Adiabatic scale factor approach

Another adiabatic method to solve the Wheeler-DeWitt equation (4.4) is to treat the scale factor α as a slowly varying variable.

Once more, we reexpress the scale factor α by $v \equiv e^{3\alpha}$, implying that we consider the Wheeler-DeWitt equation (4.5) in the following. Considering v as the adiabatic variable means that we neglect derivatives with respect to v in a first step. Then Eq. (4.5) becomes

$$\left(-\hbar^2 \frac{\partial^2}{\partial \phi^2} + \frac{\hbar^2}{9} c_1 + \frac{1}{9} m^2 \phi^2 v^2 \right) \hat{\chi}(v, \phi) = E(v) \hat{\chi}(v, \phi) \quad (4.18)$$

with $E(v)$ being a still undetermined eigenvalue. By defining

$$\omega = \frac{1}{3} m v, \quad \tilde{x} = \sqrt{\frac{\omega}{\hbar}} \phi, \quad \eta = \frac{1}{\hbar \omega} \left(E(v) - \frac{\hbar^2}{9} c_1 \right), \quad (4.19)$$

the differential equation (4.18) is simply transformed into that for the dimensionless harmonic oscillator

$$\left[\frac{d^2}{d\tilde{x}^2} + (\eta - \tilde{x}^2) \right] \tilde{y}(\tilde{x}) = 0. \quad (4.20)$$

By substituting $x = \tilde{x}^2$ and then transforming $\tilde{y}(x) = e^{-x/2} y(x)$, we eventually obtain Kummer's differential equation

$$x \frac{d^2 y(x)}{dx^2} + (\mu - x) \frac{dy(x)}{dx} - \nu y(x) = 0, \quad (4.21)$$

where in our case $\mu = 1/2$ and $\nu = (1 - \eta)/4$. This differential equation possesses the general solution

$$y(x) = A_1 F_1(\nu, \mu; x) + B_1 F_1(\nu - \mu + 1, 2 - \mu; x) x^{1-\mu}, \quad (4.22)$$

expressed with the help of the confluent hypergeometric function ${}_1F_1(a, b; x)$. A and B are constants. Thus the partial problem of finding the solution of Eq. (4.18) is done and the result is written as

⁹This is usually called adiabatic approximation with respect to ϕ .

$$\hat{\chi}_\eta(v, \phi) = e^{-\omega(v)\phi^2/2\hbar} \left[A_1 F_1[(1-\eta)/4, 1/2; \omega(v)\phi^2/\hbar] + B \sqrt{\frac{\omega(v)}{\hbar}} \phi_1 F_1[(3-\eta)/4, 3/2; \omega(v)\phi^2/\hbar] \right] \quad (4.23)$$

with $\hat{\chi}_\eta$ denoting the eigenfunction to the real, continuous eigenvalue η .

For further processing to find an adiabatic solution for the Wheeler-DeWitt equation (4.5) we make the integral ansatz

$$\tilde{\chi}(v, \phi) = \int_\eta g_\eta(v) \hat{\chi}_\eta(v, \phi) d\eta. \quad (4.24)$$

Inserting this into Eq. (4.5), we obtain (for $\epsilon=0$)

$$0 = \int_\eta \hbar^2 v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} [g_\eta(v) \hat{\chi}_\eta(v, \phi)] + g_\eta(v) \left[-\hbar^2 \frac{\partial^2}{\partial \phi^2} + \frac{1}{9} (\hbar^2 c_1 + m^2 \phi^2 v^2) \right] \hat{\chi}_\eta(v, \phi) d\eta. \quad (4.25)$$

The expression enclosed in the brackets [] in the second term is substituted by the eigenvalue

$$E(v) = \frac{1}{9} [3\hbar m \eta v - \hbar^2 (c_2 + c_3)], \quad (4.26)$$

following from Eq. (4.18) and the definitions (4.19). In contrast to the preceding approach with an adiabatic scalar field,

$E(v)$ depends explicitly on the adiabatic variable which is in this case v . Note that $E(v)$ also depends on the eigenvalues c_2 , c_3 , and η . Now we utilize the assumption of the adiabaticity with respect to v by neglecting terms containing derivatives $\partial \hat{\chi}_\eta / \partial v$ and $\partial^2 \hat{\chi}_\eta / \partial v^2$ which appear in the first term of Eq. (4.25). Furthermore, supposing that all functions $\hat{\chi}_\eta(v, \phi)$ are independent with respect to η , each integrand in Eq. (4.25) vanishes. Thus we remain with solving the differential equation

$$\left[\hbar^2 v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} + E(v) \right] g(v) = 0 \quad (4.27)$$

or, more explicitly,

$$\left[v^2 \frac{\partial^2}{\partial v^2} + v \frac{\partial}{\partial v} + \kappa_1 v - \frac{\kappa_2^2}{4} \right] g(v) = 0 \quad (4.28)$$

with $\kappa_1 = m \eta / 3\hbar$ and $\kappa_2^2 = 4(c_2 + c_3)/9$. Applying the transformation $x = 2\sqrt{\kappa_1}v$ and denoting the solution as $y(x) = g(v(x))$, Eq. (4.28) takes the same Bessel form as given in Eq. (4.8), whereas the index is κ_1 instead of Λ here. For this reason the general solution of Eq. (4.28) reads

$$g_\eta(v) = D_1 J_{4(c_2+c_3)/9}(\sqrt{4m\eta v/3\hbar}) + D_2 J_{-4(c_2+c_3)/9}(\sqrt{4m\eta v/3\hbar}) \quad (4.29)$$

with constants D_1 and D_2 .

Thus, the complete general wave function obtained with an adiabatic scale factor approach (indicated by superscript α) is found to be

$$\begin{aligned} \psi_{c_2 c_3}^{(\alpha)}(\alpha, \phi, r, s) &= [\rho_1 \exp\{\sqrt{c_2}r\} + \rho_2 \exp\{-\sqrt{c_2}r\}] (\sigma_1 \exp\{\sqrt{c_3}s\} + \sigma_2 \exp\{-\sqrt{c_3}s\}) \\ &\times \int_\eta [D_1 J_{4(c_2+c_3)/9}(\sqrt{4m\eta e^{3\alpha}/3\hbar}) + D_2 J_{-4(c_2+c_3)/9}(\sqrt{4m\eta e^{3\alpha}/3\hbar})] \\ &\times \exp\{-m\phi^2 e^{3\alpha}/6\hbar\} \left[A_1 F_1[(1-\eta)/4, 1/2; m\phi^2 e^{3\alpha}/3\hbar] \right. \\ &\left. + B \sqrt{\frac{m}{3\hbar}} \phi e^{3\alpha/2} F_1[(3-\eta)/4, 3/2; m\phi^2 e^{3\alpha}/3\hbar] \right] d\eta \end{aligned} \quad (4.30)$$

with constants ρ_1 , ρ_2 , σ_1 , σ_2 , A , B , D_1 , and D_2 to be determined via appropriate boundary conditions. The anisotropy quantum numbers c_2 and c_3 are supposed to be real at this stage. Certain boundary conditions, such as requiring the wave function to vanish for infinitely large values of α and/or $-\alpha$, lead to replacing the integral by a sum over η .

E. Visualization of the world function

Now we visualize the results of our numerical calculation of the world function (4.17). To get an impression about the

probability amplitude, we concentrate on the absolute value of the wave function which is given in Figs. 1–6 in dependence of α , the logarithm of the cosmic scale factor, and the scalar field ϕ . Each figure is printed at fixed values of the other parameters, see the corresponding explanation in the text.

For all these essentially different ranges of parameters we got essentially the same picture: The right part of the figure is the region where the WKB approximation is valid, that means, the solution is close to a sinus-shaped wave with a slowly varying amplitude. The left-hand side is the quantum




FIG. 1. The absolute value of the wave function $|\psi|$ (vertical axis) in dependence on α (axis from left to right) and ϕ (axis from front side to back side of the picture) at $r=s=1$ and $c_2=4$ and $c_3=7$.

region where no oscillations exist at all.

The range where these two behaviors go into each other, is relatively sharply defined. This range is what one usually calls the “cosmological quantum boundary,” and these pictures show up that this notion is relatively well defined.

A new feature of these pictures is the following: All of them show a period-doubling bifurcation of the frequency if one looks from the bottom to the top of these pictures.

V. DISCUSSION

A. Short review of the existing literature

Before we discuss the results of the present paper we give a short review to other papers. The original paper [6] by DeWitt was seminal to the whole development of quantum gravity. Its application to minisuperspace cosmological mod-




FIG. 2. The same as Fig. 1, now $c_2=-4$ and $c_3=7$. The different spacing of the $|\psi|$ axis should be noted.




FIG. 3. The same as Fig. 1, now $c_2=-4$ and $c_3=-7$.

els has been developed by several researchers, e.g., the authors and editors of Refs. [7] and [8]. At that time, the main discussion dealt with the closed isotropic Friedmann models, and the spatially flat Friedmann models had been included as less interesting limit cases, too.

In [9] and [10], Schön and Hajicek perform a quantization of systems with quadratic constraints and discuss the Wheeler-DeWitt equation. In Ref. [11], Landsman gives details which Hilbert space might be appropriate for the isotropic minisuperspace quantization, a topic which was not much discussed before.

In the recent preprint by Kim [12], the minimally coupled massless scalar field in an open isotropic Friedmann model has been discussed and its Wheeler-DeWitt equation be solved from the point of view that the Universe is created quantum mechanically from “nothing.”

Recently, Capozziello and Lambiase [13] investigated the connection between the Hartle criterion for selecting correlated regions in the configuration space of dynamical variables and an associated Noether symmetry. This relationship serves to classify solutions of the Wheeler-DeWitt equations in semiclassical minisuperspace models. Thus, the oscilla-





FIG. 4. The same as Fig. 1, now $c_2=0$ and $c_3=0$.

FIG. 5. The same as Fig. 1, now $c_2=1$ and $c_3=1$.

tory behavior of a subset of solutions entails the presence of Noether symmetries which, in the consequence, select classical Universe models.

The paper [14] by Schunck and Mielke is related to the models discussed here as follows: They apply the Wagoner-Bekenstein-Starobinsky-transformation for classifying inflationary solutions with scalar field as source, and this classification should also apply to the corresponding Wheeler-DeWitt equations.

In [15] and [16], a one-parameter set of minisuperspace metrics in arbitrary dimensions is considered, from which we have chosen only that one which gives classically the correct correspondence to Einstein's theory. In [15], the signature of the superspace metric in dependence on the signature of the underlying manifold is evaluated with the result that the normal-hyperbolic character of the Wheeler-DeWitt equation exists only for the Euclidean and the Lorentzian signature of the underlying manifold.

In [17], Horiguchi, Maeda, and Sakamoto perform an expansion of solutions of the Wheeler-DeWitt equation in powers of the Planck length. Vilenkin [18] compares several ap-




FIG. 6. Magnification of a subrange of Fig. 1.

proaches to quantum cosmology, Kim and Page [19] discuss quantum Friedmann models and power-law inflation, Kim [20] compares quantum Friedmann models with conformally and minimally coupled scalar fields. In Ref. [21], Bleyer and Ivashchuk discuss multidimensional cosmological models and their corresponding Wheeler-DeWitt equations. Reference [22] solves the Wheeler-DeWitt equation for scalar fields as the source.

Reference [23] represents the famous paper in which the ‘‘Hartle-Hawking boundary conditions’’ for the Wheeler-DeWitt equation have been derived. One takes a path integral over all such space times V_4 whose boundary is the prescribed spatial hypersurface V_3 .

In Ref. [24], Kiefer constructs wavepackets in minisuperspace for a Friedmann Universe. An adiabatic approach is used in the case of a massive scalar field, thereby assuming the scale parameter a to be slightly changing only. In Zeh [24], these solutions are discussed under the point of view of the definition of the direction of time. Our Eq. (4.1) coincides with the form discussed in Ref. [24] [Eq. (6.5) of Zeh, and Eq. (2.2) of Kiefer]. The approximation of our Sec. IV D was used by Kiefer [24] for solving the Wheeler-DeWitt equation of a Friedmann Universe. In contrast to Kiefer's procedure, where the harmonic oscillator has discrete eigenvalues and thus the wave function decreases for large values of the scale factor α , we allow all real eigenvalues.

Conradi [25] solves the Wheeler-DeWitt equation for Bianchi type-IX model and a massive scalar field. Grishchuk and Sidorov [26] discuss the initial conditions for the Wheeler-DeWitt equation, especially for the massive scalar field in a closed Friedmann model.

In Ref. [27], Amendola, Khalatnikov, Litterio, and Occhionero consider quantum cosmology with a complex field. Guendelmann and Kaganovich [28] discuss cosmic time in quantum cosmology. The factor-ordering problem is solved such that the kinetic term gets the form $\square + \xi R$ where $\xi \ll \xi_{conf}$.

A comparison of the minisuperspace of minimally and conformally coupled scalar fields was done by Page [29]. He solves the factor-ordering problem of the Wheeler-DeWitt equation by requiring that the kinetic term is proportional to the Laplacian in the minisuperspace metric, i.e., our variant B with $\epsilon=1$. The classical equation is similar to the geodesic equation in superspace.

Reference [30] deals with quantum cosmology from the path-integral point of view: The Wheeler-DeWitt equation can be derived in a first approximation from the corresponding path integral. Halliwell [30] solves the factor-ordering problem by requiring invariance with respect to field redefinition of both the three metric and lapse function. Jafarizadeh, Darabi, and Rastegar [30] apply the method of Duru and Kleinert to evaluate the path integral for quantum cosmology, cf. Kleinert [30].

DeWitt [30] assumes the path integral to be the more fundamental approach, but the Wheeler-DeWitt equation in the minisuperspace to remain a good approximation to it.

By considering the Wheeler-DeWitt equation

$$\left(a^{-p} \frac{\partial}{\partial a} a^p \frac{\partial}{\partial a} + a^4 \frac{\Lambda}{3} \right) \psi(a) = 0, \quad \frac{1}{2m} \Delta \psi + \tilde{V} \cdot \psi = 0,$$

Gibbons and Grishchuk [31] obtain the result that inflation is typical in the set of spatially flat Friedmann models in Einstein's theory with a Λ term. The parameter p is due to the factor-ordering ambiguity, they take $p=1$ as a preferred value. An analogous result is given by Hawking and Page [32]. Melnikov and Pevcov [33] discuss the factor-ordering problem for the Wheeler-DeWitt equation in closed and open Friedmann models and give some solutions to it.

Reuter and Schmidt [34] and Schmidt [35] derive the Wheeler-DeWitt equation of fourth-order gravity for a spatially flat Friedmann model and compare with the corresponding conformally equivalent (due to the Bicknell theorem) second-order models. The solution of Reuter-Schmidt for flat Friedmann models is generalized in Ref. [36] by Pimentel and Obregon to closed and open models. Fabris and Reuter [36] continue to generalize the results of [34] to show that the Bicknell theorem applies also at the level of the Wheeler-DeWitt equation.

Rainer [37] gives an overview on higher dimensions and discusses three types of conformal transformations of different levels for the Wheeler-DeWitt equation. Quite recently, the solutions of the Wheeler-DeWitt equation in comparison with the appearance of singularities is treated by Mongan [38]. Zhang and Shen [39] consider quantum cosmology with a complex scalar field at finite temperature. A critical discussion of the Wheeler-DeWitt equation and an alternative quantization scheme is presented in Ref. [40] by Peres.

From recent constraint calculations of Hwang and Noh [41] to an inflation model based on a nonminimally coupled massive scalar field and comparisons with observational data [Cosmic Background Explorer (COBE) Differential Microwave Radiometer (DMR)], one can state that minimal coupling is a good approximation for inflationary models. Following Futamase and Maeda [42], the coupling constant is either quite small, $\xi < 1/1000$, or negative. According to the coupling factor $(1 - \xi \phi^2)$ in front of R , critical behavior appears for $\xi = \phi^{-2}$, i.e., for positive values of ξ only.

B. Comparison with similar results by others

Let us start with explaining the misnomer with Schrödinger and Wheeler-DeWitt equation, see the footnotes to the Introduction and to Sec. III: In quantum mechanics, the classical Hamiltonian \mathbf{H} gives (for a given energy E) rise to the Schrödinger equation $\hat{\mathbf{H}}(\Psi) = E \cdot \Psi$, where Ψ is the time-dependent wave function and $\hat{\mathbf{H}}$ is the operator form of \mathbf{H} . In the most simple case and setting $\hbar = 1$ this equation reads

$$i \frac{\partial \Psi}{\partial t} = - \frac{1}{2m} \Delta \Psi + V \cdot \Psi$$

and is of first order in t . With the ansatz $\Psi = \psi \cdot e^{-iEt}$ we get an equation for the time-independent wave function ψ ; it has the following structure:

where Δ is the spatial Laplacian, i.e., the operator d^2/dx^2 in the one-dimensional case. This equation is of second order, and it is this form of the Schrödinger equation which one had in mind when one compared with the Wheeler-DeWitt equation. However, even this is a yet-to-be-explained comparison: The Laplacian Δ appearing here is an elliptic operator, whereas in the corresponding place of the Wheeler-DeWitt equation we have a differential operator of normal hyperbolic type, a point which was mentioned already many times, e.g. in Ref. [15]. So, if one wishes to make the analogy of the two equations more strict, then one has to allow a term with negative kinetic energy in the classical Hamiltonian \mathbf{H} together with all these known consequences for stability, etc. Of course, formally it is possible to achieve this negativity by an imaginary transformation of the corresponding coordinate, and this procedure does work in the search for exact solutions, but physically, of course, it really changes the system. In other words: The imaginary transformation changing from the Laplace- to the Klein-Gordon-type equation changes also the character of the set of solutions.¹⁰

There exist problems with the probability interpretation of the wave function of the universe, cf. Ref. [43]: Similarly as for the one-dimensional free particle in quantum mechanics, the world function can in general not be normalized, and one has in the result statements about relative probabilities rather than probabilities themselves, see also [44], and the recent review [45] on this subject.

In [44] it is correctly stated that the approach, which is also used here and all similar papers, has a further problem: namely that the homogeneity of the cosmological model is presumed *before* quantization. If taken literally, this would lead, see Ref. [45] (Sec. II), to a violation of the uncertainty principle. This problem should be dealt with by finding an appropriate effective potential to be added.

As we did, Ref. [44] deals with the geometry of an anisotropic Bianchi type-I model. Otherwise the paper [44] deals with the relation to $N=1$ supergravity which is not the topic of the present paper.

References [36] and [45] (Sec. IV) deal with the comparison of the Wheeler-DeWitt equations in different, but classically conformally equivalent theories, especially with fourth-order gravity theories following from a nonlinear Lagrangian $f(R)$, on the one hand, and with scalar-tensor theories, on the other hand. As a rule one can summarize up to different versions how to solve the factor-ordering problems, the conformal relation can be carried over from the classical to the quantum theory.

¹⁰The topological origin of this difference is the noncompactness of the Lorentz group as compared to the compactness of the rotation group, cf. Ref. [49]. In practice, see, e.g., [45] and the references cited there, this has the consequence, that the norm in the corresponding Hilbert space becomes a pseudonorm, i.e., it loses its positive definiteness.

The question for excited states is under debate, and our quantization condition $\hat{\mathcal{H}}\Psi=0$ concerns only the ground state. References [46] and [47] are closely related to ours, so let us compare them in more detail: There such solutions of the Wheeler-DeWitt equation have been found which can be written by a separation ansatz. So, Ref. [47], Eq. (3) (though deduced for the Brans-Dicke theory and a closed Friedmann model), which contains the linear combination of two Hankel functions in dependence on $a^2\phi$, can be compared with our solution (4.12) representing the linear combination of two Bessel functions (which is almost the same).

A further point to be discussed is the relation mentioned in [47] of the quantum cosmology with quantum wormholes; here we only mention a further approach [48] dealing with wormhole-type solutions.

C. Discussion of the present paper

Let us now summarize our results: We solved the Wheeler-DeWitt equation for the minisuperspace of a cosmological Bianchi type-I model with a minimally coupled massive scalar field ϕ as the source by generalizing the calculation of Lukash and Schmidt [1]. Contrarily to other approaches we allowed strong anisotropy.

Combining analytical and numerical methods, we applied an adiabatic approximation for ϕ , and as new feature we found a period-doubling bifurcation of the typical solutions. This bifurcation takes place near the cosmological quantum boundary, i.e., the boundary of the quasiclassical region with an oscillating ψ function where the WKB approximation is good. The numerical calculations suggest that such a notion of a ‘‘cosmological quantum boundary’’ is well defined, because sharply beyond that boundary, the WKB approximation is no longer applicable.

This result confirms the adequateness of the introduction of a cosmological quantum boundary in quantum cosmology. We applied the supercovariance principle, i.e., the underlying theory should also be covariant with respect to transformations representing a mixture between space time and matter degrees of freedom. With our figures, we tried to visualize the birth of the Universe.

ACKNOWLEDGMENTS

M.B. is supported by HSP III and the Studienstiftung des deutschen Volkes. H.-J. S. gratefully acknowledges financial support from HSP III and DFG (KL 256/31-2). We thank the colleagues of the Free University Berlin, where this work has been done, especially Professor H. Kleinert and Dr. A. Pelster, for kind hospitality, and Professor V. Gurovich and Dr. A. Kirillov for valuable comments.

APPENDIX: VARIANT A VS VARIANT B

Until Eq. (4.6) we parallelly dealt with variant A ($\epsilon=0$) and B ($\epsilon=1$), but beginning from Eq. (4.7) we simplified by restricting to variant A. Now we want to complete the calculation by showing what changes using variant B.

The purpose of this Appendix is to show in detailed calculations what has been verbally mentioned in the literature several times, namely the fact that differences in solving the factor-ordering problem do not essentially change the results. For this consideration we now set $\epsilon=1$ in Eq. (4.6) and use the definitions below that equation:

$$\left(\left[v \frac{\partial}{\partial v} v \frac{\partial}{\partial v} + v \frac{\partial}{\partial v} \right] + \gamma^2 v^2 - \Lambda^2 \right) \hat{\chi}(v, \phi) = 0. \quad (\text{A1})$$

The transformation from Eq. (4.7) to Eq. (4.8) now leads to

$$x^2 \frac{d^2 y}{dx^2} + 2x \frac{dy}{dx} + (x^2 - \Lambda^2)y = 0, \quad (\text{A2})$$

whose solutions are also Bessel functions, modified by a factor $x^{-1/2}$:

$$y(x) = x^{-1/2} [\tilde{C}_1 J_{\sqrt{\Lambda^2+1/4}}(x) + \tilde{C}_2 J_{-\sqrt{\Lambda^2+1/4}}(x)] \quad (\text{A3})$$

where \tilde{C}_1, \tilde{C}_2 are constants.

This means that the wave function (4.17) using variant A is changed by multiplying by a factor $\sqrt{3\hbar} e^{-3\alpha/m\phi}$. Moreover, the index Λ of the Bessel functions must be replaced by $\sqrt{\Lambda^2+1/4}$.

-
- [1] V. Lukash, H.-J. Schmidt, *Astron. Nachr.* **309**, 25 (1988); H.-J. Schmidt, *J. Math. Phys.* **37**, 1244 (1996).
[2] P. Amsterdamski, *Phys. Rev. D* **31**, 3073 (1985).
[3] C. Da Silva and R. Williams, *Class. Quantum Grav.* **16**, 2681 (1999).
[4] A. Pelster, Ph.D. thesis, University of Stuttgart, Germany, 1996.
[5] H. Kleinert, *Phys. Lett. B* **460**, 36 (1999).
[6] B. DeWitt, *Phys. Rev.* **160**, 1113 (1967).
[7] L. P. Grishchuk and Yu. V. Sidorov, in *Proceedings of the 4th Seminar on Quantum Gravity*, edited by M. Markov, V. Berezin, and V. Frolov (World Scientific, Singapore, 1988), p. 700.
[8] D. N. Page, in *Proceedings of the 5th Seminar on Quantum Gravity*, edited by M. Markov, V. Berezin, and V. Frolov (World Scientific, Singapore, 1991), p. 82.
[9] M. Schön and P. Hajicek, *Class. Quantum Grav.* **7**, 861 (1990).
[10] P. Hajicek, *Class. Quantum Grav.* **7**, 871 (1990).
[11] N. Landsman, in *Current Topics in Mathematical Cosmology*, edited by M. Rainer, and H.-J. Schmidt (World Scientific, Singapore, 1998), p. 256.
[12] S. P. Kim, gr-qc/9909002, 1999.
[13] S. Capozziello and G. Lambiase, *Gen. Relativ. Gravit.* **32**, 673 (2000).
[14] F. Schunck and E. Mielke, *Phys. Rev. D* **50**, 4794 (1994).
[15] H.-J. Schmidt, in *Differential Geometry and Applications*, edited by J. Janyška and D. Krupka (World Scientific, Singapore, 1990), p. 405.
[16] S. Odintsov and I. Ševčenko, *Isv. Vuzov Fizika (Univ. Tomsk)* **7**, 74 (1991); D. Giulini, *Phys. Rev. D* **51**, 5630 (1995).

- [17] T. Horiguchi, K. Maeda, and M. Sakamoto, *Phys. Lett. B* **344**, 105 (1995).
- [18] A. Vilenkin, *Phys. Rev. D* **50**, 2581 (1994).
- [19] S. Kim and D. Page, *Phys. Rev. D* **45**, R3296 (1992).
- [20] S. Kim, *Phys. Rev. D* **46**, 3403 (1992).
- [21] U. Bleyer and V. Ivashchuk, *Phys. Lett. B* **332**, 292 (1994).
- [22] D. Page, in *Quantum Concepts in Space and Time*, edited by C. Isham and R. Penrose (Oxford University Press, Oxford, 1985), p. 274.
- [23] J. Hartle and S. Hawking, *Phys. Rev. D* **28**, 2960 (1983).
- [24] C. Kiefer, *Phys. Rev. D* **38**, 1761 (1988); H. Zeh, *The Physical Basis of the Direction of Time* (Springer-Verlag, Berlin, 1992).
- [25] H. Conradi, *Phys. Rev. D* **46**, 612 (1992).
- [26] L. Grishchuk and Ju. Sidorov, *Zh. Éksp. Teor. Fiz.* **94**, 29 (1988) [*Sov. Phys. JETP* **67**, 1533 (1988)].
- [27] L. Amendola, I. Khalatnikov, M. Litterio, and F. Occhionero, *Phys. Rev. D* **49**, 1881 (1994).
- [28] E. Guendelmann and A. Kaganovich, *Int. J. Mod. Phys. D* **2**, 221 (1993).
- [29] D. Page, *J. Math. Phys.* **32**, 3427 (1991).
- [30] J. Halliwell, *Phys. Rev. D* **38**, 2468 (1988); H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, and Polymer Physics*, 2nd ed. (World Scientific, Singapore, 1995); M. Jafarizadeh, F. Darabi, and A. Rastegar, *Phys. Lett. A* **248**, 19 (1998); B. DeWitt, in *Proceedings of the 8th M. Grossmann Meeting*, Jerusalem, edited by T. Piran (World Scientific, Singapore, 1999).
- [31] G. Gibbons and L. Grishchuk, *Nucl. Phys.* **B313**, 736 (1989).
- [32] S. Hawking and D. Page, *Nucl. Phys.* **B264**, 185 (1986).
- [33] V. Melnikov and G. Pevcov, *Izv. Vuzov Fizika (Tomsk)* **4**, 45 (1985).
- [34] S. Reuter and H.-J. Schmidt, in *Proceedings of the 5th International Conference on Differential Geometry and Applications*, edited by O. Kowalski and D. Krupka (Silesian University, Opava, 1993), p. 243.
- [35] H.-J. Schmidt, *Phys. Rev. D* **49**, 6354 (1994); **54**, 7906(E) (1996).
- [36] L. Pimentel and O. Obregon, *Class. Quantum Grav.* **11**, 2219 (1994); J. Fabris and S. Reuter, *Gen. Relativ. Gravit.* (to be published).
- [37] M. Rainer, *Grav. Cosmol.* **1**, 121 (1995).
- [38] T. Mongan, *Gen. Relativ. Gravit.* **31**, 1429 (1999).
- [39] T. Zhang and Y. Shen, *Int. J. Theor. Phys.* **38**, 1969 (1999).
- [40] A. Peres, in *On Einstein's Path, Essays in Honour of E. Schücking*, edited by A. Harvey (Springer-Verlag, Berlin, 1999): When moving to the United States, Schücking changed his name to Schucking, so both spellings of his name are equally correct.
- [41] J. Hwang and H. Noh, *Phys. Rev. D* **60**, 123001 (1999).
- [42] T. Futamase and K. Maeda, *Phys. Rev. D* **39**, 399 (1989).
- [43] M. Ryan, *Hamiltonian Cosmology* (Springer, New York, 1972).
- [44] A. Macias, O. Obregon, and M. Ryan, *Class. Quantum Grav.* **4**, 1477 (1987).
- [45] A. Macias, *Gen. Relativ. Gravit.* **31**, 653 (1999).
- [46] L. Pimentel, O. Obregon, and J. Rosales, *Class. Quantum Grav.* **14**, 379 (1997).
- [47] L. Pimentel and C. Mora, in *Proceedings of the Marcel Grossmann Meeting MG8 Jerusalem*, edited by T. Piran and R. Ruffini (World Scientific, Singapore, 1999), p. 1029.
- [48] V. Dzhunushaliev and H.-J. Schmidt, *Phys. Rev. D* (to be published), gr-qc/9911080.
- [49] H.-J. Schmidt, *Int. J. Theor. Phys.* **37**, 691 (1998).