

B_c meson and the light-heavy quarkonium spectrum

M. Baldicchi and G. M. Prosperi

Dipartimento di Fisica dell'Università di Milano, and I.N.F.N., Sezione di Milano, Milano, Italy

(Received 2 August 2000; published 9 November 2000)

We compute the $c\bar{b}$ spectrum from a first principles Salpeter equation obtained in a previous paper. For comparison we report also the heavy-light quarkonium spectrum and the hyperfine separations previously presented only in a graphical form. Notice that all results are parameter free.

PACS number(s): 12.38.Aw, 11.10.St, 12.38.Lg, 12.39.Ki

The spectrum and properties of the $c\bar{b}$ systems have been calculated various times in the past in the framework of the heavy quarkonium theory [1]. However the recent experimental observation of the B_c^+ meson [2] has inspired new theoretical interest in the problem [3–5]. The mentioned spectrum has been considered again either from the potential or the lattice simulation point of view. A particularly interesting quantity should be the hyperfine splitting that as for the $c\bar{c}$ case seems to be sensitive to relativistic and subleading corrections in α_s .

For the above reasons it seems worthwhile to present in this paper a calculation of the $c\bar{b}$ spectrum based on an effective mass operator with full relativistic kinematics which we have obtained in previous works and applied with a certain success to fit the entire quarkonium spectrum: heavy-heavy, light-light, and light-heavy cases with the exception however of the $c\bar{b}$ case [6,7]. For comparison and completeness we also report numerical results for the light-heavy spectrum which we have previously given only in graphical form.

The mass operator was obtained by a three dimensional reduction of the $q\bar{q}$ Bethe-Salpeter (BS) equation introduced in [8]. It has the quadratic form $M^2 = M_0^2 + U$, with a kinetic part $M_0 = w_1 + w_2 = \sqrt{m_1^2 + \mathbf{k}^2} + \sqrt{m_2^2 + \mathbf{k}^2}$ and a ‘‘potential’’ that in terms of the instantaneous approximation of the BS kernel is given by

$$\langle \mathbf{k} | U | \mathbf{k}' \rangle = \frac{1}{(2\pi)^3} \sqrt{\frac{w_1 + w_2}{2 w_1 w_2}} \hat{t}_{\text{inst}}(\mathbf{k}, \mathbf{k}') \sqrt{\frac{w_1' + w_2'}{2 w_1' w_2'}} \quad (1)$$

with \mathbf{k} denoting the momentum of the quark in the center of mass frame, and $i = 1, 2$ the quark and the antiquark.

The BS equation was derived from QCD first principles, taking advantage of the Feynman-Schwinger path integral representation for the ‘‘second order’’ quark propagator in an external field.¹ The only assumption used consisted in writing the logarithm of the Wilson loop correlator $W = \frac{1}{3} \langle \text{Tr} P \exp(\oint dx^\mu A_\mu) \rangle$, as the sum of its perturbative expression and an area term

¹Second order propagator in the sense that it is defined by a second order differential equation; the quadratic form of the mass operator derives essentially from this fact.

$$i \ln W = i(\ln W)_{\text{pert}} + \sigma S_{\text{min}}, \quad (2)$$

with σ denoting the string tension.

An explicit expression for U is given in Ref. [7]. The perturbative part of such quantity was evaluated at the lowest order in α_s . However for α_s we have used the standard running expression

$$\alpha_s(\mathbf{Q}) = \frac{4\pi}{(11 - \frac{2}{3}N_f) \ln(\mathbf{Q}^2/\Lambda^2)} \quad (3)$$

(with $N_f = 4$ and $\Lambda = 200$ MeV) cut at a maximum value $\alpha_s(0)$, to treat properly the infrared region [9]. This amounts to include important perturbative subleading contributions.

Notice that, contrary to all the usual potential models, we have given the light quark current and not component masses in our treatment. Component masses of the usual order of magnitude can be recovered at a successive step as effective values in a semirelativistic reformulation [6]. Actually, we have fixed such masses on typical values, $m_u = m_d = 10$ MeV, $m_s = 200$ MeV, which are not adjusted in the fit (the results depend essentially on $\langle k \rangle$ and are affected very little by the precise value of the light quark masses). The other parameters of the theory are assumed to be: $m_c = 1.394$ GeV, $m_b = 4.763$ GeV, $\sigma = 0.2$ GeV², and $\alpha_s(0) = 0.35$. The first two are chosen in order to reproduce cor-

TABLE I. $b\bar{c}$ quarkonium systems. Experimental B_c mass equal to $6.40 \pm 0.39 \pm 0.13$ GeV.

States	Quadratic formalism (GeV)	Linear formalism (GeV)	Fulcher (GeV)	Lattice (GeV)
1 1S_0	6.258	6.293	6.286	6.280 ± 0.200
1 3S_1	6.334	6.355	6.341	6.321 ± 0.200
2 1S_0	6.841	6.848	6.882	6.960 ± 0.080
2 3S_1	6.883	6.881	6.914	6.990 ± 0.080
3 1S_0	7.222	7.221		
3 3S_1	7.254	7.245		
1 P	6.772	6.762	6.754	6.764 ± 0.030
2 P	7.154	7.138		
1 D	7.043	7.025	7.028	
2 D	7.367	7.346		

TABLE II. Light-heavy quarkonium systems.

States		Experimental values (MeV)	Linear formalism (MeV)	Quadratic formalism (MeV)
$u\bar{c}$				
1^1S_0	$\begin{cases} D^\pm \\ D^0 \end{cases}$	$\begin{cases} 1869.3 \pm 0.5 \\ 1864.5 \pm 0.5 \end{cases}$	1890	1875
1^3S_1	$\begin{cases} D^*(2010)^\pm \\ D^*(2007)^0 \end{cases}$	$\begin{cases} 2010.0 \pm 0.5 \\ 2006.7 \pm 0.5 \end{cases}$	2001	2020
2^1S_0	D'	2580	2556	2525
2^3S_1	$D^{*'} $	2637 ± 8	2615	2606
$1 P$	$\begin{cases} D_2^*(2460)^\pm \\ D_2^*(2460)^0 \\ D_1(2420)^\pm \\ D_1(2420)^0 \end{cases}$	$\begin{cases} 2459 \pm 4 \\ 2458.9 \pm 2.0 \\ 2427 \pm 5 \\ 2422.2 \pm 1.8 \end{cases}$	2442	2475
Δ_{avg}			12	21
$u\bar{b}$				
1^1S_0	$\begin{cases} B^\pm \\ B^0 \end{cases}$	$\begin{cases} 5278.9 \pm 1.8 \\ 5279.2 \pm 1.8 \end{cases}$	5282	5273
1^3S_1	B^*	5324.8 ± 1.8	5341	5339
2^1S_0			5878	5893
2^3S_1	$B^{*'} $	5906 ± 14	5916	5933
$1 P$		5825 ± 14	5777	5792
Δ_{avg}			34	19
$s\bar{c}$				
1^1S_0	D_s^\pm	1968.5 ± 0.6	1999	1982
1^3S_1	$D_s^{*\pm}$	2112.4 ± 0.7	2107	2120
2^1S_0			2667	2617
2^3S_1			2729	2698
$1 P$	$\begin{cases} D_{sJ}(2573)^\pm \\ D_{s1}(2536)^\pm \end{cases}$	$\begin{cases} 2573.5 \pm 1.7 \\ 2535.35 \pm 0.34 \end{cases}$	2528	2548
Δ_{avg}			21	9
$s\bar{b}$				
1^1S_0	B_s^0	5369.3 ± 2.0	5373	5364
1^3S_1	B_s^*	5416.3 ± 3.3	5433	5429
2^1S_0			5974	5985
2^1S_0			6014	6024
$1 P$	$B_{sJ}^*(5850)$	5853 ± 15	5848	5859
Δ_{avg}			5	4

rectly the J/ψ and the $\Upsilon(1S)$ masses, the string tension to give the correct slope for the Regge ρ trajectory, $\alpha_s(0) = 0.35$ to give the right $J/\psi - \eta_c$ splitting. Notice that, consequently, the results reported in this paper are completely parameter free, with the exception of the $c\bar{c}(1S)$ hyperfine

splitting.

We have also used in our calculations the more conventional ‘‘linear mass’’ operator (or center of mass relativistic Hamiltonian) $M = M_0 + V$ (where V is defined by $U = M_0 V + VM_0 + V^2$) which makes an easier comparison with the

TABLE III. Hyperfine splitting (MeV).

	$u\bar{c}$	$s\bar{c}$	$u\bar{b}$	$s\bar{b}$	$c\bar{c}$	$c\bar{b}$	$b\bar{b}$	$u\bar{u}$	$u\bar{s}$	$s\bar{s}$
1 S	145	138	66	65	115	77	86	349	298	259
Exp	141(1)	144(1)	46(3)	47(4)	117(2)	-	-	630.5(0.6)	393.92(0.24)	335.3(0.1)
2 S	81	81	40	39	67	42	35	135	130	127
Exp	57	-	-	-	92(5)	-	-	165(103)	-	-
Δ_{avg}	21	19	9	4	20	-	10	19	48	18

usual phenomenological models. If we neglect the V^2 term, V is obtained from Eq. (1) simply by the kinematical replacement

$$\sqrt{\frac{w_1 + w_2}{2w_1w_2}} \sqrt{\frac{w'_1 + w'_2}{2w'_1w'_2}} \rightarrow \frac{1}{4\sqrt{w_1w_2w'_1w'_2}}. \quad (4)$$

This is the form we have used in Ref. [6] (for some state however $\langle V^2 \rangle$ is not negligible). In the calculations based on this linear formalism we have used the same values for the light quark masses as before, a fixed coupling constant $\alpha_s = 0.363$ and taken $m_c = 1.40$ GeV, $m_b = 4.81$ GeV, and $\sigma = 0.175$ GeV².

Details on the numerical treatment of the eigenvalue equation are given in [6] and [7]. In Table I we have reported the $c\bar{b}$ spectrum as obtained by the quadratic and the linear formalism, together with the values presented in Refs. [4] and [5]. The observed mass $M(B_c) = 6.40 \pm 0.39 \pm 0.13$ GeV has to be referred to the 1^1S_0 state. For such a state, all calculations give very close results and reproduce the experimental value equally well within the errors. Larger discrepancies among the various methods occur for the excited states.

In Table II we have reported the spectrum for light-heavy mesons obtained by our formalism in numerical form. We have considered the hyperfine structure but omitted the fine one. We have also reported the quantity Δ_{avg} defined as the

average of the deviations of the theoretical values from the experimental data diminished by the experimental errors. Obviously Δ_{avg} provides a measure of the accuracy in reproducing the data and gives an idea of the precision one can expect in the $c\bar{b}$ case.

Finally, in Table III, we have reported the hyperfine splitting for the $1S$ and $2S$ states as obtained in the quadratic formalism and the Δ_{avg} quantity even for the channels for which we do not reproduce the full results here.

Notice the strong discrepancies with the data in the hyperfine splittings of the $1S$ light-light cases. This is obviously due to the chiral symmetry breaking problem and the related inadequacy of replacing the quark full propagator in the BS equation with the free form, as implied in the three-dimensional reduction. For the rest, the agreement is good for the states involving light and c quarks, while the theoretical value tends to be too large for states involving b quarks.

For comparison we can mention that in the linear formalism the hyperfine splitting turns out less agreeable, being, e.g., 97 MeV for $c\bar{c}(1S)$, 111 MeV for $u\bar{c}(1S)$, and 108 MeV for $c\bar{s}(1S)$. Such a difference has to be ascribed to relativistic and α_s subleading effects, taken into account in the quadratic formalism in Eq. (3).

In conclusion, let us mention explicitly that Δ_{avg} as reported in Table III does not include the states $c\bar{c}(4S)$ and $b\bar{b}(6S)$ (which are largely above threshold) and the 1^1S_0 and 1^1P_1 light-light states for the reasons recalled above.

[1] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985); A. Martin, in *Proceedings in Heavy Flavours and High-Energy Collisions in the 1-100 TeV Range*, edited by A. Ali and L. Cifarelli (Plenum, New York, 1989), p. 141; W. Kwong and J.L. Rosner, Phys. Rev. D **44**, 212 (1991); E.J. Eichten and C. Quigg, *ibid.* **49**, 5845 (1994); E. Bagan, H.G. Dosch, P. Godzinsky, S. Narison, and J.M. Richard, Z. Phys. C **64**, 57 (1994); R. Roncaglia, A. Dzierba, D.B. Lichtenberg, and E. Predazzi, Phys. Rev. D **51**, 1248 (1995).

[2] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **81**, 2432 (1998); CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **58**, 112004 (1998); OPAL Collaboration, K. Ackerstaff *et al.*, Phys. Lett. B **420**, 157 (1998); ALEPH Collaboration, R. Barate *et al.*, *ibid.* **402**, 231 (1997); DELPHI Collaboration, P. Abreu *et al.*, *ibid.* **398**, 207 (1997).

[3] N. Brambilla and A. Vairo, Phys. Rev. D **62**, 094019 (2000); P. Colangelo and F. De Fazio, *ibid.* **61**, 034012 (2000).

[4] L.P. Fulcher, Phys. Rev. D **60**, 074006 (1999).

[5] C.T.H. Davis, K. Hornbostel, G.P. Lepage, A.J. Lidsey, J. Shigemitsu, and J. Sloan, Phys. Lett. B **382**, 131 (1996).

[6] M. Baldicchi and G.M. Prosperi, Phys. Lett. B **436**, 145 (1998).

[7] M. Baldicchi and G.M. Prosperi, Fiz. B **8**, 251 (1999).

[8] N. Brambilla, E. Montaldi, and G.M. Prosperi, Phys. Rev. D **54**, 3506 (1996).

[9] Yu.L. Dokshitzer, A. Lucenti, G. Marchesini, and G.P. Salam, J. High Energy Phys. **05**, 003 (1998).