

Semileptonic form factors: A model-independent approach

Todd Coleman and M. G. Olsson

Department of Physics, University of Wisconsin, 1150 University Avenue, Madison, Wisconsin 53706

Siniša Veseli

Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, Illinois 60510

(Received 15 September 2000; published 11 January 2001)

We demonstrate that the $B \rightarrow D^{(*)} l \nu$ form factors can be accurately predicted given the slope parameter ρ^2 of the Isgur-Wise function. Only weak assumptions, consistent with lattice results, on the wave function for the light degrees of freedom are required to establish this result. We observe that the QCD and $1/m_Q$ corrections can be systematically represented by an effective Isgur-Wise function of shifted slope. This greatly simplifies the analysis of semileptonic B decay. We also investigate what the available semileptonic data can tell us about lattice QCD and heavy quark effective theory. A rigorous identity relating the form factor slope difference $\rho_{D^*}^2 - \rho_{A_1}^2$ to a combination of form factor intercepts is found. The identity provides a means of checking theoretically evaluated intercepts with experiment.

DOI: 10.1103/PhysRevD.63.032006

PACS number(s): 13.20.He

I. OUTLINE

We obtain a nearly model-independent description of the Isgur-Wise (IW) function [1] $\xi(w)$ in terms of a single measurable parameter—the slope at zero-recoil, ρ^2 . Only modest assumptions about the shape of the heavy-light wave function are required, and these are consistent with an established lattice result. We obtain a simple functional form for the IW function in terms of ρ^2 :

$$\xi(w, \rho^2) = \frac{2(w+1)}{\left[w+1 + \left(\rho^2 - \frac{1}{2} \right) (w-1) \right]^2}. \quad (1.1)$$

We also demonstrate that the effect of radiative QCD corrections and $1/m$ corrections can be described as the product of a term linear in w and the IW function. The resulting function is also approximately an IW function of shifted slope. Thus, fitting the above functional form to the measured form factors provides an accurate description of the extrapolation to zero-recoil. Over the range of expected slopes, our result is in good agreement with the work of [2–5] which is based on dispersion relations.

Recent analyses [6,7] of $B \rightarrow D l \nu$ and $B \rightarrow D^* l \nu$ decays extract two physical form factors and two ratios of physical form factors. We show that this information can be directly related to QCD predictions.

In Sec. II we provide some background formalism behind form factors for semileptonic B -decay and the Isgur-Wise function. We present our description of the IW function in terms of the slope parameter, ρ^2 , in Sec. III. In Sec. IV we discuss the effects of corrections to the heavy-quark limit and their equivalence to slope-shifted IW functions. The relationship of heavy quark effective theory (HQET) and QCD lattice simulation to semileptonic data is explored in Sec. V.

II. INTRODUCTION

As the observed sample of semileptonic B decays accumulates, the need for a rigorous method of analysis becomes more pressing. Even in the heavy quark limit, a universal form factor is required for each light degree of freedom state of the meson. To compute the Isgur-Wise (IW) functions $\xi(w)$, where $w = v^\mu v'_\mu$, requires considerable knowledge of the heavy-light meson dynamics. For this reason, the IW function is usually thought of as “model-dependent” and not susceptible to reliable calculation. In the first portion of this paper we seek a balance between rigorous constraints and phenomenological usefulness. We will demonstrate, from heavy quark symmetry and some qualitative lattice results, that the IW function is accurately determined by specifying only the slope parameter, ρ^2 .

Experimentalists commonly adopt the straightforward procedure of expanding form factors about the meson zero-recoil point ($w = 1$):

$$\xi(w) = 1 - \rho^2(w-1) + c(w-1)^2 + \dots \quad (2.1)$$

Burdman [8] has pointed out that the effect of the curvature term c is significant but that, for statistical reasons, much predictive power is lost when it is a free parameter. A “model-independent” relation between c and the slope parameter ρ^2 has been proposed by Boyd, Grinstein and Lebed [2]. This method has been modified by Caprini and Neubert [3] and by Caprini, Lellouch, and Neubert in expanded form [4]. This result has been criticized however by Boyd, Grinstein, and Lebed [5]. These latter authors propose a similar but weaker relation between c and ρ^2 . In Sec. III, we propose a rigorous one-parameter expression for the IW function.

The second observation we make here concerns the relation of the IW function to the actual (physical) form factors with $1/m_Q$ and QCD radiative corrections. We show in Sec. IV that all of these corrections can be distilled into a new effective IW function with a shift in slope. The analysis of

experimental data is thereby greatly simplified. The slopes appearing in $B \rightarrow D$ and $B \rightarrow D^*$ will be different and this difference can be compared to the theoretical predictions.

The decay rate for $B \rightarrow D^{(*)} l \bar{\nu}$ can be written [9] as

$$\frac{d\Gamma}{dw} = \frac{G_F^2 |V_{cb}|^2}{48\pi^3} m_B^2 m_{D^{(*)}}^3 \sqrt{w^2 - 1} f^{(*)} |F^{(*)}|^2, \quad (2.2)$$

where, for $B \rightarrow D l \bar{\nu}$,

$$\begin{aligned} f &= (w^2 - 1)(1 + r)^2, \\ r &= \frac{m_D}{m_B} \approx 0.35, \end{aligned} \quad (2.3)$$

and for $B \rightarrow D^* l \bar{\nu}$

$$\begin{aligned} f^* &= (w + 1)[(w + 1)(1 - r^*)^2 + 4w(1 - 2wr^* + r^{*2})], \\ r^* &= \frac{m_{D^*}}{m_B} \approx 0.38. \end{aligned} \quad (2.4)$$

The two form factors $F(w)$ and $F^*(w)$ can be expressed [10] in terms of the fundamental form factors h_+ , h_- , h_V , h_{A_1} , h_{A_2} , and h_{A_3} . These form factors can in turn be related to the Isgur-Wise function through

$$h_i(w) = [\alpha_i + \beta_i(w) + \gamma_i(w)] \xi(w), \quad (2.5)$$

where $\alpha_+ = \alpha_V = \alpha_{A_1} = \alpha_{A_3} = 1$ and $\alpha_- = \alpha_{A_2} = 0$. The $\beta_i(w)$ describe perturbative radiative corrections and are in principle predictable within the heavy quark effective theory. The $\gamma_i(w)$ represent the power ($1/m_Q$) deviations from heavy quark symmetry and require further theoretical assumptions [10]. In Sec. IV we will point out that, for those form factors which do not vanish in the heavy quark limit, the pre-factors in Eq. (2.5) can nevertheless be approximately absorbed into an effective IW function which has a flavor and spin-dependent slope

$$h_i(w) \approx h_i(1) \xi(w, \rho_i^2). \quad (2.6)$$

III. MODEL INDEPENDENT PARAMETRIZATION OF THE ISGUR-WISE FUNCTION

The IW function appears to require detailed information on the nature of heavy-light dynamics. In this section, we propose a method to distill most of this knowledge into a single parameter—the slope ρ^2 . Almost all model dependence will be seen to vanish once the slope is specified.

In the heavy-light limit, heavy quark symmetry provides a unique prescription [11,12] relating the IW function to the wave function and light degrees of freedom energy E ,

$$\xi(w) = \frac{2}{w+1} \left\langle j_0 \left(2Er \sqrt{\frac{w-1}{w+1}} \right) \right\rangle. \quad (3.1)$$

The expectation value may involve a multi-component wave function such as from the Dirac equation. For simplicity of

notation, we assume a single component wave function $\psi(\mathbf{r}) = R(r) Y_l^m(\theta, \phi)$. The expectation value is consequently defined by

$$\langle A(r) \rangle \equiv \int_0^\infty dr r^2 R^2(r) A(r). \quad (3.2)$$

Using $\lim_{y \ll 1} j_0(y) = 1 - \frac{1}{6}y^2 + \dots$, we may expand Eq. (3.1) about the zero recoil point $w = 1$ to yield

$$\xi(w) \approx \left[1 - \frac{1}{2}(w-1) + \dots \right] \left[\langle 1 \rangle - \frac{4E^2}{6} \left(\frac{w-1}{w+1} \right) \langle r^2 \rangle + \dots \right] \quad (3.3)$$

$$\xi(w) \approx 1 - \rho^2(w-1) + \dots \quad (3.4)$$

where

$$\rho^2 = \frac{1}{2} + \frac{E^2}{3} \langle r^2 \rangle. \quad (3.5)$$

We now use Eq. (3.5) to eliminate the energy E in the general expression (3.1) to obtain

$$\xi(w) = \frac{2}{w+1} \left\langle j_0 \left(\frac{r}{r_{rms}} \sqrt{\frac{12 \left(\rho^2 - \frac{1}{2} \right) (w-1)}{w+1}} \right) \right\rangle, \quad (3.6)$$

where $r_{rms} = \sqrt{\langle r^2 \rangle}$.

In the above Eq. (3.6), if we know the slope (ρ^2) and the radial ground state heavy quark wave function, we can compute the IW function for all w . We can make the wave function dependence more explicit by introducing the dimensionless quantities

$$x \equiv \frac{r}{r_0},$$

$$R(r) = R_0 f \left(\frac{r}{r_0} \right) = R_0 f(x), \quad (3.7)$$

where r_0 is a hadronic scale factor. We define the n^{th} moment of f^2 as

$$N_n \equiv \int_0^\infty dx x^{n+2} f^2(x) \quad (3.8)$$

so that wave function normalization and $\langle r^2 \rangle$ become

$$\begin{aligned} r_0^3 R_0^2 N_0 &= 1, \\ \langle r^2 \rangle &= r_0^5 R_0^2 N_2 = r_0^2 \frac{N_2}{N_0}. \end{aligned} \quad (3.9)$$

The IW function (3.6) is then




FIG. 1. The lattice data [13] for the heavy-light wave function along with curves corresponding to the exponential parametrization (3.11) with $k=0.5$, $k=1.0$, and $k=2.0$. It is readily seen that $k=1.0$, which corresponds to a simple exponential, provides a good fit.

$$\xi(w) = \frac{2}{w+1} \frac{1}{N_0} \int_0^\infty dx x^2 f^2(x) j_0 \times \left(x \sqrt{\frac{12 \left(\rho^2 - \frac{1}{2} \right) (w-1) N_0}{(w+1) N_2}} \right). \quad (3.10)$$

It should be noted that both the normalization constant R_0 and the hadronic scale parameter r_0 do not appear in the above expression for the IW function. The IW function thus depends only on the dimensionless slope parameter ρ^2 and one dimensionless function $f(x)$ which is essentially the heavy-light wave function.

A few years ago, the heavy-light wave function was evaluated in quenched lattice simulation [13]. The result for the ground state is shown in Fig. 1. One may observe that the

lattice wave function closely resembles a simple exponential. We use this lattice result as a guide and parametrize the wave function as [14]

$$f(x) = e^{-x^k}. \quad (3.11)$$

When $k=1$ the wave function is a simple exponential and the lattice simulation is closely reproduced. In Fig. 1, we also show that by choosing $k=0.5$ and $k=2$ we very conservatively bracket the observed wave function. Assuming the heavy-light wave function is given by Eq. (3.11) with

$$0.5 \leq k \leq 2, \quad (3.12)$$

the IW function is determined for all w once the slope is specified. In Fig. 2 we show the predicted IW function with $\rho^2=0.9$. The central curve corresponds to $k=1$ with a cor-




FIG. 2. Wave function sensitivity: The corridor for the Isgur-Wise function as determined using $k=0.5$ and $k=2.0$. The solid line corresponds to $k=1.0$. All curves have the same slope parameter, $\rho^2=0.9$.




FIG. 3. Equivalence of a ‘physical’ form factor to an Isgur-Wise function of a shifted slope. The ‘shifted’ IW function $\xi(w, \rho^2=0.9)$ and the physical form factor approximation $\xi(w, \rho^2=1.0)[1+0.1(w-1)]$ differ only about 2% at $w=1.5$.

ridor implied by the limits of Eq. (3.12). We note that at $w=1.5$, which is approximately the largest allowed for $B \rightarrow D^{(*)} l \nu$ decay, the corridor width is only about 0.02, or 3%. This small uncertainty shows that most of the IW shape is well-determined by the slope alone.

A particularly important special case is when $k=1$. Straightforward evaluation of Eq. (3.10) with $k=1$ in Eq. (3.11) yields the IW function

$$\xi(w, \rho^2) = \frac{2(w+1)}{\left[w+1 + \left(\rho^2 - \frac{1}{2} \right) (w-1) \right]^2}. \quad (3.13)$$

The above Isgur-Wise function will be accurate within the corridor shown in Fig. 2.

IV. THE PHYSICAL FORM FACTORS

An important observation from specific theoretical models [15] is that the subleading Isgur-Wise form factors which characterize the $1/m_Q$ corrections are to a good approximation linear in w . In addition, the radiative QCD corrections are monotonic and vary slowly with w , so that they too can be approximated linearly. Hence each of the $h_i(w)$ form factors (2.5) can be written as

$$h_i(w) \simeq [\alpha_i + \lambda_i + \mu_i(w-1)] \xi(w), \quad (4.1)$$

where both λ_i and μ_i are small, dimensionless constants. The advantage of this observation is that an analysis of the experimental form factor can now be carried out without *a priori* knowledge of the λ_i and μ_i coefficients.

To take advantage of the above, we start with the expansion (3.3) and note that for small $(w-1)$, IW functions of slopes $\rho^2 + \Delta\rho^2$ and ρ^2 are related by

$$\xi(w, \rho^2 + \Delta\rho^2) \simeq \xi(w, \rho^2) - \Delta\rho^2(w-1). \quad (4.2)$$

Over a wider range of w a more accurate expansion is

$$\xi(w, \rho^2 + \Delta\rho^2) \simeq \xi(w, \rho^2) [1 - \Delta\rho^2(w-1)]. \quad (4.3)$$

To illustrate the above approximation, we show in Fig. 3 a simple physical form factor (with $\mu=0.1$),

$$h(w) = \xi(w, 1.0) [1 + 0.1(w-1)]. \quad (4.4)$$

Also on Fig. 3 is a shifted IW function $\xi(w, 0.9)$ with slope chosen by the above prescription. We note that the two form factors differ by less than 2% at $w=1.5$. We now apply this result to the form factors $h_i(w)$. Comparing Eq. (4.3) to Eq. (4.1) in the case where $\alpha_i=1$ we see that

$$h_i(w) \simeq \xi(w, \rho^2 + \Delta\rho^2) + \lambda_i \xi(w, \rho^2), \quad (4.5)$$

$$\begin{aligned} &\simeq \xi(w, \rho^2 + \Delta\rho^2) + \lambda_i \xi(w, \rho^2 + \Delta\rho^2) \\ &\quad \times [1 + \Delta\rho^2(w-1)]. \end{aligned} \quad (4.6)$$

We drop the small product $\lambda_i \Delta\rho^2$ to obtain

$$h_i(w) \simeq h_i(1) \xi(w, \rho_i^2), \quad (4.7)$$

with

$$\rho_i^2 = \rho^2 - \mu_i,$$

$$h_i(1) = 1 + \lambda_i. \quad (4.8)$$

We may therefore conclude that the physical form factors are nearly equivalent to IW form factors with shifted slopes and normalizations.

As pointed out by Neubert [4,15], the sub-leading form factor χ_1 contributes identically to all of the h_i which remain in the heavy quark limit. We therefore simplify the analysis by absorbing this χ_1 contribution into a new IW type function which maintains spin-symmetry, but now has flavor dependence. It is this slope ρ^2 which we refer to as the IW slope in the following.

The absorption of χ_1 into the IW function is allowed as long as χ_1 remains small. The evidence for this is ambigu-




FIG. 4. The form factor $F(w)/F(1)$ from CLN [4] along with an IW function (1.1) of the same slope parameter, $\rho^2=1.0$.

ous. A QCD sum rule evaluation [15] gives a wide range of possible values for χ_1 , some of which are large. Evidence that χ_1 is small comes, indirectly, from the dispersive analysis of Caprini, Lellouch and Neubert (CLN) [4] where the $B \rightarrow D l \nu$ is found for a given slope. In Fig. 4 we show the CLN prediction for $\rho^2=1$ compared to our $k=1$ prediction (3.13) for the same slope. The two curves are nearly identical, which is expected if χ_1 and other corrections are small and can be absorbed into an effective IW function.

V. WHAT CAN WE LEARN FROM EXPERIMENTAL FORM FACTORS?

The semileptonic B decays yield definite but limited information about QCD corrections. It is important to understand exactly what in a theoretical framework is being tested by experiment. In this section, we examine each observable quantity to see which aspect of QCD is being tested. We consider the semileptonic decays separately.

A. $B \rightarrow D l \nu$

For $B \rightarrow D l \bar{\nu}$, the form factor is [4]

$$F(w) \equiv V_1(w) = h_+(w) + \frac{1-r}{1+r} h_-(w) \quad (5.1)$$

$$= \xi(w, \rho^2) [1 + \lambda_D + \mu_D(w-1) + \dots] \quad (5.2)$$

where

$$\lambda_D = \lambda_{h_+} + \frac{1-r}{1+r} \lambda_{h_-} \quad (5.3)$$

$$\mu_D = \mu_{h_+} + \frac{1-r}{1+r} \mu_{h_-} \quad (5.4)$$

and hence

$$\rho_D^2 = \rho^2 - \mu_D. \quad (5.5)$$

The only measurable parameter here is the physical form factor slope, ρ_D^2 , which has been investigated at CLEO [7]. If the data improves sufficiently, the difference between ρ_D^2 and the slope of a different physical form factor (such as F^*) could provide information about QCD. To gain a rough idea of the sizes of λ_D and μ_D we compute the Wilson coefficients and substitute the QCD sum rule approximations for the actual values of the subleading IW form factors. The results are included in Table II of Appendix A.

There are some directly applicable tests of HQET provided by QCD lattice simulations. Accurate lattice simulations are currently possible at the zero recoil point ($w=1$). A recent calculation [16] obtains

$$\lambda_{h_+} = 0.01, \quad (5.6)$$

$$\lambda_{h_-} = -0.11. \quad (5.7)$$

In HQET, these coefficients are due to the radiative corrections and $1/m$ corrections [4]. These are enumerated in Appendix A. We observe that λ_{h_+} has no $1/m$ corrections as required by Luke's theorem [17]. The result from the Wilson coefficients alone of $\lambda_{h_+} = 0.02$ is consistent with the lattice result (5.6).

The comparison with λ_{h_-} of Eq. (5.7) is more interesting. The $1/m$ correction here contains the subleading form factor η . This form factor has been estimated by QCD sum rules [18], by the updated Isgur-Scora-Grinstein-Wise (ISGW2) method [19], and by a Salpeter equation method [20]. While these various calculations yield reasonably consistent results for the subleading form factor $\chi_2(w)$, the expectations for $\eta(w)$ vary widely. From Appendix A we see that

$$\lambda_{h_-} = h_-(1) = (C_2 - C_3) + \left(\frac{1}{2m_b} + \frac{1}{2m_c} \right) (2\eta - 1) \bar{\Lambda}. \quad (5.8)$$

Using the calculated Wilson coefficients and the parameters $m_b = 4.80$ GeV, $m_c = 1.45$ GeV, and $\bar{\Lambda} = 0.5$ GeV we can evaluate the previous expression to yield

$$\lambda_{h_-} = -0.07 + 0.12(2\eta - 1). \quad (5.9)$$

The lattice result in Eq. (5.7) suggests that the second term above is negative, i.e. $\eta < 0.5$. This prediction is not completely rigorous as the size of the effect is of order a few percent and second-order $1/m$ corrections could be of that order. In addition, the lattice calculation for $h_-(1)$ has currently been calculated only to tree level, but a 1-loop calculation is underway [16]. However, the preliminary indication is that η is less than the central value of 0.62 which comes from QCD sum rules. At the same time however, the much smaller values of η which come from ISGW2 and the Salpeter model are also disfavored because they produce a result for $h_-(1)$ which is too negative.

B. $B \rightarrow D^* l \nu$

In this case, there is additional information provided by the $D^* \rightarrow D\pi$ decay distribution [6]. As suggested by CLN [4] and followed by CLEO [6] this decay can be analyzed in terms of the h_{A_1} form factor and two ratios of form factors.

The CLEO experiment, which to date has the best measurements of the $B \rightarrow D^*$ form factor parameters, has adopted the convention of fitting the slope of h_{A_1} , $\rho_{A_1}^2$ and two form factor ratios:

$$R_1(w) = \frac{h_V(w)}{h_{A_1}(w)},$$

$$R_2(w) = \frac{h_{A_3}(w) + r^* h_{A_2}(w)}{h_{A_1}(w)}, \quad (5.10)$$

where $r^* = m_{D^*}/m_B \approx 0.38$. These ratios are expected to be nearly independent of w so they are treated as constants in the fit. The form factor F^* can be expressed in terms of these parameters by combining Eq. (A3) from the Appendix and the definitions (5.10). We would also like to write $F^*(w)$ as a shifted IW function,

$$F^*(w) = \xi(w, \rho^2) [1 + \lambda_{D^*} + \mu_{D^*}(w-1) + \dots], \quad (5.11)$$

and hence

$$F^*(w) = F^*(1) \xi(w, \rho_{D^*}^2), \quad (5.12)$$

where $\rho_{D^*}^2 = \rho^2 - \mu_{D^*}$. Since $|F^*|^2$ is the sum of h_i contributions and kinematical factors, the λ_{D^*} and μ_{D^*} are the result of an expansion about $w = 1$. The result is

$$\lambda_{D^*} = \lambda_{A_1}, \quad (5.13)$$

$$\mu_{D^*} = \mu_{A_1} + \frac{1}{3} \lambda_V - \frac{1}{3(1-r^*)} [\lambda_{A_3} + r^* \lambda_{A_2} - r^* \lambda_{A_1}]. \quad (5.14)$$

As in the $B \rightarrow D l \nu$ case, we can obtain rough approximations of μ_{D^*} , λ_{D^*} , μ_{A_1} and λ_{A_1} . The results are included in Table II of Appendix A.

The above expression (5.14) has a remarkable consequence. The slope difference $\rho_{D^*}^2 - \rho_{A_1}^2 = \mu_{A_1} - \mu_{D^*}$ can be related to the zero-recoil quantities, λ_i above. It is therefore possible in the near future to test (for the first time) the ‘‘intercept’’ quantities λ_i by direct comparison to experiment. At this point only λ_{A_1} has been computed in lattice simulation [21], but the corresponding values of λ_{A_2} , λ_{A_3} , and λ_V could be evaluated. At present the experimental situation [6] is also not precise enough:

$$\rho_{D^*}^2 - \rho_{A_1}^2 = 0.08 \pm 0.15, \quad (5.15)$$

but this result can also be considerably sharpened. We emphasize that the above prediction (5.14) is an important check of the values of the intercepts, λ_i , which are in turn of critical importance in the extraction of the Cabibbo-Kobayashi-Maskawa (CKM) parameter V_{cb} . We should point out that the relation (5.14) is implicitly contained in CLN’s Eq. (35).

In the heavy quark limit, $R_1 = R_2 = 1$. These ratios’ deviation from unity, experimentally and theoretically, indicates deviation from heavy quark symmetry and, consequently, can test theoretical predictions for the symmetry-breaking corrections. The measurements of R_1 and R_2 (currently with large error bars) agree with theoretical predictions, and thus roughly confirm the predictions of HQET. Here we investigate the exact nature of the predictions and their value as tests of HQET.

Treating the form factor ratios as constants, while reasonable theoretically and necessary for the fit, makes the tests of symmetry-breaking corrections less informative. $R_1(w)$ and $R_2(w)$ are essentially reduced to $R_1(1)$ and $R_2(1)$, and because many of the corrections vanish at $w = 1$, they cannot be tested. For example, the subleading Isgur-Wise function $\chi_3(w)$ which has only been calculated up to the errors inherent to QCD sum rules is completely lost in this procedure. If we apply the expressions given in Appendix A to the various form factors at $w = 1$, eliminate those corrections which vanish at zero-recoil, and express the results in terms of the Wilson coefficients and subleading IW form factors, we obtain

$$R_1(1) = \frac{C_1(1) + \frac{1}{2m_c}\bar{\Lambda} - \frac{1}{2m_b}(2\eta - 1)\bar{\Lambda}}{C_1^5(1)} \quad (5.16)$$

$$R_2(1) = \frac{C_1^5(1) + C_2^5(1) + r^*C_2^5(1) - \eta\bar{\Lambda}\left[\frac{1}{m_b} + \frac{(1+r^*)}{2m_c}\right] + \frac{4\chi_2(1)}{2m_c}(r^* - 1) + \bar{\Lambda}\left(\frac{1}{2m_b} - \frac{r^*}{2m_c}\right)}{C_1^5(1)}. \quad (5.17)$$

If we use the theoretical estimates as rough guides to the size of each term, we can determine what predictions are actually being tested.

1. What do we learn from R_1 ?

The subleading combination $2\eta - 1$ is expected to be small on the basis of the λ_{h_-} considerations and is further reduced by $1/m_b$. Consequently, after evaluating the Wilson coefficients C_i (see Table I of Appendix A), R_1 must be greater than unity and provides a fairly direct probe of the HQET parameter $\bar{\Lambda}$, which is the light degrees of freedom energy to this order.

2. What do we learn from R_2 ?

In the expression for R_2 , χ_2 is generally agreed to be quite small and $1/2m_b - r^*/2m_c \approx 0.03$. Thus after calculating the Wilson coefficients, a measurement of R_2 provides an additional probe of the value of the subleading form factor η . Using the Wilson coefficients in the Appendix, as well as $m_b = 4.8$ GeV, $m_c = 1.45$ GeV, $r^* = 0.38$ and $\bar{\Lambda} = 0.5$ GeV, the expression for R_2 becomes

$$R_2 = 1.00 - 0.34\eta(1) - 0.84\chi_2(1). \quad (5.18)$$

The term containing η is the dominant correction as χ_2 is small, so a precise measurement of R_2 effectively measures η .

VI. CONCLUSIONS

We have addressed the question of how to analyze semileptonic B decay while minimizing both the amount of theoretical assumption and the number of parameters. We first establish that specifying the slope parameter ρ^2 accurately determines the entire Isgur-Wise function. To do so, we need only an approximation to the light degrees of freedom wave function provided by a QCD simulation. Our result (1.1) is

$$\xi(w, \rho^2) = \frac{2(w+1)}{\left[w+1 + \left(\rho^2 - \frac{1}{2}\right)(w-1)\right]^2}. \quad (6.1)$$

The form factors $h_i(w)$ either vanish or approach the IW function in the heavy quark limit. These amplitudes can be accurately written as

$$h_i(w) = [\alpha_i + \lambda_i + \mu_i(w-1) + \dots] \xi(w, \rho^2), \quad (6.2)$$

where α_i is 0 or 1 and λ_i and μ_i are small dimensionless constants. We observe that by altering the IW slope ρ^2 to $\rho_i^2 = \rho^2 - \mu_i$ the physical form factors can be expressed as

$$h_i(w) = (\alpha_i + \lambda_i) \xi(w, \rho_i^2), \quad (6.3)$$

where $\rho_i^2 = \rho^2 - \mu_i$. That this works well is shown by Fig. 3.

From the above we conclude that semileptonic data are parametrized by an intercept value $V_{cb}(1 + \lambda_i)$ and an effective Isgur-Wise slope parameter ρ_i^2 . To find the CKM element V_{cb} one must use a theoretical estimate of λ_i .

For $B \rightarrow D^* l \nu$ decay the parameters are $V_{cb}(1 + \lambda_{A_1})$, $\rho_{A_1}^2$, R_1 , and R_2 . In this case one has a consistency condition (5.14) relating the λ_i and the difference between actual D^* slope and $\rho_{A_1}^2$. We further point out that R_1 is nearly model-independent while R_2 depends sensitively on various estimates of the subleading form factor $\eta(1)$. Also, the value of the form factor $h_-(w)$ at zero-recoil, λ_{h_-} , appears to offer an additional probe of the value $\eta(1)$.

Finally, one might ask, what is the essential advantage of the slope shift scheme described here? The answer is, economy of parameters and a decoupling from theoretical assumptions. As seen in Fig. 3, one of the effective IW function slopes is equivalent to an (unknown) IW slope with subleading corrections. If these corrections were securely known our scheme offers no advantage. On the other hand, if one tried to fit both the IW slope and HQET μ parameters, a hopeless parameter correlation would arise.

We believe that it is best to rely as little as possible on theory and to use a direct phenomenological approach. As pointed out previously [8], one must have some theoretical constraint on the curvature term in Eq. (2.1). We show here that the shape of the form factor is specified once the slope parameter is given. Later, after the decay distributions have been parametrized, the fitted parameters can be compared to predictions.

ACKNOWLEDGMENTS

This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

TABLE I. Wilson coefficient linear fits for $\alpha(\mu)=0.12\pi$ [15].

C_i	$C_i(1)$	$C'_i(1)$
C_1	1.13	-0.08
C_1^5	1.00	-0.04
C_2	-0.086	0.037
C_2^5	-0.11	0.038
C_3	-0.019	0.005
C_3^5	0.040	-0.013

APPENDIX

The fundamental form factors h_i are defined [4,10] by

$$\frac{\langle D(v') | \bar{c} \gamma^\mu b | B(v) \rangle}{\sqrt{m_B m_D}} = (v+v')^\mu h_+(w) + (v-v')^\mu h_-(w),$$

$$\frac{\langle D^*(v', \epsilon) | \bar{c} \gamma^\mu b | B(v) \rangle}{\sqrt{m_B m_{D^*}}} = i \epsilon^{\mu\nu\alpha\beta} \epsilon_\nu^* v'_\alpha v_\beta h_V(w),$$

$$\frac{\langle D^*(v', \epsilon) | \bar{c} \gamma^\mu \gamma^5 b | B(v) \rangle}{\sqrt{m_B m_{D^*}}} = (w+1) \epsilon^{*\mu} h_{A_1}(w) - \epsilon^{*\nu} v (v^\mu h_{A_2} + v'^\mu h_{A_3}). \quad (\text{A1})$$

The matrix elements for $B \rightarrow D l \bar{\nu}$ in Eq. (2.4) can be expressed in terms of $\{h_i\}$ as

$$F(w) = h_+(w) - \left(\frac{1-r}{1+r} \right) h_-(w) \quad (\text{A2})$$

and the squared matrix element for $B \rightarrow D^* l \bar{\nu}$ in Eq. (2.4) as

$$f^* |F^*(w)|^2 = 2(w+1)(1-2wr^* + r^{*2}) [(w+1)h_{A_1}^2 + (w-1)h_V^2] + (w+1)^2 [(w-r^*)h_{A_1} - (w-1)(r^*h_{A_2} + h_{A_3})]^2. \quad (\text{A3})$$

The h_i can, as indicated in Eq. (2.5), be expressed by the short-range corrections β_i and the $1/m_Q$ corrections γ_i together as

$$h_V(w) = \xi(w) [C_1 + \epsilon_c(L_2 - L_5) + \epsilon_b(L_1 - L_4)]$$

$$h_{A_3}(w) = \xi(w) [C_1^5 + C_3^5 + \epsilon_c(L_2 - L_3 - L_5 + L_6) + \epsilon_b(L_1 - L_4)]$$

$$h_{A_2}(w) = \xi(w) [C_2^5 + \epsilon_c(L_3 + L_6)]$$

TABLE II. Estimates of the λ and μ coefficients for the various relevant form factors using the Wilson coefficients of Table I and QCD sum rules approximations [4,18] as summarized in Eq. (A5).

Form factor	λ_i	μ_i
h_{A_1}	0.004	0.050
$F_D = V_1$	0.041	0.072
$F^* = \mathcal{F}$	0.004	0.253

$$h_{A_1}(w) = \xi(w) \left[C_1^5 + \epsilon_c \left(L_2 - \frac{w-1}{w+1} L_5 \right) + \epsilon_b \left(L_1 - \frac{w-1}{w+1} L_4 \right) \right]$$

$$h_+(w) = \xi(w) \left[C_1 + \frac{w+1}{2} (C_2 + C_3) + (\epsilon_c + \epsilon_b) L_1 \right]$$

$$h_-(w) = \xi(w) \left[\frac{w+1}{2} (C_2 - C_3) + (\epsilon_c + \epsilon_b) L_4 \right] \quad (\text{A4})$$

where $\epsilon_{c,b} = (2m_{c,b})^{-1}$, the C_i are the Wilson coefficients—which are discussed extensively in [15]—and the L_i are combinations of the sub-leading Isgur Wise form factors and have been approximated by QCD sum rules [4]:

$$L_1 = -4(w-1) \frac{\chi_2}{\xi} + 12 \frac{\chi_3}{\xi} \approx 0.72(w-1) \bar{\Lambda}$$

$$L_2 = -4 \frac{\chi_3}{\xi} \approx -0.16(w-1) \bar{\Lambda}$$

$$L_3 = 4 \frac{\chi_2}{\xi} \approx 0.24 \bar{\Lambda}$$

$$L_4 = (2\eta - 1) \bar{\Lambda} \approx 0.24 \bar{\Lambda}$$

$$L_5 = -\bar{\Lambda}$$

$$L_6 = -\frac{2}{w+1} (\eta+1) \bar{\Lambda} \approx -\frac{3.24}{w+1} \bar{\Lambda} \quad (\text{A5})$$

where $\bar{\Lambda} \approx 0.5$ GeV is the ‘‘binding energy’’ of a heavy meson.

Using the detailed expressions given in [4] and [15] and the expressions above, we can extract λ_i and μ_i for each form factor h_i which appears in Eq. (4.1). They appear in Table II. We use the values $m_b = 4.8$ GeV and $m_c = 1.45$ GeV and calculate the Wilson coefficients at the scale $\alpha_s(\mu) = 0.12\pi$ (see Tables I and II).

- [1] N. Isgur and M.B. Wise, Phys. Lett. B **232**, 113 (1989); **237**, 527 (1990).
- [2] G.C. Boyd, B. Grinstein, and R.F. Lebed, Phys. Lett. B **353**, 306 (1995).
- [3] I. Caprini and M. Neubert, Phys. Lett. B **380**, 396 (1996).
- [4] I. Caprini, L. Lellouch, and M. Neubert, Nucl. Phys. **B530**, 151 (1998).
- [5] C.G. Boyd, B. Grinstein, and R.F. Lebed, Phys. Rev. D **56**, 6895 (1997).
- [6] J.E. Duboscq *et al.*, Phys. Rev. Lett. **76**, 3898 (1996).
- [7] J. Bartelt *et al.*, Phys. Rev. Lett. **82**, 3746 (1999).
- [8] G. Burdman, Phys. Lett. B **284**, 133 (1992).
- [9] E.g., S. Veseli and M.G. Olsson, Phys. Rev. D **54**, 886 (1996).
- [10] A. Falk and M. Neubert, Phys. Rev. D **47**, 2965 (1993).
- [11] M. Sadzikowski and K. Zalewski, Z. Phys. C **59**, 677 (1993); H. Høggassen and M. Sadzikowski, *ibid.* **64**, 427 (1994); M.G. Olsson and S. Veseli, Phys. Rev. D **51**, 2224 (1995); M.R. Ahmady, R.R. Mendel, and J.D. Talman, *ibid.* **52**, 254 (1995).
- [12] S. Veseli and M.G. Olsson, Phys. Lett. B **367**, 302 (1996).
- [13] A. Duncan, E. Eichten, and H. Thacker, Phys. Lett. B **303**, 109 (1993); Phys. Rev. D **51**, 5101 (1995).
- [14] M.G. Olsson and S. Veseli, Phys. Lett. B **397**, 263 (1997).
- [15] M. Neubert, Phys. Rep. **245**, 259 (1994).
- [16] S. Hashimoto *et al.*, Phys. Rev. D **61**, 014502 (2000).
- [17] M. Luke, Phys. Lett. B **252**, 447 (1990).
- [18] Z. Ligeti, Y. Nir, and M. Neubert, Phys. Rev. D **49**, 1302 (1994).
- [19] D. Scora and N. Isgur, Phys. Rev. D **52**, 2783 (1995).
- [20] A. Abd El-Hady *et al.*, Phys. Rev. D **51**, 5245 (1995).
- [21] J.N. Simone *et al.*, Nucl. Phys. B (Proc. Suppl.) **73**, 399 (1999).