

Photons from quantized electric flux representations

Madhavan Varadarajan*

Raman Research Institute, Bangalore 560 080, India

(Received 17 April 2001; published 28 September 2001)

The quantum theory of $U(1)$ connections admits a diffeomorphism invariant representation in which the electric flux through any surface is quantized. This representation is the analog of the representation of quantum $SU(2)$ theory used in loop quantum gravity. We investigate the relation between this representation, in which the basic excitations are “polymerlike,” and the Fock representation, in which the basic excitations are wavelike photons. We show that normalizable states in the Fock space are associated with “distributional” states in the quantized electric flux representation. This work is motivated by the question of how wavelike gravitons in linearized gravity arise from polymerlike states in nonperturbative loop quantum gravity.

DOI: 10.1103/PhysRevD.64.104003

PACS number(s): 04.60.Ds

I. INTRODUCTION

The loop quantum gravity approach [1,2] is based on a Hamiltonian description of classical general relativity in which the basic configuration variable is a connection [3,4] and its conjugate is a triad field. The spatial 3-metric on a Cauchy slice is constructed from the triad and the connection contains information about the extrinsic curvature of the slice as embedded in the spacetime, thus establishing contact with the usual Arnowitt-Deser-Misner formulation. In the quantum theory, the basic connection dependent operators are holonomies of the connection around loops in the Cauchy slice [5,6].

In recent years the approach has been put on a firm mathematical footing with the following key features. A (quantum) configuration space, $\overline{\mathcal{A}/\mathcal{G}}$, of (generalized) $SU(2)$ connections modulo gauge and a canonical diffeomorphism invariant measure, $d\mu_0$ (also called the Ashtekar-Lewandowski measure [7]), on $\overline{\mathcal{A}/\mathcal{G}}$ have been constructed [7–11]. The space $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$ provides a kinematical Hilbert space on which the (self-adjoint) $SU(2)$ holonomy operators act by multiplication. This kinematical Hilbert space is spanned by an orthonormal set of “spin network” states, each associated with an oriented, closed graph whose edges are labeled by representations of $SU(2)$ [12–14].

Every spin network state is an eigenstate of operators corresponding to the area of 2-surfaces in the Cauchy slice—roughly speaking, the area of a surface gets a contribution of $\sqrt{j(j+1)}$ units of Planck area from each edge labeled by spin j which intersects the surface transversely. Since the area operator is constructed from the triad field [or equivalently the $SU(2)$ electric field], the edges of the graph may be thought of as carrying quanta of non-Abelian electric flux. Since the edges are 1 dimensional, the intuitive picture of states in $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$ is of “polymerlike” quantum excitations.

As noted earlier, $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$ is a kinematical structure. The dynamics of general relativity in its canonical description is encoded in the diffeomorphism and Hamiltonian constraints. Physical states in the quantum theory are in the ker-

nel of the corresponding quantum constraint operators. The representation of holonomies, functionals of the triad, as well as the unitary action of diffeomorphisms on $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$ provide a representation of the diffeomorphism constraints [15], and in a key breakthrough by Thiemann, the Hamiltonian constraint [16].¹ It turns out that elements of the kernel of the constraints are “too distributional” to be normalizable states in $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$. Rather, they are expressible as non-normalizable, infinite, sums of spin network states.

Thus, a nonperturbative physical state of the quantum gravitational field is a distributional sum of kinematic states, each of which is associated with 1 dimensional, polymer-like excitations. Given such quantum states, a key open question is: how do classical configurations of the gravitational field arise? In particular, how does flat spacetime (and small perturbations around it) arise from non-perturbative quantum states of the gravitational field?

The latter question is particularly interesting for the following reason. Small perturbations about flat spacetime correspond to solutions of linearized gravity. Quantum states of linearized gravity lie in the familiar graviton Fock space on which the conventional perturbative approaches to quantum gravity are based. Such approaches seem to fail due to non-normalizability problems. Thus, an understanding of the relation between the quantum states of linearized gravity and states in full nonperturbative loop quantum gravity would shed light on the reasons behind the failure of perturbative methods.

There are many aspects of this yet-to-be-understood relation between perturbative and nonperturbative states and we shall focus on only one of them, namely, the dramatically different nature of the basic excitations in Fock space and in loop quantum gravity: whereas Fock space gravitons can be thought of as *propagating 3D wave-packets*, states in loop

¹Note that it is not clear if Thiemann’s quantization has general relativity as its classical limit [17]. It has also been realized that there are a large family of distinct Thiemann-like regularizations of the Hamiltonian constraint [18]. On the other hand, recent results of Bojowald [19] in the context of minisuperspace quantization may be taken as evidence that some elements of Thiemann’s procedure could lead to avoidance of singularities in quantum theory.

*Email address: madhavan@rri.ernet.in

quantum gravity are associated with *1D polymer-like excitations*.

This aspect of the relation between perturbative and non-perturbative states in quantum gravity can be isolated and studied in the simpler yet highly instructive context of source-free Maxwell theory. To this end, consider a $U(1)$ connection on a spatial slice diffeomorphic to R^3 . Starting from the Poisson brackets algebra of $U(1)$ holonomies around loops in R^3 , it can be seen that exact counterparts of the $SU(2)$ constructions of loop quantum gravity exist [8,7,20,21]. Thus, a canonical diffeomorphism invariant measure exists on the quantum configuration space of generalized $U(1)$ connections modulo gauge [we shall continue to denote these by $d\mu_0$ and $\overline{\mathcal{A}/\mathcal{G}}$ —it will be clear from the context whether these symbols refer to $SU(2)$ or $U(1)$], holonomies act as unitary operators by multiplication on $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$, and $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$ is spanned by an orthonormal basis, each element of which is associated with a closed, oriented graph and a labeling of edges of the graph by representations of $U(1)$. Since such representations are labeled by integers called charges, we refer to these states as “charge network” states. Each such state is an eigenfunction of electric flux operators associated with surfaces in R^3 and the edges of the graph underlying the state may be thought of as carrying quanta of electric flux. We refer to this diffeomorphism invariant representation, in which the basic excitations are, once again, 1 dimensional and “polymer-like” as the *quantized electric flux (QEF) representation*.

In sharp contrast to this, is the usual Poincaré invariant quantum theory of a $U(1)$ connection on a fixed flat space-time with flat spatial slices diffeomorphic to R^3 . Here, the connection and its conjugate electric field are represented as operator valued distributions on the Fock space of photons where the basic quantum excitations are 3D and wave-like.

Thus quantum $U(1)$ theory presents an excellent arena to discuss the question raised earlier, namely *how can the 3D wave-like excitations of Fock space arise from underlying 1D polymer-like excitations?*

At first sight, the Fock and the QEF representations are very different. In the Fock representation, the connection is an operator valued distribution which needs to be smeared in 3 dimensions to obtain a well defined operator. Holonomies of this operator valued distribution involve smearings only over the 1 dimension provided by the loop, and are *not* well defined. In contrast, in the QEF representation, holonomies *are* well defined operators on $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$. In Sec. II we review our previous results [22] which point to a way around this apparently insurmountable obstacle to relating the two representations.

In Sec. III we present our main result, namely that states in the Fock representation are associated with distributional sums of charge network states in the QEF representation. This mirrors the mathematical structure of loop quantum gravity in the sense that, there, physical states are associated with distributional sums of spin network states in the kinematic Hilbert space. Section IV contains a discussion of our results and some concluding remarks.

We shall use units in which $\hbar = c = 1$.

II. THE ISSUE OF SMEARING

Since the Abelian Poisson brackets algebra of holonomies is the primary structure from which the QEF representation is obtained [21,22], we would like to relate the Fock representation to this holonomy algebra. As noted earlier, the usual Fock representation is not a representation of the holonomy algebra because the connection operators in this representation are too singular for their holonomies to be defined. Instead, it is possible to obtain the usual Fock representation as the representation of a related algebra of “smeared holonomies.” By using the smeared holonomy algebra as a link between the algebra of holonomies and the Fock representation we shall be able to relate the two.

In order to define the smeared holonomies we recall the following definitions from [23,28,22]. \vec{x} denotes Cartesian coordinates of a point on R^3 . \mathcal{A} is the space of smooth $U(1)$ connections, $A_a(\vec{x})$,² [on the trivial $U(1)$ bundle on R^3] whose Cartesian components are functions of rapid decrease at infinity. \mathcal{L}_{x_0} is the space of unparametrized, oriented, piecewise analytic loops on R^3 with basepoint \vec{x}_0 . Composition of a loop α with a loop β is denoted by $\alpha \circ \beta$. Given a loop $\alpha \in \mathcal{L}_{x_0}$, the holonomy of $A_a(x)$ around α is $H_\alpha(A) := \exp(i\oint_\alpha A_a dx^a)$. The holonomy can equivalently be defined as

$$H_\alpha(A) = \exp i \int_{R^3} X_\gamma^a(\vec{x}) A_a(\vec{x}) d^3x, \quad (1)$$

with

$$X_\gamma^a(\vec{x}) := \oint_\gamma ds \delta^3(\vec{\gamma}(s), \vec{x}) \dot{\gamma}^a, \quad (2)$$

where s is a parametrization of the loop γ , $s \in [0, 2\pi]$. $X_\gamma^a(\vec{x})$ is called the form factor of γ . The Gaussian smeared form factor [23] is defined as

$$X_{\gamma(r)}^a(\vec{x}) := \int_{R^3} d^3y f_r(\vec{y} - \vec{x}) X_\gamma^a(\vec{y}) = \oint_\gamma ds f_r[|\vec{\gamma}(s) - \vec{x}|] \dot{\gamma}^a \quad (3)$$

where

$$f_r(\vec{x}) = \frac{1}{(2\pi)^{3/2} r^3} e^{-x^2/2r^2} \quad x := |\vec{x}| \quad (4)$$

approximates the Dirac delta function for small r .

Then, the smeared holonomy is defined as

$$H_{\gamma(r)}(A) = \exp i \int_{R^3} X_{\gamma(r)}^a(\vec{x}) A_a(\vec{x}) d^3x. \quad (5)$$

²As noted in [20], to get an object with the dimensions of a $U(1)$ connection, we need to divide the usual magnetic potential by a parameter, q_0 , which has dimensions of electric charge. This reflects in the Poisson brackets (7) and the definition (31).

As shown in [22], the Fock representation is a representation of the Poisson brackets algebra generated by the smeared holonomies, $H_{\gamma(r)}(A)$ and the electric field $E^a(\vec{x})$:

$$\begin{aligned} \{H_{\gamma(r)}, H_{\alpha(r)}\} &= \{E^a(\vec{x}), E^b(\vec{y})\} = 0, \\ \{H_{\gamma(r)}, E^a(\vec{x})\} &= \frac{i}{q_0} X_{\gamma(r)}^a(\vec{x}) H_{\gamma(r)}. \end{aligned} \quad (6)$$

These Poisson brackets are generated from the elementary Poisson brackets

$$\{A_a(\vec{x}), E^b(\vec{y})\} = \frac{1}{q_0} \delta_a^b \delta(\vec{x}, \vec{y}). \quad (7)$$

Here, q_0 is a parameter with the units of electric charge [20]. In the Fock representation, the smeared holonomies, $\hat{H}_{\gamma(r)}$, are unitary operators and the electric field, $\hat{E}^a(\vec{x})$, is an operator valued distribution.

How is the above algebra involving smeared holonomies and the electric field related to the holonomy algebra? To this end we define the classical Gaussian smeared electric field $E_r^a(\vec{x})$ by

$$E_r^a(\vec{x}) := \int d^3y f_{r, \vec{x}}(\vec{y} - \vec{x}) E^a(\vec{y}). \quad (8)$$

The Poisson brackets algebra generated by the (unsmeared) holonomies and the Gaussian smeared electric field is

$$\begin{aligned} \{H_\gamma, H_\alpha\} &= \{E_r^a(\vec{x}), E_r^b(\vec{y})\} = 0, \\ \{H_\gamma, E_r^a(\vec{x})\} &= \frac{i}{q_0} X_{\gamma(r)}^a(\vec{x}) H_\gamma. \end{aligned} \quad (9)$$

Then, as is hinted by Eqs. (6) and (9) and proved in detail in [22], the abstract algebraic structures underlying the Poisson brackets algebras generated by $(H_{\gamma(r)}(A), E^a(\vec{x}))$ and $(H_\gamma(A), E_r^a(\vec{x}))$ are *identical*. In other words, $E^a(\vec{x})$ and $E_r^a(\vec{x})$, and $H_{\gamma(r)}(A)$ and $H_\gamma(A)$ may be identified with the same abstract objects as far as the algebraic structure of the two Poisson brackets algebras is concerned. Therefore, any representation of the Poisson brackets algebra generated by $(H_{\gamma(r)}(A), E^a(\vec{x}))$ defines a representation of the Poisson brackets algebra generated by $(H_\gamma(A), E_r^a(\vec{x}))$. In particular, the Fock representation of the $(H_{\gamma(r)}(A), E^a(\vec{x}))$ algebra defines a representation of the $(H_\gamma(A), E_r^a(\vec{x}))$ algebra. We shall call this representation, the “ r -Fock representation” of the $(H_\gamma(A), E_r^a(\vec{x}))$ Poisson brackets algebra. Thus, in the r -Fock representation the holonomies, \hat{H}_γ are well defined operators.

We can see this explicitly as follows. The Fock representation, or for that matter, any representation with a cyclic “vacuum” state, can be reconstructed from the vacuum expectation values of the algebra of operators. Thus, we may

specify the r -Fock representation via the vacuum expectation values of the Fock representation as follows. The Fock representation is a cyclic representation generated from the Fock vacuum, $|0\rangle$ with vacuum expectation values³

$$\langle 0 | \hat{H}_{\gamma(r)} | 0 \rangle = \exp\left(-\int \frac{d^3k}{4q_0^2 k} |X_{\gamma(r)}^a(\vec{k})|^2\right), \quad (10)$$

$$\begin{aligned} \langle 0 | \hat{H}_{\alpha(r)} \hat{E}^a(\vec{x}) \hat{H}_{\beta(r)} | 0 \rangle &= \frac{X_{\beta(r)}^a(\vec{x}) - X_{\alpha(r)}^a(\vec{x})}{2q_0} \\ &\times \exp\left(-\int \frac{d^3k}{4q_0^2 k} |X_{\alpha\circ\beta(r)}^a(\vec{k})|^2\right), \end{aligned} \quad (11)$$

where $X_{\gamma(r)}^a(\vec{k})$ denotes the Fourier transform of $X_{\gamma(r)}^a(\vec{x})$. This defines the r -Fock representation as the cyclic representation generated from the r -Fock “vacuum” $|0_r\rangle$ with vacuum expectation values

$$\langle 0_r | \hat{H}_\gamma | 0_r \rangle = \exp\left(-\int \frac{d^3k}{4q_0^2 k} |X_{\gamma(r)}^a(\vec{k})|^2\right). \quad (12)$$

$$\begin{aligned} \langle 0_r | \hat{H}_\alpha \hat{E}_r^a(\vec{x}) \hat{H}_\beta | 0_r \rangle &= \frac{[X_{\beta(r)}^a(\vec{x}) - X_{\alpha(r)}^a(\vec{x})]}{2q_0} \\ &\times \exp\left(-\int \frac{d^3k}{4q_0^2 k} |X_{\alpha\circ\beta(r)}^a(\vec{k})|^2\right). \end{aligned} \quad (13)$$

The fact that the holonomies are well defined operators in the r -Fock representation allows us to relate the r -Fock representation and the QEF representation in a fairly direct manner, as we show in the next section.

To summarize, although it is not possible to relate the QEF representation with the usual Fock representation directly, it is possible to relate the QEF representation with the r -Fock representation, since both provide representations of the Poisson bracket algebra generated by $(H_\gamma(A), E_r^a(\vec{x}))$.

The only remaining question is of the relation between the Fock and the r -Fock representations. The mathematical relation between the two, in terms of representations of two different realizations of the same algebraic structure, is clear and is true for *any* $r > 0$. So the only question is of the *physical* relation between r -Fock and Fock representations. We argue below that for certain measurements which are of physical relevance in the context of our motivations from quantum gravity, the two representations are physically indistinguishable for sufficiently small r .

Since we shall phrase our argument in terms of measurements of Fourier modes, we first discuss the behavior of Fourier modes under Gaussian smearing. Given any function $h(\vec{x})$, its Fourier transform is

³Our conventions for the Fock space representation are displayed in Sec. III B.

$$h(\vec{k}) = \frac{1}{(2\pi)^{3/2}} \int_{R^3} d^3x h(\vec{x}) e^{-i\vec{k}\cdot\vec{x}} \quad (14)$$

and its Gaussian smeared version is

$$h_r(\vec{x}) = \int_{R^3} d^3y f_r(\vec{y}-\vec{x}) h(\vec{y}). \quad (15)$$

It follows that

$$h_r(\vec{k}) = e^{-k^2 r^2/2} h(\vec{k}). \quad (16)$$

In particular we have

$$X_{\gamma(r)}^a(\vec{k}) = e^{-k^2 r^2/2} X_{\gamma}^a(\vec{k}) \quad (17)$$

and

$$E_r^a(\vec{k}) = e^{-k^2 r^2/2} E^a(\vec{k}). \quad (18)$$

From Eqs. (17) and (5) it follows, in obvious notation, that

$$H_{\gamma}(A(\vec{k})) = H_{\gamma(r)}(e^{k^2 r^2/2} A(\vec{k})). \quad (19)$$

Consider measurements of quantities at or above a length scale L . More precisely, let the measurements be of Fourier modes, $E^a(\vec{k}), A_a(\vec{k})$ of the electric and connection fields (in, say, the Coulomb gauge) for $k \leq 1/L$. Further, let the accuracy of the measurement process be characterized by the (small) positive number δ . If $\Delta E^a(\vec{k}), \Delta A_a(\vec{k})$ are the accuracies to which $E^a(\vec{k}), A_a(\vec{k})$ are measured, then δ is defined through $\Delta E^a(\vec{k}) = E^a(\vec{k}) \delta$ and $\Delta A_a(\vec{k}) = A_a(\vec{k}) \delta$. Then if r is chosen small enough that the condition $r^2/2L^2 < \delta$ holds, it follows that the measurements cannot distinguish between the modes $E^a(\vec{k})$ and $e^{-k^2 r^2/2} E^a(\vec{k})$, and $A_a(\vec{k})$ and $e^{k^2 r^2/2} A_a(\vec{k})$. Thus, the measurements cannot distinguish between unsmeared fields and their “ r -counterparts” for sufficiently small r . Since the primary operators in the Fock and r -Fock representations are related by Eqs. (18) and (19), it is straightforward to see that for any state in the Fock space there exists a state in the r -Fock space such that the above type of measurement can never distinguish between the two.⁴

We loosely interpret this statement to mean that, given measurements at some length scale performed to some finite accuracy, there is always a sufficiently small r such that the r -Fock representation is experimentally indistinguishable from the usual Fock representation. Although for Maxwell theory the introduction of a length scale seems arbitrary, in the context of linearized gravity it is necessary to restrict attention to length scales much larger than the Planck scale or else the linearized approximation will not be physically valid.

⁴In the language and notation of [22], the state in the r -Fock space is the image of the state in the Fock space via the map I_r .

Hence, for the remainder of the paper we may alter our original question to: *how can the 3D wave-like excitations of r -Fock space arise from underlying 1D polymer-like excitations?*

III. r -FOCK STATES AS DISTRIBUTIONS IN THE QEF REPRESENTATION

In this section the r -Fock space representation is derived from the QEF representation by identifying the r -Fock vacuum as a distributional state in the QEF representation. We show that the r -Fock vacuum can be written as a formal non-normalizable sum of charge network states and that it resides in the algebraic dual to the space of finite linear combinations of charge network states.

In Sec. III A we briefly review the properties of charge network states as well as the definition of the algebraic dual representation. In Sec. III B 1 we encode the Poincaré invariance of the Fock vacuum in a relation between smeared holonomy and electric field operators. In Sec. III B 2 we show how this relation implies the identification of the r -Fock vacuum with a distributional sum of charge network states.

A. Charge network states

Straightforward repetitions of constructions for the $SU(2)$ case of loop quantum gravity [12,11,10,13] lead to the following results for $U(1)$ charge networks.

(i) The charge network states constitute an (uncountable) orthonormal spanning set in $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_0)$. Each such state is labeled by a closed, oriented graph whose edges carry non-trivial representations of $U(1)$. Representations of $U(1)$ are labeled by integers called “charges” [20], hence the name “charge network states.” $U(1)$ gauge invariance implies that the sum of charges at each vertex vanishes. We denote the normalized state labeled by the graph “ γ ,” with N edges carrying the charges $(p_1 \dots p_N)$ as $|\gamma, \{p\}\rangle$. Orthonormality implies

$$\langle \alpha, \{q\} | \gamma, \{p\} \rangle = \delta_{(\alpha, \{q\}), (\gamma, \{p\})}, \quad (20)$$

i.e., the inner product vanishes unless $\alpha = \gamma$ and $q_i = p_i$, $i = 1 \dots N$.

(ii) The electric field operator acts as

$$\hat{E}^a(\vec{x}) |\gamma, \{p\}\rangle = \frac{X_{\gamma, \{p\}}^a(\vec{x})}{q_0} |\gamma, \{p\}\rangle. \quad (21)$$

Here

$$X_{\gamma, \{p\}}^a(\vec{x}) := \sum_{i=1}^N p_i \int_{e_i} ds_i \delta^3(\vec{e}_i(s_i), \vec{x}) \dot{e}_i^a. \quad (22)$$

In this equation e_i denotes the i th edge of γ and is parametrized by the parameter s_i so that $\vec{e}_i(s_i)$ is the coordinate of the point on the edge e_i at parameter value s_i .

The electric flux operator associated with a surface S with surface normal n_a acts as

$$\int_S \hat{E}^a n_a d^2 s |\gamma, \{p\}\rangle = \frac{\sum_{i_S} p_{i_S} \kappa_{i_S}}{q_0} |\gamma, \{p\}\rangle. \quad (23)$$

Here the index i_S ranges only over the edges of γ which intersect the surface S . $\kappa_{i_S} = 1$ iff $e_{i_S}^a n_a > 0$, $\kappa_{i_S} = -1$ iff $e_{i_S}^a n_a < 0$ and $\kappa_{i_S} = 0$ iff $e_{i_S}^a n_a = 0$. This action may be derived rigorously via a regularization along the lines of [24].

From Eq. (23), every charge network state $|\gamma, \{p\}\rangle$ is an eigenstate of the electric flux operator and the edges of γ may be pictured as carrying quanta of electric flux in multiples of q_0^{-1} . Although physical intuition for (and, indeed, the naming of) the QEF representation arises from this property of the electric flux, the electric flux operator itself will not play a role in the considerations of this work. Instead, it is the Gaussian smeared electric field operator which will play a key role and we now exhibit its action on charge network states.

In the notation of Eq. (3), it follows that Eq. (21) implies

$$\hat{E}_r^a(\vec{x}) |\gamma, \{p\}\rangle = \frac{X_{\gamma, \{p\}(r)}^a(\vec{x})}{q_0} |\gamma, \{p\}\rangle. \quad (24)$$

In addition to (i) and (ii), the Abelian nature of $U(1)$ implies the following.

(iii) Every charge network state $|\gamma, \{p\}\rangle$ can be obtained from the “vacuum” state $|\Omega\rangle$ [i.e. the state $\Omega(A) = 1$, $\Omega \in L^2(\overline{\mathcal{A}\mathcal{G}}, d\mu_0)$] via the action of the holonomy operator around a suitably defined loop, β , so that

$$|\gamma, \{p\}\rangle = \hat{H}_\beta |\Omega\rangle. \quad (25)$$

Here $\beta = \beta_1^{p_1} \circ \beta_2^{p_2} \cdots \circ \beta_N^{p_N}$ and $\beta_i^{p_i}$ denotes the loop obtained by traversing p_i times around β_i . β_i , $i = 1 \dots N$ are defined by the construction (3.2) of [7] as

$$\beta_i = Q(v_i^+) \circ e_i \circ Q(v_i^-) \quad (26)$$

where v_i^\pm are the vertices of γ which constitute the beginning and end points of e_i and $Q(v)$ is a path from the base point \vec{x}_0 to the point v such that $Q(v)$ intersects γ at most at a finite number of isolated points. It can be verified that

$$X_{\gamma, \{p\}}^a(\vec{x}) = X_\beta^a(\vec{x}) \quad (27)$$

and that Eq. (25) holds. For this reason we denote \hat{H}_β by $\hat{H}_{\gamma, \{p\}}$.

Conversely, it can be checked that for any loop $\beta \in \mathcal{L}_{x_0}$, $X_\beta^a(\vec{x}) = X_{\gamma, \{q\}}^a(\vec{x})$, where the closed, oriented graph γ is the union of the edges which comprise β with the orientations of the edges in γ chosen arbitrarily and the labeling $\{q\}$, given such a choice of orientation is as follows. Let the number of times an edge e_i is traversed in β , in the same direction as its orientation in γ , be q_{1i} . Let the number of times the edge e_i is traversed in β , in the opposite direction to its orientation in γ , be q_{2i} . Then e_i is labeled by $q_i = q_{1i} - q_{2i}$. Henceforth,

we shall use the labeling of holonomies by their associated charge networks (i.e. $H_{\gamma, \{q\}}$) interchangeably with their labeling by loops (i.e. H_β). Thus, if there is no charge labeling in the subscript to H , the label is to be understood as a loop; else as an associated charge network.

(iv) Consider the holonomy operator associated with charge network label $(\alpha, \{q\})$.⁵ Then $\hat{H}_{\alpha, \{q\}}$ maps $|\gamma, \{p\}\rangle$ to a new charge network state based on the graph $\gamma \cup \alpha$ consisting of the union of the sets of edges belonging to γ and α .⁶ The edges of $\gamma \cup \alpha$ are oriented and labeled with charges as follows. Edges which are not shared by γ and α retain their orientations and charge labels. Any shared edge labeled by the charge p in γ retains its orientation from γ and has charge $p + q$ if it has the same orientation in α and charge $p - q$ if it has opposite orientation in α . We denote this new state by $|\gamma \cup \alpha, \{p \cup q\}\rangle$. Thus

$$\hat{H}_{\alpha, \{q\}} |\gamma, \{p\}\rangle = |\gamma \cup \alpha, \{p \cup q\}\rangle. \quad (28)$$

It can be checked that Eqs. (20), (24) and (28) define a representation of the Poisson bracket algebra of Eq. (9) such that \hat{H}_γ is unitary and $\hat{E}_r^a(\vec{x})$ is self-adjoint.

Next, we review the construction of the “dual” representation on the space of algebraic duals. Let $\mathcal{D} \subset L^2(\overline{\mathcal{A}\mathcal{G}}, d\mu_0)$ be the (dense) set of finite linear combinations of charge network states. Let \mathcal{D}^* be the space of algebraic duals to \mathcal{D} i.e. every $\Phi \in \mathcal{D}^*$ is a complex linear map on \mathcal{D} . Define the action of $\hat{E}_r^a(\vec{x}), \hat{H}_\alpha$ on \mathcal{D}^* through

$$\begin{aligned} [\hat{E}_r^a(\vec{x})\Phi](|\gamma, \{p\}\rangle) &:= \Phi(\hat{E}_r^a(\vec{x})^\dagger |\gamma, \{p\}\rangle) \\ &= \Phi(\hat{E}_r^a(\vec{x}) |\gamma, \{p\}\rangle), \end{aligned} \quad (29)$$

$$\begin{aligned} (\hat{H}_\alpha \Phi)(|\gamma, \{p\}\rangle) &:= \Phi(\hat{H}_\alpha^\dagger |\gamma, \{p\}\rangle) \\ &= \Phi(\hat{H}_{\alpha^{-1}} |\gamma, \{p\}\rangle). \end{aligned} \quad (30)$$

Equations (29) and (30) provide an (anti-)representation of the Poisson bracket algebra (9). Note that \mathcal{D}^* merely provides a linear representation space for the dual representation—it does not inherit any natural inner product from \mathcal{D} .

In Sec. III B we shall see that the natural arena to discuss the relation between the r -Fock representation and the QEF representation is the space of algebraic duals, \mathcal{D}^* .

⁵Strictly speaking the discussion should and can be framed in terms of holonomically equivalent labels [i.e. $(\alpha, \{p\})$ is equivalent to $(\beta, \{q\})$ iff $X_{\alpha, \{p\}}^a = X_{\beta, \{q\}}^a$]. We gloss over this subtlety in the interest of pedagogy.

⁶It is assumed that the edges of α, γ overlap only if they are identical and that intersections of α, γ occur only at vertices of α, γ . This entails no loss of generality, since we can always find graphs which are holonomically equivalent to α, γ and for which the assumption holds.

B. The condition of Poincaré invariance

Even in linear quantum field theory there is no analog of the Stone–von Neuman uniqueness theorem for quantum mechanics on a vector space. Hence there are infinitely many inequivalent representations of the Poisson brackets algebra of smeared holonomies [see Eq. (6)]. The Fock space representation is singled out by the additional requirement of Poincaré invariance, which in turn, is encoded in the specific choice of complex structure (i.e. the positive-negative frequency decomposition) for the Fock representation. This choice is equivalent to the requirement that the Fock vacuum be a zero eigenstate of the Fock space annihilation operators. In Sec. III B 1 we shall express this requirement as a relation between the action of the smeared holonomy and electric field operators on the Fock vacuum [see Eq. (37)].

The image of this relation in the r -Fock representation [see Eq. (38)] may be thought of as a condition which picks out the r -Fock vacuum. In other words, the r -Fock vacuum may be thought of as a solution to the condition (38). Since this condition relates the action of the holonomy and the smeared electric field operators, it is well defined in *any* representation of the Poisson brackets algebra of Eq. (9). In Sec. III B 2 we impose the condition (38) in the (dual-) QEF representation and show that it has a *unique* (up to a multiplicative constant) solution. Since this condition is deduced from the requirement of Poincaré invariance, we may interpret the solution of Eq. (38) as the r -Fock vacuum expressed as a state in the (dual-) QEF representation.

1. Poincaré invariance in terms of smeared holonomies

Our conventions for the Fock space representation are as follows. The expansions of the field operators in the Coulomb gauge are

$$\hat{A}_a(\vec{x}) = \frac{1}{q_0(2\pi)^{3/2}} \int \frac{d^3k}{\sqrt{k}} \left(e^{i\vec{k}\cdot\vec{x}} \frac{\hat{a}_a(\vec{k})}{\sqrt{2}} + \text{Hermitian conjugate} \right), \quad (31)$$

$$\hat{E}_a(\vec{x}) = \frac{1}{(2\pi)^{3/2}} \int d^3k \sqrt{k} \left(-i e^{i\vec{k}\cdot\vec{x}} \frac{\hat{a}_a(\vec{k})}{\sqrt{2}} + \text{Hermitian conjugate} \right) \quad (32)$$

$$\Rightarrow \hat{H}_{\alpha, \{q\}(r)} = \exp \left(\frac{i}{q_0} \int \frac{d^3k}{\sqrt{2k}} X_{\alpha, \{q\}(r)}^a(\vec{k}) \times [\hat{a}_a(\vec{k}) + \hat{a}_a^\dagger(\vec{k})] \right). \quad (33)$$

The commutation relation between the annihilation and creation operators is

$$[\hat{a}_a(\vec{k}), \hat{a}_b^\dagger(\vec{l})] = \left(\delta_{ab} - \frac{k_a k_b}{k^2} \right) \delta(\vec{k}, \vec{l}). \quad (34)$$

Using these equations, the vacuum expectation value of the smeared holonomy operator [also called, in the language of [22], “the Fock positive linear functional” (PLF) and denoted by Γ_F] evaluates to⁷

$$\begin{aligned} \Gamma_F([\alpha, \{q\}]) &:= \langle 0 | H_{\alpha, \{q\}(r)} | 0 \rangle \\ &= \exp \left(-\frac{1}{4q_0^2} \int \frac{d^3k}{k} |X_{\alpha, \{q\}(r)}^a(\vec{k})|^2 \right). \end{aligned} \quad (35)$$

With Eq. (34), Eqs. (32) and (33) provide a representation of the Poisson brackets algebra (6). As discussed above, the condition of Poincaré invariance implies

$$\hat{a}_a(\vec{k}) | 0 \rangle = 0. \quad (36)$$

This, in turn, implies the following relation between the smeared holonomy and the electric field operators:

$$\begin{aligned} &[\Gamma_F([\alpha, \{q\}])]^2 \\ &\times \exp \left[-\int \frac{d^3x}{q_0} \left(\frac{1}{\sqrt{-(\partial^c \partial_c)}} X_{\alpha, \{q\}(r)}^a(\vec{x}) \hat{E}_a(\vec{x}) \right) \right] | 0 \rangle \\ &= \hat{H}_{\alpha, \{q\}(r)}^\dagger | 0 \rangle. \end{aligned} \quad (37)$$

This equation holds for *every* closed oriented graph α and encodes the condition of Poincaré invariance in terms of elements of the algebra of smeared holonomies and the electric field.

2. Poincaré invariance in terms of holonomies

The image of condition (37) in the r -Fock representation is

$$\begin{aligned} &[\Gamma_F([\alpha, \{q\}])]^2 \exp \left[-\int \frac{d^3x}{q_0} \left(\frac{1}{\sqrt{-(\partial^c \partial_c)}} X_{\alpha, \{q\}(r)}^a(\vec{x}) \right) \right. \\ &\left. \times \hat{E}_{ra}(\vec{x}) \right] | 0_r \rangle = \hat{H}_{\alpha, \{q\}}^\dagger | 0_r \rangle. \end{aligned} \quad (38)$$

We impose this condition in the QEF representation on the space \mathcal{D}^* of algebraic duals defined in Sec. III A. Thus, the following equation is to be solved for some $\Phi_0 \in \mathcal{D}^*$:

$$\begin{aligned} &[\Gamma_F([\alpha])]^2 \exp \left[-\int \frac{d^3x}{q_0} \left(\frac{1}{\sqrt{-(\partial^c \partial_c)}} X_{\alpha, \{q\}(r)}^a(\vec{x}) \right) \right. \\ &\left. \times \hat{E}_{ra}(\vec{x}) \right] \Phi_0 = \hat{H}_{\alpha, \{q\}}^\dagger \Phi_0. \end{aligned} \quad (39)$$

⁷Note that the expression for Γ_F in [22] is incorrect. The correct expression is Eq. (35) above and differs from the expression in [22] by a factor in its exponent. Also, in [22] the parameter q_0^{-1} was written as e , but factors of e appeared in [22] often in the wrong places. We have corrected the erroneous expressions of [22] in this work. With appropriate corrections regarding these factors of q_0 , all the results of [22] continue to hold.

Any element of \mathcal{D}^* can be written as a formal sum over all charge network states as follows. If $\Phi_0(|\gamma, \{p\}\rangle) = c_{\gamma, \{p\}}$ where $c_{\gamma, \{p\}}$ is a complex number, it follows that Φ_0 can be written as

$$\Phi_0 = \sum_{\gamma, \{p\}} c_{\gamma, \{p\}} |\gamma, \{p\}\rangle. \quad (40)$$

Substitution of this in Eq. (39) and projection of the resulting equation onto the ket $|\beta, \{t\}\rangle$ yields

$$\begin{aligned} & [\Gamma_F([\alpha, \{q\}])]^2 \sum_{\gamma, \{p\}} c_{\gamma, \{p\}} \langle \gamma, \{p\} | \exp \left[- \int \frac{d^3x}{q_0} \right. \\ & \left. \times \left(\frac{1}{\sqrt{-(\partial^c \partial_c)}} X_{\alpha, \{q\}(r)}^a(\vec{x}) \right) \hat{E}_{ra}(\vec{x}) \right] |\beta, \{t\}\rangle \\ & = \sum_{\gamma, \{p\}} c_{\gamma, \{p\}} \langle \gamma, \{p\} | \hat{H}_{\alpha, \{q\}} |\beta, \{t\}\rangle. \end{aligned} \quad (41)$$

Orthonormality of the charge network states, together with Eqs. (24) and (28), implies that

$$\begin{aligned} & [\Gamma_F([\alpha, \{q\}])]^2 \exp \left[- \int \frac{d^3x}{q_0^2} \left(\frac{1}{\sqrt{-(\partial^c \partial_c)}} X_{\alpha, \{q\}(r)}^a(\vec{x}) \right) \right. \\ & \left. \times X_{\beta, \{t\}(r)}(\vec{x}) \right] c_{\beta, \{t\}} = c_{\beta \cup \alpha, \{t \cup q\}} \end{aligned} \quad (42)$$

This equation can now be solved for the coefficients $c_{\gamma, \{p\}}$; of course, since the equation is linear and homogeneous the solution will be ambiguous by an overall constant. We fix this ambiguity by setting the coefficient labeled by the trivial graph $\gamma=0$, $\vec{0}(s)=\vec{x}_0$, to be unity. Then setting $\beta=0$ in Eq. (42) yields

$$c_{\alpha, \{q\}} = [\Gamma_F([\alpha, \{q\}])]^2 \quad (43)$$

for every charge network $|\alpha, \{q\}\rangle$. It may be verified that, miraculously, this also provides a solution to Eq. (42). Thus

$$\Phi_0 = \sum_{\gamma, \{p\}} [\Gamma_F([\gamma, \{p\}])]^2 |\gamma, \{p\}\rangle. \quad (44)$$

is the *unique* (up to an overall constant) solution to the condition (39). We identify Φ_0 as the state corresponding to the r -Fock vacuum.

As shown in [22] the action of the smeared holonomy operators on the Fock vacuum generates a dense subset of the Fock space. It follows that the action of the holonomy operators on the r -Fock vacuum generates a dense set of the r -Fock space. Therefore, we can use the dual representation of the holonomy operator [see Eq. (30)] on \mathcal{D}^* to generate the corresponding set of states from Φ_0 . Call this set \mathcal{L}^* . Thus any element of \mathcal{L}^* is of the form $\sum_{l=1}^N a_l \hat{H}_{\gamma^l, \{p^l\}} \Phi_0$ for some complex a_l and N finite.

As noted earlier, \mathcal{D}^* (and hence \mathcal{L}^*) is not equipped with an inner product. Therefore an inner product on \mathcal{L}^* must be

chosen which implements the classical ‘‘reality conditions,’’ namely $H_{\gamma, \{p\}}^* = H_{\gamma, \{-p\}}$ and $E_r(\vec{x})^* = E_r(\vec{x})$,⁸ on the corresponding quantum operators. It can be verified that the following inner product (naturally extendible to all of \mathcal{L}^*) implements the reality conditions:

$$\begin{aligned} & (\hat{H}_{\alpha, \{p\}} \Phi_0, \hat{H}_{\beta, \{q\}} \Phi_0) \\ & = \exp \left(- \frac{1}{4q_0^2} \int \frac{d^3k}{k} |X_{\alpha \cup \beta, \{-p \cup q\}(r)}^a(\vec{k})|^2 \right). \end{aligned} \quad (45)$$

It follows that the Cauchy completion of \mathcal{L}^* with respect to this inner product results in a Hilbert space which can be identified as r -Fock space and that the representation given by Eqs. (29) and (30) is exactly the r -Fock representation.

IV. CONCLUDING REMARKS

In this work we have related the diffeomorphism invariant, non-separable quantized electric flux representation for quantum $U(1)$ theory to its standard Poincaré invariant Fock space representation. This relation is based on the fact that the $U(1)$ holonomies play an important role in the construction of the QEF representation.

Since the holonomy operators are well defined in the QEF representation but *not* in the Fock representation, we first constructed a 1 parameter family of representations in which the holonomy operators *are* well defined and which are physically indistinguishable from the standard Fock representation. More precisely, the new representations are labeled by a positive parameter, r , with dimensions of length. For finite accuracy measurements at distance scales much larger than r , these ‘‘ r -Fock representations’’ are indistinguishable from the standard Fock representation.

Next, we related the r -Fock representation (for any fixed r) to the QEF representation by identifying the r -Fock vacuum as a (distributional) state in the (dual) QEF representation. This identification was achieved by solving, in the QEF representation, Eq. (39) inspired by Poincaré invariance, which enforced the condition that the annihilation operator of the r -Fock representation kill its vacuum state. The QEF representation is built on the property of diffeomorphism invariance and ‘‘knows’’ nothing about Poincaré invariance and hence we find it truly remarkable that the condition (39) which arises from Poincaré invariance of the Fock vacuum can be solved essentially uniquely in the (dual) QEF representation. Once the r -Fock vacuum was identified as a state in the dual QEF representation, we constructed states corresponding to a dense set in r -Fock space by the (dual) action of the holonomy operators on the r -Fock vacuum. Finally, the inner product was obtained on this set of states by requiring that the classical reality conditions be implemented as adjointness conditions on the corresponding quantum operators. Thus, in the qualitative language of the

⁸Rather than using these reality conditions directly for the operator $\hat{E}_r(\vec{x})$, it is simpler to use them to induce adjointness relations on the operators $\exp[-\int (d^3x/q_0) [(1/\sqrt{-(\partial^c \partial_c)} X_{\alpha, \{q\}(r)}^a(\vec{x})) \hat{E}_a(\vec{x})]$.

Introduction, we may say that non-normalizable infinite superpositions of 1 dimensional, polymer like excitations conspire, in collusion with the inner product (45), to acquire the character of 3D wavelike excitations in Fock space.

From the point of view of quantum gravity, we think that we have unraveled an important set of structures which will help relate the Fock space of gravitons of linearized gravity to appropriate semiclassical states in loop quantum gravity. In loop quantum gravity also, it is the *dual* representation to the kinematic spin network representation which serves as the “home” for physical, dynamically relevant quantum states of the gravitational field. Although there is no consensus on the exact physical states of the theory, it is still true that the *structure* of the physical states is that of distributions on the finite span of spin network states and that quantum operators act on these distributions via dual action. Further, the issue of the correct inner product on the space of physical states is still open and this inner product may have very different properties from the kinematical one in the context of solutions to the Hamiltonian constraint. Since the r -Fock representation for $U(1)$ theory has been obtained, in this work, as the dual representation on distributions to the finite span of charge network states with the “physical” inner product (45) unrelated to the “kinematic” inner product (20), we feel that our results will play an important role in relating gravitons to physical states in loop quantum gravity. In the loop quantum gravity case there are other complications such as the “linearization” of the non-Abelian gauge group to 3 copies of $U(1)$ [23], as well as the identification of a state corresponding to flat spacetime. These issues are currently under investigation. It is also of interest to ask if there is a generalization of our r -Fock constructions to the non-Abelian case. In this regard, Ashtekar and Lewandowski [25] have shown that a suitable reformulation of our results using “heat kernel” ideas admits a generalization to the non-Abelian context.

Apart from potential applications to quantum gravity, it would be of interest to understand the r -Fock representations in their own right. In this regard one question which comes to mind is that of dynamics in the r -Fock representation. The dynamics of the r -Fock representation is governed by the “ r -image” of the standard Maxwell Hamiltonian operator. The classical Hamiltonian for free Maxwell theory is

$$H = \int d^3k [E^a(\vec{k})E_a(-\vec{k}) + B^a(\vec{k})B_a(-\vec{k})], \quad (46)$$

where $B^a(\vec{k})$ is the Fourier transform of the magnetic field, $\vec{B}(\vec{x}) := q_0 \vec{\nabla} \times \vec{A}(\vec{x})$. It follows that the classical function corresponding to the Hamiltonian operator for the r -Fock representation is given by

$$H_r = \int d^3k \{ E_r^a(\vec{k}) E_{ar}(-\vec{k}) + [e^{k^2 r^2/2} B^a(\vec{k})] \times [e^{k^2 r^2/2} B_a(-\vec{k})] \}. \quad (47)$$

From considerations similar to those at the end of Sec. II, it can be argued that the dynamics generated by H_r , though

different, is physically indistinguishable from the usual Maxwell dynamics generated by H in the context of finite accuracy measurements at distance scales much larger than r .

We digress here to describe an alternative viewpoint⁹ on the issue of physical indistinguishability of the r -Fock representation. This is the viewpoint that only *algebraic* properties of functions on phase space are physically measurable. With this viewpoint, there is no way of asserting whether the pair $(\hat{H}_{\gamma(r)}, \hat{E}^a(\vec{x}))$ are being measured with dynamics generated by \hat{H} in the Fock representation or the pair $(\hat{H}_\gamma, \hat{E}_r^a(\vec{x}))$ are being measured with dynamics generated by \hat{H}_r in the r -Fock representation. With such a viewpoint, at least in the context of source-free Maxwell theory, the physics of the r -Fock representation for *any* positive r is (exactly, not approximately) *identical* to that of the usual Fock representation. The difference between this viewpoint and the one in the preceding paragraph is that the latter tacitly assumes that there is some property other than purely algebraic properties of the pair $(\hat{H}_{\gamma(r)}, \hat{E}^a(\vec{x}))$ by virtue of which the measuring apparatus measures them rather than $(\hat{H}_\gamma, \hat{E}_r^a(\vec{x}))$. This ends our brief digression.

Another question of interest intrinsic to the r -Fock representation is the one we raised in [22] as to whether the r -Fock representation could be realized as an $L^2(\overline{\mathcal{A}/\mathcal{G}}, d\mu_{F(r)})$ representation for some measure $d\mu_{F(r)}$ on $\overline{\mathcal{A}/\mathcal{G}}$. As we show in the Appendix, the answer to this question is in the affirmative and it would be of interest to understand the properties of this new r -Fock measure. In this regard the recent work of Velhinho [26] contains a detailed analysis of the properties of the r -Fock measures. Whether these new representations have applications outside of loop quantum gravity remains to be seen.

ACKNOWLEDGMENTS

I gratefully acknowledge helpful discussions on this matter with Abhay Ashtekar. I am indebted to Marcus van Bers for sharing his insights with me.

APPENDIX

1. Existence of the r -Fock measure on $\overline{\mathcal{A}/\mathcal{G}}$

In what follows we shall freely use results and notation from [22] as well as from previous sections.¹⁰ We shall be brief—the interested reader may work out the details. The r -Fock measure exists on $\overline{\mathcal{A}/\mathcal{G}}$ iff the r -Fock positive linear functional,

$$\Gamma_{F(r)} \left(\sum_{I=1}^N a_I [\alpha_I] \right) = \sum_{I=1}^N a_I \exp \left(- \frac{1}{4q_0^2} \int \frac{d^3k}{k} |X_{a_{I(r)}}^a(\vec{k})|^2 \right), \quad (A1)$$

⁹I learned of this viewpoint from Abhay Ashtekar.

¹⁰We shall use the corrected expressions of [22]—see footnote 7 in this regard.

is continuous with respect to the C^* norm, $\|\sum_{I=1}^N a_I[\alpha_I]\| := \sup_{A \in \mathcal{A}} |\sum_{I=1}^N a_I H_{\alpha_I}(A)|$. From [22] we have that

$$\left\| \sum_{I=1}^N a_I[\alpha_I] \right\| = \sup_{A \in \mathcal{A}} \left| \sum_{I=1}^N a_I H_{\alpha_I}(A) \right|. \quad (\text{A2})$$

Also, from Eq. (10) we have that

$$\langle 0 | \sum_{I=1}^N a_I \hat{H}_{\alpha_I}(r) | 0 \rangle = \int_{A_a \in \mathcal{S}^*} d\mu_G(A) \sum_{I=1}^N a_I H_{\alpha_I}(r)(A) \quad (\text{A4})$$

$$\Rightarrow \left| \langle 0 | \sum_{I=1}^N a_I \hat{H}_{\alpha_I}(r) | 0 \rangle \right| \leq \sup_{A \in \mathcal{S}^*} \left| \sum_{I=1}^N a_I H_{\alpha_I}(r)(A) \right| \int_{A_a \in \mathcal{S}^*} d\mu_G(A) = \sup_{A \in \mathcal{S}^*} \left| \sum_{I=1}^N a_I H_{\alpha_I}(r)(A) \right|. \quad (\text{A5})$$

Since $X_{\alpha(r)}^a(\vec{x})$ is in Schwartz space, it follows that every $A_a \in \mathcal{S}^*$ defines a homeomorphism, h , from $\mathcal{H}G_r$ to $U(1)$ [the element of $U(1)$ corresponding to a loop α is just $\exp i \int_{R^3} X_{\alpha(r)}^a(\vec{x}) A_a(\vec{x}) d^3x$]. It follows from the considerations of [22] [see especially Eq. (A15) of [22]] that

$$\sup_{A \in \mathcal{S}^*} \left| \sum_{I=1}^N a_I H_{\alpha_I}(r)(A) \right| \leq \sup_{A \in \mathcal{A}} \left| \sum_{I=1}^N a_I H_{\alpha_I}(r)(A) \right|. \quad (\text{A6})$$

$$\Gamma_{F(r)} \left(\sum_{I=1}^N a_I[\alpha_I] \right) = \langle 0 | \sum_{I=1}^N a_I \hat{H}_{\alpha_I}(r) | 0 \rangle. \quad (\text{A3})$$

Using the standard $L^2(\mathcal{S}^*, d\mu_G)$ [22,27] representation of Fock space where \mathcal{S}^* is an appropriate space of tempered distributions and $d\mu_G$ is the standard, unit volume, Gaussian measure, we have

It follows from Eqs. (A2), (A3), (A5) and (A6) that

$$\left| \Gamma_{F(r)} \left(\sum_{I=1}^N a_I[\alpha_I] \right) \right| \leq \left\| \sum_{I=1}^N a_I[\alpha_I] \right\|. \quad (\text{A7})$$

This implies that $\Gamma_{F(r)}$ is continuous with respect to $\|\cdot\|$ and hence that an r -Fock measure, $d\mu_{F(r)}$ exists on \mathcal{A}/\mathcal{G} .

- [1] T. A. Schilling, "Bibliography of publications related to Classical and Quantum Gravity in terms of the Ashtekar variables," gr-qc/9409031.
- [2] C. Rovelli, *Living Rev. Relativ.* **1**, 1 (1998).
- [3] A. Ashtekar, *Phys. Rev. D* **36**, 1587 (1987).
- [4] J. Fernando Barbero G., *Phys. Rev. D* **51**, 5507 (1995).
- [5] C. Rovelli and L. Smolin, *Nucl. Phys.* **B331**, 80 (1990).
- [6] T. Jacobson and L. Smolin, *Nucl. Phys.* **B299**, 295 (1988).
- [7] A. Ashtekar and J. Lewandowski, in *Quantum Gravity and Knots*, edited by J. Baez (Oxford University Press, Oxford, 1994).
- [8] A. Ashtekar and C. J. Isham, *Class. Quantum Grav.* **9**, 1433 (1992).
- [9] D. Marolf and J. Mourão, *Commun. Math. Phys.* **170**, 583 (1995).
- [10] J. Baez, *Lett. Math. Phys.* **31**, 213 (1994).
- [11] A. Ashtekar and J. Lewandowski, *J. Geom. Phys.* **17**, 19 (1995).
- [12] C. Rovelli and L. Smolin, *Phys. Rev. D* **52**, 5743 (1995).
- [13] J. Baez, in *The Interface of Knots and Physics*, edited by L. Kauffman (American Mathematical Society, Providence, 1996).
- [14] R. de Pietri, *Class. Quantum Grav.* **14**, 53 (1997); R. de Pietri and C. Rovelli, *Phys. Rev. D* **54**, 2664 (1996).
- [15] A. Ashtekar, J. Lewandowski, D. Marolf, J. Mourão, and T. Thiemann, *J. Math. Phys.* **36**, 6456 (1995).
- [16] T. Thiemann, *Class. Quantum Grav.* **15**, 839 (1998).
- [17] L. Smolin, gr-qc/9609034; J. Lewandowski and D. Marolf, *Int. J. Mod. Phys. D* **7**, 299 (1998).
- [18] M. Gaul and C. Rovelli, *Class. Quantum Grav.* **18**, 1593 (2001); Roberto de Pietri (private communication).
- [19] M. Bojowald, *Phys. Rev. Lett.* **86**, 5227 (2001).
- [20] A. Corichi and K. V. Krasnov, *Mod. Phys. Lett. A* **13**, 1339 (1998).
- [21] A. Ashtekar and C. J. Isham, *Phys. Lett. B* **274**, 393 (1992).
- [22] M. Varadarajan, *Phys. Rev. D* **61**, 104001 (2000).
- [23] A. Ashtekar, C. Rovelli, and L. Smolin, *Phys. Rev. D* **44**, 1740 (1991).
- [24] A. Ashtekar and J. Lewandowski, *Class. Quantum Grav.* **14**, A55 (1997).
- [25] A. Ashtekar and J. Lewandowski, gr-qc/0107043.
- [26] J. M. Velhinho, math-ph/0107002.
- [27] J. Glimm and A. Jaffe, *Quantum Physics* (Springer-Verlag, New York, 1987).
- [28] A. Ashtekar and A. Corichi, *Class. Quantum Grav.* **14**, A43 (1997).