

Nonresonant contributions in $B \rightarrow \rho \pi$ decay

Jusak Tandean*

Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506-0055

S. Gardner†

*Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506-0055‡
and Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309*

(Received 16 April 2002; published 20 August 2002)

We consider nonresonant contributions in the Dalitz-plot analysis of $B \rightarrow \rho \pi \rightarrow \pi^+ \pi^- \pi^0$ decay and their potential impact on the extraction of the Cabibbo-Kobayashi-Maskawa parameter α . In particular, we examine the role of the heavy mesons B^* and B_0 , via the process $B \rightarrow \pi(B^*, B_0) \rightarrow \pi^+ \pi^- \pi^0$, and their interference with resonant contributions in the ρ -mass region. We discuss the inherent uncertainties and suggest that the effects may be substantially smaller than previously indicated.

DOI: 10.1103/PhysRevD.66.034019

PACS number(s): 13.25.Hw, 11.30.Er, 12.15.Hh, 14.40.Nd

I. INTRODUCTION

The recent observation of CP violation in the B -meson system, realized through the measurement of a nonzero, time-dependent, CP -violating asymmetry in the process $B^0(\bar{B}^0) \rightarrow J/\psi K_S$ (and related ones) [1], heralds a new era of discovery. The result yields a value of $\sin(2\beta)$ in accord with standard model (SM) expectations [2], where β , defined by $\exp(i\beta) \equiv -V_{cb}^* V_{cd} / (V_{ub}^* V_{ud})$, is an angle of the unitarity triangle, V_{ij} being an element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [3]. Ascertaining the presence of physics beyond the SM thus demands the determination of all the angles of the unitarity triangle.

In this paper, we consider the decays $B^0(\bar{B}^0) \rightarrow \rho \pi \rightarrow \pi^+ \pi^- \pi^0$, as a Dalitz-plot analysis of the possible $\rho \pi$ final states, under the assumption of isospin symmetry, permits the determination of the CKM parameter α [4], where $\alpha = \pi - \beta - \gamma$ and $\exp(i\gamma) \equiv -V_{ub}^* V_{ud} / (V_{cb}^* V_{cd})$. Our interest is in assessing the size of the nonresonant contributions which could possibly obscure the analysis, and in ameliorating their impact. Indeed, the strategy for the extraction of α relies, in part, on the assumption that the ρ mesons dominate the 3π final state. There are, however, empirical indications that this assumption may not always be warranted. For example, combining the CLEO measurements of the branching fractions, $\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm) = (27.6^{+8.4}_{-7.4} \pm 4.2) \times 10^{-6}$ and $\mathcal{B}(B^- \rightarrow \rho^0 \pi^-) = (10.4^{+3.3}_{-3.4} \pm 2.1) \times 10^{-6}$ [5], with the BABAR result $\mathcal{B}(B^0 \rightarrow \rho^\pm \pi^\mp) = (28.9 \pm 5.4 \pm 4.3) \times 10^{-6}$ [6] yields

$$\mathcal{R} = \frac{\mathcal{B}(\bar{B}^0 \rightarrow \rho^\mp \pi^\pm)}{\mathcal{B}(B^- \rightarrow \rho^0 \pi^-)} = 2.7 \pm 1.2, \quad (1)$$

where we have added the errors in quadrature and ignored correlations. These ratios are small [7] with respect to simple

theoretical estimates, which give $\mathcal{R} \sim 6$ [8]. An interesting possibility for the resolution of this discrepancy has been suggested in Refs. [9,10], whose authors investigate the possible backgrounds to $B \rightarrow \rho \pi \rightarrow 3\pi$ decay which arise from contributions mediated by other resonances. They find that the light σ resonance, a broad $I=J=0$ enhancement in $\pi\pi$ scattering, as well as the heavy-meson resonances B^* ($J^P = 1^-$) and B_0 ($J^P = 0^+$), can modify the $B \rightarrow 3\pi$ branching ratios in the ρ -mass region and give rise to values of \mathcal{R} crudely compatible with the empirical value of Eq. (1), given its large error. In particular, the contribution of $B^- \rightarrow \sigma \pi^-$ decay significantly enhances the effective $B^- \rightarrow \rho^0 \pi^-$ branching ratio and lowers the value of \mathcal{R} . Analogously, the σ modestly impacts the $B^0 \rightarrow \rho^0 \pi^0$ branching ratio [11]; let us consider the issues.

The analysis of $B^0(\bar{B}^0) \rightarrow \rho \pi \rightarrow \pi^+ \pi^- \pi^0$ decay posits a two-step process, that is, that the amplitude for $B^0 \rightarrow \pi^+ \pi^- \pi^0$ decay can be written as

$$A(B^0 \rightarrow \pi^+ \pi^- \pi^0) = f_+ a_{+-} + f_- a_{-+} + f_0 a_{00}, \quad (2)$$

where $a_{ij} \equiv A(B^0 \rightarrow \rho^i \pi^j)$ and f_i is the vector form-factor describing $\rho^i \rightarrow \pi\pi$ [4]. An analogous construct can be made for $B^0(\bar{B}^0) \rightarrow \sigma \pi \rightarrow \pi^+ \pi^- \pi^0$ decay, which contains the scalar form-factor describing $\sigma \rightarrow \pi^+ \pi^-$. It is evident that the manner in which the σ populates the ρ phase-space will depend on the amplitude for $B^0 \rightarrow \sigma \pi^0$ decay, as well as on the accompanying scalar form-factor. The σ is a state of definite CP , so that the isospin analysis of Ref. [4] can be enlarged to include it [11]; nevertheless, the analysis relies on the form factors adopted for the $\rho^i \rightarrow \pi\pi$ and $\sigma \rightarrow \pi\pi$ processes. The resonances of interest are broad, so that Breit-Wigner form-factors are generally insufficient: they do not satisfy general theoretical constraints, such as analyticity and unitarity, over the $\pi\pi$ invariant-mass interval needed. As discussed in detail in Ref. [11], the differences are striking for the scalar form-factor, and the resulting numerical impact on $B \rightarrow 3\pi$ decay is sizable. In contrast, the numerical differences for the vector form-factor are not large.

*Electronic address: jtandean@pa.uky.edu

†Electronic address: gardner@pa.uky.edu

‡Permanent address.




FIG. 1. Diagrams contributing to $B \rightarrow 3\pi$, decay with each square denoting a weak vertex.

The purpose of this paper is to extend the work of Ref. [11], which deals exclusively with the ρ and σ contributions. We incorporate the B^* and B_0 contributions suggested in Ref. [9], as the effects they find in the $B^0 \rightarrow \rho^0 \pi^0$ channel are considerable. In this paper, however, we show that the off-shell nature of the B^* and B_0 weak and strong vertices adds considerably to the uncertainty of the estimate of Ref. [9] and may well reduce these contributions significantly. Nevertheless, we also explore kinematical cuts which would be useful in reducing the impact of these effects in the ρ -mass region.

We begin in Sec. II with the weak, effective Hamiltonian and the matrix elements pertinent to our calculations. Subsequently, in Sec. III, we derive the amplitudes associated with the various contributions of interest in the ρ -mass region of $B \rightarrow 3\pi$ decay. We discuss our numerical results in Sec. IV and conclude in Sec. V.

II. EFFECTIVE HAMILTONIAN AND MATRIX ELEMENTS

The effective, $|\Delta B| = 1$ Hamiltonian for $b \rightarrow dq\bar{q}$ decay is given by [12]

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left[\lambda_u (C_1 O_1^u + C_2 O_2^u) + \lambda_c (C_1 O_1^c + C_2 O_2^c) - \lambda_t \sum_{i=3}^{10} C_i O_i \right], \quad (3)$$

where G_F is the Fermi coupling constant, $\lambda_q \equiv V_{qb} V_{qd}^*$ are CKM factors, C_i are Wilson coefficients, and O_i are four-quark operators. The expressions for C_i and O_i are detailed in Ref. [12], though we interchange $C_1 O_1^q \leftrightarrow C_2 O_2^q$, so that $C_1 \sim 1$ and $C_1 > C_2$. We neglect the electroweak-penguin operators $O_{7, \dots, 10}$ because their coefficients $C_{7, \dots, 10}$ are smaller than the others. In the decay amplitudes that we derive, the C_i enter through the combinations $a_i = C_i + C_{i+1}/N_c$ if i is odd and $a_i = C_i + C_{i-1}/N_c$ if i is even, where $N_c = 3$ is the number of colors.

The diagrams contributing to the $B \rightarrow 3\pi$ amplitudes considered here, as shown in Fig. 1, each have a strong vertex and a weak vertex, where the latter describes the transition $M_b \rightarrow M_1 M_2$, in which M_b is a heavy meson containing a b quark and $M_{1,2}$ are light mesons. The amplitude corresponding to the weak vertex is given by

$$A(M_b \rightarrow M_1 M_2) = \langle M_1 M_2 | \mathcal{H}_{\text{eff}} | M_b \rangle. \quad (4)$$

To evaluate this, we adopt the naive factorization approximation, following earlier calculations [9–11] to which we compare.

The relevant matrix elements are

$$\langle \pi^-(p) | \bar{d} \gamma^\mu L u | 0 \rangle = \sqrt{2} \langle \pi^0(p) | \bar{u} \gamma^\mu L u | 0 \rangle = i f_\pi p^\mu, \quad (5)$$

$$\langle \rho^-(p, \varepsilon) | \bar{d} \gamma^\mu u | 0 \rangle = \sqrt{2} \langle \rho^0(p, \varepsilon) | \bar{u} \gamma^\mu u | 0 \rangle = f_\rho \varepsilon^{*\mu},$$

$$q^\mu \langle \rho^+(p, \varepsilon) | \bar{u} \gamma_\mu L b | \bar{B}^0(k) \rangle = -2i A_0^{B^0 \rightarrow \rho}(q^2) M_\rho \varepsilon^* \cdot q,$$

$$\begin{aligned} \langle \pi^+(p) | \bar{u} \gamma^\mu L b | \bar{B}^0(k) \rangle &= (k+p)^\mu F_1^{B^0 \rightarrow \pi}(q^2) \\ &+ \frac{M_B^2 - M_\pi^2}{q^2} q^\mu [F_0^{B^0 \rightarrow \pi}(q^2) - F_1^{B^0 \rightarrow \pi}(q^2)], \end{aligned} \quad (6)$$

$$q^\mu \langle \sigma(p) | \bar{d} \gamma_\mu L b | \bar{B}^0(k) \rangle = -i(M_B^2 - M_\sigma^2) F_0^{B^0 \rightarrow \sigma}(q^2),$$

$$\begin{aligned} q^\mu \langle \pi^+(p) | \bar{u} \gamma_\mu L b | \bar{B}^{*0}(k, \varepsilon_{B^*}) \rangle \\ = 2i \sqrt{2} A_0^{B^* \rightarrow \pi}(q^2) M_{B^*} \varepsilon_{B^*} \cdot q, \end{aligned} \quad (7)$$

$$\begin{aligned} q^\mu \langle \pi^+(p) | \bar{u} \gamma^\mu L b | \bar{B}_0^0(k) \rangle \\ = -i(M_{B_0}^2 - M_\pi^2) F_0^{B_0 \rightarrow \pi}(q^2), \end{aligned}$$

where f_π and f_ρ are the usual decay constants, $q \equiv k - p$, and $L \equiv 1 - \gamma_5$. The various $A_0(q^2)$ and $F_{0,1}(q^2)$ are form factors. Other meson-to-meson matrix elements can be determined using isospin symmetry. In our phase convention, the meson flavor wave functions are given by $\pi^+ = u\bar{d}$, $\sqrt{2}\pi^0 = u\bar{u} - d\bar{d}$, $\pi^- = d\bar{u}$, $\bar{B}^0 = b\bar{d}$, $B^- = b\bar{u}$, and similarly for the ρ , B^* , and B_0 . This implies that we have, for example, $\langle \pi^+ | \bar{u} \gamma_\mu b | \bar{B}^0 \rangle = -\sqrt{2} \langle \pi^0 | \bar{d} \gamma_\mu b | \bar{B}^0 \rangle = +\sqrt{2} \langle \pi^0 | \bar{u} \gamma_\mu b | B^- \rangle = \langle \pi^- | \bar{d} \gamma_\mu b | B^- \rangle$. We now employ these matrix elements to realize amplitudes for $B \rightarrow 3\pi$ decays.

III. AMPLITUDES

Practical considerations drive our interest in the $\pi^+ \pi^- \pi^0$ and $\pi^\mp \pi^\mp \pi^\pm$ decay modes; we shall not consider the $\pi^0 \pi^0 \pi^\pm$ ones. We write the amplitude for $\bar{B}^0 \rightarrow \pi^+(p_+) \pi^-(p_-) \pi^0(p_0)$ decay as a coherent sum of the ρ , σ , B^* , and B_0 amplitudes, namely,

$$A^{+-0} = A_\rho^{+-0} + A_\sigma^{+-0} + A_{B^*}^{+-0} + A_{B_0}^{+-0}. \quad (8)$$

For $B^- \rightarrow \pi^-(p_1) \pi^-(p_2) \pi^+(p_+)$, the amplitude A^{--+} can be constructed in an analogous manner.

We consider first the $B \rightarrow \rho \pi \rightarrow 3\pi$ contributions, represented by the diagram denoted by “ ρ ” in Fig. 1(a). For each ρ^i diagram and 3π state, the amplitude is written as a product of an amplitude for the $B \rightarrow \rho^i \pi^j$ weak transition and a vertex function $\Gamma_{\rho\pi\pi}$ describing the $\rho^i \rightarrow \pi\pi$ form factor. Were the ρ a narrow resonance, the Breit-Wigner (BW) form

$$\Gamma_{\rho\pi\pi}^{\text{BW}}(s) = \frac{g_\rho}{s - M_\rho^2 + i\Gamma_\rho M_\rho} \quad (9)$$

would suffice, where \sqrt{s} is the invariant mass of the 2π system and g_ρ is the $\rho \rightarrow \pi\pi$ coupling constant. However, since the ρ is not narrow—its width is some 20% of its mass—this form must be generalized to accommodate known theoretical constraints over the region in s for which it is appreciable. For example, unitarity and time-reversal invariance compel the phase of $\Gamma_{\rho\pi\pi}(s)$ to be that of $L=1, I=1$ $\pi\pi$ scattering for $s \leq (M_\pi + M_\omega)^2$, for which the scattering is elastic. Moreover, the imaginary part of $\Gamma_{\rho\pi\pi}(s)$ must vanish below physical threshold, $s = 4M_\pi^2$. For a detailed discussion with references to earlier work, see Refs. [11,13]. Following Ref. [13], we have

$$\Gamma_{\rho\pi\pi}(s) = \frac{-F_\rho(s)}{f_{\rho\gamma}}, \quad (10)$$

where $F_\rho(s)$ is the vector form-factor of the pion and $f_{\rho\gamma}$ is the ρ - γ coupling constant. The parameters in $F_\rho(s)$ are determined by fitting to $e^+e^- \rightarrow \pi^+\pi^-$ data; what is important is that the parametrization itself is consistent with theoretical constraints. The value of $f_{\rho\gamma}$ is determined from the $\rho \rightarrow e^+e^-$ width, which, in turn, is extracted from the $e^+e^- \rightarrow \pi^+\pi^-$ cross section at $s = M_\rho^2$ [13,14]. The overall sign is chosen so that Eq. (10) is equivalent to the BW form, Eq. (9), as $s \rightarrow M_\rho^2$. At $s = M_\rho^2$ the BW form is compatible with the various theoretical constraints. In our numerical analysis, we adopt the “solution B” fit of Ref. [13] for F_ρ , for which $f_{\rho\gamma} = 0.122 \pm 0.001 \text{ GeV}^2$ [14]. Alternatively, a BW form with a running width $\Gamma_\rho(s)$, chosen to be compatible with the form of the $\pi\pi$ phase shift (in the crossed channel) as $s \rightarrow 4M_\pi^2$, is given in Ref. [15]. However, the numerical differences between this form and the one we have chosen are small [11].

For the decay amplitudes, after summing over the ρ polarizations, we find

$$\begin{aligned} A_\rho^{+-0} &= \eta^-(s_{+-} - s_{+0})\Gamma_{\rho\pi\pi}(s_{-0}) \\ &\quad + \eta^+(s_{-0} - s_{+-})\Gamma_{\rho\pi\pi}(s_{+0}) \\ &\quad - \eta^0(s_{-0} - s_{+0})\Gamma_{\rho\pi\pi}(s_{+-}), \end{aligned} \quad (11)$$

$$\begin{aligned} A_\rho^{-+} &= -\bar{\eta}^0[(s_{12} - s_{1+})\Gamma_{\rho\pi\pi}(s_{2+}) \\ &\quad + (s_{12} - s_{2+})\Gamma_{\rho\pi\pi}(s_{1+})], \end{aligned}$$

where $s_{kl} \equiv (p_k + p_l)^2$, with

$$\begin{aligned} \eta^- &= \frac{G_F}{\sqrt{2}}(\lambda_u a_1 - \lambda_t a_4) f_\rho F_1^{B\pi}, \\ \eta^+ &= \frac{G_F}{\sqrt{2}}[\lambda_u a_1 - \lambda_t(a_4 - a_6 R_q)] f_\pi M_\rho A_0^{B\rho}, \\ \eta^0 &= \frac{-G_F}{2\sqrt{2}}\{[\lambda_u a_2 + \lambda_t(a_4 - a_6 R_q)] f_\pi M_\rho A_0^{B\rho} \\ &\quad + (\lambda_u a_2 + \lambda_t a_4) f_\rho F_1^{B\pi}\}, \\ \bar{\eta}^0 &= \frac{G_F}{2}\{[\lambda_u a_1 - \lambda_t(a_4 - a_6 R_q)] f_\pi M_\rho A_0^{B\rho} \\ &\quad + (\lambda_u a_2 + \lambda_t a_4) f_\rho F_1^{B\pi}\}. \end{aligned} \quad (12)$$

Here $A_0^{B\rho} \equiv A_0^{B \rightarrow \rho}(M_\pi^2)$ and $F_1^{B\pi} \equiv F_1^{B \rightarrow \pi}(M_\rho^2)$, whereas $R_q \equiv M_\pi^2 / [(m_b + \hat{m})\hat{m}]$ —note that we work in the isospin-symmetric limit, for which $\hat{m} = m_u = m_d$. The relative signs between the different terms in Eq. (11) follow from the $\rho\pi\pi$ couplings¹

$$\begin{aligned} \langle \pi^0(p_0) \pi^\pm(p_\pm) | \rho^\pm \rangle &= \pm g_\rho \varepsilon_\rho \cdot (p_\pm - p_0), \\ \langle \pi^+(p_+) \pi^-(p_-) | \rho^0 \rangle &= g_\rho \varepsilon_\rho \cdot (p_- - p_+), \end{aligned} \quad (13)$$

which follow, in turn, from the phase conventions we have chosen for the flavor wave functions: $|\pi^\pm\rangle = \mp |I=1, I_3 = \pm 1\rangle$ and $|\pi^0\rangle = |I=1, I_3 = 0\rangle$, and similarly for the ρ states. Our A_ρ amplitudes agree with those of earlier calculations [9,11,16].

We turn next to the σ “meson” contributions, represented by the diagram denoted by “ σ ” in Fig. 1(a). We use the σ to denote a two-pion state with total isospin $I=0$ and total angular-momentum $J=0$; it need not be a “pre-existing” resonance, but, rather, can be generated dynamically by the strong pionic final-state interactions in this channel [17]. The peak of the broad enhancement associated with the σ is close to the ρ in mass, so that the decay $B \rightarrow \sigma\pi \rightarrow 3\pi$ can populate the $B \rightarrow \rho\pi\pi$ phase space [10]. As in the ρ case, the amplitudes for $B \rightarrow \sigma\pi \rightarrow 3\pi$ decays are written as a product of an amplitude for the $B \rightarrow \sigma\pi$ weak transition and a vertex function $\Gamma_{\sigma\pi\pi}$ describing the $\sigma \rightarrow \pi\pi$ form factor. We write [11]

$$\Gamma_{\sigma\pi\pi}(s) = \chi \Gamma_1^{n*}(s), \quad (14)$$

where Γ_1^n is defined as

$$\langle 0 | \bar{d}d | \pi^+(p_+) \pi^-(p_-) \rangle = \sqrt{\frac{2}{3}} \Gamma_1^n(s_{+-}) B_0. \quad (15)$$

We note that $B_0 \equiv M_\pi^2 / (2\hat{m})$ is the vacuum quark condensate and χ is a normalization constant, to be discussed shortly.

¹We use the notation $\langle M_2 M_3 | M_1 \rangle \equiv \langle M_2 M_3 | \mathcal{H}_{\text{strong}} | M_1 \rangle$.

For our numerical work in the next section, we adopt the $\Gamma_1^n(s)$ as derived in Ref. [18], after Refs. [17,19,20]. The calculated form factor is realized in a chiral, unitarized, coupled-channel approach; at low energies, the form factor is matched to the one-loop-order expression in chiral perturbation theory [18,21]. The resulting form factor is consistent with low-energy constraints and is comparable to the scalar form-factor which emerges from the dispersion analysis of Ref. [22]; however, it is notably different from the Breit-Wigner form adopted in Refs. [10,23] to study the role of the σ in B and D decays into the 3π final state. That is,

$$\Gamma_{\sigma\pi\pi}^{\text{BW}}(s) = \frac{g_{\sigma\pi\pi}}{s - M_\sigma^2 + i\Gamma_\sigma(s)M_\sigma}, \quad (16)$$

$$\Gamma_\sigma(s) = \frac{M_\sigma\Gamma_\sigma}{\sqrt{s}} \sqrt{\frac{s - 4M_\pi^2}{M_\sigma^2 - 4M_\pi^2}},$$

where the coupling $g_{\sigma\pi\pi} \equiv \langle \pi^+\pi^- | \sigma \rangle$ is determined from the $\sigma \rightarrow \pi\pi$ decay rate. For $B \rightarrow 3\pi$ decay, the numerical

changes arising from the use of $\Gamma_{\sigma\pi\pi}(s)$ in place of the BW expression are significant [11], as we will see here as well. We determine the normalization χ by requiring that [11]

$$\chi |\Gamma_1^n(M_\sigma^2)| = \frac{g_{\sigma\pi\pi}}{\Gamma_\sigma(M_\sigma^2)M_\sigma}, \quad (17)$$

which equates $|\Gamma_{\sigma\pi\pi}(s)|$ to its BW counterpart at $s = M_\sigma^2$. The values of M_σ and Γ_σ are extracted from fits of $\Gamma_{\sigma\pi\pi}^{\text{BW}}(s)$ to $D \rightarrow 3\pi$ decays [23]. The normalization condition is motivated by noting that the modulus of $\Gamma_1^n(s)$ is peaked near $s = M_\sigma^2$, whereas the normalization of $\Gamma_1^n(s)$ is sensitive to the values of certain, poorly known low-energy constants [11]. We emphasize that M_σ and Γ_σ appear merely in the normalization of $\Gamma_{\sigma\pi\pi}$.

The resulting decay amplitudes are then

$$A_\sigma^{+-0} = \eta_\sigma^0 \Gamma_{\sigma\pi\pi}(s_{+-}), \quad (18)$$

$$A_\sigma^{-+} = \bar{\eta}_\sigma^0 [\Gamma_{\sigma\pi\pi}(s_{1+}) + \Gamma_{\sigma\pi\pi}(s_{2+})],$$

where

$$\eta_\sigma^0 = \frac{G_F}{2} \left\{ [\lambda_u a_2 + \lambda_t (a_4 - a_6 R_q)] (M_B^2 - M_\sigma^2) f_\pi F_0^{B\sigma} - \lambda_t a_6 \frac{2\langle \sigma | \bar{d}d | 0 \rangle}{m_b - \hat{m}} (M_B^2 - M_\pi^2) F_0^{B\pi} \right\}, \quad (19)$$

$$\bar{\eta}_\sigma^0 = \frac{G_F}{\sqrt{2}} \left\{ [\lambda_u a_1 - \lambda_t (a_4 - a_6 R_q)] (M_B^2 - M_\sigma^2) f_\pi F_0^{B\sigma} + \lambda_t a_6 \frac{2\langle \sigma | \bar{d}d | 0 \rangle}{m_b - \hat{m}} (M_B^2 - M_\pi^2) F_0^{B\pi} \right\},$$

with $F_0^{B\pi} \equiv F_0^{B \rightarrow \pi}(M_\sigma^2)$ and $F_0^{B\sigma} \equiv F_0^{B \rightarrow \sigma}(M_\pi^2)$. From Eqs. (14) and (15), it follows that $\langle \sigma | \bar{d}d | 0 \rangle = M_\pi^2 / (\sqrt{6}\chi\hat{m})$. We agree with the weak amplitudes of Ref. [11], but disagree with those of Ref. [10] in that our η_σ^0 and $\bar{\eta}_\sigma^0$, neglecting penguin terms, are smaller and larger, respectively, than theirs by a factor of $\sqrt{2}$.

We now evaluate the B^* and B_0 contributions, whose diagrams are shown in Fig. 1(b); we suppose that other excited B -meson states could also contribute, but we expect that their larger masses ought to make them less important [9]. Presently, no reliable data exist on the widths of these heavy mesons, so that their values have to be calculated. Recent estimates [24,25] suggest that the B^* is a very narrow resonance, whereas the B_0 is less so, its width being some 6% of its mass. Nevertheless, the resonances are sufficiently narrow that it is reasonable to adopt a Breit-Wigner representation for the propagators of these

mesons, as in Ref. [9]. In the combined heavy-quark and chiral limit [26], the strong couplings connecting the (B^*, B_0) , B , and π mesons are [9,24,27]²

$$\langle B^-(p') \pi^+(p) | \bar{B}^{*0}(k, \varepsilon) \rangle = -\frac{2g\sqrt{M_B M_{B^*}}}{f_\pi} \varepsilon \cdot p, \quad (20)$$

$$\langle B^-(p') \pi^+(p) | \bar{B}_0^0(k) \rangle = \frac{h\sqrt{M_B M_{B_0}}}{f_\pi} \frac{k^2 - M_B^2}{M_{B_0}}. \quad (21)$$

Using isospin symmetry, we derive

²We note that $\langle B^{*-}(k, \varepsilon) \pi^+ | \bar{B}^0 \rangle = -\langle B^- \pi^+ | \bar{B}^{*0}(k, \varepsilon) \rangle$ and $\langle B_0^-(k) \pi^+ | \bar{B}^0 \rangle = -\langle B^- \pi^+ | \bar{B}_0^0(k) \rangle$.

$$\langle B^- \pi^+ | \bar{B}^{*0} \rangle = -\sqrt{2} \langle \bar{B}^0 \pi^0 | \bar{B}^{*0} \rangle = \langle \bar{B}^0 \pi^- | B^{*-} \rangle \quad (22)$$

and analogous relations for $\langle B \pi | B_0 \rangle$. We then obtain

$$A_{B^*}^{+-0} = \frac{1}{\sqrt{2}} \frac{K \Pi(s_{-0}, s_{+-}) + K_1 \Pi(s_{-0}, s_{+0})}{s_{-0} - M_{B^*}^2 + i \Gamma_{B^*} M_{B^*}} - \frac{1}{\sqrt{2}} \frac{K \Pi(s_{+-}, s_{-0})}{s_{+-} - M_{B^*}^2 + i \Gamma_{B^*} M_{B^*}}, \quad (23)$$

$$A_{B^*}^{--0} = \frac{K \Pi(s_{1+}, s_{12})}{s_{1+} - M_{B^*}^2 + i \Gamma_{B^*} M_{B^*}} + \frac{K \Pi(s_{2+}, s_{12})}{s_{2+} - M_{B^*}^2 + i \Gamma_{B^*} M_{B^*}},$$

$$A_{B_0}^{+-0} = \left(\frac{\tilde{K}^0 + \tilde{K}^{cc}}{s_{-0} - M_{B_0}^2 + i \Gamma_{B_0} M_{B_0}} - \frac{\tilde{K}^0}{s_{+-} - M_{B_0}^2 + i \Gamma_{B_0} M_{B_0}} \right) \frac{(M_{B_0}^2 - M_\pi^2)}{\sqrt{2}}, \quad (24)$$

$$A_{B_0}^{--0} = \left(\frac{\tilde{K}^0}{s_{1+} - M_{B_0}^2 + i \Gamma_{B_0} M_{B_0}} + \frac{\tilde{K}^0}{s_{2+} - M_{B_0}^2 + i \Gamma_{B_0} M_{B_0}} \right) (M_{B_0}^2 - M_\pi^2),$$

where

$$K = -4 G_F [\lambda_u a_1 - \lambda_t (a_4 - a_6 R_q)] \times g M_{B^*} \sqrt{M_B M_{B^*}} A_0^{B^* \pi}, \quad (25)$$

$$K_1 = -2 \sqrt{2} G_F [\lambda_u a_2 + \lambda_t (a_4 - a_6 R_q)] \times g M_{B^*} \sqrt{M_B M_{B^*}} A_0^{B^* \pi},$$

$$\tilde{K}^0 = \frac{G_F}{\sqrt{2}} [\lambda_u a_1 - \lambda_t (a_4 - a_6 R_q)] \times \frac{M_{B_0}^2 - M_B^2}{M_{B_0}} h \sqrt{M_B M_{B_0}} F_0^{B_0 \pi}, \quad (26)$$

$$\tilde{K}^{cc} = \frac{G_F}{\sqrt{2}} [\lambda_u a_2 + \lambda_t (a_4 - a_6 R_q)] \times \frac{M_{B_0}^2 - M_B^2}{M_{B_0}} \sqrt{M_B M_{B_0}} F_0^{B_0 \pi},$$

and the sum over B^* polarizations yields

$$\Pi(u, v) = \frac{(M_B^2 - M_\pi^2 - u)u}{4M_{B^*}^2} + M_\pi^2 - \frac{v}{2}. \quad (27)$$

Note that $A_0^{B^* \pi} \equiv A_0^{B^* \rightarrow \pi}(M_\pi^2)$ and $F_0^{B_0 \pi} \equiv F_0^{B_0 \rightarrow \pi}(M_\pi^2)$. Our expressions for A^{+-0} in Eqs. (23) and (24) disagree with those in Ref. [9] in that the factors of $1/\sqrt{2}$ are missing in their formulas, and that the minus sign in the middle of the big brackets in Eq. (24) is opposite to theirs. However, our expressions for A^{--0} in Eqs. (23) and (24) agree with theirs.

IV. NUMERICAL RESULTS AND DISCUSSION

We begin by listing the parameters that we use; we conform with the parameter choices of Refs. [9,11], in order to realize a crisp comparison with their results. In specific, the Wilson coefficients we use are

$$C_1 = 1.100, \quad C_2 = -0.226, \quad C_3 = 0.012, \quad (28)$$

$$C_4 = -0.029, \quad C_5 = 0.009, \quad C_6 = -0.033.$$

For the CKM factors, we adopt the Wolfenstein parametrization [28], retaining terms of $\mathcal{O}(\lambda^3)$ in the real part and of $\mathcal{O}(\lambda^5)$ in the imaginary part, to wit,

$$V_{ud} = 1 - \lambda^2/2, \quad V_{ub} = A \lambda^3 [\rho - i \eta (1 - \lambda^2/2)], \quad (29)$$

$$V_{td} = A \lambda^3 (1 - \rho - i \eta), \quad V_{tb} = 1,$$

and using

$$\lambda = 0.2196, \quad \rho = 0.05, \quad \eta = 0.36, \quad A = 0.806. \quad (30)$$

For decay constants, light meson masses, and resonance parameters, we have

$$f_\pi / \sqrt{2} = 92.4 \text{ MeV}, \quad M_\pi = 139.57 \text{ MeV},$$

$$f_\rho = 0.15 \text{ GeV}^2, \quad M_\rho = 769.3 \text{ MeV},$$

$$\Gamma_\rho = 150 \text{ MeV}, \quad g_\rho = 5.8, \quad (31)$$

$$M_\sigma = 478 \text{ MeV}, \quad \Gamma_\sigma = 324 \text{ MeV},$$

$$g_{\sigma \pi \pi} = 2.52 \text{ GeV}.$$

The decay constants f_π and f_ρ are associated with π^\pm and ρ^\pm decay, respectively. We neglect isospin-violating effects throughout, so that $M_{\pi^\pm} = M_{\pi^0} = M_\pi$, $M_{\rho^\pm} = M_{\rho^0} = M_\rho$, as well as $M_{\bar{B}^0} = M_{B^-} = M_B$. Moreover, $\hat{m} = 6 \text{ MeV}$. The B^* and B are degenerate in the heavy-quark limit, so that we neglect their mass difference as well. We also neglect the lifetime difference between the \bar{B}^0 and B^- , setting $\tau_{\bar{B}^0} = \tau_{B^-} = \tau_B$. For the B and related mesons, we have

TABLE I. Effective branching ratios for $B \rightarrow \rho\pi$ decays, as per Eq. (35), with $\delta=0.3$ GeV. Breit-Wigner form factors are used throughout, noting Eqs. (9) and (16) for the ρ and σ contributions, respectively. All branching ratios are reported in units of 10^{-6} .

Decay mode	ρ	σ	B^*	B_0	$\rho+B^*$	$\rho+B^*+B_0$	$\rho+\sigma$	$\rho+\sigma+B^*$	$\rho+\sigma+B^*+B_0$
$\bar{B}^0 \rightarrow \rho^- \pi^+$	16.0	0.0003	0.54	0.009	16.5	16.3	16.0	16.4	16.3
$\bar{B}^0 \rightarrow \rho^+ \pi^-$	4.76	0.0003	0.13	0.020	4.98	4.98	4.78	5.00	5.00
$\bar{B}^0 \rightarrow \rho^0 \pi^0$	0.91	0.045	0.39	0.016	1.43	1.29	0.93	1.59	1.43
$B^- \rightarrow \rho^0 \pi^-$	4.10	5.18	2.71	0.107	7.42	8.45	8.83	7.67	7.92
\mathcal{R}	5.1				2.9	2.5	2.3	2.8	2.7

$$M_B = 5.279 \text{ GeV}, \quad \tau_B = 1.6 \times 10^{-12} \text{ s}, \quad m_b = 4.6 \text{ GeV},$$

$$\Gamma_{B^*} = 0.2 \text{ keV}, \quad M_{B_0} = 5.697 \text{ GeV}, \quad \Gamma_{B_0} = 0.36 \text{ GeV}, \quad (32)$$

and use

$$g = 0.6, \quad h = -0.7. \quad (33)$$

The heavy-to-light transition form factors are given by

$$A_0^{B\rho} = 0.29, \quad F_0^{B\rho} = 0.37, \quad F_1^{B\rho} = 0.37, \quad F_0^{B\sigma} = 0.46, \quad (34)$$

$$A_0^{B^*\pi} = 0.16, \quad F_0^{B^*\pi} = -0.19.$$

Finally, for the vector and scalar form-factors, $\Gamma_{\rho\pi\pi}(s)$ and $\Gamma_{\sigma\pi\pi}(s)$, respectively, we follow the treatment of Ref. [11]. The $F_\rho(s)$ parametrization we adopt was fit to $e^+e^- \rightarrow \pi^+\pi^-$ data in the elastic region [13], $2M_\pi \leq \sqrt{s} \leq M_\pi + M_\omega$, only, so that for larger values of s we use a Breit-Wigner form, matched to the value of $\Gamma_{\rho\pi\pi}(s)$ at $\sqrt{s} = M_\pi + M_\omega$. That is, for $\sqrt{s} \geq 923$ MeV we employ $\Gamma_{\rho\pi\pi}(s) = [c_r(M_\rho^2 - s) + ic_i\Gamma_\rho M_\rho]g_\rho / [(M_\rho^2 - s)^2 + \Gamma_\rho^2 M_\rho^2]$, with $c_r \approx 0.929$ and $c_i \approx 1.29$. For the scalar form-factor, we employ the $\Gamma_1^{\sigma\pi\pi}(s)$ derived in Ref. [18], which is valid for $\sqrt{s} \leq 1.2$ GeV. The normalization of Eq. (17) implies that $\chi = 20.0 \text{ GeV}^{-1}$. For $\sqrt{s} > 1.2$ GeV, we match to the asymptotic form of $\Gamma_{\sigma\pi\pi}(s)$ [22], as detailed in Ref. [11].

To obtain branching ratios for $B \rightarrow 3\pi$ decay in the ρ -mass region, we integrate over the region of phase space satisfying the requirement that two of the three pions reconstruct the ρ mass within an interval of 2δ , as was done in Refs. [9,11]. This amounts in each case to calculating the effective width

$$\Gamma_{\text{eff}}(B \rightarrow \rho(p_1 + p_2)\pi(p_3)) = \Gamma(B \rightarrow \pi(p_1)\pi(p_2)\pi(p_3))|_{(M_\rho - \delta)^2 \leq s_{12} \leq (M_\rho + \delta)^2}. \quad (35)$$

We choose $\delta=0.3$ GeV, following earlier work [9,11].

For crisp comparison with Ref. [9], we begin by computing the effective branching ratios arising from the use of Breit-Wigner forms, as in Eqs. (9) and (16) for the ρ and σ , respectively, throughout. The various contributions, reflective of the enumerated terms in Eq. (8), are reported in Table I. There are differences between our results for the ρ , $\rho+B^*$, and $\rho+B^*+B_0$ contributions and the corresponding ones in Ref. [9]. The differences are, however, not large and arise in part from missing factors in the formulas for the B^* and B_0 amplitudes, which we delineated in the last section. In contrast, as pointed out in Ref. [11], the σ effect on the B^- decay is much bigger than that found in Ref. [10], because our σ amplitude is larger than theirs by a factor of $\sqrt{2}$. This is evident in the $\rho+\sigma$ and $\rho+\sigma+B^*$ columns. Our results agree with those in Ref. [11], to the extent that they are applicable; we note that Ref. [11] neglects penguin contributions altogether and deals exclusively with the ρ and σ contributions. The last column of Table I contains the sum of all the contributions, $\rho+\sigma+B^*+B_0$. Overall, it is apparent that the effect of the B_0 is smaller than that of the other contributions, although it is not negligible. Finally, in the last row, we collect the ratios of branching ratios \mathcal{R} defined in Eq. (1). These results show that the inclusion of the σ and B^* , either individually or together, makes the estimated value of \mathcal{R} consistent with the empirical one, given its large error.

We now proceed to compute the effective branching ratios with the ρ and σ form-factors, Eqs. (10) and (14), which we advocate. These results are presented in Table II. The results

TABLE II. Effective branching ratios for $B \rightarrow \rho\pi$ decays, as per Eq. (35), with $\delta=0.3$ GeV. We adopt the ρ and σ form factors, Eqs. (10) and (14), respectively, which we have advocated. All branching ratios are reported in units of 10^{-6} .

Decay mode	ρ	σ	B^*	B_0	$\rho+B^*$	$\rho+B^*+B_0$	$\rho+\sigma$	$\rho+\sigma+B^*$	$\rho+\sigma+B^*+B_0$
$\bar{B}^0 \rightarrow \rho^- \pi^+$	16.0	0.001	0.54	0.009	16.6	16.4	15.9	16.5	16.3
$\bar{B}^0 \rightarrow \rho^+ \pi^-$	4.76	0.001	0.13	0.020	4.90	4.93	4.80	4.94	4.98
$\bar{B}^0 \rightarrow \rho^0 \pi^0$	0.86	0.065	0.39	0.016	1.35	1.21	0.91	1.47	1.33
$B^- \rightarrow \rho^0 \pi^-$	4.06	7.66	2.71	0.107	7.20	8.25	11.1	11.9	12.7
\mathcal{R}	5.1				3.0	2.6	1.9	1.8	1.7




FIG. 2. The B^* and B_0 contributions to $\bar{B}^0 \rightarrow \pi^+(p_+) \pi^-(p_-) \pi^0(p_0)$ decay, specifically, $|A_{B^*}^{+-0} + A_{B_0}^{+-0}|^2$ (in dimensionless units) as a function of its arguments s_{+0} and s_{-0} , both in units of GeV^2 .

without the σ contributions change little, as the vector form-factor is not terribly different from its BW counterpart [11]. In the presence of the σ , this similarity persists for the \bar{B}^0 decays, but, in contrast, the B^- branching ratios are significantly increased compared to the corresponding ones in Table I. This effect also tends to diminish the relative impact of the B^* and B_0 contributions on the $\rho^0 \pi^-$ mode, though the heavy mesons persist in making a substantial impact on the effective branching ratio for the $\rho^0 \pi^0$ mode.

Were the heavy-meson contributions to the $\rho^0 \pi^0$ mode seen in Table II as large as we have estimated, the impact on the Dalitz-plot analysis to extract α from $\bar{B}^0(B^0) \rightarrow \pi^+ \pi^- \pi^0$ decays would be significant [9]. Since the B^* and B_0 masses lie outside the phase-space region of $B \rightarrow 3\pi$, their effects behave as part of the nonresonant background, but are not uniform and obviously interfere with other contributions. The manner in which the contributions are distributed throughout the Dalitz plot is shown for $\bar{B}^0 \rightarrow \pi^+ \pi^- \pi^0$ decay in Fig. 2; the heavy-meson contributions preferentially populate the edges of the Dalitz plot, in which the ρ contributions lie as well. In $B^- \rightarrow \pi^+ \pi^- \pi^-$ decay, the distribution of the heavy-meson contributions is somewhat more uniform, as illustrated in Fig. 3.

We now proceed to consider the reliability of the estimates we have effected. Let us first note that the parameters g and h of the strong heavy-meson couplings in Eqs. (20), (21) assume the values given in Eq. (33)—these reflect the upper limits of their estimated ranges [9,24].³ Thus, the results we find with these parameters can be regarded as extremal estimates (although variations in other numerical inputs, such as the form factors, do introduce further uncertainties). Choosing central values of g and h in their estimated ranges decreases the heavy-meson effects by up to

³We also note, however, that the g value in Eq. (33) is, by virtue of heavy-quark symmetry, favored by the recent measurement of the $D^* \rightarrow D\pi$ width [29], which yields $g = 0.59 \pm 0.01 \pm 0.07$.




FIG. 3. The B^* and B_0 contributions to $B^- \rightarrow \pi^-(p_1) \pi^-(p_2) \pi^+(p_+)$ decay, specifically, $|A_{B^*}^{--0} + A_{B_0}^{--0}|^2$ (in dimensionless units) as a function of its arguments s_{1+} and s_{2+} , both in units of GeV^2 .

some 50% [9], as explicitly shown in Table III.

Moreover, the relative signs chosen for the ρ , σ , and heavy-meson contributions will impact the numerical values of the effective branching ratios. As noted by Ref. [9], the relative sign of the B^* and B_0 contributions is fixed in the heavy-quark and chiral limits. The relative signs of the heavy-meson, ρ , and σ contributions, however, are less clear. We define the $\rho \rightarrow \pi\pi$ coupling as per Eq. (13), after Refs. [9,11], though we note that a chiral Lagrangian analysis suggests that the relations of Eq. (13) should possess an additional overall sign. With this modification, the branching ratios for the $\rho + B^* + B_0$ combination in Table II typically become smaller by no more than 15%. However, the $\rho + \sigma$ results in $\bar{B}^0 \rightarrow \rho^0 \pi^0$ and $B^- \rightarrow \rho^0 \pi^-$ become some 3% and 10% larger, respectively. The impact on the $\rho + \sigma + B^* + B_0$ results is mixed, leading to a suppression of about 10% in the $\rho^0 \pi^0$ mode and an enhancement of 2% in the $\rho^0 \pi^-$ mode.

Kinematical cuts can mitigate the impact of the heavy-meson and σ contributions. Since the $\rho^\pm \pi^\mp$ modes are little affected by these notions, we evaluate only the $\rho^0 \pi^0$ and $\rho^0 \pi^-$ modes. We try two different sets of kinematical cuts. For the first one, we set $\delta = 0.15 \text{ GeV} = \Gamma_\rho$ and report our results in Table IV. The relative suppression of the heavy-meson and σ contributions is quite modest, if it exists at all. For the second set, we impose not only a δ cut but also a cut on $\cos \theta$, where θ is the helicity angle, defined as the angle between the direction of one member of a pion pair from ρ decay and the direction of the parent B -meson evaluated in the pair's rest-frame. Since the ρ contribution has a $\cos^2 \theta$ distribution in $B \rightarrow \pi^+ \pi^- \pi^0$ decay [4], larger values of $|\cos \theta|$ enhance the ρ contribution. Interference effects in the $B^- \rightarrow \pi^+ \pi^- \pi^-$ channel will make this cut less effective. We set $\delta = 0.3 \text{ GeV}$ and $|\cos \theta| > 0.4$, and collect the results in Table V. Comparing to Table II, the θ cut is seen to decrease the relative size of the σ background, as discussed in Ref. [11]. The helicity-angle cut only modestly reduces the σ contribution in $B^- \rightarrow \rho^0 \pi^-$ decay; however, an assumption of ρ

TABLE III. Effective branching ratios for $B \rightarrow \rho \pi$ decays, as in Table II, except that $g=0.40$ and $h=-0.54$ have been used.

Decay mode	ρ	B^*	B_0	$\rho+B^*$	$\rho+B^*+B_0$	$\rho+\sigma$	$\rho+\sigma+B^*$	$\rho+\sigma+B^*+B_0$
$\bar{B}^0 \rightarrow \rho^- \pi^+$	16.0	0.24	0.005	16.3	16.1	15.9	16.2	16.1
$\bar{B}^0 \rightarrow \rho^+ \pi^-$	4.76	0.06	0.012	4.82	4.87	4.80	4.86	4.91
$\bar{B}^0 \rightarrow \rho^0 \pi^0$	0.86	0.17	0.009	1.10	1.03	0.91	1.20	1.13
$B^- \rightarrow \rho^0 \pi^-$	4.06	1.20	0.064	5.55	6.12	11.1	11.0	11.4
\mathcal{R}	5.1			3.8	3.4	1.9	1.9	1.8

dominance is only needed when one employs an isospin analysis to extract $\sin(2\alpha)$ from $B^0(\bar{B}^0) \rightarrow \pi^+ \pi^- \pi^0$ decay, as detailed in Ref. [4]. For the $\rho^0 \pi^0$ mode, the chosen cut does significantly reduce an already small contribution. Were the σ contribution to the $\rho^0 \pi^0$ mode much larger than we estimate, then a full partial-wave analysis to separate the s - and p -wave contributions could be both practicable and necessary.

Finally, we must discuss a tacit assumption we have made in the estimation of the B^* and B_0 contributions, which is made in Ref. [9] as well. That is, in realizing the diagrams of Fig. 1(b), we have treated the strong $(B^*, B_0)B\pi$ and weak $(B^*, B_0) \rightarrow \pi\pi$ vertices as if the B^* and B_0 mesons were on their mass shell. This assumption is compatible with the assumed use of the combined heavy-quark and chiral limits in the treatment of the strong $(B^*, B_0)B\pi$ vertices. However, neither assumption is appropriate for $B \rightarrow \rho\pi$ decay. That is, for the $B \rightarrow 3\pi$ decays of interest, we require that two of the three pions have an invariant mass \sqrt{s} comparable to that of the ρ meson. This implies that in most of the relevant phase-space region the mediating heavy-mesons carry s values much smaller than their squared masses—they are highly off-mass-shell. Moreover, the bachelor π is never soft in this kinematical region. Thus the combined heavy-quark and chiral limits are used beyond their range of validity. These effects modify the vertices we have assumed in Eqs. (20), (21) and Eq. (7). Unfortunately, the needed off-shell extrapolations cannot be done reliably, although we would generically expect this effect to suppress the numerical importance of the B^* and B_0 contributions. For example, the form factors of Eqs. (7) now depend on both q^2 and k^2 ; the vertices are only “half” off-shell, so that p^2 does not enter, as the final-state π is on its mass shell. Moreover, additional form factors appear. To illustrate, we note that the general parametrization

$$\begin{aligned}
& \langle \pi^+(p) | \bar{u} \gamma^\mu L b | \bar{B}_0^0(k) \rangle \\
&= -i F_0^{B_0 \rightarrow \pi}(k^2, p^2, q^2) \frac{(M_{B_0}^2 - M_\pi^2) q^\mu}{q^2} \\
&\quad - i F_1^{B_0 \rightarrow \pi}(k^2, p^2, q^2) \left[p^\mu + k^\mu - (M_{B_0}^2 - M_\pi^2) \frac{q^\mu}{q^2} \right]
\end{aligned} \tag{36}$$

predicated by an assumption of Lorentz invariance yields

$$\begin{aligned}
& q^\mu \langle \pi^+(p) | \bar{u} \gamma^\mu L b | \bar{B}_0^0(k) \rangle \\
&= -i (M_{B_0}^2 - M_\pi^2) F_0^{B_0 \rightarrow \pi}(q^2, k^2) \\
&\quad + i (M_{B_0}^2 - k^2) F_1^{B_0 \rightarrow \pi}(q^2, k^2)
\end{aligned} \tag{37}$$

for the half-off-shell matrix element of interest. The matrix element is a linear combination of signed, uncertain contributions, so that its sign is ultimately unclear. Similar considerations apply to the $B^* \rightarrow \pi$ matrix element, as well as to the strong vertices of Eqs. (20), (21). In the treatment of Ref. [30], an off-shell extrapolation of Eq. (20), in the kinematic region of interest, is effected through the replacement $\sqrt{M_B M_{B^*}} \rightarrow \sqrt{M_B} \sqrt{s}$. To assess the impact of these considerations on the numerical results we have reported, we shall adopt a similarly *ad hoc* prescription. Thus, we perform the replacement

$$M_{B^*}^{3/2} \rightarrow s^{3/4} \tag{38}$$

TABLE IV. Effective branching ratios for $B \rightarrow \rho\pi$ decays, as in Table II, except that $\delta=0.15$ GeV has been used.

Decay mode	ρ	σ	B^*	$\rho+B^*$	$\rho+B^*+B_0$	$\rho+\sigma$	$\rho+\sigma+B^*$	$\rho+\sigma+B^*+B_0$
$\bar{B}^0 \rightarrow \rho^0 \pi^0$	0.33	0.029	0.20	0.55	0.49	0.36	0.59	0.53
$B^- \rightarrow \rho^0 \pi^-$	3.36	3.46	1.38	4.82	5.39	6.46	7.56	8.17

TABLE V. Effective branching ratios for $B \rightarrow \rho\pi$ decays, as in Table II, but with the additional kinematic cut $|\cos \theta| > 0.4$ as explained in the text.

Decay mode	ρ	σ	B^*	$\rho+B^*$	$\rho+B^*+B_0$	$\rho+\sigma$	$\rho+\sigma+B^*$	$\rho+\sigma+B^*+B_0$
$\bar{B}^0 \rightarrow \rho^0 \pi^0$	0.84	0.039	0.27	1.22	1.11	0.87	1.29	1.18
$B^- \rightarrow \rho^0 \pi^-$	3.81	4.79	1.71	5.92	6.58	7.97	8.63	9.13

in the numerator of the B^* amplitudes in Eq. (23), so that the ‘‘off-shellness’’ of both the strong and weak vertices is taken into account. We neglect the B_0 in this simple numerical estimate, as its effect was rather small to start with. We calculate the corresponding branching ratios and collect the results in Table VI. Our simple prescription leads to a dramatic reduction of the B^* contributions, as a comparison with Table II makes clear. Note that the computed values of \mathcal{R} are still consistent with the empirical ones, as a reduction in \mathcal{R} is still realized through the σ contributions. Although we cannot draw firm conclusions from this simple exercise, it serves to illustrate that neglecting the off-shell nature of the heavy-meson vertices in the kinematic region of interest could easily lead to a considerable overestimate of their effects.

V. CONCLUSIONS

We have examined resonant and nonresonant backgrounds to $B \rightarrow \rho\pi \rightarrow 3\pi$ decays which can potentially impact the extraction of α from a Dalitz plot analysis of $B \rightarrow \pi^+ \pi^- \pi^0$ decays [4], as well as the value of the ratio of branching ratios we term \mathcal{R} , as defined in Eq. (1). In particular, we have evaluated the effects of nonresonant contributions mediated by the heavy mesons B^* and B_0 , as well as the contributions from the light σ resonance via $B \rightarrow \sigma\pi \rightarrow 3\pi$ decay, in the ρ -mass region. In this, our analysis parallels that of Refs. [9,10], though it differs fundamentally in two points. Firstly, we use the vector and scalar form-factors of Ref. [11], which are consistent with low-energy theoretical constraints and thus are suitable for the description of broad resonant structures such as the ρ and the σ . The scalar form factor, in particular, is quite different from the Breit-Wigner form adopted in other analyses [10,23] and leads to differing results [11]. Secondly, in the kinematics of interest, the B^* and B_0 are highly off-mass-shell, impacting the strong and weak vertices which mediate the $B \rightarrow (B^*, B_0)\pi$

$\rightarrow \pi\pi\pi$ decay. We find that these effects can reduce the heavy-meson contributions substantially.

Our numerical results show, were we to neglect the off-shell effects we have mentioned, that the $B \rightarrow \rho^\pm \pi^\mp$ decay modes are little affected by the σ and heavy-meson backgrounds, whereas the $\rho^0 \pi^0$ mode receives large contributions from the latter. In contrast, the $B^- \rightarrow \pi^+ \pi^- \pi^-$ decay mode contains large contributions from both the σ and B^* , though the σ contributions numerically dominate. Effecting a simple model of off-shell effects, we find that the B^* effects are substantially reduced. The off-shell extrapolation of interest cannot be effected with certainty; nevertheless, our estimates indicate that the neglect of this effect may lead to a substantial overestimate of the B^* contributions in $B \rightarrow 3\pi$ decay. The role of the σ in lowering the theoretical value of \mathcal{R} and yielding a favorable comparison with experiment persists despite these considerations.

Note added. Since the submission of this paper for publication, a report by the BABAR Collaboration has appeared [31], giving the experimental bound $\mathcal{B}(B^\pm \rightarrow \pi^+ \pi^- \pi^\pm) < 15 \times 10^{-6}$ at 90% C.L. This can be used to constrain the contribution of the B^* - and B_0 -pole diagrams. Using the g and h values as in Table II, we find $\mathcal{B}(B^- \rightarrow \pi^+ \pi^- \pi^-) = 24.8 \times 10^{-6}$ for the combined $\rho + \sigma + B^* + B_0$ contribution, where we have integrated over all the allowed phase-space. Were we to use the intermediate values of g and h given in Table III, though such a g is not favored by data [29], we would obtain $\mathcal{B}(B^- \rightarrow \pi^+ \pi^- \pi^-) = 18.7 \times 10^{-6}$. If we use our off-shell extrapolation (neglecting the small B_0 contribution) and the parameters of Table II, we find $\mathcal{B}(B^- \rightarrow \pi^+ \pi^- \pi^-) = 15.4 \times 10^{-6}$. This comparison supports our assertion: the treatment of the B^* vertices in Ref. [9] tends to yield an overestimate of their contribution to $B \rightarrow 3\pi$ decay. On a related note, the failure to confront the empirical bound on $\mathcal{B}(B^- \rightarrow K^+ K^- \pi^-)$ decay has been described in recent work by Cheng and Yang [32].

TABLE VI. Effective branching ratios for $B \rightarrow \rho\pi$ decays, as in Table II, except that the off-shellness of the B^* meson is included as explained in the text.

Decay mode	ρ	σ	B^*	$\rho+B^*$	$\rho+\sigma$	$\rho+\sigma+B^*$
$\bar{B}^0 \rightarrow \rho^- \pi^+$	16.0	0.001	0.03	16.0	15.9	16.0
$\bar{B}^0 \rightarrow \rho^+ \pi^-$	4.76	0.001	0.01	4.85	4.80	4.88
$\bar{B}^0 \rightarrow \rho^0 \pi^0$	0.86	0.065	0.02	0.88	0.91	0.95
$B^- \rightarrow \rho^0 \pi^-$	4.06	7.66	0.25	4.43	11.1	10.7
\mathcal{R}	5.1			4.7	1.9	1.9

ACKNOWLEDGMENTS

We thank Ulf-G. Meißner for helpful discussions and J.A. Oller for the use of his scalar form factor program. The work of J.T. is supported by the U.S. Department of Energy under

contract DE-FG01-00ER45832. S.G. thanks the SLAC Theory Group for hospitality and is supported by the U.S. Department of Energy under contracts DE-FG02-96ER40989 and DE-AC03-76SF00515.

-
- [1] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 091801 (2001); Belle Collaboration, K. Abe *et al.*, *ibid.* **87**, 091802 (2001).
- [2] See, e.g., Y. Nir, hep-ph/0109090; A.J. Buras, hep-ph/0109197.
- [3] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [4] H.J. Lipkin, Y. Nir, H.R. Quinn, and A. Snyder, Phys. Rev. D **44**, 1454 (1991); A.E. Snyder and H.R. Quinn, *ibid.* **48**, 2139 (1993); H.R. Quinn and J.P. Silva, *ibid.* **62**, 054002 (2000).
- [5] CLEO Collaboration, C.P. Jessop *et al.*, Phys. Rev. Lett. **85**, 2881 (2000).
- [6] BABAR Collaboration, B. Aubert *et al.*, hep-ex/0107058.
- [7] CLEO Collaboration, Y. Gao and F. Wurthwein, hep-ex/9904008.
- [8] M. Bauer, B. Stech, and M. Wirbel, Z. Phys. C **34**, 103 (1987).
- [9] A. Deandrea *et al.*, Phys. Rev. D **62**, 036001 (2000); A. Deandrea, hep-ph/0005014.
- [10] A. Deandrea and A.D. Polosa, Phys. Rev. Lett. **86**, 216 (2001).
- [11] S. Gardner and U.-G. Meißner, Phys. Rev. D **65**, 094004 (2002).
- [12] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [13] S. Gardner and H.B. O’Connell, Phys. Rev. D **59**, 076002 (1999).
- [14] S. Gardner and H.B. O’Connell, Phys. Rev. D **57**, 2716 (1998); **62**, 019903(E) (2000).
- [15] BABAR Collaboration, P. F. Harrison and H. R. Quinn, “The BaBar Physics Book,” SLAC-R-0504, <http://www.slac.stanford.edu/pubs/slacreports/slac-r-504.html>
- [16] A. Ali, G. Kramer, and C.D. Lu, Phys. Rev. D **58**, 094009 (1998).
- [17] J.A. Oller and E. Oset, Phys. Rev. D **60**, 074023 (1999).
- [18] U.-G. Meißner and J.A. Oller, Nucl. Phys. **A679**, 671 (2001).
- [19] J.A. Oller and E. Oset, Nucl. Phys. **A620**, 438 (1997); **A652**, 407(E) (1997).
- [20] O. Babelon, J.L. Basdevant, D. Caillerie, and G. Mennessier, Nucl. Phys. **B113**, 445 (1976).
- [21] J. Gasser and H. Leutwyler, Ann. Phys. (N.Y.) **158**, 142 (1984).
- [22] J.F. Donoghue, J. Gasser, and H. Leutwyler, Nucl. Phys. **B343**, 341 (1990).
- [23] E791 Collaboration, E.M. Aitala *et al.*, Phys. Rev. Lett. **86**, 770 (2001).
- [24] R. Casalbuoni *et al.*, Phys. Rep. **281**, 145 (1997), references therein.
- [25] D. Becirevic and A. Le Yaouanc, J. High Energy Phys. **03**, 021 (1999).
- [26] G. Burdman and J.F. Donoghue, Phys. Lett. B **280**, 287 (1992); M.B. Wise, Phys. Rev. D **45**, 2188 (1992); T.-M. Yan *et al.*, *ibid.* **46**, 1148 (1992); **55**, 5851(E) (1992).
- [27] A.F. Falk, Nucl. Phys. **B378**, 79 (1992); A.F. Falk and M.E. Luke, Phys. Lett. B **292**, 119 (1992); U. Kilian, J.G. Korner, and D. Pirjol, *ibid.* **288**, 360 (1992).
- [28] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
- [29] CLEO Collaboration, S. Ahmed *et al.*, Phys. Rev. Lett. **87**, 251801 (2001).
- [30] N.G. Deshpande, G. Eilam, X.-G. He, and J. Trampetic, Phys. Rev. D **52**, 5354 (1995).
- [31] BABAR Collaboration, B. Aubert *et al.*, hep-ex/0206004.
- [32] H.-Y. Cheng and K.-C. Yang, hep-ph/0205133.