

Higgs boson production cross section as a precision observable?

S. Dawson* and S. Heinemeyer†

Physics Department, Brookhaven National Laboratory, Upton, New York 11973

(Received 7 March 2002; published 23 September 2002)

We investigate what can be learned at a linear collider about the sector of electroweak symmetry breaking from a precise measurement of the Higgs boson production cross section through the process $e^+e^- \rightarrow hZ$. We focus on deviations from the standard model arising in its minimal supersymmetric extension. The analysis is performed within two realistic future scenarios, taking into account all prospective experimental errors on supersymmetric particle masses as well as uncertainties from unknown higher order corrections. We find that information on $\tan\beta$ and M_A could be obtained from a cross section measurement with a precision of $0.5 - 1\%$. Alternatively, information could be obtained on the gaugino mass parameters M_2 and μ if they are relatively small, $M_2, \mu \approx 200$ GeV.

DOI: 10.1103/PhysRevD.66.055002

PACS number(s): 14.80.Cp, 12.15.Lk, 14.80.Ly

I. INTRODUCTION

One of the fundamental problems facing particle physics is understanding the nature of electroweak symmetry breaking. If this symmetry breaking is due to a light Higgs boson, then the Higgs boson will certainly be discovered at the Fermilab Tevatron [1] or the CERN Large Hadron Collider (LHC) [2–4]. The remaining challenge will then be to understand whether this new object is the Higgs boson of the standard model (SM) or some more exotic particle. In the SM, the couplings of the Higgs boson to all particles are completely fixed once the mass is known and so the validity of the SM can be confirmed by measuring Higgs boson production and decay rates and eventually the Higgs potential itself [5]. In alternative models, the Higgs couplings can be quite different from the SM values and so can potentially be used to distinguish between models.

A linear collider (LC) with an energy in the range $\sqrt{s} \sim 350 - 500$ GeV has the capability of performing precision measurements of both Higgs boson production and decay rates [6–8], provided that the Higgs boson mass, M_h , lies in the range predicted by electroweak precision observables, $M_h \lesssim 200$ GeV [9,10]. The dominant production mechanism for such a light Higgs boson is $e^+e^- \rightarrow hZ$ [11], with the largest decay channel being $h \rightarrow b\bar{b}$ or $h \rightarrow WW^*$. The measurements must be interpreted in terms of SM expectations or some model of physics beyond the SM. The goal is then to use the experimental data to disentangle the underlying structure of the model. An important question is thus the required experimental precision for production rates and branching ratios in order to distinguish it from the SM and perhaps to measure the parameters of the new theory.

An integrated luminosity of $\mathcal{L} \sim 500 \text{ fb}^{-1}$ and $\sqrt{s} = 350$ GeV is expected to produce measurements of the various Higgs branching ratios with precisions in the $2 - 10\%$ range at an e^+e^- collider [6–8]. The precision will be less at $\sqrt{s} = 500$ GeV, primarily due to the reduced rate [6,12]. The

LHC can measure some, but not all, Higgs branching ratios, with a precision which is typically less than that obtainable at a LC [13].

Supersymmetric (SUSY) theories [14] are widely considered as the theoretically most appealing extension of the SM. They are consistent with the approximate unification of the three gauge coupling constants at the grand unified theory (GUT) scale and provide a way to cancel the quadratic divergences in the Higgs sector, hence stabilizing the huge hierarchy between the GUT and the Fermi scales. Furthermore, in SUSY theories the breaking of the electroweak symmetry is naturally induced at the Fermi scale, and the lightest supersymmetric particle can be neutral, weakly interacting and absolutely stable, providing a natural solution for the dark matter problem. Therefore the implications of the measurements of Higgs boson branching ratios have been extensively studied within the context of the minimal supersymmetric model (MSSM) [15,16]. This model is extremely predictive and so is useful for comparing the experimental reach achievable in various channels.

In this paper we address the question of whether the total $e^+e^- \rightarrow hZ$ cross section, σ_{hZ} , can be used as a precision observable to help determine the structure of the electroweak sector of the MSSM. The measurement of the $e^+e^- \rightarrow hZ$ Higgsstrahlung production cross section is expected to achieve a $2 - 3\%$ accuracy at $\sqrt{s} = 350$ GeV [6]. This assumes $\mathcal{L} = 500 \text{ fb}^{-1}$ and the analysis of the $Z \rightarrow l^+l^-$ events only. From this measurement, some restrictions can be inferred about the parameters of the MSSM, which we investigate here. For the Higgsstrahlung process, the complete next-to-leading order corrections (involving SM and SUSY particles), including all vertex and box corrections have been calculated [17]. More recently also the leading two-loop corrections have been included [18]. Since these corrections are significant, their inclusion is crucial for drawing conclusions about the underlying model.

Our study differs considerably from previous studies of the Higgs branching ratios in that we investigate plausible future scenarios and estimate uncertainties from all relevant sources. We assume that some MSSM particle masses and mixing angles have been determined at the LHC and/or the LC, and vary *all* inputs accordingly within realistic errors,

*Email address: dawson@bnl.gov

†Email address: Sven.Heinemeyer@physik.uni-muenchen.de

instead of fixing all parameters and then varying just one or two. Furthermore, the anticipated theory errors from unknown higher order corrections in the MSSM Higgs sector are taken into account in a consistent manner. We then ask what can be learned about the remaining unknown parameters of the model. These assumptions try to represent *possible* future scenarios and thus give an idea of what might be inferred from a precise σ_{hZ} measurement.

The rest of this paper is organized as follows: In Sec. II we review the necessary MSSM input parameters and existing higher order corrections in the Higgs sector. Our approach to the investigation, with emphasis on the attempt to look into realistic future scenarios, is explained in detail in Sec. III. Section IV contains our analysis and the corresponding results, while conclusions can be found in Sec. V.

II. THE MSSM: BASICS

The Higgs sector of the MSSM consists of two Higgs doublets, \mathcal{H}_1 and \mathcal{H}_2 [19]. After electroweak symmetry breaking, there remain 5 physical Higgs bosons: h , H , A , and H^\pm . In this paper, we will be concerned only with the production of the lightest Higgs boson, h . The Higgs sector is described at tree level by two additional parameters (besides the SM parameters), which are usually chosen to be $\tan\beta$, the ratio of the Higgs vacuum expectation values (VEVs), and M_A , the mass of the pseudoscalar Higgs boson. The mass eigenstates of the neutral scalar Higgs bosons are obtained from the interaction eigenstates ϕ_1 and ϕ_2 by the rotation

$$\begin{pmatrix} H \\ h \end{pmatrix} = \begin{pmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad (1)$$

where at tree level

$$\mathcal{M}_{\tilde{t}}^2 = \begin{pmatrix} M_{\tilde{t}_L}^2 + m_t^2 + \cos 2\beta \left(\frac{1}{2} - \frac{2}{3} s_W^2 \right) M_Z^2 & m_t X_t \\ m_t X_t & M_{\tilde{t}_R}^2 + m_t^2 + \frac{2}{3} \cos 2\beta s_W^2 M_Z^2 \end{pmatrix}, \quad (4)$$

$$\mathcal{M}_{\tilde{b}}^2 = \begin{pmatrix} M_{\tilde{b}_L}^2 + m_b^2 + \cos 2\beta \left(-\frac{1}{2} + \frac{1}{3} s_W^2 \right) M_Z^2 & m_b X_b \\ m_b X_b & M_{\tilde{b}_R}^2 + m_b^2 - \frac{1}{3} \cos 2\beta s_W^2 M_Z^2 \end{pmatrix}, \quad (5)$$

where $s_W^2 = 1 - c_W^2 = 1 - M_W^2/M_Z^2$ and

$$m_t X_t = m_t (A_t - \mu \cot\beta),$$

$$m_b X_b = m_b (A_b - \mu \tan\beta). \quad (6)$$




FIG. 1. Feynman diagram for the lowest order contribution to $e^+e^- \rightarrow hZ$.

$$\tan 2\alpha = \tan 2\beta \left(\frac{M_A^2 + M_Z^2}{M_A^2 - M_Z^2} \right). \quad (2)$$

At tree level, the mass of the lightest Higgs boson is completely fixed in terms of M_Z , M_A and $\tan\beta$.

The process $e^+e^- \rightarrow hZ$ proceeds (at the tree-level) via the Feynman diagram shown in Fig. 1 and is hence sensitive to the ZZh coupling. At the tree level, the ZZh coupling in the MSSM is altered from the SM value:

$$g_{ZZh}^{\text{SUSY}} = g_{ZZh}^{\text{SM}} \sin(\beta - \alpha). \quad (3)$$

For $M_A \gg M_Z$, $\sin(\beta - \alpha) \rightarrow 1$ and the coupling of the lightest MSSM Higgs boson to the Z boson approaches that of the SM. We therefore expect that $\sigma_{hZ}^{\text{SUSY}}$ will be sensitive to small M_A .

There are two important effects which arise when going beyond the tree level. The first is that the Higgs boson mass prediction is significantly increased by radiative corrections, leading to an upper bound at the two-loop level [20–22] of $M_h \lesssim 135$ GeV [21]. The most important corrections are those in the t/\tilde{t} sector [23] and for large $\tan\beta$ also those in the b/\tilde{b} sector. The mass matrices in the basis of the current eigenstates \tilde{t}_L, \tilde{t}_R and \tilde{b}_L, \tilde{b}_R are given by

Here A_t denotes the trilinear Higgs-boson–top-squark coupling, A_b is the Higgs-boson–sbottom coupling, and μ is the Higgs mixing parameter. SU(2) gauge invariance leads to the relation

$$M_{\tilde{t}_L} = M_{\tilde{b}_L}. \quad (7)$$

The two mass matrices (4),(5) are diagonalized by the angles $\theta_{\tilde{t}}$ and $\theta_{\tilde{b}}$, respectively. The physical squark masses are $m_{\tilde{t}_1}$, $m_{\tilde{t}_2}$, $m_{\tilde{b}_1}$ and $m_{\tilde{b}_2}$. Specifying $m_{\tilde{t}_1}$, $m_{\tilde{t}_2}$, and $\theta_{\tilde{t}}$, along with μ and $\tan\beta$ therefore implicitly fixes the tri-linear mixing parameter A_t , and similarly in the b -squark sector. The radiatively corrected value for the lightest MSSM Higgs boson mass, M_h , depends sensitively on the parameters of the top squark mass matrix (4).

The second important effect of going beyond the tree level is that the SUSY particles enter into loop corrections. The complete set of one-loop corrections to the process $e^+e^- \rightarrow hZ$ has been computed in Ref. [17]. In addition, the leading two loop corrections have been included [18]. The effects of including these corrections have been discussed in detail in Ref. [18] and are seen to be large. This applies in particular for the two-loop corrections. Our analysis includes therefore all one-loop SM and SUSY corrections, along with the leading two-loop corrections. From the analysis in Ref. [18] one can infer a theoretical uncertainty due to unknown higher order corrections for the prediction of σ_{hZ} of $\sim 5\%$.

III. CONCEPT OF THE ANALYSIS

The focus here is to determine in the context of SUSY what new information can be obtained from a precision measurement of σ_{hZ} , beyond the direct measurement of the lightest Higgs boson mass. At the time of a σ_{hZ} measurement at the LC, SUSY (if it exists at a low mass scale) will have been discovered at the LHC and possibly confirmed by the LC. Therefore some SUSY parameters will be known with high precision from the LC measurements, while others (e.g. masses beyond the kinematic reach of the LC) will be known with lesser precision from the LHC data. In the Higgs sector it is possible that only the lightest MSSM Higgs boson will have been measured (e.g. for $M_A \gtrsim 300$ and moderate $\tan\beta$ values, $\tan\beta \sim 10$) [2,3,6]. Only for relatively small masses, $M_H, M_A \lesssim \sqrt{s}/2$, will the heavy Higgs bosons be visible at the LC.

In a realistic analysis at the time of the LC the following has to be taken into account: (i) uncertainties of the measured SM parameters, (ii) uncertainties of the measured MSSM parameters, (iii) intrinsic uncertainties on the theoretical prediction of the MSSM Higgs sector parameters (M_h, σ_{hZ}, \dots) from unknown higher order corrections, (iv) bremsstrahlung, (v) beamstrahlung, and (vi) other machine related uncertainties, e.g. due to the luminosity measurement, detector smearing etc.

A full simulation clearly goes beyond the scope of this exploratory analysis. However, we try to give a realistic impression about the information which can be obtained from a σ_{hZ} measurement. To this end we include the following.

(i) We include the relevant SM uncertainties arising from the m_t measurement.

(ii) We take into account *all* uncertainties on the MSSM parameters from their measurement at the LHC [2,3] and/or

the LC [6–8].¹ To study the dependence of the cross section on the parameters, we vary *all* parameters within their expected precisions and include effects of SUSY particles beyond the leading order as described in the previous section.

(iii) We assume a future theoretical uncertainty in the prediction of M_h from the other SUSY input parameters of 0.5 GeV (which affects mainly the connection of the different SUSY parameters to each other). For the theoretical prediction of σ_{hZ} an uncertainty of 1% is assumed from unknown higher order corrections. However, the Higgs boson mass value that will be used in the future will be determined to ± 0.05 GeV (see below) and thus will have a negligible error. [Numerically the uncertainty of σ_{hZ} is taken into account by allowing a variation of the Higgs boson mass as an input parameter in the σ_{hZ} evaluation by ± 0.5 GeV. This (by numerical coincidence) reproduces the “desired” theoretical uncertainty in σ_{hZ} of $\sim 1\%$.]

(iv) We do not include beamstrahlung, bremsstrahlung, or detector effects, which are beyond the scope of this paper. While the latter can only be realized in a full simulation, the former mostly induce a shift in the numerical results, but have a much smaller effect on the errors.

(v) We neglect luminosity errors. Concerning these, it might be helpful not to investigate σ_{hZ} directly, but to consider e.g. σ_{hZ}/σ_{ZZ} , since in this ratio many uncertainties cancel out. However, the idea of this analysis is to show the possible potential of a precise cross section measurement, which can already be obtained from an analysis of σ_{hZ} alone.

Taking into account the relevant uncertainties in the above manner necessarily weakens the potential of a precise σ_{hZ} measurement, see Sec. IV. This approach is contrary to existing analyses [16]. In these previous analyses, all parameters, except for the one under investigation, are fixed. Furthermore, all theoretical uncertainties for the evaluation of the Higgs sector observables are neglected. The potentially measured effect is then attributed solely to the one parameter under investigation, whereas part of the effect could be due to other sources, such as variations in one of the parameters held fixed (within the corresponding experimental errors) or due to the theoretical uncertainties. In this way the sensitivity to the investigated parameters is incorrectly enhanced. Our approach, on the other hand, results in a smaller sensitivity, but constitutes a more realistic scenario for the investigation of LC analyses.

For this analysis we assume that σ_{hZ} is measured at $\sqrt{s} = 350$ GeV with $\mathcal{L} = 500 \text{ fb}^{-1}$.² In all the investigated scenarios, we assume that the Higgs boson mass will have been measured to an experimental accuracy of [6–8]

¹The errors are similar to those used in Ref. [24], where besides the pure experimental resolution also the anticipated theoretical uncertainty entering the extraction of the parameters has been taken into account.

²Possible LC run scenarios have been investigated in Ref. [25]. They usually assume first some high(er)-energy run and afterwards several shorter runs at lower energies, which we summarize here as one run at $\sqrt{s} = 350$ GeV with $\mathcal{L} = 500 \text{ fb}^{-1}$.

$$M_h^{\text{exp}} = 115 \pm 0.05 \text{ GeV}. \quad (8)$$

However, as mentioned above, within the MSSM this experimental error will always be dominated by the theoretical uncertainty on the prediction of M_h due to unknown higher order corrections. While the current uncertainty in the M_h prediction is estimated to be $\sim 3 \text{ GeV}$ [26], we assume for the future uncertainty

$$\delta M_h^{\text{theo}} (\text{future}) = \pm 0.5 \text{ GeV}. \quad (9)$$

Also the dependence of M_h on the top quark mass is very strong, $\delta m_t / \delta M_h \approx 1$. However, m_t will be determined to an accuracy better than $\sim 130 \text{ MeV}$ at a LC [6,27], so that the parametric uncertainty is smaller than the theoretical uncertainty. It is, however, taken into account.

Since the value of M_h in the MSSM is not a free parameter, but depends on the other SUSY parameters, they have to be chosen such that the value of $M_h = 115 \pm 0.5 \text{ GeV}$ emerges. The numerical evaluation of the MSSM Higgs sector (including M_h and σ_{hZ}) is based on the code FeynHiggsXS [28,18].

IV. ANALYSIS AND RESULTS

In order to make progress in understanding the sensitivity of the total cross section to the input parameters, the approach explained above has been applied to two *possible* future scenarios. In both scenarios we make assumptions about what parameters will be measured and what parameters are left free. This choice, since it involves the unknown MSSM parameters and their detectability, is of course subject to personal opinions. However, the scenarios certainly reflect the possible strength of the σ_{hZ} measurement as explained in the previous section.

A. The Higgs sector scenario

In the first scenario we assume that the gaugino and squark masses and mixing angles have been measured at the LHC [2,3] and/or the LC [6–8]. For our analysis, the most important input parameter is the top quark mass and its associated error. Here we assume

$$m_t^{\text{exp}} = 175 \pm 0.1 \text{ GeV}, \quad (10)$$

which is the anticipated precision from a high energy linear collider [27].

In the \tilde{t} sector, we chose

$$m_{\tilde{t}_1} = 500 \pm 2 \text{ GeV}$$

$$m_{\tilde{t}_2} = 700 \pm 10 \text{ GeV}$$

$$\sin \theta_{\tilde{t}} = -.69 \pm 0.014. \quad (11)$$

This precision for $m_{\tilde{t}_1}$ and $\sin \theta_{\tilde{t}}$ could most probably only be realized with an LC measurement at an energy of $\sqrt{s} = 1 \text{ TeV}$. A more conservative choice would be $\delta m_{\tilde{t}_1} = 10 \text{ GeV}$ and an error on $\sin \theta_{\tilde{t}}$ of up to 10%, which can be

achieved at the LHC [2]. In the analysis we will first investigate the implications of a LC precision, but comment also on the LHC precision results as well.

With the above measurements, A_t is given implicitly in terms of $m_{\tilde{t}_1}$, $m_{\tilde{t}_2}$, $\sin \theta_{\tilde{t}}$, μ and $\tan \beta$. We furthermore fix

$$\mu = 200 \pm 1 \text{ GeV}. \quad (12)$$

We assume approximate unification of the trilinear Higgs-sfermion couplings and take

$$A_b = A_t \pm 10\% \quad (13)$$

$$A_l = A_t \pm 1\%. \quad (14)$$

In addition, we assume the relationship between gaugino masses predicted in many unified models. The specific values we take are

$$M_2 = 400 \pm 2 \text{ GeV}$$

$$M_1 = \frac{5}{3} \frac{s_W^2}{c_W^2} M_2 \pm 1 \text{ GeV}$$

$$m_{\tilde{g}} = M_3 = 500 \pm 10 \text{ GeV}. \quad (15)$$

Finally, for the remaining sfermion sector we choose

$$M_{\tilde{b}_R} = M_{\tilde{t}_R} \pm 10\% \quad (16)$$

$$m_{\tilde{e}_1}, m_{\tilde{e}_2} = 200 \pm 2 \text{ GeV}, \quad (17)$$

where the selectron masses enter in the vertex and box corrections. The uncertainties chosen above are consistent with those given in Refs. [2,3,6], see Sec. III. Equations (13) and (16) reflect the assumed future measurement of the scalar bottom sector. However, the b/\tilde{b} sector plays only a minor role here, since (as will be shown below) either μ or $\tan \beta$ (or both) do not reach large values. This, however, is necessary to have large corrections from b/\tilde{b} loops to the MSSM Higgs sector. With the above choices, the only remaining free parameters are M_A and $\tan \beta$, which we assume to be only poorly known in this scenario. Our procedure is to pick a value for M_A and $\tan \beta$ and check that the chosen parameters generate $M_h = 115 \pm 0.5 \text{ GeV}$, which cuts out a slice of the $M_A - \tan \beta$ -plane. For the above set of parameters, we then calculate σ_{hZ} and compare with the value obtained for the SM, $\sigma_{hZ}^{\text{SM}} = 0.1530 \text{ pb}$. The resulting variations of the cross section from the SM value are shown in Fig. 2. Since the measurement of σ_{hZ} is a missing mass experiment, our results are independent of the Higgs boson decay channel.

The different panels of Fig. 2 show the regions where the rate differs from the SM prediction by a specified amount. This includes a theoretical uncertainty in the SM rate which we approximate by varying M_h within the range, $M_h = 115 \pm 0.5 \text{ GeV}$ (as described in Sec. III). The cross section is quite sensitive to $\tan \beta$. A measurement which differs from the SM prediction by 1.4% or less will restrict $\tan \beta < 10$. Concerning the indirect M_A determination, a 1.4% measure-




FIG. 2. The deviation of $\sigma_{hZ}^{\text{SUSY}}$ from $\sigma_{hZ}^{\text{SM}} = 0.1530$ pb is shown in the $M_A - \tan \beta$ -plane for $M_h = 115 \pm 0.5$ GeV at $\sqrt{s} = 350$ GeV with $\mathcal{L} = 500 \text{ fb}^{-1}$. For the \tilde{t} sector we have assumed the LC errors in Eq. (11).

ment would only be sensitive to $M_A \lesssim 200$ GeV. However, a measurement at the 0.5% level, finding a deviation larger than 0.8%, can be realized only for $M_A \lesssim 300$ GeV. A smaller deviation from the SM value can be realized for all M_A values with $M_A \gtrsim 300$ GeV. Thus a weak upper bound might be established; in case of a direct observation (which will be possible for such small M_A values), the cross section measurement can confirm the direct M_A measurement. Interestingly, this could also happen for values where the LHC can see only the lightest MSSM Higgs boson (in the so-called “LHC wedge region”). The currently envisaged accuracy on σ_{hZ} of 2–3 % is unfortunately not sufficient for σ_{hZ} to be used as a such precision determination.

In this scenario it is important to keep the uncertainties of the \tilde{t} sector in mind, which up to now we have assumed to come partially from the LC and partially from the LHC, see Eq. (11). If the more conservative assumption of LHC errors is made, the cut-out region in the $M_A - \tan \beta$ -plane is visibly enlarged. In particular the band is widened to larger $\tan \beta$ values by about 2, depending somewhat on M_A . The obtained results from the cross section measurement for M_A are

affected in a two-fold way. The lower M_A bound is hardly affected at all. The upper M_A bound is weakened by ~ 50 GeV in the relevant M_A region, $M_A \sim 300$ GeV, but without spoiling the possible determination of an upper bound as explained in the previous section.

B. The gaugino scenario

To demonstrate the possible amount of information that σ_{hZ} might deliver on μ and M_2 , in this scenario we make the assumption that M_A and $\tan \beta$ will have been measured,

$$M_A = 250 \pm 10 \text{ GeV}$$

$$\tan \beta = 4 \pm 0.5, \quad (18)$$

but we leave the gaugino mass parameters M_2 and μ as free parameters (the scan stops at an upper bound of 1 TeV). The other MSSM parameters are assumed to have the same values as in Sec. IV A, together with their corresponding uncertainties. As in the previous section, all experimental and theoretical errors are fully taken into account.




FIG. 3. The deviation of $\sigma_{hZ}^{\text{SUSY}}$ from $\sigma_{hZ}^{\text{SM}} = 0.1530$ pb is shown in the M_2 - μ -plane for $M_h = 115 \pm 0.5$ GeV at $\sqrt{s} = 350$ GeV with $\mathcal{L} = 500 \text{ fb}^{-1}$.

Figure 3 shows the dependence of $\sigma_{hZ}^{\text{SUSY}}$ on μ and M_2 . It is obvious that a reasonable sensitivity only appears for $M_2 \approx 200$ GeV or $\mu \approx 200$ GeV, where $\sigma_{hZ}^{\text{SUSY}}$ has a minimum. It is very unlikely that these two parameters, if they possess such a low value, will not have been measured directly, see e.g. [6] and references therein. Thus, in this scenario σ_{hZ} can only offer complementary information which can verify the internal consistency of the MSSM (see Ref. [24] for a detailed discussion on this subject).

V. CONCLUSIONS AND OUTLOOK

We have investigated whether a precise measurement of the Higgs boson production cross section, $\sigma(e^+e^- \rightarrow hZ)$, offers additional information to pin down the unknown parameters of the MSSM. We have chosen two possible future scenarios. We have explained in detail what uncertainties will be present at the time of a σ_{hZ} measurement and how we take them into account. This includes realistic assumptions for all mass parameters together with the expected uncertainties obtainable at the LHC and/or LC. We also took into

account realistic assumptions on the theoretical uncertainties for the predictions in the MSSM Higgs sector.

We find that the total rate needs to be measured to a 0.5 – 1 % accuracy in order to be useful as a precision observable. Then additional information on $\tan \beta$ or M_A (if it is not too high, $M_A \lesssim 500$ GeV) may be obtainable. The dependence of σ_{hZ} on the gaugino parameters μ and M_2 shows a strong enough dependence to be useful only for very low values, $\mu, M_2 \approx 200$ GeV. Hence, in this case σ_{hZ} could only test the internal consistency of the MSSM.

The required precision for σ_{hZ} at the 1% level, as compared to the envisaged 2–3 %, could possibly be achieved by either accumulating a higher integrated luminosity (also at different center of mass energies) and/or by taking other than the leptonic Z decay modes into account.

ACKNOWLEDGMENTS

This work was supported by the Department of Energy under contract DE-AC02-98CH10886.

- [1] Tevatron Higgs Working Group, M. Carena *et al.*, hep-ph/0010338.
- [2] ATLAS Collaboration, CERN/LHCC/99-15 (1999), see atlasinfo.cern.ch/Atlas/GROUPS/PHYSICS/TDR/access.html.
- [3] CMS Collaboration, see cmsinfo.cern.ch/Welcome.html/CMSdocuments/CMSplots/.
- [4] J. Ellis, S. Heinemeyer, K. Olive, and G. Weiglein, Phys. Lett. B **515**, 348 (2001).
- [5] A. Djouadi, W. Kilian, M. Mühlleitner, and P.M. Zerwas, Eur. Phys. J. C **10**, 45 (1999); F. Boudjema and A. Semenov, hep-ph/0201219.
- [6] ECFA/DESY LC Physics Working Group Collaboration, J. Aguilar-Saavedra *et al.*, “TESLA Technical Design Report Part III: Physics at an e^+e^- Linear Collider,” hep-ph/0106315, see tesla.desy.de.
- [7] American LC Working Group Collaboration, T. Abe *et al.*, “Resource Book for Snowmass 2001,” hep-ex/0106056.
- [8] ACFA LC Working Group Collaboration, K. Abe *et al.*, hep-ph/0109166.
- [9] LEP Electroweak Working Group, see lepewwg.web.cern.ch/LEPEWWG/Welcome.html
- [10] U. Baur, R. Clare, J. Erler, S. Heinemeyer, D. Wackeroth, G. Weiglein, and D. Wood, contribution to the P1-WG1 report of the workshop “The Future of Particle Physics,” Snowmass, Colorado, 2001, hep-ph/0111314.
- [11] W. Kilian, M. Krämer, and P.M. Zerwas, Phys. Lett. B **373**, 135 (1996).
- [12] J. Brau, C. Potter, and M. Iwasaki, in *Physics and Experiments with Future Linear e^+e^- Colliders: LCWS 2000*, Proceedings, Batavia, Illinois, 2000, edited by A. Para and H.E. Fisk, AIP Conf. Proc. No. 578 (AIP, Melville, NY, 2001), p. 267.
- [13] D. Zeppenfeld, R. Kinnunen, A. Nikitenko, and E. Richter-Was, Phys. Rev. D **62**, 013009 (2000).
- [14] H.P. Nilles, Phys. Rep. **110**, 1 (1984); H.E. Haber and G.L. Kane, *ibid.* **117**, 75 (1985); R. Barbieri, Riv. Nuovo Cimento **11**, 1 (1988).
- [15] K. Babu and C. Kolda, Phys. Lett. B **451**, 77 (1999); S. Heinemeyer, W. Hollik, and G. Weiglein, Eur. Phys. J. C **16**, 139 (2000); S. Heinemeyer and G. Weiglein, hep-ph/0102117.
- [16] J. Guasch, W. Hollik, and S. Peñaranda, Phys. Lett. B **515**, 367 (2001); M. Carena, H. Haber, H. Logan, and S. Mrenna, Phys. Rev. D **65**, 055005 (2002); A. Curiel, M. Herrero, D. Temes, and J. De Troconiz, *ibid.* **65**, 075006 (2002).
- [17] P.H. Chankowski, S. Pokorski, and J. Rosiek, Nucl. Phys. **B423**, 437 (1994); **B423**, 497 (1994); V. Driesen and W. Hollik, Z. Phys. C **68**, 485 (1995); V. Driesen, W. Hollik, and J. Rosiek, *ibid.* **71**, 259 (1996).
- [18] S. Heinemeyer, W. Hollik, J. Rosiek, and G. Weiglein, Eur. Phys. J. C **19**, 535 (2001); S. Heinemeyer and G. Weiglein, Nucl. Phys. B (Proc. Suppl.) **89**, 210 (2000).
- [19] J. Gunion, H. Haber, G. Kane, and S. Dawson, *The Higgs Hunter’s Guide* (Addison-Wesley, Reading, MA, 1990).
- [20] M. Carena, J. Espinosa, M. Quirós, and C. Wagner, Phys. Lett. B **355**, 209 (1995); M. Carena, M. Quirós, and C. Wagner, Nucl. Phys. **B461**, 407 (1996); H. Haber, R. Hempfling, and A. Hoang, Z. Phys. C **75**, 539 (1997); R. Zhang, Phys. Lett. B **447**, 89 (1999); J. Espinosa and R. Zhang, J. High Energy Phys. **03**, 026 (2000); Nucl. Phys. **B586**, 3 (2000); G. Degrandi, P. Slavich, and F. Zwirner, *ibid.* **B611**, 403 (2001); A. Brignole, G. Degrandi, P. Slavich, and F. Zwirner, *ibid.* **B631**, 195 (2002).
- [21] S. Heinemeyer, W. Hollik, and G. Weiglein, Phys. Rev. D **58**, 091701 (1998); Phys. Lett. B **440**, 296 (1998); Eur. Phys. J. C **9**, 343 (1999).
- [22] M. Carena, H. Haber, S. Heinemeyer, W. Hollik, C. Wagner, and G. Weiglein, Nucl. Phys. **B580**, 29 (2000).
- [23] H. Haber and R. Hempfling, Phys. Rev. Lett. **66**, 1815 (1991); Y. Okada, M. Yamaguchi, and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991); J. Ellis, G. Ridolfi, and F. Zwirner, Phys. Lett. B **257**, 83 (1991); **262**, 477 (1991); R. Barbieri and M. Frigeni, *ibid.* **258**, 395 (1991).
- [24] J. Erler, S. Heinemeyer, W. Hollik, G. Weiglein, and P.M. Zerwas, Phys. Lett. B **486**, 125 (2000).
- [25] J. Barron *et al.*, contribution to the workshop “The Future of Particle Physics,” Snowmass, Colorado, 2001, hep-ph/0201177.
- [26] M. Frank, S. Heinemeyer, W. Hollik, and G. Weiglein, hep-ph/0202166.
- [27] A. Hoang *et al.*, Eur. Phys. J. C **3**, 1 (2000); M. Martinez, talk at the Linear Collider Workshop, Cracow, Poland, 2001, see webnt.physics.ox.ac.uk/lc/ecfadesy_topqcd.htm
- [28] S. Heinemeyer, W. Hollik, and G. Weiglein, Comput. Phys. Commun. **124**, 76 (2000); hep-ph/0002213. The codes are accessible at www.feynhiggs.de.