

Neutrino observatories can characterize cosmic sources and neutrino properties

Gabriela Barenboim*

Theoretical Physics Department, Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, Illinois 60510

Chris Quigg†

*Theoretical Physics Department, Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, Illinois 60510
and Kavli Institute for Theoretical Physics, University of California, Santa Barbara, California 93106*

(Received 26 January 2003; published 29 April 2003)

Neutrino telescopes that measure relative fluxes of ultrahigh-energy ν_e, ν_μ, ν_τ can give information about the location and characteristics of sources, about neutrino mixing, and can test for neutrino instability and for departures from *CPT* invariance in the neutrino sector. We investigate the consequences of neutrino mixing for the neutrino flux arriving at Earth, and consider how terrestrial measurements can characterize distant sources. We contrast mixtures that arise from neutrino oscillations with those signaling neutrino decays. We stress the importance of measuring ν_e, ν_μ, ν_τ fluxes in neutrino observatories.

DOI: 10.1103/PhysRevD.67.073024

PACS number(s): 96.40.Tv, 11.30.Er, 13.35.Hb, 14.60.Pq

I. INTRODUCTION

Neutrino telescopes promise to probe the deepest reaches of stars, galaxies, and exotic structures in the cosmos [1–4]. Unlike charged particles, neutrinos arrive on a direct line from their source, undeflected by magnetic fields. Unlike photons, neutrinos interact weakly, so they can penetrate thick columns of matter [20]. It is plausible that ultrahigh-energy extraterrestrial neutrinos will emerge from the atmospheric-neutrino background at energies between 1 and 10 TeV. Prospecting for extraterrestrial neutrino sources leads the science agenda for neutrino observatories, with characterizing the sources and the processes that operate within them to follow. The fluxes of extraterrestrial neutrinos may, in addition, offer clues to the properties of neutrinos themselves.

The task of developing diagnostics for neutrino sources by measuring relative fluxes of electron, muon, and tau neutrinos is complicated by neutrino oscillations. Prominent among expected sources is the diffuse flux of neutrinos produced in the jets of active galactic nuclei such as the TeV-gamma sources Mkn 421 and 501, which are some 140 Mpc distant from Earth. The vacuum oscillation length, $L_{\text{osc}} = 4\pi E_\nu / |\Delta m^2|$, is short compared with such intergalactic distances, so neutrinos oscillate many times between source and detector. For $|\Delta m^2| = 10^{-5} \text{ eV}^2$, for example, the oscillation length is $L_{\text{osc}} \approx 2.5 \times 10^{-24} \text{ Mpc} \cdot (E_\nu / 1 \text{ eV})$, a fraction of a megaparsec even for 10^{20} -eV neutrinos. The fluxes $\Phi = \{\varphi_e, \varphi_\mu, \varphi_\tau\}$ that arrive at Earth are not identical to the source fluxes $\Phi^0 = \{\varphi_e^0, \varphi_\mu^0, \varphi_\tau^0\}$, and the transfer matrix \mathcal{X} that maps Φ^0 to Φ is not, in general, invertible [21]. Moreover, over their long flight paths, cosmic neutrinos are vulnerable to decay processes that would have gone undetected in terrestrial or solar experiments. Should the neutrino sector exhibit departures from *CPT* invariance, the pattern of neutrino mixing could be considerably richer—and less circum-

scribed by experiment—than conventional wisdom holds.

In this paper, we investigate the consequences of neutrino mixing for the cosmic neutrino flux Φ at Earth, and explore how measurements on Earth can characterize the source flux Φ^0 . We contrast the fluxes that result from neutrino mixing with those that might arise from simple decay scenarios, and we look at the unconventional consequences that might obtain if *CPT* symmetry were violated. We first carry out an idealized analysis, taking our cue from current experiments; then we take into account the uncertainties of existing experimental constraints; finally we ask what we will know after the next round of neutrino oscillation experiments. Our aim is to examine the scientific potential of neutrino observatories in light of current information about neutrino properties, and to project the situation five years hence.

The basic operational goal of neutrino telescopes is the detection of energetic—hence, long-range—muons generated in charged-current interactions $(\nu_\mu, \bar{\nu}_\mu)N \rightarrow (\mu^-, \mu^+) + \text{anything}$. Efficient, well-calibrated detection of $(\nu_e, \bar{\nu}_e)$ interactions is also required to make neutrino observatories incisive tools for the investigation of cosmic sources and neutrino properties. Good detection of $(\nu_\tau, \bar{\nu}_\tau)$ interactions would test the expectation that muon neutrinos and tau neutrinos arrive at Earth in nearly equal numbers. The normalized electron-neutrino flux at Earth emerges as a promising diagnostic for the character of cosmic sources and for non-standard neutrino properties. If neutrinos behave as expected—mixing according to the standard three-generation picture—it should prove possible, over time, to infer something about the flavor mix of neutrinos produced in distant sources. As is by now well known, neutrino oscillations—conventionally understood—map the standard neutrino mixture at the source, $\Phi_{\text{std}}^0 = \{\frac{1}{3}, \frac{2}{3}, 0\}$ into a mixture at Earth that approximates $\{\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\}$. We explore the uncertainties that attach to this expectation in light of recent improvements in the experimental constraints on neutrino-mixing parameters, and we compute the mixture at Earth to be expected for other source mixtures. Neutrino decays over the long path from astrophysical sources could distort the flavor mixture of neu-

*Email address: gabriela@fnal.gov

†Email address: quigg@fnal.gov

trinos arriving at Earth. Current speculations about *CPT* violation in the neutrino sector imply small, and perhaps undetectable, modifications to the conventional oscillation scenario for ultrahigh-energy neutrinos, but would lead to striking consequences for antineutrino decay.

II. THE INFLUENCE OF NEUTRINO OSCILLATIONS

Neutrino emission from active galactic nuclei (AGN) may constitute the dominant diffuse flux at energies above a few TeV, where cosmic sources should emerge from the background of atmospheric neutrinos. With luminosities on the order of $10^{45\pm 3}$ erg/s, AGNs are the most powerful radiation sources in the universe. They are cosmic accelerators powered by the gravitational energy of matter falling in upon a supermassive black hole. Protons accelerated to very high energies within an AGN may interact with ultraviolet photons in the bright jets along the rotation axis or with matter in the accretion disk. The resulting pp or $p\gamma$ collisions yield approximately equal numbers of π^+ , π^0 , π^- that decay into $\mu^+\nu_\mu$, $\gamma\gamma$, or $\mu^-\bar{\nu}_\mu$. The subsequent muon decays yield

additional muon neutrinos and electron neutrinos, so the final count from $\pi^+ + \pi^0 + \pi^-$ is $2\gamma + 2\nu_\mu + 2\bar{\nu}_\mu + 1\nu_e + 1\bar{\nu}_e$, for a normalized neutrino flux

$$\Phi_{\text{std}}^0 = \left\{ \varphi_e^0 = \frac{1}{3}, \varphi_\mu^0 = \frac{2}{3}, \varphi_\tau^0 = 0 \right\} \quad (1)$$

at the source. This standard mixture—equally divided between neutrinos and antineutrinos—is the canonical expectation from most hypothesized sources of ultrahigh-energy neutrinos.

In the standard, three-generation description of neutrino oscillations, the flavor eigenstates ν_α are related to the mass eigenstates ν_i through $\nu_\alpha = \sum_i U_{\beta i} \nu_i$, or

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}. \quad (2)$$

A canonical form for the neutrino mixing matrix is [5]

$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}, \quad (3)$$

where $s_{ij} = \sin \theta_{ij}$, $c_{ij} = \cos \theta_{ij}$ and δ is a *CP*-violating phase. A useful idealized form of the mixing matrix that holds for $\sin \theta_{13} = 0$, and maximal atmospheric mixing, $\sin 2\theta_{23} = 1$, is

$$U_{\text{ideal}} = \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12}/\sqrt{2} & c_{12}/\sqrt{2} & 1/\sqrt{2} \\ s_{12}/\sqrt{2} & -c_{12}/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}. \quad (4)$$

We can express the oscillation (or survival) probabilities in a transfer matrix \mathcal{X} that maps an initial mixture of flavors at the source into the observed mixture at the detector,

$$\begin{pmatrix} \varphi_e \\ \varphi_\mu \\ \varphi_\tau \end{pmatrix} = \mathcal{X} \begin{pmatrix} \varphi_e^0 \\ \varphi_\mu^0 \\ \varphi_\tau^0 \end{pmatrix}. \quad (5)$$

Averaged over many oscillations, \mathcal{X} takes the form

$$\mathcal{X}_{\beta\alpha} = \delta_{\alpha\beta} - 2\Re \sum_{i>j} U_{\alpha i}^* U_{\beta i} U_{\alpha j} U_{\beta j}^* = \sum_j |U_{\alpha j}|^2 |U_{\beta j}|^2, \quad (6)$$

which does not depend on the phase δ . The idealized form that corresponds to Eq. (4) is

$$\mathcal{X}_{\text{ideal}} = \begin{pmatrix} 1-2x & x & x \\ x & \frac{1}{2}(1-x) & \frac{1}{2}(1-x) \\ x & \frac{1}{2}(1-x) & \frac{1}{2}(1-x) \end{pmatrix}, \quad (7)$$

where $x = \sin^2 \theta_{12} \cos^2 \theta_{12}$. Because the second and third rows are identical, the ν_μ and ν_τ fluxes that result from any source mixture Φ^0 are equal: $\varphi_\mu = \varphi_\tau$.

It is noteworthy that $\mathcal{X}_{\text{ideal}}$ maps the standard source mixture Φ_{std}^0 into

$$\Phi_{\text{std}} = \left\{ \varphi_e = \frac{1}{3}, \varphi_\mu = \frac{1}{3}, \varphi_\tau = \frac{1}{3} \right\}, \quad (8)$$

independent of x , i.e., independent of θ_{12} . Because ν_μ and ν_τ are fully mixed, the result is more general: *any source* with $\varphi_e^0 = \frac{1}{3}$ is mapped into Φ_{std} , independent of x . The source mixture $\Phi_Z^0 = \{\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\}$ that arises from Z^0 decay is yet more special because it is unchanged by neutrino oscillations: *any three-generation transfer matrix* maps it into Φ_{std} . For an arbitrary value of the ν_e fraction at the source, $\mathcal{X}_{\text{ideal}}$ leads to $\varphi_e = \varphi_e^0(1-3x) + x$, with $\varphi_\mu = \varphi_\tau = \frac{1}{2}(1-\varphi_e)$. The variation of φ_e with the ν_e source fraction φ_e^0 is shown as a sequence of small black squares (for $\varphi_e^0 = 0, 0.1, \dots, 1$) in Fig. 1 for the value $x = 0.21$, which corresponds to θ_{12}




FIG. 1. (Color online only.) Ternary plots of the neutrino flux Φ at Earth, showing the implications of current (left panel) and future (right panel) knowledge of neutrino mixing. The small black squares indicate the ν_e fractions produced by the idealized transfer matrix $\mathcal{X}_{\text{ideal}}$ as φ_e^0 varies from 0 to 1 in steps of 0.1. A crossed circle marks the standard mixed spectrum at Earth, $\Phi_{\text{std}}^0 = \{\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\}$; for comparison, a red dot marks the standard source spectrum, $\Phi_{\text{std}}^0 = \{\frac{1}{3}, \frac{2}{3}, 0\}$. Colored swaths delimit the fluxes at Earth produced by neutrino oscillations from the source mixtures $\Phi_0^0 = \{0, 1, 0\}$ (pink), Φ_{std}^0 (red), and $\Phi_1^0 = \{1, 0, 0\}$ (orange), using 95% C.L. ranges for the oscillation parameters. Black crosses (\times) show the mixtures at Earth that follow from neutrino decay, assuming normal ($\varphi_e \approx 0.7$) and inverted ($\varphi_e \approx 0$) mass hierarchies. The blue bands show how current and future uncertainties blur the predictions for neutrino decays. The violet tripod indicates how *CPT*-violating oscillations shape the mix of antineutrinos that originate in a standard source mixture, and the violet cross averages that $\bar{\nu}$ mixture with the standard neutrino mixture. The brown squares denote consequences of *CPT* violation for antineutrino decays.

$= 0.57$, the central value in a recent global analysis [6]. The ν_e fraction at Earth ranges from 0.21, for $\varphi_e^0 = 0$, to 0.59, for $\varphi_e^0 = 1$.

The simple analysis based on $\mathcal{X}_{\text{ideal}}$ is useful for orientation, but it is important to explore the range of expectations implied by global fits to neutrino-mixing parameters. The recent KamLAND data [7] on the reactor- $\bar{\nu}_e$ rate and spectral shape, taken together with solar-neutrino data, select the large-mixing-angle (LMA) solution, with a mixing angle bounded at 95% C.L. to lie in the interval $0.49 < \theta_{12} < 0.67$ [6]. Observations of atmospheric neutrinos favor maximal mixing; we take $\pi/4 \times 0.8 < \theta_{23} < \pi/4 \times 1.2$ [8], also at 95% C.L. Experimental evidence favors $\theta_{13} \ll 1$; informed by the nonobservation of oscillations in the CHOOZ and Palo Verde reactor experiments [9,10], we take $0 < \theta_{13} < 0.1$.

With current uncertainties in the oscillation parameters, a standard source spectrum, $\Phi_{\text{std}}^0 = \{\frac{1}{3}, \frac{2}{3}, 0\}$, is mapped by oscillations onto the (red) boomerang near $\Phi_{\text{std}}^0 = \{\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\}$ in the left panel of Fig. 1. Given that $\mathcal{X}_{\text{ideal}}$ maps $\Phi_{\text{std}}^0 \rightarrow \Phi_{\text{std}}^0$ for any value of θ_{12} , it does not come as a great surprise that the target region is of limited extent, and this is a familiar result [11]. The variation of θ_{23} away from $\pi/4$ breaks the identity $\varphi_\mu \equiv \varphi_\tau$ of the idealized analysis.

One goal of neutrino observatories will be to characterize cosmic sources by determining the source mix of neutrino flavors. It is therefore of interest to examine the outcome of different assumptions about the source. We show in the left

panel of Fig. 1 the mixtures at Earth implied by current knowledge of the oscillation parameters for source fluxes $\Phi_0^0 = \{0, 1, 0\}$ (the purple band near $\varphi_e \approx 0.2$) and $\Phi_1^0 = \{1, 0, 0\}$ (the orange band near $\varphi_e \approx 0.6$) [12,22]. For the Φ_{std}^0 and Φ_1^0 source spectra, the uncertainty in θ_{12} is reflected mainly in the variation of φ_e , whereas the uncertainty in θ_{23} is expressed in the variation of $\varphi_\mu / \varphi_\tau$. For the Φ_0^0 case, the influence of the two angles is not so orthogonal. For all the source spectra we consider, the uncertainty in θ_{13} has little effect on the flux at Earth. The extent of the three regions, and the absence of a clean separation between the regions reached from Φ_{std}^0 and Φ_0^0 indicates that characterizing the source flux will be challenging, in view of the current uncertainties of the oscillation parameters.

Over the next five years—roughly the time scale on which large-volume neutrino telescopes will come into operation—we can anticipate improved information on θ_{12} and θ_{23} from KamLAND [13] and the long-baseline accelerator experiments at Soudan [14] and Gran Sasso [15]. We base our projections for the future on the ranges $0.54 < \theta_{12} < 0.63$ and $\pi/4 \times 0.9 < \theta_{23} < \pi/4 \times 1.1$, still with $0 < \theta_{13} < 0.1$. The results are shown in the right panel of Fig. 1. The (purple) target region for the source flux Φ_0^0 shrinks appreciably and separates from the (red) region populated by Φ_{std}^0 , which is now tightly confined around Φ_{std}^0 . The (orange) region mapped from the source flux Φ_1^0 by oscillations shrinks by about a factor of two in the φ_e and $\varphi_\mu - \varphi_\tau$ dimensions.

If CPT invariance is not respected in the neutrino sector [16], the neutrino and antineutrino mixing matrices need not coincide. The KamLAND experiment's confirmation of solar-like oscillations of $\bar{\nu}_e$ restricts the space of CPT -violating mixing matrices, but Barenboim, Borissov, and Lykken [17] have found two illustrative antineutrino solutions that respect all existing constraints:

$$\text{solution 1: } \begin{pmatrix} \theta_{12}=0.6 \\ \theta_{23}=0.5 \\ \theta_{13}=0.08 \end{pmatrix}, \quad \text{solution 2: } \begin{pmatrix} \theta_{12}=0.785 \\ \theta_{23}=0.52 \\ \theta_{13}=0.08 \end{pmatrix}. \quad (9)$$

These two solutions lead to essentially identical transfer matrices [cf. Eq. (6)] that map a standard source mixture of antineutrinos into $\bar{\Phi}_{\text{CPTV}} = \{0.42, 0.36, 0.22\}$, which is plotted as a violet tripod in Fig. 1. Observing $\varphi_e / \bar{\varphi}_e = \frac{4}{5} \neq 1$ would be manifest evidence of CPT violation. Although in principle it might be possible to distinguish the flux of ν_e and $\bar{\nu}_e$ at Earth by observing the resonant formation $\bar{\nu}_e e \rightarrow W^- \rightarrow \text{hadrons}$, the $\nu_e + \bar{\nu}_e$ flux is more likely to be the available observable. In the presence of CPT -violating mixing, the expectation from a standard flux at the source is $\langle \Phi_{\text{CPTV}} \rangle = \frac{1}{2}(\Phi_{\text{std}} + \bar{\Phi}_{\text{CPTV}}) = \{0.375, 0.35, 0.275\}$, which is plotted as a violet cross in Fig. 1. Distinguishing this value from $\Phi_{\text{std}} = \{\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\}$ would be extraordinarily challenging.

III. RECONSTRUCTING THE NEUTRINO MIXTURE AT THE SOURCE

What can observations of the blend Φ of neutrinos arriving at Earth tell us about the source? Inferring the nature of the processes that generate cosmic neutrinos is more complicated than it would be in the absence of neutrino oscillations. Because ν_μ and ν_τ are fully mixed—and thus enter identically in $\mathcal{X}_{\text{ideal}}$ —it is not possible fully to characterize Φ^0 . We can, however, reconstruct the ν_e fraction at the source as

$$\varphi_e^0 = \frac{\varphi_e - x}{1 - 3x}. \quad (10)$$

The reconstructed source flux φ_e^0 is shown in Fig. 2 as a function of the ν_e flux at Earth. The heavy solid line represents the best-fit value for θ_{12} ; the light blue lines and thin solid lines indicate the current and future 95% C.L. bounds on θ_{12} .

A possible strategy for beginning to characterize a source of cosmic neutrinos might proceed by measuring the ν_e / ν_μ ratio and estimating φ_e under the plausible assumption—later to be checked—that $\varphi_\mu = \varphi_\tau$. Let us first note that very large ($\varphi_e \geq 0.65$) or very small ($\varphi_e \leq 0.15$) ν_e fluxes cannot be accommodated in the standard neutrino-oscillation picture. Observation of an extreme ν_e fraction would implicate unconventional physics.

As we have already seen from Fig. 1, constraining the source flux sufficiently to test the nature of the neutrino production process will require rather precise determinations of




FIG. 2. (Color online only.) The source flux φ_e^0 of electron neutrinos reconstructed from the ν_e flux φ_e at Earth, using the ideal transfer matrix $\mathcal{X}_{\text{ideal}}$ of Eq. (7). The heavy solid line refers to our chosen central value, $\theta_{12}=0.57$. The light blue lines refer to the current experimental constraints (at 95% C.L.), and the thin solid lines refer to our projection of future experimental constraints.

the neutrino flux at Earth. Suppose we want to test the idea that the source flux has the standard composition of Eq. (1). With today's uncertainty on θ_{12} , a 30% measurement that locates $\varphi_e = 0.33 \pm 0.10$ implies only that $0 \leq \varphi_e^0 \leq 0.68$. For a measured flux in the neighborhood of $\frac{1}{3}$, the uncertainty in the solar mixing angle is of little consequence: the constraint that arises if we assume the central value of θ_{12} is not markedly better: $0.06 \leq \varphi_e^0 \leq 0.59$. A 10% measurement of the ν_e fraction, $\varphi_e = 0.33 \pm 0.033$, would make possible a rather restrictive constraint on the nature of the source. The central value for θ_{12} leads to $0.26 \leq \varphi_e^0 \leq 0.43$, blurred to $0.22 \leq \varphi_e^0 \leq 0.45$ with current uncertainties.

Measured ν_e fractions that depart significantly from the canonical $\varphi_e = \frac{1}{3}$ would suggest nonstandard neutrino sources. An observed flux $\varphi_e = 0.5 \pm 0.1$ points to a source flux $0.47 \leq \varphi_e^0 \leq 1$, with current uncertainties, whereas $\varphi_e = 0.25 \pm 0.10$ indicates $0 \leq \varphi_e^0 \leq 0.35$.

IV. INFLUENCE OF NEUTRINO DECAYS

Beacom, Bell, and collaborators [18,19] have observed that the decays of unstable neutrinos over cosmic distances can lead to mixtures at Earth that are incompatible with the oscillations of stable neutrinos. The candidate decays are transitions between mass eigenstates by emission of a very light particle, $\nu_i \rightarrow (\nu_j, \bar{\nu}_j) + X$. Dramatic effects occur when the decaying neutrinos disappear, either by decay to invisible products or by decay into active neutrino species so degraded in energy that they contribute negligibly to the total flux at the lower energy. If the lifetimes of the unstable mass eigenstates are short compared with the flight time from source to Earth, decay of the unstable neutrinos will be complete, and

the (unnormalized) flavor ν_α flux at Earth is given by

$$\tilde{\varphi}_\alpha(E_\nu) = \sum_{i=\text{stable}} \sum_{\beta} \varphi_\beta^0(E_\nu) |U_{\beta i}|^2 |U_{\alpha i}|^2, \quad (11)$$

with $\varphi_\alpha = \tilde{\varphi}_\alpha / \sum_{\beta} \tilde{\varphi}_\beta$.

If only the lightest neutrino survives, the flavor mix of neutrinos arriving at Earth is determined by the flavor composition of the lightest mass eigenstate, *independent of the flavor mix at the source*. (The flavor mix at the source of course determines the relative fluxes of the mass eigenstates at the source, and so the rate of surviving neutrinos at Earth.) For a normal mass hierarchy $m_1 < m_2 < m_3$, the ν_α flux at Earth is $\varphi_\alpha = |U_{\alpha 1}|^2$. Accordingly, the neutrino flux at Earth is $\Phi_{\text{normal}} = \{|U_{e1}|^2, |U_{\mu 1}|^2, |U_{\tau 1}|^2\} \approx \{0.70, 0.17, 0.13\}$ for our chosen central values of the mixing angles. If the mass hierarchy is inverted, $m_1 > m_2 > m_3$, the lightest (hence, stable) neutrino is ν_3 , so the flavor mix at Earth is determined by $\varphi_\alpha = |U_{\alpha 3}|^2$. In this case, the neutrino flux at Earth is $\Phi_{\text{inverted}} = \{|U_{e3}|^2, |U_{\mu 3}|^2, |U_{\tau 3}|^2\} \approx \{0.05, 0.5, 0.5\}$. Both Φ_{normal} and Φ_{inverted} , which are indicated by crosses (\times) in Fig. 1, are very different from the standard flux $\Phi_{\text{std}} = \{\varphi_e = \frac{1}{3}, \varphi_\mu = \frac{1}{3}, \varphi_\tau = \frac{1}{3}\}$ produced by the ideal transfer matrix from a standard source. Observing either mixture would represent a departure from conventional expectations.

The fluxes that result from neutrino decays *en route* from the sources to Earth are subject to uncertainties in the neutrino-mixing matrix. The expectations for the two decay scenarios are indicated by the blue regions in Fig. 1, where we indicate the consequences of 95% C.L. ranges of the mixing parameters now and in the future. With current uncertainties, the normal hierarchy populates $0.61 \leq \varphi_e \leq 0.77$, and allows considerable departures from $\varphi_\mu = \varphi_\tau$. The normal-hierarchy decay region based on current knowledge overlaps the flavor mixtures that oscillations produce in a pure- ν_e source, shown in orange. (It is, however, far removed from the standard region that encompasses Φ_{std} .) With the projected tighter constraints on the mixing angles, the range in φ_e swept out by oscillation from a pure- ν_e source or decay from a normal hierarchy shrinks by about a factor of two. Neutrino decay then populates $0.65 \leq \varphi_e \leq 0.74$, and is separated from the oscillations. The degree of separation between the region populated by normal-hierarchy decay and the one populated by mixing from a pure- ν_e source depends on the value of the solar mixing angle θ_{12} . For the seemingly unlikely value $\theta_{12} = \pi/4$, both mechanisms yield $\Phi = \{\frac{1}{2}, \frac{1}{4}, \frac{1}{4}\}$.

With current uncertainties, the inverted hierarchy spans the range $(\varphi_\mu, \varphi_\tau) = (0.34, 0.66)$ to $(0.66, 0.34)$, always with $\varphi_e \approx 0$. With the improvements we project in the knowledge of mixing angles, the range will decrease to $(\varphi_\mu, \varphi_\tau) = (0.42, 0.58)$ to $(0.58, 0.42)$, with $\varphi_e \approx 0$. In both cases, these mixtures are well separated from the mixtures that would result from neutrino oscillations, for any conceivable source at cosmic distances.

Should *CPT* not be a good symmetry for neutrinos, the mass hierarchies and mixing matrices can be different for neutrinos and antineutrinos. Different numbers of stable ν

and $\bar{\nu}$ may reach the Earth, so we do not know how to combine neutrino and antineutrino results to compare with neutrino-telescope measurements. The analysis of neutrino decays does not change if we relax *CPT* invariance, because neutrino mixing is already highly constrained by experiment. The Barenboim-Borissov-Lykken antineutrino mixing parameters [17] [cf. Eq. (9)] provide examples of what can be expected for antineutrino decays if *CPT* is violated. Consider first the normal mass hierarchy, in which ν_2 and ν_3 both decay completely before reaching Earth. The resulting antineutrino fluxes at Earth are $\bar{\Phi}_{\text{CPTV1}} = \{0.68, 0.28, 0.04\}$ and $\bar{\Phi}_{\text{CPTV2}} = \{0.50, 0.42, 0.08\}$, both characterized by unusually large ratios of $\bar{\varphi}_\mu / \bar{\varphi}_\tau$ that place them outside the range of *CPT*-conserving decays and oscillations alike [23]. In the case of an inverted mass hierarchy, both BBL parameter sets predict $\bar{\Phi}_{\text{CPTvinv}} = \{0.01, 0.23, 0.76\}$, an unusually small ratio of $\bar{\varphi}_\mu / \bar{\varphi}_\tau$. Conventional oscillations—or *CPT*-conserving decays—populate the ternary plots of Fig. 1 along the line $\varphi_\mu = \varphi_\tau$. *CPT*-violating decays offer an example of exotic physics with a very specific signature.

Beacom and Bell [19] have shown that observations of solar neutrinos set the most stringent plausible lower bound on the reduced lifetime of a neutrino of mass m as $\tau/m \geq 10^{-4}$ s/eV. This rather modest limit opens the possibility that some neutrinos do not survive the journey from astrophysical sources, with consequences we have just explored. The energies of neutrinos that may be detected in the future from AGNs and other cosmic sources range over several orders of magnitude, whereas the distances to such sources vary over perhaps one order of magnitude. The neutrino energy sets the neutrino lifetime in the laboratory frame; more energetic neutrinos survive over longer flight paths than their lower-energy companions [24]. Under propitious circumstances of reduced lifetime, path length, and neutrino energy, it might be possible to observe the transition from more energetic survivor neutrinos to less energetic decayed neutrinos.

If decay is not complete, the (unnormalized) flavor ν_α flux arriving at Earth from a source at distance L is given by

$$\tilde{\varphi}_\alpha(E_\nu) = \sum_i \sum_{\beta} \varphi_\beta^0(E_\nu) |U_{\beta i}|^2 |U_{\alpha i}|^2 e^{-(L/E_\nu)(m_i/\tau_i)}, \quad (12)$$

with the normalized flux $\varphi_\alpha(E_\nu) = \tilde{\varphi}_\alpha(E_\nu) / \sum_{\beta} \tilde{\varphi}_\beta(E_\nu)$. An idealized case will illustrate the possibilities for observing the onset of neutrino decay and estimating the reduced lifetime. Assume a normal mass hierarchy, $m_1 < m_2 < m_3$, and let $\tau_3/m_3 = \tau_2/m_2 \equiv \tau/m$. For a given path length L , the neutrino energy at which the transition occurs from negligible decays to complete decays is determined by τ/m . We show in the left panel of Fig. 3 the energy evolution of the normalized neutrino fluxes arriving from a standard source; the energy scale is appropriate for the case $\tau/m = 1$ s/eV and $L = 100$ Mpc. The result shown there is actually universal—within our simplifying assumptions—provided that the neutrino energy scale is renormalized as




FIG. 3. (Color online only.) Energy dependence of normalized ν_e , ν_μ , and ν_τ fluxes, for the two-body decay of the two upper mass eigenstates, with the neutrino source at $L=100$ Mpc from Earth and $\tau/m=1$ s/eV. The left panel shows the result for a normal mass hierarchy; the right panel shows the result for an inverted mass hierarchy. With suitable rescaling of the neutrino energy [cf. Eq. (13)], these plots apply for any combination of path length and reduced lifetime.

$$E_\nu = \varepsilon \left(\frac{1 \text{ s/eV}}{\tau/m} \right) \left(\frac{L}{100 \text{ Mpc}} \right). \quad (13)$$

We show in the left panel of Fig. 4 that over about one decade in energy, the flavor mix of neutrinos arriving at

Earth changes from the normal decay flux, Φ_{normal} , to the standard oscillation flux, Φ_{std} . For the case at hand, the transition takes place between 3×10^{15} eV and 3×10^{16} eV. The corresponding results for the inverted mass hierarchy are depicted in the right panels of Figs. 3 and 4, which show a rapid transition from Φ_{inverted} to Φ_{std} .




FIG. 4. (Color online only.) Ternary plots showing the energy dependence of normalized ν_e , ν_μ , and ν_τ fluxes, for the two-body decay of the two upper mass eigenstates, with the neutrino source at $L=100$ Mpc from Earth and $\tau/m=1$ s/eV. The left panel shows the result for a normal mass hierarchy; the right panel shows the result for an inverted mass hierarchy. Each unlabeled step multiplies ε by $10^{1/4}$. With suitable rescaling of the neutrino energy [cf. Eq. (13)], these plots apply for any combination of path length and reduced lifetime.

If we locate the transition from survivors to decays at neutrino energy E^* , then we can estimate the reduced lifetime in terms of the distance to the source as

$$\tau/m \approx 100 \text{ s/eV} \left(\frac{L}{1 \text{ Mpc}} \right) \left(\frac{1 \text{ TeV}}{E^*} \right). \quad (14)$$

In practice, ultrahigh-energy neutrinos are likely to arrive from a multitude of sources at different distances from Earth, so the transition region will be blurred [25]. Nevertheless, it would be rewarding to observe the decay-to-survival transition, and to use Eq. (14) to estimate—even within one or two orders of magnitude—the reduced lifetime. If no evidence appears for a flavor mix characteristic of neutrino decay, then Eq. (14) provides a lower bound on the neutrino lifetime. For that purpose, the advantage falls to large values of L/E^* , and so to the lowest energies at which neutrinos from distant sources can be observed. Observing the standard flux, $\Phi_{\text{std}} = \{\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\}$, which is incompatible with neutrino decay, would strengthen the current bound on τ/m by some seven orders of magnitude, for 10-TeV neutrinos from sources at 100 Mpc [19].

V. ASSESSMENT

The conventional mechanism for ultrahigh-energy neutrino production in astrophysical sources, when modulated by the standard picture of neutrino oscillations, leads to approximately equal fluxes of ν_e , ν_μ , and ν_τ at a terrestrial detector. Observing a strong deviation from this expectation will indicate that we must revise our conception of neutrino sources—or that neutrinos behave in unexpected ways on the long journey from source to Earth. With the more precise understanding of neutrino mixing that we expect will develop over the next five years, it may become possible to characterize the neutrino mixture at the source. On the other hand, detecting equal fluxes of the three neutrino flavors can strengthen existing bounds on neutrino lifetimes quite dra-

matically. If neutrinos do decay between source and Earth, the flavor mixture at Earth will be incompatible with the consequences of the standard source mixture. As constraints improve on the mixing angles θ_{12} and θ_{23} , the flavor mix that results from neutrino decay will separate decisively from what can be generated by any source mix plus oscillations. In the best imaginable case for neutrino decays, the neutrino lifetime might reveal itself in an energy dependence of the flavor mix observed at Earth. If neutrinos decay, the fluxes at Earth are very different for normal and inverted mass hierarchies.

Conventional neutrino oscillations—and conventional decays—lead to nearly equal fluxes of ν_μ and ν_τ , and so populate only a portion of the $\varphi_e - \varphi_\mu - \varphi_\tau$ ternary plot. A neutrino mixture at Earth that is far from the $\varphi_\mu = \varphi_\tau$ line points to unconventional neutrino physics. *CPT* violation offers one example of novel behavior, but its predictions are tested most cleanly by observing differences between neutrinos and antineutrinos, for which the experimental prospects are distinctly limited at ultrahigh energies.

The analysis presented here offers further motivation to develop techniques for identifying interactions of ultrahigh-energy ν_e , ν_μ , and ν_τ —and for measuring the neutrino energies—in cubic-kilometer-scale neutrino observatories. If this can be achieved, prospects for learning about the properties of neutrinos and about the nature of cosmic neutrino sources will be greatly enhanced.

ACKNOWLEDGMENT

We thank Stephen Parke for a helpful observation, and thank John Beacom and Nicole Bell for lively discussions. Fermilab is operated by Universities Research Association Inc. under Contract No. DE-AC02-76CH03000 with the U.S. Department of Energy. One of us (C.Q.) is grateful for the hospitality of the Kavli Institute for Theoretical Physics during the program “Neutrinos: Data, Cosmos, and Planck Scale.” This research was supported in part by the National Science Foundation under Grant No. PHY99-07949.

-
- [1] J.G. Learned and K. Mannheim, *Annu. Rev. Nucl. Part. Sci.* **50**, 679 (2000).
 [2] F. Halzen and D. Hooper, *Rep. Prog. Phys.* **65**, 1025 (2002).
 [3] F. Halzen, “High-Energy Neutrino Astronomy: Science and First Results,” astro-ph/0301143.
 [4] R. Gandhi, C. Quigg, M.H. Reno, and I. Sarcevic, *Astropart. Phys.* **5**, 81 (1996); *Phys. Rev. D* **58**, 093009 (1998).
 [5] P. Fisher, B. Kayser, and K.S. McFarland, *Annu. Rev. Nucl. Part. Sci.* **49**, 481 (1999).
 [6] P.C. de Holanda and A.Yu. Smirnov, “LMA MSW Solution of the Solar Neutrino Problem and First KamLAND Results,” hep-ph/0212270.
 [7] KamLAND Collaboration, K. Eguchi *et al.* *Phys. Rev. Lett.* **90**, 021802 (2003).
 [8] M.C. Gonzalez-Garcia, “Theory of Neutrino Masses and Mixing,” Plenary report at the 2003 International Conference on High Energy Physics, Amsterdam, hep-ph/0210359.
 [9] CHOOZ Collaboration, M. Apollonio *et al.*, *Eur. Phys. J. C* **27**, 331 (2003).
 [10] F. Boehm *et al.*, *Phys. Rev. D* **64**, 112001 (2001).
 [11] J.G. Learned and S. Pakvasa, *Astropart. Phys.* **3**, 267 (1995); H. Athar, M. Jezabek, and O. Yasuda, *Phys. Rev. D* **62**, 103007 (2000). O. Yasuda, “Neutrino oscillations in high energy cosmic neutrino flux,” hep-ph/0005135; L. Bento, P. Keranen, and J. Maalampi, *Phys. Lett. B* **476**, 205 (2000); G.J. Gounaris and G. Moultaqa, “The Flavor Distribution of Cosmic Neutrinos,” hep-ph/0212110.
 [12] J.P. Rachen and P. Mészáros, *Phys. Rev. D* **58**, 123005 (1998).
 [13] V.D. Barger, D. Marfatia, and B.P. Wood, *Phys. Lett. B* **498**, 53 (2001); see also <http://www.awa.tohoku.ac.jp/KamLAND/>
 [14] <http://www.numi.fnal.gov>
 [15] <http://proj-cngs.web.cern.ch/proj-cngs/>
 [16] G. Barenboim, L. Borissov, J. Lykken, and A.Yu. Smirnov, *J. High Energy Phys.* **10**, 001 (2002); G. Barenboim, L. Borissov,

- and J. Lykken, Phys. Lett. B **534**, 106 (2002).
- [17] G. Barenboim, L. Borissov, and J. Lykken, "CPT Violating Neutrinos in the Light of KamLAND," hep-ph/0212116.
- [18] J.F. Beacom, N.F. Bell, D. Hooper, S. Pakvasa, and T.J. Weiler, "Decay of High-Energy Astrophysical Neutrinos," hep-ph/0211305.
- [19] J.F. Beacom and N.F. Bell, Phys. Rev. D **65**, 113009 (2002).
- [20] The interaction length of a 1-TeV neutrino is about 2.5 million kilometers of water, or 250 kton/cm², whereas high-energy photons are blocked by a few hundred g/cm².
- [21] We use the symbols Φ and φ to denote the sum of neutrinos and antineutrinos except when we discuss implications of possible CPT violation, where Φ, φ refer to neutrinos, and $\bar{\Phi}, \bar{\varphi}$ to antineutrinos.
- [22] Muon cooling within the source, as discussed in Ref. [12], would deplete the source flux of ν_e . We have no candidate mechanism to produce an ultrahigh-energy flux like Φ_1^0 . In view of the current puzzlement over the origins of ultrahigh-energy cosmic rays, we think it prudent to keep an open mind about the range of possibilities.
- [23] The barred quantities refer to antineutrinos.
- [24] A similar phenomenon is familiar for cosmic-ray muons.
- [25] Our assumption that $\tau_3/m_3 = \tau_2/m_2 \equiv \tau/m$ is also a special case.