Temperature and entropy of Schwarzschild-de Sitter space-time

S. Shankaranarayanan*

DCTD, University of Azores, 9500 Ponta Delgada, Portugal (Received 23 January 2003; published 24 April 2003)

In the light of recent interest in quantum gravity in de Sitter space, we investigate semiclassical aspects of four-dimensional Schwarzschild-de Sitter space-time using the method of complex paths. The standard semiclassical techniques (such as Bogoliubov coefficients and Euclidean field theory) have been useful to study quantum effects in space-times with single horizons; however, none of these approaches seem to work for Schwarzschild-de Sitter space-time or, in general, for space-times with multiple horizons. We extend the method of complex paths to space-times with multiple horizons and obtain the spectrum of particles produced in these space-times. We show that the temperature of radiation in these space-times is proportional to the effective surface gravity-the inverse harmonic sum of surface gravity of each horizon. For the Schwarzschild-de Sitter space-time, we apply the method of complex paths to three different coordinate systems-spherically symmetric, Painlevé, and Lemaître. We show that the equilibrium temperature in Schwarzschild-de Sitter space-time is the harmonic mean of cosmological and event horizon temperatures. We obtain Bogoliubov coefficients for space-times with multiple horizons by analyzing the mode functions of the quantum fields near the horizons. We propose a new definition of entropy for space-times with multiple horizons, analogous to the entropic definition for space-times with a single horizon. We define entropy for these space-times to be inversely proportional to the square of the effective surface gravity. We show that this definition of entropy for Schwarzschild-de Sitter space-time satisfies the D-bound conjecture.

DOI: 10.1103/PhysRevD.67.084026

PACS number(s): 04.70.Dy, 04.62.+v

I. INTRODUCTION

Over the last three decades, quantum field theory in de Sitter (dS) space has been a subject of growing interest. In the 1970s, the attention was due to the large symmetry group of dS space, which made the field theory in dS space less ambiguous than, for example, in Schwarzschild space-time. In the 1980s, the focus was due to the role it played during inflation-accelerated expansion in the universe's distant past. Recent attention to dS space and asymptotic dS spacetimes is motivated by two aspects. (i) Observations [1] suggest that the universe might be currently asymptotic dS and approach a pure dS space; (ii) the success of the AdS/ conformal field theory (CFT) correspondence [2] has led to the intense study of the quantum gravity of de Sitter space [3]. The focus has been to obtain an analogue of the ADS/ CFT correspondence in dS space [3-5]. For recent attempts on the semiclassical aspects of dS and asymptotic dS spacetime in the light of the dS/CFT correspondence, please see Refs. [6-10]. [The authors in Refs. [6,7] extended the method introduced in Ref. [11] to the Painlevé coordinates of (Schwarzschild) dS space-time.]

Even though there has been an extensive study of the semiclassical aspects of dS space-time (for a recent review, see Ref. [12]) very little has been understood in the case of Schwarzschild-de Sitter (SdS) space-time. (An incomplete list of references, with regard to the semiclassical aspect, is given in Refs. [13–16].) The fundamental difference between SdS and dS (also Schwarzschild) space-times is the existence of multiple horizons. SdS space-time has two—cosmological and event—horizons, while dS (and Schwarzs-

child) space-time has only one horizon.

Various semiclassical approaches or techniques (such as Bogoliubov coefficients, particle detectors, effective action, Euclidean field theory) have been used in the literature to study quantum effects in space-times with a single horizon (like dS and Schwarzschild space-times). All the approaches conclude that the notion of temperature (and entropy) of the space-time is associated with the horizon. In the case of Bogoliubov coefficients, one uses the mode functions to obtain the spectrum of particles, while using Euclidean field theory, one obtains temperature using periodicity arguments. Even though these approaches work well for space-times with a single horizon, none of them work for SdS or, in general, space-times with multiple horizons. A naive extension of these approaches to SdS space-time leads us to the conclusion that the SdS space-time has two different temperatures associated with the two horizons. Using this extension, it has been argued [6,17] that the SdS space will inevitably evolve toward an empty de Sitter space, indicating that SdS spacetime may never be in thermodynamic equilibrium with a single temperature associated with the space-time.

The above argument seems to be in contradiction with the well known case of a Schwarzschild black hole in thermal equilibrium with radiation in a bounded box. In this case, the black hole has a negative specific heat while the radiation has a positive specific heat. The two will be in thermal equilibrium if the box is bounded, in other words compact. On the contrary, if the box is unbounded the black hole evaporates completely. The situation is identical to the case of our interest—black holes in de Sitter space. The de Sitter space is compact with no notion of spatial infinity. In addition, it has a positive specific heat similar to the above mentioned case. The specific heat of de Sitter space is given by [12,14]

^{*}Email address: shanki@notes.uac.pt

$$C_V = \frac{\partial E}{\partial T} = \frac{1}{4\pi T_{dS}^2} = \pi l^2 = S_{dS}, \qquad (1)$$

where S_{dS} (T_{dS}) is the entropy (temperature) of de Sitter space. However, the difference between de Sitter space and the bounding box is that the de Sitter has a (cosmological) horizon while the bounding box, by construction, does not possess a horizon. The similarity of the two systems strongly suggests that we should be able to obtain a temperature for SdS space corresponding to a system in thermal equilibrium.

Given this, one would like to ask the following question: Can one obtain a temperature for SdS space which corresponds to a system in thermal equilibrium using semiclassical techniques? The purpose of this paper is an attempt in this direction. As mentioned in earlier paragraphs, standard quantum field theoretic techniques have not proven useful for space-times with multiple horizons. In this paper, we extend the method of complex paths to space-times with multiple horizons and obtain the spectrum of particles produced in these space-times. (The method of complex paths has proved to be useful in obtaining the temperature associated with a quantum field propagating in a spherically symmetric coordinate space-times with single horizon [18-21].) We show that the temperature of radiation in these space-times is proportional to the effective surface gravity-the inverse harmonic sum of the surface gravity of each horizon. In the case of Schwarzschild-de Sitter space time we apply the method of complex paths to three different coordinate systemsspherically symmetric, Painlevé, and Lemaitre. We show that the equilibrium temperature in Schwarzschild-de Sitter space time is the harmonic mean of the cosmological and event horizon temperatures. We obtain Bogoliubov coefficients for space-times with multiple horizons by analyzing the mode functions of the quantum fields near the horizons.

We propose a new definition of entropy for space-times with multiple horizons analogous to the entropic definition for space-times with a single horizon. We define the entropy for these space-times to be inversely proportional to the square of the effective surface gravity. We show that this definition of entropy for SdS space-time satisfies the *D*-bound conjecture [22].

The paper is organized as follows. In Sec. II A, we discuss the general properties of spherically symmetric space-times. A brief description of SdS geometry is given in Sec. II B. In Sec. III A, we apply the method of complex paths to general spherically symmetric space-times and show that the equilibrium temperature is proportional to the inverse harmonic sum of the surface gravity of each horizon. In Secs. III B and III C, we apply the method of complex paths to three coordinate systems—spherically symmetric, Painlevé, and Lemaître—of SdS space-time. In Sec. V, we propose a new definition of entropy for space-times with multiple horizons and discuss its implications for SdS space-time. Finally, in Sec. VI, we discuss the results.

Throughout this paper, the metric signature we shall adopt is (-, +, +, +). We use greek letters for (3+1)D and lower case latin letters for (1+1)D. The quantum field is a massless, minimally coupled scalar field (Φ) .

II. SPHERICALLY SYMMETRIC SPACE-TIMES

A. General analysis

The line element for an interval in a spherically symmetric space-time can be written in the following form:

$$ds^{2} \equiv g_{\mu\nu}^{(4)} dx^{\mu} dx^{\nu} = g_{ab}^{(2)} dx^{a} dx^{b} + \exp[-2\phi(x^{a})] d\Omega^{2}$$
(2)
= $-\exp[\nu(x^{0}, x^{1})] (dx^{0})^{2} + \exp[\lambda(x^{0}, x^{1})] (dx^{1})^{2}$ + $\exp[-2\phi(x^{0}, x^{1})] d\Omega^{2}$, (3)

where $d\Omega^2$ is the two-dimensional angular line element. As discussed in Ref. [23] (see also Refs. [19]), the space-time structure of spherically symmetric space-times can be understood via the *R* and *T* regions. If at the given event in the coordinate system (3) the inequality

$$\exp[\nu - \lambda] > \left(\frac{\partial \phi}{\partial x^0} \middle/ \frac{\partial \phi}{\partial x^1}\right)^2 \tag{4}$$

is satisfied, then the event is defined as in the *R* region. (If the above inequality is satisfied at a certain world point then by virtue of the continuity $[\exp(\nu-\lambda), \frac{\partial \phi}{\partial x^0}, \frac{\partial \phi}{\partial x^1}$ cannot be discontinuous] it is satisfied in some neighborhood of this point. Thus, the points in the neighborhood of this system of coordinates satisfy the above inequality and are *R* points and a set of them is an *R* region.) If the opposite inequality is satisfied, the event is in a *T* region. The definitions of *R* and *T* regions can be shown to be coordinate invariant.

Spherically symmetric coordinate. Choosing the Schwarzschild gauge, the line element (3) can be written as

$$ds^{2} = -g(r)dt^{2} + \frac{dr^{2}}{g(r)} + r^{2} d\Omega^{2},$$
 (5)

where g(r) is an arbitrary (continuous, differentiable) function of r. For space-times with a single horizon (like Schwarzschild, dS), g(r) vanishes at one point, say, $r=r_0$. Near r_0 , g(r) can be expanded as

$$g(r) = R(r_0)(r - r_0), \tag{6}$$

where $R(r_0)$ is twice the surface gravity (κ) of the horizon.

For space-times with multiple horizons (like SdS), g(r) vanishes at more than one point, say, $r=r_i$, where i = 1, 2, ..., n. In general, g(r) can be written in the following form

$$g(r) = a \frac{(r-r_1)(r-r_2)(r-r_3)\cdots(r-r_n)}{r^m},$$
 (7)

where *a* is a constant, m < n, $r_n > r_{n-1} > \cdots > r_1$, and all r_i 's are assumed to be positive. Around each of these points one can expand $g(r) = R(r_i) (r - r_i)$ where $R(r_i)/2$ is the surface gravity (κ_i) of each of these horizons.

The R(T) region in the spherically symmetric coordinate system satisfies the inequality condition g(r) > 0 [g(r)] <0]. For space-times with multiple horizons, since g(r) has multiple zeros, there are multiple *R* and *T* regions.

Painlevé coordinate. In order to obtain a line element which is regular at the horizon, we define a new time coordinate (t_P) which is related to the static time coordinate (t) by the relation

$$t = t_P \pm f(r), \tag{8}$$

where *f* is required to be a function of *r* alone to ensure that the metric remains stationary [6,7]. The form of f(r) can be obtained by imposing the condition that the resulting metric be regular at the horizon. This can be realized by demanding that the constant-time slices be flat, i.e.,

$$\frac{1}{g(r)} - g(r) \left[\frac{df}{dr} \right]^2 = 1 \quad \Rightarrow \quad \frac{df}{dr} = \frac{\sqrt{1 - g(r)}}{g(r)}.$$
 (9)

Substituting the expressions for f(r) and t in Eq. (5), we get

$$ds^{2} = -g(r)dt_{P}^{2} \pm 2\sqrt{1-g(r)} dr dt_{P} + dr^{2} + r^{2} d\Omega^{2}.$$
(10)

We will refer to the above line element as the *Painlevé co*ordinate. The above line element is a stationary—but not a static—system. The + sign in the cross term corresponds to the ingoing null geodesic while the - sign corresponds to an outgoing null geodesic.

For both ingoing and outgoing null geodesics, the inequality condition for the *R* region is given by g(r) > 0, implying that the whole of space-time is doubly mapped with respect to spherically symmetric coordinate system [19].

Lemaître coordinate. We can get rid of the cross term in line element (10) by performing a transformation of the radial coordinate (r). This can be achieved by demanding that a set of curves $[t_P, r(t_P)]$ have tangent vectors $(1, dr/d\tau)$ orthogonal to the surface of constant Painlevé time. This gives

$$R + t_P = \int \frac{dr}{\sqrt{1 - g(r)}}.$$
 (11)

Substituting for r in Eq. (10), we get

$$ds^{2} = -dt_{P}^{2} + [1 - g(r)]dR^{2} + r^{2} d\Omega^{2}.$$
 (12)

We will refer to the above line element as the *Lemaître co*ordinate. The line element is explicitly dependent on time and test particles at rest relative to the reference system are particles moving freely in the given field. The coordinate system can be modeled as that natural to a freely infalling/ outfalling observer whose velocity at radial infinity is zero. [The - sign in Eq. (11) corresponds to the infalling observer while the + sign in Equ. (11) corresponds to an outfalling observer.]

In terms of the spherically symmetric coordinate time (t), we have

$$R \mp t = \int \frac{dr}{g(r)} \frac{1}{\sqrt{1 - g(r)}}.$$
(13)

From Eq. (11) it can be seen that

$$r =$$
function of $(R \mp t_P)$. (14)

For the upper sign in Eq. (11), defining 1-g(r) = F(U) and $r^2 = G(U)$, the Lemaître line element (12) in terms of the light cone coordinates (U,V) $(U \equiv R - t_P, V \equiv R - t_P)$ is

$$ds^{2} = \left(\frac{F(U) - 1}{4}\right) (dV^{2} + dU^{2}) + 2\left(\frac{F(U) + 1}{4}\right) dU \, dV + G(U) d\Omega^{2}.$$
 (15)

For the line element corresponding to an infalling observer, the *R* region is given by F(U)>0. For the *T* region, the inequality is opposite. For an outfalling observer, the *R* region is given by the inequality F(V)>0.

In the *T* region there is an asymmetry in the direction of flow and hence the *T* regions corresponding to the infalling and outfalling observer will be different while the *R* regions will be the same. (For an elaborate discussion on this aspect, please refer to Refs. [19].) Hence, the *T* region in the Lemaître coordinates is a doubly mapped *T* region of spherically symmetric coordinates [19].

B. Schwarzschild-de Sitter space-time

The spherically symmetric coordinate of SdS space-time is given by the line element (5), where

$$g(r) = \left(1 - \frac{2M}{r} - \frac{r^2}{l^2}\right),$$
 (16)

M is the mass of the black hole, and l^2 is related to the positive cosmological constant. The space-time has more than one horizon if 0 < y < 1/27 where $y = M^2/l^2$. The black hole horizon (r_h) and the cosmological horizon (r_c) are located, respectively, at

1

$$r_h = \frac{2M}{\sqrt{3y}} \cos\frac{\pi + \psi}{3},\tag{17}$$

$$r_c = \frac{2M}{\sqrt{3y}} \cos\frac{\pi - \psi}{3},\tag{18}$$

where

$$\psi = \cos^{-1}(3\sqrt{3y}). \tag{19}$$

In the limit of $y \rightarrow 0$, we get $r_h \rightarrow 2M$ and $r_c \rightarrow l$. (Note that $r_h < r_c$, i.e., the event horizon is the smallest positive root.) The space-time is dynamic for $r < r_h$ and $r > r_c$. In the limit $y \rightarrow 1/27$, the two horizons—event and cosmological—coincide and is the well known Nariai space-time. If y > 1/27, the space-time is dynamic for all r > 0.

The surface gravity of these horizons is given by [24]

$$\kappa_h = \alpha \left| \frac{M}{r_h^2} - \frac{r_h}{l^2} \right|,\tag{20}$$



FIG. 1. (A) is the Penrose diagram of the de Sitter space where the left and right sides are identified. The spherically symmetric coordinate system of de Sitter space covers region I while the Painlevé and Lemaître coordinate systems cover regions I and II. (B) is the Penrose diagram illustrating the causal structure of the Schwarzschild–de Sitter space-time. Here again, the left and right sides are identified. The spherically symmetric coordinate system covers regions I and II. The Painlevé coordinate system covers regions II and III.

$$\kappa_c = \alpha \left| \frac{M}{r_c^2} - \frac{r_c}{l^2} \right|,\tag{21}$$

where

$$\alpha = \frac{1}{\sqrt{1 - (27y)^{1/3}}}.$$
(22)

In the limit of $y \rightarrow 0$, we get $\kappa_h \rightarrow 1/(4M)$ and $\kappa_c \rightarrow 1/l$. The ranges of κ_h and κ_c are

$$\frac{3\sqrt{3}}{4l} < \kappa_h < \frac{\sqrt{3}}{l} \Rightarrow \frac{l}{\sqrt{3}} < \frac{1}{\kappa_h} < \frac{4l}{3\sqrt{3}}, \tag{23}$$

$$\frac{1}{l} < \kappa_c < \frac{\sqrt{3}}{l} \Rightarrow \frac{l}{\sqrt{3}} < \frac{1}{\kappa_c} < l.$$
(24)

[The above ranges are obtained by setting $y \rightarrow 0$ and $y \rightarrow (1/27)$ in Eqs. (20),(21).]

Figure 1(B) illustrates one version of the Penrose diagram of SdS space-time. (For an easy comparison, we have also provided the Penrose diagram of de Sitter space.) The static coordinate system covers regions I and II of the Penrose diagram. The boundaries of the static region consist of (i) past and future cosmological horizons, and (ii) past and future event horizons.

The Painlevé line element of SdS space-time is given by

$$ds^{2} = -g(r)dt_{P}^{2} \pm 2\sqrt{\frac{2M}{r} + \frac{r^{2}}{l^{2}}} dr dt_{P} + dr^{2} + r^{2} d\Omega^{2}.$$
(25)

where t_P is related to t by the relation

$$t = t_P \pm \int dr \frac{\sqrt{2M/r + r^2/l^2}}{(1 - 2M/r - r^2/l^2)}.$$
 (26)

The above line element is a stationary—but not a static system and it covers regions II and III of the Penrose diagram. The + sign in the above chart is suitable for studying the physical experience of observers falling freely and radially into the hole. (The + sign in the cross term corresponds to the ingoing null geodesic while the - sign corresponds to an outgoing null geodesic.) For both ingoing and outgoing null geodesics, the inequality condition for the *R* region [23] is given by g(r) > 0, implying that the whole of space-time is doubly mapped with respect to the spherically symmetric coordinate system (5).

The Lemaitre line element is given by

$$ds^{2} = -dt_{p}^{2} + \left(\frac{2M}{r} + \frac{r^{2}}{l^{2}}\right)dR^{2} + r^{2} d\Omega^{2}, \qquad (27)$$

where

$$2r^{3/2} = (2Ml^2) \exp\left[-\frac{3(R \pm t_p)}{2l}\right] - \exp\left[\frac{3(R \pm t_p)}{2l}\right].$$
(28)

The line element is explicitly dependent on time and test particles at rest relative to the reference system are particles moving freely in the given field. $R - t_p$ in the right-hand side (RHS) of the above equation corresponds to the infalling observer while the $R + t_p$ in the RHS of the above expression corresponds to an outfalling observer. Like the Painlevé, the Lemaître line element also covers regions II and III of the Penrose diagram.

For the line element corresponding to an infalling observer, the *R* region [23] is given by F(U)>0 ($U=R-t_P$). For the *T* region, the inequality is opposite. For an outfalling observer, the *R* region is given by the inequality F(V)>0.

III. PARTICLE PRODUCTION IN SPACE-TIMES WITH MULTIPLE HORIZONS

In Refs. [18–20] the method of complex paths was applied to space-times with a single horizon. The temperature associated with the quantum field (in the three coordinate systems—spherically symmetric, Painlevé, and Lemaître) in these space-times were shown to be consistent with the temperatures obtained by using other quantum field theoretic techniques like Bogoliubov coefficients, Euclidean field theory, and the effective action. As discussed in the Introduction, the standard quantum field theoretic techniques do not work well for space-times with multiple horizons. In this

section, we extend the method of complex paths to spacetimes with multiple horizons and in particular to Schwarzschild-de Sitter space-time. We show that the effective temperature for the quantum fields in these space-times is proportional to the inverse harmonic sum of the surface gravity at each horizon.

A. General spherically symmetric space-time

In this subsection, we obtain the effective temperature in a general spherically symmetric space-time with multiple horizons. (The result in this subsection is from Ref. [26] and is summarized here for completeness.)

Consider a quantum field propagating in a general spherically symmetric space-time (5) with multiple horizons. The wave equation is given by

$$\partial_{\mu}(\sqrt{-gg^{\mu\nu}}\partial_{\nu})\Phi = 0,$$

$$\frac{r^{2}}{g(r)}\frac{\partial^{2}\Psi}{\partial t^{2}} - \frac{\partial}{\partial r}\left(r^{2}g(r)\frac{\partial\Psi}{\partial r}\right) + L^{2}\Psi = 0.$$
(29)

where $\Phi(x^{\mu}) = \Psi(t,r)Y_{lm}(\theta,\phi)$. The semiclassical wave functions satisfying the above are obtained by making the ansatz

$$\Psi(r,t) = \exp\left[\frac{i}{\hbar}S(r,t)\right],\tag{30}$$

where *S* is a functional which will be expanded in powers of \hbar . Expanding *S* in a power series of \hbar ,

$$S(r,t) = S_0(r,t) + \hbar S_1(r,t) + \hbar^2 S_2(r,t) + \cdots, \quad (31)$$

and substituting the resulting expression in Eq. (29) (neglecting terms of order \hbar or higher), we have

$$-\frac{1}{g(r)}\left(\frac{\partial S_0}{\partial t}\right)^2 + g(r)\left(\frac{\partial S_0}{\partial r}\right)^2 + \frac{L^2}{r^2} = 0.$$
(32)

This is simply the Hamilton-Jacobi equation satisfied by a massless particle moving in a general spherically symmetric coordinate. Making the ansatz $S_0 = -Et + B(r)$, we get

$$B(r) = \pm \int^{r} \frac{dr}{g(r)} \sqrt{E^{2} - (L^{2}/r^{2})g(r)}$$

= $\pm \frac{E}{a} \int^{r} dr \frac{r^{m}}{(r - r_{1})(r - r_{2}) \cdots (r - r_{n})}.$ (33)

(The last expression is obtained by setting L=0 and noticing that near the horizon the presence of the L^2 term can be neglected. It can also be noted that the above expression is valid for an arbitrary dimension and hence the following result is true for a spherically symmetric space-time in an arbitrary dimension.) Rewriting the RHS in the above expression with partial fractions, i.e.,

$$\frac{1}{a} \frac{r^m}{(r-r_1)(r-r_2)\cdots(r-r_n)} = \sum_{i=1}^n \frac{A_i}{(r-r_i)}, \quad (34)$$

where

$$A_{i} = \frac{r_{i}^{m}}{a} \prod_{j=1}^{n} \frac{1}{(r_{i} - r_{j})}$$
$$= \frac{1}{(m-1)\kappa_{i}}, \quad j \neq i,$$
(35)

and κ_i is the surface gravity of each of the horizons. Substituting the above expressions in Eq. (33), we get

$$B(r_{s}, r_{f}) = \pm \frac{E}{m-1} \sum_{i=1}^{n} \frac{1}{\kappa_{i}} \int_{r_{s}}^{r_{f}} \frac{dr}{r-r_{i}}.$$
 (36)

The sign ambiguity is related to the *outgoing* $(\partial S_0 / \partial r > 0)$ or *ingoing* $(\partial S_0 / \partial r < 0)$ nature of the particle. Unlike space-times with a single horizon [18,19], space-times with multiple horizons have many possible ways of choosing the initial (r_1) and final (r_2) points. However, we will consider a particular situation of the particle crossing all the horizons, i.e., $r_s < r_1$ and $r_f > r_n$. (This result can be used to obtain other possible cases.)

For an outgoing particle $(\partial S_0 / \partial r > 0)$ with respect to the horizon (r_1) , the contribution to S_0 is

$$S_{0}(\text{emission}) = -\frac{E}{(m-1)\kappa_{1}} \int_{r_{1}-\epsilon}^{r_{1}+\epsilon} \frac{dr}{r-r_{1}}$$
$$-\frac{E}{(m-1)\kappa_{2}} \int_{r_{2}+\epsilon}^{r_{2}-\epsilon} \frac{dr}{r-r_{2}} - \cdots$$
$$-\frac{E}{(m-1)\kappa_{n}} \int_{r_{n}+\epsilon}^{r_{n}-\epsilon} \frac{dr}{r-r_{n}} + \text{real part}$$
$$= i \frac{\pi E}{(m-1)\kappa_{\text{eff}}} + \text{real part}, \qquad (37)$$

where

$$\frac{1}{\kappa_{\rm eff}} = \sum_{i=1}^{n} \frac{1}{\kappa_i},\tag{38}$$

and the - sign in front of the integrals corresponds to the condition that $\partial S_0 / \partial r > 0$ at $r = r_s < r_1$ and $\partial S_0 / \partial r < 0$ at $r = r_f > r_n$ and the integrals are evaluated by taking the contours to lie in the upper complex plane. It should be noted that the above expression can also be considered as the action for the ingoing particles with respect to the horizon r_n . [Note that we have assumed all r_i 's to be positive, and hence S_0 will have an *imaginary* contribution from all of them. For SdS space-time, which we consider in detail, one of the roots of g(r) is negative and will not contribute to S_0 .]

Using the above procedure, the action for an ingoing particle $(\partial S_0 / \partial r < 0)$ with respect to the horizon r_1 is given by

$$S_{0}(\text{absorption}) = -\frac{E}{(m-1)\kappa_{1}} \int_{r_{1}+\epsilon}^{r_{1}-\epsilon} \frac{dr}{r-r_{1}}$$
$$-\frac{E}{(m-1)\kappa_{2}} \int_{r_{2}-\epsilon}^{r_{2}+\epsilon} \frac{dr}{r-r_{2}} - \dots$$
$$-\frac{E}{(m-1)\kappa_{n}} \int_{r_{n}-\epsilon}^{r_{n}+\epsilon} \frac{dr}{r-r_{n}} + \text{real part}$$
$$= -i \frac{\pi E}{(m-1)\kappa_{\text{eff}}} + \text{real part.}$$
(39)

The last expression is obtained by evaluating the integrals with the contours taken in the lower half plane. Taking the modulus square of Eqs. (37),(39), we get

$$P[\text{emission}] = \exp\left[-\frac{4\pi E}{(m-1)\kappa_{\text{eff}}}\right] P[\text{absorption}]. \quad (40)$$

The exponential dependence on the energy allows one to give a *thermal* interpretation for the above result. The temperature of the emission spectrum is given by

$$\beta^{-1} = \frac{(m-1)}{4\pi} \kappa_{\text{eff}}.$$
(41)

The above result shows that the spectrum of particles created in a spherically symmetric space-time with multiple horizons is thermal, and the temperature is proportional to the inverse harmonic sum of the surface gravity of each horizon. The above result can be interpreted as a particle propagating from inside the horizon (r_1) to outside the horizon (r_n) and picking up $\exp(-\beta_i E)$ at each horizon, resulting in $\exp[-\beta E)$. (Note that we have neglected the backscattering of the particles from the horizon.) The above analysis can be performed in Painlevé and Lemaître coordinates for these multiple-horizon space-times as discussed in Appendix A. In the next section, we explicitly demonstrate the above result for three different coordinate systems of SdS space-time.

B. Spherically symmetric SdS coordinate

For a quantum field propagating in the spherically symmetric coordinate system of SdS space-time, the field equation is given by Eq. (29) where g(r) is given by Eq. (16). In the previous section we assumed that all the roots of g(r) are positive and hence S_0 has contributions from all r_i 's. However, for SdS space-time, one of the roots of g(r) is negative and the results in the previous section do not follow automatically.

Making the usual semiclassical ansatz for Ψ , and expanding *S* in powers of \hbar , we get

$$B(r) = \pm \int^{r} \frac{dr}{g(r)} \sqrt{E^{2} - L^{2}/r^{2}g(r)}$$
$$= \pm E \int^{r} \frac{dr}{1 - 2M/r - r^{2}/l^{2}},$$
(42)

where the last expression is obtained by setting L=0. After a bit of lengthy algebra, it can be shown that

$$\left(1 - \frac{2M}{r} - \frac{r^2}{l^2}\right)^{-1} = \frac{\alpha}{2\kappa_h(r - r_h)} - \frac{\alpha}{2\kappa_c(r - r_c)} - \frac{\alpha}{2\kappa_c(r - r_c)} - \frac{\alpha}{2}\left(\frac{1}{\kappa_h} - \frac{1}{\kappa_c}\right)\frac{1}{r + r_c + r_h}, \quad (43)$$

where κ_h , κ_c are given by expressions (20),(21) and r_h , r_c are given by Eqs. (17),(18). Substituting the above expression in Eq. (42), we get

$$B(r) = \pm \frac{E\alpha}{2\kappa_h} \int^r \frac{dr}{r - r_h} \mp \frac{E\alpha}{2\kappa_c} \int^r \frac{dr}{r - r_c}$$
$$\mp \frac{E\alpha}{2} \left(\frac{1}{\kappa_h} - \frac{1}{\kappa_c}\right) \int^r \frac{dr}{r + r_c + r_h}.$$
(44)

The sign ambiguity is related to the *outgoing* $(\partial S_0 / \partial r > 0)$ or *ingoing* $(\partial S_0 / \partial r < 0)$ nature of the particle. Unlike spacetimes with a single horizon [18,19], SdS has many possible ways of choosing the initial (r_1) and the final (r_2) points. The following scenarios will be investigated.

(1) $r_1 < r_h$ and $r_2 > r_c$: The first two integrals in the RHS of Eq. (44) do not exist.

(2) $r_1 < r_h$ and $r_2 < r_c$: The first integral in the RHS of the above expression does not exist since the denominator vanishes at $r = r_h$.

(3) $r_1 > r_h$ and $r_2 > r_c$: The second integral in the RHS of the above expression does not exist since the denominator vanishes at $r = r_h$.

In what follows, we shall obtain the spectrum of particles produced in the first scenario. (The other two cases follows automatically from this discussion.)

In the first case, both the points are outside the region (r_h, r_c) . Hence, the contribution to the probability amplitude for emission and absorption will be due to both the horizons—event and cosmological. For an outgoing particle $(\partial S_0 / \partial r > 0)$ with respect to the event horizon (r_h) , the contribution to S_0 is

$$S_{0}(\text{emission}) = -\frac{E\alpha}{2\kappa_{h}} \int_{r_{h}-\epsilon}^{r_{h}+\epsilon} \frac{dr}{r-r_{h}} - \frac{E\alpha}{2\kappa_{c}} \int_{r_{c}+\epsilon}^{r_{c}-\epsilon} \frac{dr}{r-r_{c}} + \text{real part}$$
(45)

$$=i\frac{\pi E\alpha}{2\kappa_{\rm eff}} + \text{real part},\tag{46}$$

where

$$\kappa_{\rm eff} = \left[\frac{1}{\kappa_h} + \frac{1}{\kappa_c}\right]^{-1},\tag{47}$$

and the - sign in front of the integrals corresponds to the condition that $\partial S_0 / \partial r > 0$ at $r = r_1 < r_h$ and $\partial S_0 / \partial r < 0$ at $r = r_2 > r_c$ and the integrals are evaluated by taking the contours to lie in the upper complex plane. It should be noted

that the above expression can also be considered as the action for the ingoing particles with respect to the cosmological horizon (r_h) .

Using the above procedure, the action for an ingoing particle $(\partial S_0 / \partial r < 0)$ with respect to the event horizon (r_c) is

$$S_{0}(\text{absorption}) = -\frac{E\alpha}{2\kappa_{h}} \int_{r_{h}+\epsilon}^{r_{h}-\epsilon} \frac{dr}{r-r_{h}} - \frac{E\alpha}{2\kappa_{c}} \int_{r_{c}-\epsilon}^{r_{c}+\epsilon} \frac{dr}{r-r_{c}} + \text{real part}$$
(48)

$$= -i\frac{\pi E\,\alpha}{2\,\kappa_{\rm eff}} + \text{real part},\tag{49}$$

where the integrals are evaluated by taking the contour in the lower half plane. Taking the modulus squared, we get

$$P[\text{emission}] = \exp\left[-\frac{2\pi E\,\alpha}{\kappa_{\text{eff}}}\right] P[\text{absorption}].$$
(50)

The exponential dependence on the energy allows one to give a *thermal* interpretation for the above result. The temperature of the emission spectrum is given by

$$\beta^{-1} = \frac{1}{2\pi\alpha} \frac{\kappa_h \kappa_c}{\kappa_h + \kappa_c}.$$
(51)

The following features are noteworthy regarding this result.

(i) The equilibrium temperature in SdS space-time is inversely proportional to the harmonic mean of the two horizons. [The range of κ_{eff} is $(1/l) < \kappa_{\text{eff}} < (\sqrt{3}l)$.]

(ii) In this case, radiation is propagating inward from the cosmological horizon and outward from the black-hole horizon. (We have neglected the scattering of the radiation from these horizons.) Thus, there is a constant flux of radiation flowing between these horizons and a static observer is in a thermal bath of radiation with the above temperature. (This result was obtained in Ref. [15] from the Euclidean quantum gravity point of view.)

(iii) In the limit of $1/l \rightarrow 0$, it reproduces the Hawking temperature for a Schwarzschild black hole in asymptotically flat space-time.

(iv) In the limit of $M \rightarrow 0$, this reproduces the Gibbons-Hawking temperature for a vacuum de Sitter space-time.

(v) In the Nariai limit $(y \rightarrow 1/27)$, the event and cosmological horizons coincide. Thus, the results obtained for space-times with multiple horizons cannot be applied. In the Nariai limit, using the analysis performed in Ref. [18] and Appendix A, we get [24]

$$\beta_{\text{Nariai}}^{-1} = \frac{\sqrt{3}}{2\pi l}.$$
(52)

C. Painlevé and Lemaître SdS coordinate

Let us now consider a scalar field propagating in the Painlevé coordinate system of SdS space-time. Retracing the steps as discussed in Appendix A, we get (for L=0)

$$B(r) = E \int^{r} dr \frac{\sqrt{2M/r + r^2/l^2 \pm 1}}{1 - 2M/r - r^2/l^2}.$$
 (53)

As discussed in Sec. II B, Painlevé coordinates cover regions II and III. In the Painlevé coordinate system, as opposed to spherically symmetric coordinates, there are only two possible ways of choosing the initial (r_1) and final (r_2) points: (i) both the points are on one side of r_h and (ii) the two points lie on the opposite sides of the event horizon.

For an outgoing particle $(\partial S_0 / \partial r > 0)$ with respect to the event horizon (r_c) , the contribution to S_0 is

$$S_0[\text{emission}] = -\frac{E\alpha}{\kappa_h} \int_{r_h-\epsilon}^{r_h+\epsilon} \frac{dr}{r-r_h} + \text{real terms}$$
$$= i\frac{\pi E\alpha}{\kappa_h} + \text{real terms.}$$
(54)

In order to obtain the action for absorption of particles corresponding to the ingoing particles $[(\partial S_0/\partial x) < 0]$, we have to repeat the above calculation for the lower sign of the metric given in Eq. (25). In this case, it is easy to see that the only singular solution corresponds to ingoing particles and we get

$$S_0[\text{absorption}] = -\frac{\pi E \alpha}{\kappa_h} + \text{real terms.}$$
 (55)

Constructing the semiclassical propagator in the usual manner and taking the modulus squared we obtain the probability. From the earlier discussion, we know that in calculating the probability of absorption/emission there is an extra contribution to the probability from four sets of complex paths satisfying the semiclassical ansatz. Taking this into account, we get

$$\beta^{-1} = \frac{\kappa_h}{2\pi\alpha}.$$
(56)

In the case of Lemaître coordinates, repeating the above procedure it is easy to show that the temperature of the radiation is the same as that obtained in the case of Painlevé coordinates and case II of spherically symmetric coordinates. The Painlevé and Lemaître observers will be in thermal equilibrium with the above temperature.

IV. BOGOLIUBOV COEFFICIENTS FOR SPACE-TIMES WITH MULTIPLE HORIZONS

In this section, we extend the analysis of Appendix B and obtain Bogoliubov coefficients for space-times with multiple horizons. The equation of motion for the scalar field in the spherically symmetric coordinate is given by Eq. (29). Near each of the horizons, the solution to the wave equation is given by

$$\Psi = C_1^i \exp(-iEt)(x_i)^{iE/R(r_i)},$$
(57)

where C_1^i is an arbitrary constant. Ψ^* is also a solution to the scalar field equation (29). For each horizon, we can write the two sets of modes as

$$\Psi(x_i < 0) = C_1^i \psi(-x_i) + C_2^i \psi^*(-x_i),$$

$$\Psi(x_i > 0) = C_a^i \psi(x_i) + C_\beta^i \psi^*(x_i),$$
(58)

where C_1^i , C_2^i , C_{α}^i , and C_{β}^i are constants to be determined. We obtain a relation between the different constants similar to that in the single horizon case. They are given by

$$C_{\alpha}^{i} = C_{1}^{i} \exp[\pi E/R(r_{i})], \qquad (59)$$

$$C_{\beta}^{i} = C_{2}^{i} \exp[-\pi E/R(r_{i})].$$
 (60)

This leaves us with identifying the relation between the two constants (C_1^i, C_2^i) to the left of the horizon r_i . For space-times with a single horizon, we assumed that the net current to the left of the horizon is zero. However, for space-times with multiple horizons, the above assumption is not valid.

In order to see this explicitly, let us consider a quantum field propagating in spherically symmetric coordinates of a SdS space-time which has two horizons—event and cosmological. Let r_i be the cosmological horizon. To the left of the cosmological horizon, the net current is nonzero due to the presence of an event horizon which produces thermal radiation. (Here we are assuming that the backscattering of the particles from the event horizons can be neglected.) In this case, C_1^i and C_2^i are related by

$$|C_1^i|^2 = \exp(-2\pi\alpha E/\kappa_h)|C_2^i|^2.$$
(61)

This gives

$$\left|\frac{C_{\beta}^{i}}{C_{\alpha}^{i}}\right|^{2} = \exp[-2\pi\alpha E/\kappa_{\text{eff}}].$$
(62)

Interpreting $|C_{\alpha}^{i}|^{2} \equiv |\alpha_{E}^{i}|^{2}$ and $|C_{\beta}^{i}|^{2} \equiv |\beta_{E}^{i}|^{2}$ as Bogoliubov coefficients and using the unitary condition, we have

$$|\alpha_{E}^{i}|^{2} = \frac{1}{1 - \exp[-2\pi\alpha E/\kappa_{\text{eff}}]},$$
$$|\beta_{E}^{i}|^{2} = \frac{1}{\exp[2\pi\alpha E/\kappa_{\text{eff}}] - 1}.$$
(63)

The following features are noteworthy regarding this result.

(1) $|\beta_E^i|^2$ is a Planck spectrum with a temperature given by Eq. (51).

(2) The Bogoliubov coefficients at each of the horizons are the same implying that the space-time is in thermal equilibrium with temperature proportional to κ_{eff} .

(3) The condition in Eq. (61) has been crucial in obtaining the result. The condition implies that a particle propagating from inside the horizon (r_h) to outside the horizon (r_c) picks up $\exp[-\beta_h E]$ and $\exp[-\beta_c E]$, resulting in $\exp[-\beta E)$.

(4) In the Nariai limit, the condition (61) is not valid and we need to impose the condition (B6) (which implies that the net current to the left of the horizon is zero). In this case, the Bogoliubov coefficients are given by Eqs. (B8) where $\kappa = l/\sqrt{3}$.

V. ENTROPY OF SdS

In this section, we propose a new definition of entropy for space-times with multiple horizons analogous to the entropic definition for space-times with a single horizon and discuss the implications in the context of SdS space-time.

For space-times with a (compact) single horizon, the entropy is given by

$$S(\text{single horizon}) = \frac{A}{4} = c \frac{\pi}{\kappa^2}, \tag{64}$$

where c = 1 and 1/4 for dS and Schwarzschild space-times, respectively. Assuming that the relation between *S* and κ holds true for space-times with multiple horizons, the entropy for space-times with multiple horizons can be written as

$$S(\text{multiple horizons}) = \frac{\pi}{\kappa_{\text{eff}}^2},$$
 (65)

where κ_{eff} is given by Eq. (38). The following features are noteworthy regarding this expression.

(i) The above expression can be treated as an entropy associated with a single (effective) horizon for space-times with multiple horizon. For SdS space-times, this corresponds to the entropy of the system in thermal equilibrium with the temperature (51).

(ii) In the literature, the entropy of space-times with multiple horizons is defined as the sum of the entropies of individual horizons. Using our definition, the entropy will have extra nonzero contributions. In the case of SdS space-times, we have

$$S_{SdS} = \frac{\pi}{\kappa_h^2} + \frac{\pi}{\kappa_c^2} + \frac{2\pi}{\kappa_c \kappa_h}.$$
 (66)

If we set

$$\frac{\pi}{\kappa_h^2} = S_h, \quad \frac{\pi}{\kappa_c^2} = S_c, \tag{67}$$

then the expression for the entropy of SdS space-times can be written as

$$S_{SdS} = S_h + S_c + 2\sqrt{S_c S_h},\tag{68}$$

where S_h , S_c are the entropies of event and cosmological horizons, respectively.

(iii) The entropy of the SdS space-time obtained in Eq. (66) satisfies the *D*-bound conjecture [22].

Following the discussion in Sec. II B, we know that the range of κ_{eff} is $(1/l, \sqrt{3}/l)$. [The range of $1/\kappa_{\text{eff}}$ is $(l/\sqrt{3}, l)$.] Substituting the range of κ_{eff} in the expression for entropy, we get

$$S_{SdS} < \pi l^2 \equiv S_{dS} \,. \tag{69}$$

(iv) The form of the entropy in Eq. (68) is similar to Eq. (5.1) of Ref. [17]. In Ref. [17], the authors showed that the final values of the entropies of the event (S_B) and cosmological (S_C) horizons satisfy the inequality

$$S_B + S_C + \sqrt{S_B S_C} \le \pi l^2. \tag{70}$$

In our case, the inequality (69) is satisfied for all values of entropies of the event and cosmological horizons. The *D* bound we have obtained in Eq. (69) is a stronger inequality than that of (70).

VI. CONCLUSIONS

In this paper, we have studied the spectrum of created particles in SdS space-time for a linear, massless scalar field using the method of complex paths. We have shown that it is possible to obtain a temperature for SdS space-time which corresponds to a system in thermal equilibrium. The equilibrium temperature is the harmonic mean of the event and cosmological horizon temperatures. We have also obtained the same result by calculating the Bogoliubov coefficients using the wave modes near the horizons.

It has been assumed in the literature that the entropy of space-times with multiple horizons is the sum of the entropies of individual horizons. In this paper, we have proposed a new definition of entropy for space-times with multiple horizons, analogous to the entropic definition of entropy for space-times with a single horizon. We have defined the entropy for these space-times to be inversely proportional to the square of the effective surface gravity. Using this definition, we have shown that the entropy of SdS space-time satisfies the *D*-bound conjecture [22]. In Ref. [25], the authors have proposed a definition for entropy outside the cosmological horizon in SdS space-time which satisfies the *D* bound. It is interesting to look for the connection between the two proposed entropies.

The above result brings attention to the following interesting questions.

(1) For space-times with a single horizon, there is a unique way of obtaining a global coordinate system. For space-times with multiple horizons, it has not been possible to obtain a global coordinate system. Is it possible to obtain a global coordinate system whose Euclidean metric has periodicity proportional to κ_{eff} ?

(2) Can we obtain stress tensors for SdS space-time which correspond to the thermal state with the temperature proportional to κ_{eff} ?

(3) What kind of quantum vacuum state will this correspond to?

We hope to return to study some of these problems in the near future.

ACKNOWLEDGMENTS

The author would like to thank T. Padmanabhan and A. J. M. Medved for useful discussions. The author gratefully acknowledges support from Fundação para a Ciência e a Tecnologia (Portugal) under the grant SFRH/BI/9622/2002.

APPENDIX A

In this appendix, we generalize the results of Ref. [19] to two nonsingular coordinate systems—Painlevé and Lemaître—of a general spherically symmetric space-time (with a single horizon). We show that the temperature associated with the radiation in these two coordinate systems, *for space-times with a single horizon*, is the same as in the spherically symmetric coordinate system. These results can be extended to space-times with multiple horizons where g(r) is given by Eq. (7).

In the context of the method of complex paths, these two coordinate systems possess the interesting feature of double mapping of the paths. In the following, we briefly describe the multiple mapping of space-time and the measure of these paths. In Refs. [19], it was argued that the family of complex paths used to calculate the emission/absorption probability takes into account all paths irrespective of the multiple mapping of that part/whole of the space-time.

Let us consider a coordinate system like the Painlevé system where the whole of the space-time is doubly mapped with respect to the spherically symmetric coordinate. The space-time has two distinct R and T regions. Hence, the complex paths will have *equal* contributions from both of these, which do not have any point in common. Hence, the contribution to the amplitude of emission/absorption by these two paths will be mutually exclusive.

In the case of coordinate systems where part of the spacetime is doubly mapped with respect to the spherically symmetric coordinate (as in the case of Lemaître coordinates), it is always possible to find one point that is common to the paths contributing to absorption/emission. Hence, these paths are not mutually exclusive. These paths, on the other hand, will be mutually exclusive when one considers the probability amplitude—which is the important quantity in our approach. Hence, the action we obtain by regularizing the singularity and the resulting probability amplitude, for absorption/emission, will have equal contributions from both these paths.

1. Painlevé coordinate system

Let us consider a quantum field propagating in the Painlevé line element (10) with the + signature in the cross term. The equation for a scalar field propagating in general Painlevé coordinates is given by

$$r^{2} \frac{\partial^{2} \Psi}{\partial t_{p}^{2}} + 2r^{2} \sqrt{1 - g(r)} \frac{\partial^{2} \Psi}{\partial r \partial t_{p}} + \frac{d}{dr} [r^{2} g(r)] \frac{\partial \Psi}{\partial r}$$
$$- r^{2} g(r) \frac{\partial^{2} \Psi}{\partial r^{2}} + \frac{d}{dr} [r^{2} \sqrt{1 - g(r)}] \frac{\partial \Psi}{\partial t_{p}} = -L^{2} \Psi,$$
(A1)

where $\Phi(x^{\mu}) = \Psi(t_P, r) Y_{lm}(\theta, \phi)$. Making the standard semi-classical ansatz for Ψ and expanding *S* in powers of \hbar as discussed in Sec. III A, we get, to the lowest order,

$$-\left(\frac{\partial S_0}{\partial t_P}\right)^2 + 2\sqrt{1-g(r)}\left(\frac{\partial S_0}{\partial r}\right)\left(\frac{\partial S_0}{\partial t_P}\right) + g(r)\left(\frac{\partial S_0}{\partial r}\right)^2 + \frac{L^2}{r^2} = 0.$$
(A2)

The above equation is the Hamilton-Jacobi equation of a massless particle propagating in a general Painlevé line element (10). Substituting the ansatz $S_0 = -Et_P + B(r)$ in the above expression, we get

$$\frac{dB}{dr} = \frac{E\sqrt{1-g(r)}}{g(r)}$$
$$\pm \frac{\sqrt{4E^2[1-g(r)]-4g(r)[E^2-L^2/r^2]}}{g(r)}.$$
 (A3)

It is easy to see that, near the horizon, the presence of the L^2 term can be neglected since it is multiplied by g(r). Thus for L=0 we have

$$B(r) = E \int^{r} dr \frac{\sqrt{1 - g(r)} \pm 1}{g(r)}$$
$$= E \int^{r} dr \frac{\sqrt{1 - g(r)}}{g(r)} \pm E \int^{r} \frac{dr}{g(r)}.$$
 (A4)

For space-times with a single horizon, one can expand g(r) around the horizon as $g(r) = R(r_0)(r-r_0)$. The numerator in the first integral (in the RHS) of the above expression can be approximated to unity near the horizon. Noticing that the denominator is singular at $r = r_0$ only for the positive sign, we get the action for the outgoing particle as

$$S_0[\text{emission}] = -2E \int_{r_0-\epsilon}^{r_0+\epsilon} \frac{dr}{g(r)} + \text{real part}$$
$$= +\frac{2i\pi E}{R(r_0)} + \text{real part.}$$
(A5)

In order to obtain the action for absorption of particles corresponding to the ingoing particles $[(\partial S_0/\partial r) < 0]$, we have to repeat the above calculation for the metric (10) with a – sign in the cross term. In this case, it is easy to see that the only singular solution corresponds to ingoing particles and so

$$S_0[\text{absorption}] = -\frac{2i\pi E}{R(r_0)} + \text{real part.}$$
 (A6)

Constructing the semiclassical propagator in the usual manner and taking the modulus squared we obtain the probability. Extending the double mapping of the paths to the generalized Painlevé coordinates and squaring the modulus to get the probability, we get the temperature associated with the quantum fields propagating in generalized Painlevé coordinates as

$$\beta^{-1} = \frac{R(r_0)}{4\pi} = \frac{\kappa(r_0)}{2\pi}.$$
 (A7)

For dS space-time, $g(r) = (1 - r^2/l^2)$, $\kappa = (1/l)$, and hence the temperature associated with the radiation is $1/(2\pi l)$.

2. Lemaître coordinate system

Consider a quantum field propagating in the general spherically symmetric space-times described by the Lemaître line element (12). The scalar field equation [for the - sign in the expression (11)] is given by

$$-\frac{\partial}{\partial t_{P}}\left[r^{2}\sqrt{1-g(r)}\frac{\partial\Psi}{\partial t_{P}}\right] + \frac{\partial}{\partial R}\left[\frac{r^{2}}{\sqrt{1-g(r)}}\frac{\partial\Psi}{\partial R}\right]$$
$$-\sqrt{1-g(r)}L^{2}\Psi = 0, \qquad (A8)$$

where $\Phi(x^{\mu}) = \Psi(t_P, R) Y_{lm}(\theta, \phi)$. Noting from Eq. (11) that *r* is related to $R - t_P \equiv U$, the above equation rewritten in terms of the light-cone coordinates (U, V) translates into

$$\frac{\partial}{\partial V} \left[(F(U) - 1)G(U) \frac{\partial \Psi}{\partial V} \right] + \frac{\partial}{\partial U} \left[(F(U) - 1)G(U) \frac{\partial \Psi}{\partial U} \right] \\ + \frac{\partial}{\partial V} \left[(F(U) + 1)G(U) \frac{\partial \Psi}{\partial V} \right] \\ + \frac{\partial}{\partial U} \left[(F(U) + 1)G(U) \frac{\partial \Psi}{\partial V} \right] \\ - 2 \frac{\sqrt{F(U)}}{G(U)} L^2 \Psi = 0.$$
 (A9)

Making the standard semiclassical ansatz for Ψ and expanding S in powers of \hbar , we get, to the lowest order of S_0 ,

$$\left(\frac{\partial S_0}{\partial U}\right)^2 + \left(\frac{\partial S_0}{\partial V}\right)^2 + 2\left(\frac{1+F(U)}{1-F(U)}\right)\frac{\partial S_0}{\partial U}\frac{\partial S_0}{\partial V} - \left(\frac{L^2}{1-F(U)}\right) = 0.$$
(A10)

The above equation is the Hamilton-Jacobi equation of a massive particle propagating in the Lemaître line element (12). Substituting the ansatz $S_0 = -EV + B(U)$ in the above expression, we get (for L=0)

$$B(U) = E \int^{U} dU \frac{1 + F(U) \pm 2\sqrt{F(U)}}{1 - F(U)}.$$
 (A11)

[Near the horizon the contribution due to *L* can be neglected since it is multiplied by g(r).] Notice that the denominator is singular at U=1 only for the positive sign. Our interest in this exercise is to obtain the principal part of the action near the horizon $r=r_0$. The transformation relation (11) near the horizon translates into

$$U = -\frac{\sqrt{1 - R(r_0)(r - r_0)}}{R(r_0)},$$
 (A12)

thus giving

$$F(U) = 1 - R(r_0)(r - r_0) = \left(\frac{R(r_0)U}{2}\right)^2.$$
 (A13)

Substituting the above expression in Eq. (A11), we get

$$S_0[\text{emission}] = -\frac{4E}{R(r_0)} \int_{1-\epsilon}^{1+\epsilon} dU \frac{U}{1-U} + \text{real part}$$
$$= \frac{4i\pi E}{R(r_0)} + \text{real part}, \qquad (A14)$$

where U is rescaled as $[R(r_0)U/2] \rightarrow U$.

In order to obtain the action for absorption of particles corresponding to the ingoing particles $[(\partial S_0/\partial r) < 0]$, we have to repeat the above calculation for the metric (12) with the + sign of expression (11). In this case, it is easy to see that the only singular solution corresponds to ingoing particles and so

$$S_0[\text{absorption}] = -\frac{4i\pi E}{R(r_0)} + \text{real part.}$$
 (A15)

Constructing the semiclassical propagator in the usual manner and taking the modulus squared we obtain the probability. Now, extending the double mapping of the paths to the generalized Lemaître coordinates and squaring the modulus, we get the temperature associated with the quantum scalar fields propagating in the generalized Lemaître coordinates as

$$\beta^{-1} = \frac{R(r_0)}{4\pi} = \frac{\kappa(r_0)}{2\pi},$$
 (A16)

which is the same as the expression obtained in generalized Painlevé coordinates. For dS space-time, we get $\beta^{-1} = 1/(2\pi l)$.

APPENDIX B

In this appendix, we recover Bogoliubov coefficients (for space-times with a single horizon) solely by the mode functions of the scalar field near the horizon [27]. Let us consider a quantum field propagating in a spherically symmetric coordinate (5). The equation of motion of the scalar field $\Phi = \Psi(t,r)Y_{lm}(\theta,\phi)$ is

$$\frac{r^2}{g(r)}\frac{\partial^2\Psi}{\partial t^2} - \frac{\partial}{\partial r}\left(r^2g(r)\frac{\partial\Psi}{\partial r}\right) + L^2\Psi = 0.$$
(B1)

Since our interest is near the horizon, we can transform the above expression in terms of a new variable $x \equiv r - r_0$. The solution to the scalar field, close to the horizon x=0, is given by

$$\Psi = C_1 \exp(-iEt) x^{iE/R(0)}, \tag{B2}$$

where C_1 is an arbitrary constant. Note that Ψ^* is also a solution to the scalar field equation.

In the usual field theory description, Bogoliubov transformations relate two distinct orthonormal sets of modes that are solutions to the equation of motion. In the case of the method of complex paths, the two sets of modes correspond to $\Psi(x<0)$ and $\Psi(x>0)$. We can write the two sets of modes as

$$\Psi(x<0) = C_1 \psi(-x) + C_2 \psi^*(-x),$$

$$\Psi(x>0) = C_\alpha \psi(x) + C_\beta \psi^*(x),$$
(B3)

where C_1 , C_2 , C_{α} , and C_{β} are constants to be determined. In the region x < 0, $\psi(-x)$ represents an outgoing particle from the left of the horizon. Implementing the method of complex paths by rotating in the *upper* complex plane, we get

$$C_{\alpha} = C_1 \exp[\pi E/R(0)]. \tag{B4}$$

In the region x < 0, $\psi^*(-x)$ represents an ingoing particle to the left of the horizon. Here again, implementing the method of complex paths by rotating in the *lower* complex plane, we obtain

$$C_{\beta} = C_2 \exp[-\pi E/R(0)].$$
 (B5)

This leaves us with identifying a relation between C_1 and C_2 . In order to do this, let us consider the physical situation of particle production near the black-hole horizon. In this case, a virtual particle-antiparticle pair to the left of the horizon gets converted to a real particle-antiparticle pair to the right of the horizon by the tidal action of the gravitational field, implying that the total current to the left of the horizon is zero. Thus, we get

$$|C_1|^2 = |C_2|^2. (B6)$$

This implies

$$|C_{\beta}|^{2} = |C_{\alpha}|^{2} \exp[-2\pi E/\kappa].$$
 (B7)

Interpreting $|C_{\alpha}|^2 \equiv |\alpha_E|^2$ and $|C_{\beta}|^2 \equiv |\beta_E|^2$ as the Bogoliubov coefficients and using the unitarity condition, we get

$$|\alpha_{E}|^{2} = \frac{1}{1 - \exp[-2\pi E/\kappa]},$$
$$|\beta_{E}|^{2} = \frac{1}{\exp[2\pi E/\kappa] - 1}.$$
(B8)

These are the well known relations for the Bogoliubov coefficients.

- S. Perlmutter *et al.*, Astrophys. J. **483**, 565 (1997); B. Schmidt *et al.*, *ibid.* **507**, 46 (1998); A. J. Riess *et al.*, Astron. J. **116**, 1009 (1998).
- [2] J. Maldacena, Adv. Theor. Math. Phys. 2, 231 (1998); E. Witten, *ibid.* 2, 253 (1998); S. Gubser, I. Klebanov, and A. Polyakov, Phys. Lett. B 428, 105 (1998); O. Aharony, S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, Phys. Rep. 323, 183 (2000).
- [3] E. Witten, "Quantum gravity in de Sitter space," hep-th/0106109.
- [4] A. Strominger, J. High Energy Phys. 10, 034 (2001).
- [5] C. M. Hull, J. High Energy Phys. 07, 021 (1998); Mu-In Park, Phys. Lett. B 440, 275 (1998); Nucl. Phys. B544, 377 (1999);
 I. Antoniadis, P. Mazur, and E. Mottola, astro-ph/9705200; A. Volovich, hep-th/0101176; V. Balasubramanian, P. Horava, and D. Minic, J. High Energy Phys. 05, 043 (2001).
- [6] A. J. M. Medved, Phys. Rev. D 66, 124009 (2002).
- [7] M. Parikh, Phys. Lett. B 546, 189 (2002).
- [8] R. Bousso, A. Maloney, and A. Strominger, Phys. Rev. D 65, 104039 (2002).
- [9] Kyung-Seok Cha, Bum-Hoon Lee, and Chanyong Park, "dS/ CFT Correspondence from the Brick Wall Method," hep-th/0207194.
- [10] S. Nojiri and S. Odintsov, J. High Energy Phys. 12, 033 (2001); Phys. Lett. B 523, 165 (2001).
- [11] M. Parikh and F. Wilczek, Phys. Rev. Lett. 85, 5042 (2000).
- [12] M. Spradlin, A. Strominger, and A. Volovich, "Les Houches Lectures on De Sitter Space," hep-th/0110007.
- [13] G. W. Gibbons and S. W. Hawking, Phys. Rev. D 15, 2738 (1977); W. Hiscock, *ibid.* 39, 1067 (1989); D. Markovic and W. Unruh, *ibid.* 43, 332 (1991); Zhong Chao Wu, Gen. Relativ. Gravit. 32, 1823 (2000).
- [14] T. Padmanabhan, Class. Quantum Grav. 19, 5387 (2002); T.

Padmanabhan, "Entropy and Energy of a Class of Spacetimes with Horizon: A General Derivation," gr-qc/0202080.

- [15] Feng-Li Line and Chopin Soo, Class. Quantum Grav. 16, 551 (1999).
- [16] S. Nojiri and S. D. Odintsov, Phys. Rev. D 59, 044026 (1999);
 A. Bytsenko, S. Nojiri, and S. Odintsov, Phys. Lett. B 444, 121 (1998);
 S. Nojiri and S. D. Odintsov, Int. J. Mod. Phys. A 14, 1293 (1999);
 15, 989 (2000).
- [17] K. Maeda, T. Koike, M. Narita, and A. Ishibashi, Phys. Rev. D 57, 3503 (1998).
- [18] K. Srinivasan and T. Padmanabhan, Phys. Rev. D 60, 24007 (1999).
- [19] S. Shankaranarayanan, K. Srinivasan, and T. Padmanabhan, Mod. Phys. Lett. A 16, 571 (2001); S. Shankaranarayanan, T. Padmanabhan, and K. Srinivasan, Class. Quantum Grav. 19, 2671 (2002).
- [20] S. Shankaranarayanan, Ph.D. thesis, IUCAA, Pune.
- [21] Elias C. Vagenas, hep-th/0111047; Damien A. Easson, J. High Energy Phys. 02, 037 (2003).
- [22] R. Bousso, J. High Energy Phys. 11, 038 (2000); 04, 035 (2001).
- [23] I. D. Novikov, Commun. Shternberg State Astron. Inst. 132, 3-42 (1964).
- [24] R. Bousso and S. W. Hawking, Phys. Rev. D 54, 6312 (1996);
 57, 2436 (1998); J. Niemeyer and R. Bousso, *ibid.* 62, 023503 (2000).
- [25] R. Mann and M. Ghezelbash, J. High Energy Phys. 01, 005 (2002).
- [26] S. Shankaranarayanan and T. Padmanabhan (paper in preparation).
- [27] K. Srinivasan (private communication).