

Can power corrections be reliably computed in models with extra dimensions?

J. F. Oliver, J. Papavassiliou, and A. Santamaria

Departament de Física Teòrica and IFIC, Universitat de València-CSIC, Dr. Moliner 50, E-46100 Burjassot (València), Spain

(Received 11 February 2003; published 11 June 2003)

We critically reexamine the issue of power-law running in models with extra dimensions. The analysis is carried out in the context of a higher-dimensional extension of QED, with the extra dimensions compactified on a torus. It is shown that a naive β function, which simply counts the number of modes, depends crucially on the way the thresholds of the Kaluza-Klein modes are crossed. To solve these ambiguities we turn to the vacuum polarization, which, due to its special unitarity properties, guarantees the physical decoupling of the heavy modes. This latter quantity, calculated in the context of dimensional regularization, is used for connecting the low energy gauge coupling with the coupling of the D -dimensional effective field theory. We find that the resulting relation contains only logarithms of the relevant scales, and no power corrections. If, instead, hard cutoffs are used to regularize the theory, one finds power corrections, which could be interpreted as an additional matching between the effective higher-dimensional model and some unknown, more complete theory. The possibility of estimating this matching is examined in the context of a toy model. The general conclusion is that, in the absence of any additional physical principle, the power corrections depend strongly on the details of the underlying theory. Possible consequences of this analysis for gauge coupling unification in theories with extra dimensions are briefly discussed.

DOI: 10.1103/PhysRevD.67.125004

PACS number(s): 11.10.Kk, 11.10.Hi, 12.10.Dm, 12.10.Kt

I. INTRODUCTION

The study of models with extra dimensions has received a great deal of attention recently [1–4], mainly because of the plethora of theoretical and phenomenological ideas associated with them, and the flexibility they offer for realizing new, previously impossible, field-theoretic constructions. One of the most characteristic features of such models is that of “early unification:” the running of gauge couplings is supposed to be modified so strongly by the presence of the tower of Kaluza-Klein (KK) modes that instead of being logarithmic it becomes linear, quadratic, etc., depending on the number of extra dimensions [5–21]. Specifically, it has been widely argued that the gauge couplings run as μ^δ , where δ is the number of compact extra dimensions. Thus, if the extra dimensions are sufficiently large, such behavior of the couplings could allow for their unification at accessible energies, of the order of a few TeV, clearly an exciting possibility.

The assertion that gauge couplings display power-law running is based on rather intuitive arguments: In modified minimal subtraction ($\overline{\text{MS}}$) schemes the QED β function is proportional to the number of “active” flavors, namely the number of particles lighter than the renormalization scale. Using this argument, and just counting the number of modes lighter than μ , one easily finds that the “ β function” of QED in models with extra dimensions grows as μ^δ . This behavior is also justified by explicit calculations of the vacuum polarization of the photon using hard cutoffs; since the cutoff cannot be removed, due to the non-renormalizability of the theory, it is finally identified with the renormalization scale, a procedure which eventually leads to a similar conclusion [6,7] (but with the final coefficient adjusted by hand in order to match the naive expectation in $\overline{\text{MS}}$).

Even though these arguments are plausible, the importance of their consequences requires that they should be scrutinized more carefully [22]. In particular, the argument based on $\overline{\text{MS}}$ running is rather tricky. As is well known, the $\overline{\text{MS}}$ scheme, because of its mass independence, does not satisfy decoupling, already at the level of four-dimensional theories. Instead, decoupling has to be *imposed* by hand every time a threshold is passed: one builds an effective theory below the threshold, m , and matches it to the theory above the threshold. This matching is carried out by requiring that some physical amplitude or Green’s function (i.e. the effective charge) is the same when calculated using either theory, at energies where *both* theories are reliable, namely at Q^2 much below the threshold. Then, since the renormalization scale, μ , is still a free parameter, one chooses μ around m , in order to avoid large logarithms in the matching equations. In the case of gauge couplings and $\overline{\text{MS}}$ schemes with $\text{Tr}\{I_{\text{Dirac}}\} = 4$ one finds (at one loop) that gauge couplings are continuous at $\mu = m$. This statement is, however, extremely scheme dependent: just by choosing $\text{Tr}\{I_{\text{Dirac}}\} = 2^{D/2}$ it is completely modified (see for instance [23]). In addition to these standard ambiguities, a new complication arises in the context of higher-dimensional models. In particular, the aforementioned procedure requires that the different scales be widely separated in order to avoid the result that higher dimension operators, generated in the process of matching, become important. However, the condition of having well-separated thresholds is rather marginally satisfied in the case of an infinite tower of KK modes with $M_n = nM_c$ (M_c is the compactification scale). In fact, as we will see in detail later, the results obtained for a β function that just counts the number of active modes depend very strongly on the prescription chosen for the way the various thresholds are crossed.

As has been hinted above, the deeper reason behind these additional types of ambiguities is the fact that gauge theories

in more than 4 dimensions, compactified or not, are not renormalizable. At the level of the 4-dimensional theory with an infinite number of KK modes the non-renormalizability manifests itself by the appearance of extra divergences, encountered when summing over all the modes. If the theory is not compactified the non-renormalizability is even more evident, since gauge couplings in theories with δ extra dimensions have dimension $1/M^{\delta/2}$. Therefore, gauge theories in extra dimensions should be treated as effective field theories (EFT's). Working with such theories presents several difficulties, but, as we have learned in recent years, they can also be very useful. In the case of quantum field theories in extra dimensions, there is no alternative: basic questions, such as the calculation of observables or the unification of couplings, can only be addressed in the framework of EFT's. However, before attempting to answer specific questions related to the running of couplings in the extra-dimensional theories, one should first clarify the type of EFT one is going to use, since there are, at least, two types of EFT [24]: In one type, known as the ‘‘Wilsonian EFT’’ (WEFT) [25], one keeps only momenta below some scale Λ , while all the effects of higher momenta or heavy particles are encoded in the couplings of the effective theory. This method is very intuitive and leads, by definition, to finite results at each step; however, the presence of the cutoff in all expressions makes the method cumbersome to use, and in the particular case of gauge theories difficult to reconcile with gauge invariance. The WEFT approach has already been applied to the problem of running of couplings in theories with compact extra dimensions, but only for the case of scalar theories [21]. Within the context of another type of EFT, often termed ‘‘continuum effective field theories’’ (CEFT's) (see for instance [24,26–31]), one allows the momenta of particles to vary up to infinity, but heavy particles are removed from the spectrum at low energies. As in the WEFT case the effects of heavier particles are absorbed into the coefficients of higher dimension operators. Since the momenta are allowed to be infinite, divergences appear, and therefore the CEFT needs to undergo both regularization and renormalization. In choosing the specific scheme for carrying out the above procedures particular care is needed. Whereas in principle one could use any scheme, experience has shown that the most natural scheme for studying the CEFT's is dimensional regularization with minimal subtraction [24,26–31]. CEFT's are widely used in physics: for example, when in the context of QCD one talks about 3, 4 or 5 active flavors, one is implicitly using this latter type of effective theory [27,32]. Moreover, most of the analyses of grand unification [28,33] resort to CEFT-type constructions: one has a full theory at the grand unified theory (GUT) scale, then an effective field theory below the GUT scale standard model or minimal supersymmetric standard model is built, and then yet another effective field theory below the Fermi scale (just QED + QCD). In these cases the complete theory is known, and the CEFT language is used only in order to simplify the calculations at low energies and to control the large logarithms which appear when there are widely separated scales. Nevertheless, CEFT's are useful even when the complete theory is not known, or when the connection with the complete theory cannot be worked

out; this is the case of chiral perturbation theory (χ PT) [34–37] (for more recent reviews see also [29,38,39]).

It is important to maintain a sharp distinction between the two types of EFT mentioned above, i.e. Wilsonian or continuum, because conceptually they are quite different. However, perhaps due to the fact that the language is in part common to both types of theories, it seems that they are often used interchangeably in the literature, especially when employing cutoffs within the CEFT framework. In particular, since the couplings α_i have dimensions $[\alpha_i]=M^{-n}$, when computing loops one generally obtains effects which grow as $(\Lambda^n \alpha_i)^m$, where Λ is the formal CEFT cutoff, and as such is void of physics. As a consequence, physical observables should be made as independent of these cutoffs as possible by introducing as many counterterms as needed to renormalize the answer. Not performing these renormalizations correctly, or identifying naively formal cutoffs with the physical cutoffs of the effective theory, can lead to completely nonsensical results (see for instance [40,41]). This type of pitfall may be avoided by simply using dimensional regularization, since the latter has the special property of not mixing operators with different dimensionalities.

The usual way to treat theories with compactified extra dimensions is to define them as a 4-dimensional theory with a truncated tower of KK modes at some large but otherwise arbitrary N_s , a procedure which effectively amounts to using a hard cutoff in the momenta of the extra dimensions. Thus, physical quantities calculated in this scheme depend explicitly on the cutoff N_s , which is subsequently identified with some physical cutoff. However, as already commented, N_s plays the role of a formal cutoff, and is therefore plagued with all the aforementioned ambiguities. Identification of this formal cutoff with a universal physical cutoff can give the illusion of predictability, making us forget that we are dealing with a non-renormalizable theory with an infinite number of parameters, which can be predictive only at low energies, where higher dimension operators may be neglected.

In this paper we want to analyze the question of the running of gauge couplings in theories with compact dimensions from the CEFT ‘‘canonical’’ point of view. We hasten to emphasize that even the CEFT presents conceptual problems in theories with compactified dimensions. Specifically, as mentioned above, in the CEFT approach the (virtual) momenta are allowed to vary up to infinity; however, momenta related to the compactified extra dimensions turn out to be KK masses in the 4-dimensional compactified theory, where it is supposed that one keeps only particles lighter than the relevant scale. Thus, truncating the KK series amounts to cutting off the momenta of the compactified dimensions. Therefore, in order to define a true ‘‘non-cutoff’’ CEFT scheme we are forced to keep all KK modes. Our main motivation is to seriously explore this approach, and investigate both its virtues and its limitations for the problem at hand. We hope that this study will help us identify more clearly which quantities can and which cannot be computed in effective extra-dimensional theories.

The paper is organized as follows. In Sec. II we discuss the usual arguments in favor of power-law running of gauge couplings and show that they depend crucially on the way

KK thresholds are crossed. In particular we show that a one-loop β function which simply counts the number of modes diverges for more than 5 dimensions, if the physical way of passing thresholds dictated by the vacuum polarization function (VPF) is imposed.

In Sec. III we introduce a theory with one fermion and one photon in $4 + \delta$ dimensions, with the extra δ ones compactified. This theory, which is essentially QED in $4 + \delta$ dimensions, serves as a toy model for studying the issue of power corrections and the running of the coupling in a definite framework.

In Sec. IV we study the question of decoupling KK modes in the aforementioned theory by analyzing the behavior of the VPF of the (zero-mode) photon. Since, as commented above, decoupling the KK modes one by one is problematic, we study the question of how to decouple all of them at once. To accomplish this we consider the VPF of the photon with all KK modes included, and study how it reduces at $Q^2 \ll M_c$ to the standard QED VPF with only one light mode. Since the entire KK tower is kept untruncated, the theory is of course non-renormalizable; therefore, to compute the VPF we have to regularize and renormalize it in the spirit of the CEFT, in a similar way that observables are defined in χ PT. As in χ PT, it is most convenient to use dimensional regularization with minimal subtraction, in order to maintain a better control on the mixing among different operators. However, at the level of the 4-dimensional theory the non-renormalizability manifest itself through the appearance of divergent sums over the infinite KK modes, and dimensional regularization does not seem to help in regularizing them. The dimensional regularization of the VPF is eventually accomplished by exploiting the fact that its UV behavior coincides with that found when the δ extra dimensions have not been compactified.¹ To explore this point we first resort to the standard unitarity relation (optical theorem), which relates the imaginary part of the VPF to the total cross section in the presence of the KK modes; the latter is finite because the phase space truncates the series. For $Q^2 \gg M_c^2$ the uncompactified result for the imaginary part of the VPF is rapidly reached, i.e. after passing a few thresholds. We then compute the real part of the one-loop VPF in the non-compact theory in $4 + \delta$ dimensions, where, of course, we can directly use dimensional regularization to regularize it (since no KK reduction has taken place). For later use we also present results in which the same quantity is evaluated by using hard cut-offs. Finally, we show that the UV divergences of the one-loop VPF are indeed the same in both the (torus)-compactified and uncompactified theories. Therefore, in order to regularize the VPF in the compactified theory with an infinite number of KK modes it is sufficient to split the VPF into two pieces, an ‘‘uncompactified’’ piece, corresponding to the case where the extra dimensions are treated on the same footing as the four usual ones, and a piece which

contains all compactification effects. We show that this latter piece is UV and IR finite and proceed to evaluate it, while all UV divergences remain in the former, which we evaluate using dimensional regularization.

The results of previous sections are used in Sec. V to define an effective charge $\alpha_{\text{eff}}(Q)$ which can be continuously extrapolated from $Q^2 \ll M_c$ to $Q^2 \gg M_c$. We use this effective charge to study the matching of couplings in the low energy effective theory (QED) to the couplings of the theory containing an infinite of KK modes. In the context of dimensional regularization we find that this matching contains only the standard logarithmic running from m_Z to the compactification scale M_c , with no power corrections. On the other hand, if hard cutoffs are used to regularize the VPF in the non-compact space, one does find power corrections, which may be interpreted as an additional matching between the effective $D = 4 + \delta$ dimensional field theory and some more complete theory. We discuss the possibility of estimating this matching in the EFT without knowing the details of the full theory. This point is studied in a simple extension of our original toy model, by endowing the theory considered (QED in $4 + \delta$ compact dimensions) with an additional fermion with mass $M_f \gg M_c$, which is eventually integrated out.

II. CROSSING THRESHOLDS

The simplest argument (apart from the purely dimensional ones) in favor of power-law running in theories with extra dimensions is based on the fact that in $\overline{\text{MS}}$ -like schemes the β function is proportional to the number of active modes. Theories with δ extra compact dimensions contain, in general, a tower of KK modes. In particular, if we embed QED in extra dimensions we find that electrons (also photons) have a tower of KK modes with masses $M_n^2 = (n_1^2 + n_2^2 + \dots + n_\delta^2)M_c^2$ with n_i integer values and $M_c = 1/R_c$ the compactification scale. The exact multiplicity of the spectrum depends on the details of the compactification procedure (torus, orbifold, etc.). As soon as we cross the compactification scale, the KK modes begin to contribute, and therefore one expects that the β function of this theory will start to receive additional contributions from them. In a general renormalization scheme satisfying decoupling one can naively write

$$\beta = \sum_n \beta_0 f\left(\frac{\mu}{M_n}\right), \quad (2.1)$$

where μ is the renormalization scale, β_0 is the contribution of a single mode, and $f(\mu/M)$ is a general step function that decouples the modes as μ crosses the different thresholds, namely $f(\mu/M) \rightarrow 0$, $\mu \ll M$, and $f(\mu/M) \rightarrow 1$, $\mu \gg M$. For instance in $\overline{\text{MS}}$ schemes $f(\mu/M) \equiv \theta(\mu/M - 1)$ where $\theta(x)$ is the step function. Then one finds $[\Omega_\delta = 2\pi^{\delta/2}/\Gamma(\delta/2)]$

$$\beta = \sum_{n < \mu/M_c} \beta_0 \approx \beta_0 \int d\Omega_\delta g^{n^{\delta-1}} dn = \beta_0 \frac{\Omega_\delta}{\delta} \left(\frac{\mu^2}{M_c^2}\right)^{\delta/2}. \quad (2.2)$$

¹This is in a way expected, since for very large $Q^2 \gg M_c = 1/R_c$ the compactification effects should be negligible. Note, however, that this is not always the case; a known exception is provided by the orbifold compactification [22].

This argument, simple and compelling as it may seem, cannot be trusted completely because in $\overline{\text{MS}}$ schemes the decoupling is put in by hand. Therefore, other types of schemes, in which decoupling seems natural, have been studied in the literature. For instance, in Ref. [7] the VPF of the photon at $Q^2=0$ was calculated in the presence of the infinite tower of KK modes by using a hard cutoff in proper time, and the result was used to compute the β function; in that case the modes decouple smoothly. In addition, after adjusting the cutoff by hand one can reproduce the aforementioned result obtained in $\overline{\text{MS}}$. One can easily see that this procedure is equivalent to the use of the function $f(\Lambda/M) \equiv e^{-M_n^2/\Lambda^2}$ to decouple the KK modes:

$$\beta = \sum_n \beta_0 e^{-M_n^2/\Lambda^2} \approx \beta_0 \left(\pi \frac{\Lambda^2}{M_c^2} \right)^{\delta/2}. \quad (2.3)$$

If one chooses by hand $\mu^\delta = \Gamma(1 + \delta/2)\Lambda^\delta$, the sum in Eq. (2.3) agrees exactly with the sum obtained if one uses a sharp step function. Even though this particular way of decoupling KK modes appears naturally in some string scenarios [4,42–44], it hardly appears compelling from the field theory point of view; this procedure is not any better conceptually than the sharp step-function decoupling of modes: one obtains a smooth β function because one uses a smooth function to decouple the KK modes.

These two ways of decoupling KK modes, due to the very sharp step-like behavior they impose, lead to a finite result in Eq. (2.1) for any number of extra dimensions. One is tempted to ask, however, what would happen if one were to use a more physical way of passing thresholds. In fact, heavy particles decouple naturally and smoothly in the VPF, because they cannot be produced physically. Specifically, in QED in 4 dimensions at the one-loop level, the imaginary part, $\Im\Pi(q^2)$, of the VPF $\Pi(q^2)$ is directly related, via the optical theorem, to the tree level cross sections σ for the physical processes $e^+e^- \rightarrow f^+f^-$, by

$$\Im\Pi(s) = \frac{s}{2} \sigma(e^+e^- \rightarrow f^+f^-). \quad (2.4)$$

Given a particular contribution to the spectral function $\Im\Pi(s)$, the corresponding contribution to the renormalized vacuum polarization function $\Pi_R(q^2)$ can be reconstructed via a once-subtracted dispersion relation. For example, for the one-loop contribution of the fermion f , choosing the on-shell renormalization scheme, one finds (if q is the physical momentum transfer with $q^2 < 0$, as usual we define $Q^2 \equiv -q^2$)

$$\begin{aligned} \Pi_R(Q) &= Q^2 \int_{4m_f^2}^{\infty} ds \frac{1}{s(s+Q^2)} \frac{1}{\pi} \Im\Pi(s) \\ &= \frac{\alpha}{\pi} \times \begin{cases} \frac{1}{15} \frac{Q^2}{m_f^2} + \mathcal{O}\left(\frac{Q^4}{m_f^4}\right), & Q^2/m_f^2 \rightarrow 0, \\ \frac{1}{3} \ln\left(\frac{Q^2}{m_f^2}\right) - \frac{5}{9} + \mathcal{O}\left(\frac{m_f^2}{Q^2}\right), & Q^2/m_f^2 \rightarrow \infty, \end{cases} \end{aligned} \quad (2.5)$$

where $\alpha \equiv e^2/(4\pi)$. The above properties can be extended to the QCD effective charge [45], with the appropriate modifications to take into account the non-Abelian nature of the theory, and provide a physical way for computing the matching equations between couplings in QCD at quark mass thresholds. One computes the VPF of QCD with n_f flavors and that of QCD with n_f-1 flavors, and requires that the effective charge is the same for $Q^2 \ll m_f$ in the two theories. This procedure gives the correct relation between the couplings in the two theories [23,46,47]. However, one can easily see that this cannot work for more than one extra dimension. To see that, let us consider the decoupling function $f(\mu/M)$ provided by the one-loop VPF, which, as explained, captures correctly the physical thresholds. The corresponding $f(\mu/M)$ may be obtained by differentiating $\Pi_R(Q)$ once with respect to Q^2 ; it is known [48] that the answer can be well approximated by a simpler function of the form $f(\mu/M) = \mu^2/(\mu^2 + 5M^2)$. We see immediately that if we insert this last function in Eq. (2.1) and perform the sum over all KK modes the result is convergent only for one extra dimension (with a coefficient which is different from the one obtained with the renormalization schemes mentioned earlier), while it becomes highly divergent for several extra dimensions. We conclude therefore that the physical way of decoupling thresholds provided by the VPF seems to lead to a divergent β function in more than one extra dimension. As we will see, this is due to the fact that, in order to properly define the one-loop VPF for $\delta > 1$, more than one subtraction is needed.

III. A TOY MODEL

To be definite we will consider a theory with one fermion and one photon in $4 + \delta$ dimensions, in which the δ extra dimensions are compactified on a torus of equal radii $R_c \equiv 1/M_c$. The Lagrangian is given by

$$\mathcal{L}_\delta = -\frac{1}{4} F^{MN} F_{MN} + i\bar{\psi} \gamma^M D_M \psi + \mathcal{L}_{\text{ct}}, \quad (3.1)$$

where $M=0, \dots, 3, \dots, 3+\delta$. We will also use greek letters to denote four-dimensional indices $\mu=0, \dots, 3$. $D_M = \partial_M - ie_D A_M$ is the covariant derivative with e_D the coupling in $4 + \delta$ dimensions which has dimension $[e_D] = 1/M^{\delta/2}$. After compactification, the dimensionless gauge coupling in four dimensions, e_4 , and the dimensionful $4 + \delta$ coupling are related by the compactification scale²

$$e_4 = e_D \left(\frac{M_c}{2\pi} \right)^{\delta/2}. \quad (3.2)$$

Evidently e_D is determined from the four-dimensional gauge coupling and the compactification scale, but in the uncompactified space we can regard it as a free parameter (like f_π in χ PT). Finally \mathcal{L}_{ct} represents possible gauge invariant op-

²Note that the factors 2π depend on the exact way the extra dimensions are compactified (on a circle, orbifold, etc.).

erators with dimension $2 + D$ or higher, which are in general needed for renormalizing the theory; they can be computed only if a more complete theory, from which our effective theory originates, is given. For instance, by computing the VPF we will see that an \mathcal{L}_{ct} of the form

$$\mathcal{L}_{\text{ct}} = \frac{c_1}{M_s^2} D_M F^{MK} D^N F_{NK} + \dots \quad (3.3)$$

is needed to make it finite.

The spectrum after compactification contains a photon (the zero mode of the four-dimensional components of the gauge boson), the δ extra components of the gauge boson remaining in the spectrum as δ massless real scalars, a tower of massive vector bosons with masses $M_n^2 = (n_1^2 + n_2^2 + \dots + n_\delta^2) M_c^2$, $n_i \in \mathbb{Z}$, $n_i \neq 0$, $2^{\lfloor \delta/2 \rfloor}$ massless Dirac fermions (here the symbol $\lfloor x \rfloor$ represents the closest integer to x smaller than or equal to x), and a tower of massive Dirac fermions with masses also given by the above mass formula. Note that this theory does not lead to normal QED at low energies, first because the δ extra components of the gauge boson remain in the spectrum, and second because in $4 + \delta$ dimensions the fermions have $4 \cdot 2^{\lfloor \delta/2 \rfloor}$ components, which remain as zero modes, leading at low energy to a theory with $2^{\lfloor \delta/2 \rfloor}$ Dirac fermions. In the $D = 4 + \delta$ theory these will arise from the trace of the identity of the γ matrices, which just counts the number of components of the spinors. To obtain QED as a low energy theory one should project out the correct degrees of freedom by using some more appropriate compactification (for instance, orbifold compactifications can remove the extra components of the photon from the low energy spectrum, and leave just one Dirac fermion). However this is not important for our discussion of the VPF; we just have to remember to drop the additional factors $2^{\lfloor \delta/2 \rfloor}$ to make contact with usual QED with only one fermion. Theories of this type, with all particles living in extra dimensions, are called theories with ‘‘universal extra dimensions’’ [49] and have the characteristic that all the effects of the KK modes below the compactification scale cancel at tree level due to the conservation of the KK numbers. In particular, and contrary to what happens in theories where gauge and scalar fields live in the bulk and fermions in the brane [50,51], no divergences associated with summations over KK towers appear at the tree level. Finally, the couplings of the electron KK modes to the standard zero-mode photon are universal and dictated by gauge invariance. The couplings among the KK modes can be found elsewhere [52,53]; they will not be important for the discussion of the VPF that we present here.

IV. THE VACUUM POLARIZATION IN THE PRESENCE OF KK MODES

In this section we study in detail the behavior of the one-loop VPF in the theory defined above for general values of the number δ of extra dimensions. The main problems we want to address are (i) the general divergence structure of the VPF, (ii) a demonstration that it is possible to regulate the UV divergences using dimensional regularization, (iii) the appearance of non-logarithmic (power) corrections, and (iv)

their comparison to the analogous terms obtained when resorting to a hard-cutoff regularization.

A. The imaginary part of the vacuum polarization

One can try to compute the VPF of the zero-mode photon directly in a theory with infinite KK fermionic modes. However, one immediately sees that, in addition to the logarithmic divergences that one finds in QED, new divergences are encountered when summing over the infinite number of KK modes. One can understand the physical origin of these divergences more clearly by resorting to the unitarity relation (here s denotes the center-of-mass energy available for the production process):

$$\begin{aligned} \Im \Pi^{(\delta)}(s) &= \frac{s}{e_4^2} \sum_n \sigma(e^+ e^- \rightarrow f_n^+ f_n^-) \\ &= \frac{\alpha_4}{3} \sum_{n < n_{\text{th}}} \left(1 + \frac{2M_n^2}{s} \right) \sqrt{1 - 4M_n^2/s}, \end{aligned} \quad (4.1)$$

where $n < n_{\text{th}}$ represents the sum over all the electron KK modes such that $4(n_1^2 + n_2^2 + \dots + n_\delta^2) M_c^2 < s$, and $\alpha_4 = e_4^2/(4\pi)$. This sum can be evaluated approximately for $s \gg M_c^2$ by replacing it by an integral; then we obtain

$$\Im \Pi^{(\delta)}(s) \approx \frac{\alpha_4}{2^{3+\delta}} \frac{(\delta+2)\pi^{(\delta+1)/2}}{\Gamma((\delta+5)/2)} \left(\frac{s}{M_c^2} \right)^{\delta/2}. \quad (4.2)$$

It turns out that this last result captures the behavior of the same quantity when the extra dimensions are not compact; this is so because, at high energies, the effects of the compactification can be neglected. In fact, this result may be deduced on simple dimensional grounds: as commented, the gauge coupling in $4 + \delta$ dimensions has dimension $1/M^{\delta/2}$; therefore one expects that $\Im \Pi^{(\delta)}(s)$ will grow with s as $(s/M_c^2)^{\delta/2}$, which is what we obtained from the explicit calculation. To see how rapidly one reaches this regime we can plot the exact result of $\Im \Pi^{(\delta)}(s)$ together with the asymptotic value. As we can see in Fig. 1, the asymptotic limit is reached very fast, especially for higher dimensions. For practical purposes one can reliably use the asymptotic value soon after passing the first threshold, $Q > 2M_c$, incurring errors which are below 10%.

Now we can try to obtain the real part by using a dispersion relation like the one used in 4-dimensional QED, i.e. Eq. (2.5). However, one immediately sees that it will need a number of subtractions which depends on the value of δ . Thus, for just one extra dimension, as in 4-dimensional QED, one subtraction is enough, for $\delta=2$ and $\delta=3$ two subtractions are needed, and so on. This just manifests the non-renormalizability of the theory, and in the effective field theory language, the need for higher dimension operators acting as counterterms. Even though this ‘‘absorptive’’ approach is perfectly acceptable, it would be preferable to have a way of computing the real part directly at the Lagrangian




FIG. 1. $\mathfrak{I}\Pi^{(\delta)}(Q)$ as compared with the asymptotic value ($\delta = 1, 2, 3$). Q is given in units of M_c .

level (by computing loops, for instance). As commented in the Introduction, to accomplish this we will use dimensional regularization.

B. The vacuum polarization in uncompactified $4 + \delta$ dimensions

When using dimensional regularization to compute the VPF in uncompactified space, to be denoted Π_{uc} , simple dimensional arguments suggest that one should typically obtain contributions of the form

$$\Pi_{uc}(Q) \propto e_4^2 \left(2\pi \frac{Q}{M_c} \right)^\delta,$$

since the two vertices in the loop provide a factor e_D^2 , whose dimensions must be compensated by the only available scale in the problem, namely Q^2 . In the above formula we have used the relation of Eq. (3.2) in order to trade off e_D for e_4 . The omitted coefficient in front will be generally divergent, and will be regularized by letting $\delta \rightarrow \delta - \epsilon$.

Let us compute the VPF $\Pi_{uc}^{MN}(q)$ in uncompactified space, assuming that, if necessary, the dimensions will be continued to complex values. We have that

$$\Pi_{uc}^{MN}(q) = ie_D^2 \int \frac{d^{4+\delta}k}{(2\pi)^{4+\delta}} \text{Tr} \left\{ \gamma^M \frac{1}{k} \gamma^N \frac{1}{k+q} \right\}, \quad (4.3)$$

which by gauge invariance assumes the standard form

$$\Pi_{uc}^{MN}(q) = (q^2 g^{MN} - q^M q^N) \Pi_{uc}(q).$$

If we now were to use that, in D dimensions, $\text{Tr}[\gamma^M \gamma^N] = 2^{[D/2]} g^{MN}$, we would find that the low energy limit has an extra $2^{[D/2]}$ factor, which, as commented, is an artifact of the torus compactification: there are $2^{[D/2]}$ too many fermions in the theory. Therefore we simply drop this factor by hand. Moreover, we use Eq. (3.2) and employ the proper-time parametrization in intermediate steps, thus arriving at

$$\begin{aligned} \Pi_{uc}(Q) &= \frac{e_4^2}{2\pi^2} \left(\frac{\pi}{M_c^2} \right)^{\delta/2} \int_0^1 dx x(1-x) \\ &\quad \times \int_0^\infty \frac{d\tau}{\tau^{1+\delta/2}} \exp\{-\tau x(1-x)Q^2\} \\ &= \frac{e_4^2}{2\pi^2} \frac{\pi^{\delta/2} \Gamma^2(2 + \delta/2)}{\Gamma(4 + \delta)} \Gamma\left(-\frac{\delta}{2}\right) \left(\frac{Q^2}{M_c^2} \right)^{\delta/2}. \end{aligned} \quad (4.4)$$

A simple check of this result may be obtained by computing its imaginary part. To that end we let $Q^2 \rightarrow -q^2 - i\epsilon$ with $q^2 > 0$. Then

$$\mathfrak{I}\{-q^2 - i\epsilon\}^{\delta/2} = -(q^2)^{\delta/2} \sin \frac{\delta\pi}{2}.$$

Now we can use that $\Gamma(-\delta/2)\Gamma(1 + \delta/2) = -\pi/\sin(\delta\pi/2)$ to write

$$\begin{aligned} \mathfrak{I}\Pi_{uc}(q) &= \alpha_4 \frac{2\pi^{\delta/2} \Gamma^2(2 + \delta/2)}{\Gamma(4 + \delta)\Gamma(1 + \delta/2)} \left(\frac{q^2}{M_c^2} \right)^{\delta/2} \\ &= \frac{\alpha_4}{2^{3+\delta}} \frac{(\delta+2)\pi^{(\delta+1)/2}}{\Gamma((\delta+5)/2)} \left(\frac{q^2}{M_c^2} \right)^{\delta/2} \end{aligned}$$

which agrees with our previous result of Eq. (4.2).

For odd values of δ , the one-loop $\Pi_{uc}(Q)$ computed above is finite, since the $\Gamma(-\delta/2)$ can be calculated by analytic continuation. This result is expected in a way, since in an odd number of dimensions, by Lorentz invariance, there are no appropriate gauge invariant operators able to absorb any possible infinities generated in the one-loop VPF; this would require operators which give contributions that go like Q^δ . Notice, however, that at higher orders $\Pi_{uc}(Q)$ will eventually become divergent. For instance, in five dimensions at two loops, the VPF should go as Q^2 , since there are four elementary vertices. The divergences generated by these con-

tributions could be absorbed in an operator such as the one considered in the previous section, namely $D_M F^{MK} D^N F_{NK}$. On the other hand, when δ is even, $\Gamma(-\delta/2)$ has a pole, and subtractions are already needed at one loop. To compute the divergent and finite parts in a well-defined way we will use dimensional regularization, i.e. we will assume that $\delta \rightarrow \delta - \epsilon$. Notice however that, unlike in 4 dimensions, we do not need to introduce an additional scale at this point, i.e. the equivalent of the 't Hooft mass scale μ : M_c plays the role of μ , and can be used to keep e_4 dimensionless. After expanding in ϵ we find a simple pole accompanied by the usual logarithm:

$$\Pi_{\text{uc}}(Q) \propto \left(\frac{Q^2}{M_c^2} \right)^{\delta/2} \left\{ -\frac{2}{\epsilon} + \ln(Q^2/M_c^2) + \dots \right\}. \quad (4.5)$$

Here the ellipsis represents a finite constant. Now, to renormalize this result we must introduce higher dimension operators (for instance, if $\delta=2$ the operator $D_M F^{MK} D^N F_{NK}$ will do the job) which could absorb the divergent piece. The downside of this, however, is that we also have to introduce an arbitrary counterterm, κ , corresponding to the contribution of the higher dimension operator; thus we obtain a finite quantity proportional to $\log(Q^2/M_c^2) + \kappa$. Note that, since κ is arbitrary, we can always introduce back a renormalization scale and write $\log(Q^2/M_c^2) + \kappa = \log(Q^2/\mu^2) + \kappa(\mu)$ with $\kappa(\mu) = \kappa + \log(\mu^2/M_c^2)$. It is also important to remark that, in the case of an odd number of dimensions, although at one loop we do not need any counterterm to make the VPF finite, higher dimensional operators could still be present and affect its value.

In the case of uncompactified space, it is interesting to compare the above result with that obtained by regularizing the integral using a hard cutoff. To study this it is enough to carry out the integral of Eq. (4.4), with a cutoff in $\tau_0 = 1/\Lambda^2$:

$$\begin{aligned} \Pi_{\text{uc}}(Q) &= \frac{e_4^2}{2\pi^2} \left(\frac{\pi}{M_c^2} \right)^{\delta/2} \int_0^1 dx x(1-x) \\ &\quad \times \int_{\tau_0}^{\infty} \frac{d\tau}{\tau^{1+\delta/2}} \exp\{-\tau x(1-x)Q^2\}. \end{aligned} \quad (4.6)$$

Then, for $\delta=1,2,3$ we obtain

$$\Pi_{\text{uc}}^{(1)}(Q) = \frac{e_4^2}{2\pi^2} \left(-\frac{3\pi^2 Q}{64M_c} + \frac{\sqrt{\pi}Q^2}{15M_c\Lambda} + \frac{\sqrt{\pi}\Lambda}{3M_c} \right), \quad (4.7)$$

$$\Pi_{\text{uc}}^{(2)}(Q) = \frac{e_4^2}{2\pi^2} \left[\frac{\pi\Lambda^2}{6M_c^2} + \frac{\pi Q^2}{30M_c^2} \left(\log(Q^2/\Lambda^2) + \gamma - \frac{77}{30} \right) \right], \quad (4.8)$$

$$\Pi_{\text{uc}}^{(3)}(Q) = \frac{e_4^2}{2\pi^2} \left(\frac{5\pi^3 Q^3}{768M_c^3} - \frac{\pi^{3/2} Q^2 \Lambda}{15M_c^3} + \frac{\pi^{3/2} \Lambda^3}{9M_c^3} \right). \quad (4.9)$$

As we can see, the pieces that are independent of the cutoff are exactly the same ones we obtained using dimensional regularization. But, in addition, we obtain a series of contributions that depend explicitly on the cutoff. For instance we find corrections to the gauge coupling which behave as Λ^δ , and just redefine the gauge coupling we started with [54]. In the case of 5 dimensions we also generate a term linear in Q^2 ; however it is suppressed by $1/\Lambda$, and therefore it approaches zero for large Λ . In the case of 6 dimensions we obtain the same logarithmic behavior we found with dimensional regularization, and the result can be cast in identical form, if the cutoff is absorbed in the appropriate counterterm. For 7 dimensions we also find divergent contributions which go as Q^2 . This means that, when using cutoffs, higher dimension operators in the derivative expansion (e.g. operators giving contributions varying as Q^2 or higher) are necessary to renormalize the theory and must be included. In the case of dimensional regularization this type of operator is not strictly needed at one loop; however, nothing forbids them in the Lagrangian, and they could appear as ‘‘finite counterterms.’’ If one were to identify the Λ in the above expressions with a physical cutoff, one might get the impression that, contrary to the dimensional regularization approach where arbitrary counterterms are needed, one could now obtain all types of contributions with only one additional parameter, namely Λ . This is not true however: the regulator function is arbitrary; we have simply chosen one among an infinity of possibilities. By changing the regulator function we can change the coefficients of the different contributions at will, except for those few contributions that are independent of Λ . These latter are precisely the ones we have obtained by using dimensional regularization. Thus, even when using cutoffs one has to add counterterms from higher dimension operators, absorb the cutoff, and express the result in terms of a series of unknown coefficients. The lesson is that with dimensional regularization we obtain all calculable pieces, while the non-calculable pieces are related to higher dimensional terms in the Lagrangian.

What we will demonstrate next is that the one-loop VPF in the compactified theory on a torus can be renormalized exactly like the VPF in the uncompactified theory; this will allow us to compute it for any number of dimensions, and examine its behavior for large and for small values of the Q^2 .

C. The vacuum polarization in δ compact dimensions

From the four-dimensional point of view the vacuum polarization tensor in the compactified theory is

$$\Pi^{\mu\nu}(q^2) = \sum_n i e_4^2 \int \frac{d^4 k}{(2\pi)^4} \text{Tr} \left\{ \gamma^\mu \frac{1}{\not{k} - m_n} \gamma^\nu \frac{1}{\not{k} + \not{q} - m_n} \right\}$$

with $m_n^2 = (n_1^2 + n_2^2 + \dots + n_\delta^2) M_c^2$; for simplicity we have assumed a common compactification radius $R = 1/M_c$ for all the extra dimensions. The sum over n denotes collectively the sum over all the modes $n_i = -\infty, \dots, +\infty$. The contribution of each mode to this quantity seems quadratically divergent, as in ordinary QED; however, we know that gauge

invariance converts it to only logarithmically divergent. But, in addition, the sum over all the modes makes the above expressions highly divergent. Instead of attempting to compute it directly, we will add and subtract the contribution of the vacuum polarization function of the uncompactified theory in $4 + \delta$ dimensions:

$$\begin{aligned}\Pi^{\mu\nu}(q) &= [\Pi^{\mu\nu}(q) - \Pi_{\text{uc}}^{\mu\nu}(q)] + \Pi_{\text{uc}}^{\mu\nu}(q) \\ &= \Pi_{\text{fin}}^{\mu\nu}(q) + \Pi_{\text{uc}}^{\mu\nu}(q).\end{aligned}\quad (4.10)$$

Here we have already taken into account the relation between the coupling in $4 + \delta$ dimensions and the four-dimensional coupling and have restricted the external Lorentz indices to the 4-dimensional ones. Depending on the value of δ the vacuum polarization can be highly divergent (naively as $\Lambda^{\delta+2}$, and after taking into account gauge invariance as Λ^δ). However, we can use dimensional regularization (or any other regularization scheme) to make it finite. The important point is that the quantity $\Pi_{\text{fin}}^{\mu\nu}(Q)$ is UV and IR finite and can be unambiguously computed.

Instead of doing the two calculations from scratch, we will do the following:

(i) We will first compute the compactified expression by using Schwinger's proper time, τ , to regularize the UV divergences.

(ii) We will show that the UV behavior of the compactified theory, $\tau \rightarrow 0$, is just the behavior of the uncompactified theory.

(iii) Therefore, to compute $\Pi_{\text{fin}}^{\mu\nu}(Q)$ it is sufficient to compute $\Pi^{\mu\nu}(Q)$ and then subtract its most divergent contribution when $\tau \rightarrow 0$. We will see that it is sufficient to make it finite.

After a few manipulations $\Pi^{\mu\nu}(Q)$ can be written as

$$\Pi^{\mu\nu}(q) = (q^2 g^{\mu\nu} - q^\mu q^\nu) \Pi(q),$$

where [7]

$$\begin{aligned}\Pi(Q) &= \frac{e_4^2}{2\pi^2} \sum_n \int_0^1 dx x(1-x) \\ &\quad \times \int_0^\infty \frac{d\tau}{\tau} \exp\{-\tau[x(1-x)Q^2 + m_n^2]\}.\end{aligned}$$

$\Pi(Q)$ can be written in terms of the function

$$\bar{\theta}_3(\tau) \equiv \sum_{n=-\infty}^{+\infty} e^{-n^2\tau} = \sqrt{\frac{\pi}{\tau}} \bar{\theta}_3\left(\frac{\pi^2}{\tau}\right)$$

as

$$\begin{aligned}\Pi(Q) &= \frac{e_4^2}{2\pi^2} \int_0^1 dx x(1-x) \int_0^\infty \frac{d\tau}{\tau} \\ &\quad \times \exp\left\{-\tau x(1-x) \frac{Q^2}{M_c^2}\right\} \bar{\theta}_3^\delta(\tau),\end{aligned}$$

where we have rescaled τ in order to remove M_c from the $\bar{\theta}_3(\tau)$ function. This last expression for $\Pi(Q)$ is highly divergent in the UV ($\tau \rightarrow 0$), because in that limit the $\bar{\theta}_3(\tau)$ function goes as $\sqrt{\pi/\tau}$. Then, if we define, as in Eq. (4.10), $\Pi_{\text{fin}} = \Pi - \Pi_{\text{uc}}$, we have

$$\begin{aligned}\Pi_{\text{fin}}(Q) &= \frac{e_4^2}{2\pi^2} \int_0^1 dx x(1-x) \int_0^\infty \frac{d\tau}{\tau} \exp\left\{-\tau x(1-x) \frac{Q^2}{M_c^2}\right\} \\ &\quad \times \left[\bar{\theta}_3^\delta(\tau) - \left(\frac{\pi}{\tau}\right)^{\delta/2} \right],\end{aligned}$$

which is completely finite for any number of dimensions. In fact, the last term provides a factor

$$F_\delta(\tau) \equiv \bar{\theta}_3^\delta(\tau) - \left(\frac{\pi}{\tau}\right)^{\delta/2} \xrightarrow{\tau \rightarrow 0} 2\delta \left(\frac{\pi}{\tau}\right)^{\delta/2} \exp\left\{-\frac{\pi^2}{\tau}\right\}$$

that makes the integral convergent in the UV, while for large τ this function goes to 1 quite fast. In this region the integral is cut off by the exponential of momenta; so we can think of the exponential $\exp\{-\tau x(1-x)Q^2/M_c^2\}$ as providing a cutoff for $\tau > 4M_c^2/Q^2$, and $F_\delta(\tau)$ as providing a cutoff for $\tau < \pi^2$. With this in mind, we can estimate $\Pi_{\text{fin}}(Q)$ as

$$\begin{aligned}\Pi_{\text{fin}}(Q) &\approx \frac{e_4^2}{2\pi^2} \sum_n \int_0^1 dx x(1-x) \int_{\pi^2}^{4M_c^2/Q^2} \frac{d\tau}{\tau} \\ &= -\frac{e^2}{2\pi^2} \frac{1}{6} \log \frac{Q^2 \pi^2}{4M_c^2}, \quad Q^2 < 4M_c^2/\pi^2;\end{aligned}\quad (4.11)$$

it is just the ordinary running of the zero mode. As Q^2 grows, the upper limit of integration is smaller than the lower limit, and then we expect that $\Pi_{\text{fin}}(Q)$ should vanish. In that region $\Pi(Q)$ will be dominated completely by $\Pi_{\text{uc}}(Q)$.

Let us evaluate $\Pi_{\text{fin}}(Q)$ for any number of extra dimensions. To this end we will approximate the function $F_\delta(\tau)$ as follows:

$$F_\delta(\tau) = \begin{cases} 2\delta \left(\frac{\pi}{\tau}\right)^{\delta/2} \exp\left\{-\frac{\pi^2}{\tau}\right\}, & \tau < \pi, \\ 1 + 2\delta \exp\{-\tau\} - \left(\frac{\pi}{\tau}\right)^{\delta/2}, & \tau > \pi. \end{cases}\quad (4.12)$$

The matching point at $\tau = \pi$ makes the function continuous. In Fig. 2 we display the exact function $F_\delta(\tau)$ (solid) and the approximation above (dashed) for $\delta = 1, 2, 3$. The approximation is very good except in a small region around the matching point $\tau = \pi$. This can be further improved by adding more terms from the expansions of the $\bar{\theta}(\tau)$ functions.

The approximate expression of Eq. (4.12) can be used to obtain semianalytical expansions for $\Pi_{\text{fin}}^\delta(Q)$ for small Q^2 (we define $w \equiv Q^2/M_c^2$):

$$\begin{aligned} \Pi_{\text{fin}}^{(1)}(Q) &= \frac{e_4^2}{2\pi^2} [-0.335 - 0.167\log(w) \\ &\quad + 0.463\sqrt{w} - 0.110w + \dots], \end{aligned} \quad (4.13)$$

$$\begin{aligned} \Pi_{\text{fin}}^{(2)}(Q) &= \frac{e_4^2}{2\pi^2} [-0.159 - 0.167\log(w) \\ &\quad - 0.105w(\log(w) - 1.75) + \dots], \end{aligned} \quad (4.14)$$

$$\begin{aligned} \Pi_{\text{fin}}^{(3)}(Q) &= \frac{e_4^2}{2\pi^2} [-0.0937 - 0.167\log(w) + 0.298w \\ &\quad - 0.202\sqrt{w^3} + \dots]. \end{aligned} \quad (4.15)$$

To see how good these approximate results are, we can compare with the exact results that can be obtained when $\delta = 1$. In this case we have

$$\begin{aligned} \Pi_{\text{fin}}^{(1)}(Q) &= \frac{e_4^2}{2\pi^2} \int_0^1 dx x(1-x) \sum_{n=1}^{\infty} \int_0^{\infty} \frac{d\tau}{\tau} \exp\{-\tau x(1-x)w\} 2 \left(\frac{\pi}{\tau}\right)^{1/2} \exp\left\{-\frac{n^2\pi^2}{\tau}\right\} \\ &= \frac{e_4^2}{2\pi^2} \int_0^1 dx x(1-x) \sum_{n=1}^{\infty} \frac{2}{n} \exp\{-2\pi n\sqrt{x(1-x)}w\} \\ &= \frac{e_4^2}{2\pi^2} \int_0^1 dx x(1-x) (-2\log\{1 - \exp[-2\pi\sqrt{x(1-x)}w]\}). \end{aligned} \quad (4.16)$$

This last expression can be expanded for $w \ll 1$, and the integral over x can then be performed analytically, yielding

$$\begin{aligned} \Pi_{\text{fin}}^{(1)}(Q) &\approx \frac{e_4^2}{2\pi^2} \left(\frac{1}{18} [5 - 6\log(2\pi)] - \frac{1}{6} \log w \right. \\ &\quad \left. + \frac{3\pi^2}{64} \sqrt{w} - \frac{\pi^2}{90} w + \dots \right), \end{aligned} \quad (4.17)$$

which is in excellent agreement with our approximation [see Eq. (4.13)].

Equation (4.16) can also be used to obtain the behavior of $\Pi_{\text{fin}}^{(1)}(Q)$ for $w \gg 1$. In this limit we obtain

$$\Pi_{\text{fin}}^{(1)}(Q) \approx \frac{e_4^2}{2\pi^2} \frac{3\zeta(5)}{\pi^4} \frac{M_c^4}{Q^4}, \quad Q^2 \gg M_c^2.$$

For higher dimensions things are more complicated, but the behavior is the same, and we find




FIG. 2. Exact values of $F_\delta(\tau)$ (solid) as compared with the approximation discussed in the text (dashed) for $\delta=1,2,3$.

$$\Pi_{\text{fin}}^{(\delta)}(Q) \approx \frac{e_4^2}{2\pi^2} \frac{4\delta\Gamma(2+\delta/2)}{\pi^{4+\delta/2}} K_\delta \frac{M_c^4}{Q^4}, \quad Q^2 \gg M_c^2,$$

where K_δ is of the order of unity and is determined numerically [$K_1 = \zeta(5) = 1.037$, $K_2 = 1.165$, $K_3 = 1.244$]. However, since the uncompactified contribution grows as $(Q^2/M_c^2)^{\delta/2}$ it is obvious that the contributions to $\Pi^{(\delta)}(Q)$ from $\Pi_{\text{fin}}^{(\delta)}(Q)$ will be completely irrelevant for $Q^2 \gg M_c^2$.

Adding the finite and the uncompactified contributions we find that for $Q^2 \ll M_c^2$ the uncompactified contribution exactly cancels the corresponding piece obtained from the expansion of $\Pi_{\text{fin}}^{(\delta)}(Q)$ [the \sqrt{w} piece for $\delta=1$, the $w\log(w)$ piece for $\delta=2$, or the $\sqrt{w^3}$ piece for $\delta=3$]. Then, for $Q^2 \ll M_c^2$ and choosing $\mu = M_c$ we finally find

$$\begin{aligned} \Pi^{(\delta)}(Q) &= \frac{e_4^2}{2\pi^2} \left[a_0^{(\delta)} - \frac{1}{6} \log\left(\frac{Q^2}{M_c^2}\right) + a_1^{(\delta)} \frac{Q^2}{M_c^2} + \dots \right], \\ &\quad Q^2 \ll M_c^2, \end{aligned} \quad (4.18)$$

with the coefficients for 1,2 and 3 extra dimensions given by

δ	1	2	3
$a_0^{(\delta)}$	-0.335	-0.159	-0.0937
$a_1^{(\delta)}$	-0.110	0.183	0.298

As we will see below, in general the coefficients $a_1^{(\delta)}$ can be affected by non-calculable contributions from higher di-

mension operators in the effective Lagrangian, which we have not included.

The following comments related to Eq. (4.18), which is valid only in the dimensional regularization scheme we are using, are now in order:

(i) From Eq. (4.18) we see that for small Q^2 , as expected, we recover the standard logarithm with the correct coefficient, independently of the number of extra dimensions. In addition, interestingly enough, we can also compute the constant term. Thus, although the full theory in $4 + \delta$ dimensions is non-renormalizable and highly divergent, the low energy limit of the VPF calculated in our dimensional regularization scheme is actually finite: when seen from low energies the compactified extra dimensions seem to act as an ultraviolet regulator for the theory.

(ii) When the energy begins to grow, we start seeing effects suppressed by Q^2/M_c^2 , which are finite, at one loop, for any number of dimensions except for $\delta=2$. This is so because the gauge couplings have dimensions $1/M^{\delta/2}$, and therefore, the one-loop VPF goes like $1/M^\delta$.

(iii) For $\delta=1$ one finds that, because of gauge and Lorentz invariance, there are no possible counterterms of this dimension. The VPF must be finite, and that is precisely the result one obtains with dimensional regularization. This of course changes if higher loops are considered: for instance, two-loop diagrams go like $1/M^2$, and, in general, we expect that they will have divergences, which, in turn, should be absorbed in the appropriate counterterms. In principle the presence of these counterterms could pollute our result; however, the natural size of these counterterms, arising at two loops, should be suppressed compared to the finite contributions we have computed.

(iv) For $\delta=2$ one finds that the VPF goes as $1/M^2$, already at one loop, and that the result is divergent. The divergences have to be absorbed in the appropriate counterterm coming from higher dimension operators in the higher dimensional theory. The immediate effect of this is that the coefficient of the Q^2 term in $\Pi^{(2)}(Q)$ becomes arbitrary, its value depending on the underlying physics beyond the compactification scale.

(v) For $\delta>2$ all loop contributions to the Q^2 term are finite, simply because of the dimensionality of the couplings. This, however, does not preclude the existence of finite counterterms, which could be generated by physics beyond the compactification scale, that is, contributions from operators suppressed by two powers of the new physics scale like the operator in Eq. (3.3).

For $Q^2 \gg M_c^2$ the full VPF is completely dominated by the uncompactified contribution:

$$\Pi^{(1)}(Q) = -\frac{e_4^2}{2\pi^2} \frac{3\pi^2}{64} \sqrt{\frac{Q^2}{M_c^2}},$$

$$\Pi^{(2)}(Q) = \frac{e_4^2}{2\pi^2} \frac{\pi}{30} \frac{Q^2}{M_c^2} \log\left(\frac{Q^2}{M_c^2}\right),$$

$$Q^2 \gg M_c^2,$$

$$\Pi^{(3)}(Q) = \frac{e_4^2}{2\pi^2} \frac{5\pi^3}{768} \left(\frac{Q^2}{M_c^2}\right)^{3/2}.$$

As before, the VPF could also receive non-calculable contributions from higher dimension operators which we have not included; in fact, for $\delta=2$, these are needed to renormalize the VPF. How large can these non-calculable contributions be? Since our D -dimensional theory is an effective theory valid only for $Q^2 \ll M_s^2$, even above M_c the results will be dominated by the lowest power of Q^2 . In the case of $\delta=1$, the first operator that one can write down goes as Q^2 ; therefore we expect that the one-loop contribution, of order $\sqrt{Q^2}$, that we have computed will completely dominate the result, as long as we do not stretch it beyond the applicability of the effective Lagrangian approach. For $\delta=2$, counterterms are certainly needed at order Q^2 ; still one can hope that the result will be dominated by the logarithm (as happens with chiral logarithms in the χ PT). For $\delta=3$ (and higher), the one loop result grows as $(Q^2)^{\delta/2}$; however there could be operators giving contributions of order Q^2 with unknown coefficients (in fact, although in dimensional regularization those are not needed, they must be included if cutoffs are used to regularize the theory). Therefore, unless for some reason they are absent from the theory, the result will be dominated by those operators.

V. MATCHING OF GAUGE COUPLINGS

Using the VPF constructed in the previous section we can define a higher dimensional analogue of the conventional QED effective charge [55,56], which will enter in any process involving off-shell photons, e.g.

$$\frac{1}{\alpha_{\text{eff}}(Q)} \equiv \frac{1}{\alpha_4} [1 + \Pi^{(\delta)}(Q)]|_{\overline{\text{MS}}_\delta}, \quad (5.1)$$

where $\alpha_4 = e_4^2/(4\pi)$. We remind the reader that e_4 denotes the (dimensionless) coupling of the four-dimensional theory including all KK modes; it is directly related to the gauge coupling in the theory with δ extra dimensions by Eq. (3.2). The subscript $\overline{\text{MS}}_\delta$ means that the VPF has been regularized using dimensional regularization in $D=4+\delta-\epsilon$ dimensions, and that divergences, when present, are subtracted according to the $\overline{\text{MS}}$ procedure.

To determine the relation between α_4 and the low energy coupling in QED, we have to identify the effective charge computed in the compactified theory with the low energy effective charge, at some low energy scale (for instance $Q^2 = m_Z^2 \ll M_c^2$), where both theories are valid. In that limit we can trust our approximate results of Eq. (4.18), and write

$$\frac{1}{\alpha_{\text{eff}}(m_Z)} = \frac{1}{\alpha_4} + \frac{2}{\pi} a_0^{(\delta)} - \frac{2}{3\pi} \log\left(\frac{m_Z}{M_c}\right). \quad (5.2)$$

This equation connects the low energy QED coupling with the coupling in the compactified D -dimensional theory, regularized by dimensional regularization. Note that this equation is completely independent of the way subtractions are performed to remove the poles in $1/\epsilon$. These poles appear only (and only for an even number of dimensions) in the contributions proportional to Q^δ , which vanish for $Q \rightarrow 0$. Equation (5.2) contains, apart from a finite constant, the standard logarithmic running from m_Z to the compactification scale M_c . It is interesting to notice that, in this approach, the logarithm comes from the finite piece, and should therefore be considered as an infrared (IR) logarithm. When seen from scales smaller than M_c , these logarithms appear to have an UV origin, while, when seen from scales above M_c , they appear as having an IR nature.

It is important to emphasize that, in this scheme, the gauge coupling no longer runs above the compactification scale. This seems counter-intuitive, but it is precisely what happens in χ PT when using dimensional regularization: f_π does not run, it just renormalizes higher dimensional operators [26].

Now we can use Eq. (5.2) to write the effective charge at all energies in terms of the coupling measured at low energies:

$$\frac{1}{\alpha_{\text{eff}}(Q)} \equiv \frac{1}{\alpha_{\text{eff}}(m_Z)} + \frac{1}{\alpha_4} [\Pi^{(\delta)}(Q) - \Pi^{(\delta)}(m_Z)] |_{\overline{\text{MS}}_\delta}. \quad (5.3)$$

Note that the last term is independent of α_4 due to the implicit dependence of $\Pi^{(\delta)}$ on it. Equation (5.3) has the form of a momentum-subtracted definition of the coupling; in fact, in four dimensions it is just the definition of the momentum-subtracted running coupling. For $\delta=1$ and at one loop, $\Pi^{(\delta)}(Q) - \Pi^{(\delta)}(m_Z)$ is finite, and $\alpha_{\text{eff}}(Q)$ can still be interpreted as a momentum-subtracted definition of the coupling. For $\delta>1$, however, Eq. (5.3) involves additional subtractions, a fact which thwarts such an interpretation.

For $Q^2 \ll M_c^2$ we can expand $\Pi^{(\delta)}(Q)$ and obtain

$$\frac{1}{\alpha_{\text{eff}}(Q)} \equiv \frac{1}{\alpha_{\text{eff}}(m_Z)} - \frac{2}{3\pi} \log\left(\frac{Q}{m_Z}\right) + \mathcal{O}\left(\frac{Q^2}{M_c^2}\right),$$

which is nothing but the standard expression for the effective charge in QED, slightly modified by small corrections of order Q^2/M_c^2 . However, as soon as Q^2/M_c^2 approaches unity, the effects of the compactification scale start to appear in $\alpha_{\text{eff}}(Q)$, forcing it to deviate dramatically from the logarithmic behavior, as shown in Fig. 3.

The crucial point, however, is that this effective charge *cannot* be interpreted anymore as the running coupling (as can be done in four dimensions) since it may receive contributions from higher dimension operators; in fact some of them are needed to define this quantity properly. These contributions have nothing to do with the gauge coupling which




FIG. 3. The “effective charge” against the energy scale for $\delta = 1$ (solid), $\delta = 2$ (short dash), $\delta = 3$ (long dash).

is defined as the coefficient of the operator F^2 . In particular, one should not use this quantity to study gauge coupling unification. Instead, one could use Eq. (5.2), which relates the coupling measured at low energies with the one appearing in the D -dimensional Lagrangian valid at energies $M_c < Q < M_s$. This relation involves a logarithmic correction, which is the only contribution that can be reliably computed without knowing the physics beyond M_s .

It is instructive to see what happens if instead of dimensional regularization we use hard cutoffs to regularize the uncompactified part of the VPF as in Eqs. (4.6)–(4.9). Then, when using cutoffs, one can define an “effective charge” as in Eq. (5.1):

$$\frac{1}{\alpha_{\text{eff}}(Q)} \equiv \frac{1}{\alpha_4(\Lambda)} [1 + \Pi^{(\delta)}(Q)] |_\Lambda, \quad (5.4)$$

where $\alpha_4(\Lambda)$ now is the coupling constant in the theory regularized with cutoffs and the subscript Λ indicates that the VPF has been regularized with cutoffs. The use of a Λ dependent coupling obviously implies the WEFT formulation, in which the cutoff is not removed from the theory. On the other hand, in the CEFT formulation one should renormalize the coupling constant by adding the appropriate counterterms and then take the limit $\Lambda \rightarrow \infty$. This usually brings in a new scale at which the coupling is defined, and which effectively replaces Λ in the previous equation. Notice also that for $\delta=2$ in Eq. (4.8) there are logarithmic contributions proportional to Q^2 , which cannot be removed when $\Lambda \rightarrow \infty$. The same is true for $\delta>2$, but with dependencies which are proportional to $\Lambda^{(\delta-2)}$. This just manifests the need for higher dimensional operators, as was already clear in the dispersive approach, to define the effective charge properly. As one can see, the full VPF contains a term that goes as Λ^δ and is independent of Q . This piece survives when $Q \rightarrow 0$, and thus we obtain (we assume $m_Z^2 \ll Q^2$)

$$\frac{1}{\alpha_{\text{eff}}(m_Z)} = \frac{1}{\alpha_4(\Lambda)} + \frac{2}{3\pi\delta} \left(\sqrt{\pi} \frac{\Lambda}{M_c} \right)^\delta + \frac{2}{\pi} a_0^{(\delta)} - \frac{2}{3\pi} \log\left(\frac{m_Z}{M_c}\right). \quad (5.5)$$

Since $\alpha_{\text{eff}}(m_Z)$ should be the same in the two schemes, we find the following relation between α_4 and $\alpha_4(\Lambda)$:

$$\frac{1}{\alpha_4} = \frac{1}{\alpha_4(\Lambda)} + \frac{2}{3\pi\delta} \left(\sqrt{\pi} \frac{\Lambda}{M_c} \right)^\delta. \quad (5.6)$$

If one identifies Λ with the onset of a more complete theory beyond the compactification scale, but at which the EFT treatment is still valid, i.e. if one assumes that $\Lambda \sim M_G \ll M_s$, M_G being this new scale, Eq. (5.6) could be reinterpreted as a matching equation between the coupling α_4 of our effective theory and the coupling of the theory at scales M_G , $\alpha_4(M_G)$. Equation (5.6) generically tells us that one expects corrections which go as $(M_G/M_c)^\delta$. However, without knowledge of the full theory beyond M_G , the meaning of M_G [or even $\alpha_4(M_G)$] is unclear. In particular, if the new theory is some grand unified theory in extra dimensions, M_G will be, in general, not just one single mass, but several masses of the same order of magnitude, related by different coefficients. In the case of logarithmic running those coefficients can be neglected, because they give small logarithms next to the large logarithms containing the common scale. However, in the case of contributions which depend on powers of the new physics scale the situation is completely different, and the presence of several masses could completely change the picture of unification. Cutoffs can give an indication of the presence of power corrections, but the coefficients of these corrections cannot be computed without knowing the details of the full theory.

To see this point more clearly, we add to our $4 + \delta$ dimensional theory an additional fermion with mass M_f satisfying $M_s \gg M_f \gg M_c$, such that compactification corrections may be neglected, and compute its effects on the coupling constant for $M_c^2 \ll Q^2 \ll M_f^2$, using dimensional regularization. We have

$$\begin{aligned} \Pi_f^{(\delta)}(Q) &= \frac{e_4^2}{2\pi^2} \left(\frac{\pi}{M_c} \right)^{\delta/2} \Gamma(-\delta/2) \int_0^1 dx x(1-x) \\ &\quad \times [M_f^2 + x(1-x)Q^2]^{\delta/2}. \end{aligned} \quad (5.7)$$

By expanding for $Q^2 \ll M_f^2$ and integrating over x we obtain

$$\Pi_f^{(\delta)}(Q) = \frac{e_4^2}{2\pi^2} \left(\sqrt{\pi} \frac{M_f}{M_c} \right)^\delta \Gamma\left(-\frac{\delta}{2}\right) \left(\frac{1}{6} + \frac{\delta}{60} \frac{Q^2}{M_f^2} \right). \quad (5.8)$$

For odd values of δ we can use the analytic continuation of the Γ function to obtain a finite result. For even values of δ we will allow a slight departure from the integer value in order to dimensionally regularize the integral. Clearly, integrating out the heavy fermion gives power corrections to the gauge coupling. In addition, it also generates contributions to the higher dimension operators, e.g. contributions proportional to Q^2 and higher powers. As can be seen by comparing with Eqs. (4.6)–(4.9) these power corrections are qualitatively similar to those calculated using a hard cutoff. Evidently, in the context of a more complete theory (in this case, given the existence of a heavy fermion), power corrections

may be encountered even if the dimensional regularization is employed. However, as one can easily see by setting, for example, $\delta=1$ in Eq. (5.8), the coefficients of the power corrections obtained knowing the full theory are in general different from those obtained using a hard cutoff, e.g. Eq. (4.7). In fact, no choice of Λ in Eq. (4.7) can reproduce all the coefficients appearing in Eq. (5.8).

The situation is somewhat similar to what happens when χ PT with $SU(2) \otimes SU(2)$ is matched to χ PT with $SU(3) \otimes SU(3)$. In the $SU(2) \otimes SU(2)$ theory, just by dimensional arguments, one can expect corrections like m_K^2/f_π^2 . But can one compute them reliably without even knowing that there are kaons?

VI. CONCLUSIONS

We attempted a critical discussion of the arguments in favor of power-law running of coupling constants in models with extra dimensions. We showed that the naive arguments lead to an arbitrary β function depending on the particular way chosen to cross KK thresholds. In particular, if one chooses the physical way of passing thresholds provided by the vacuum polarization function of the photon, a β function that counts the number of modes is divergent for more than 5 dimensions.

We studied the question of decoupling of KK modes in QED with $4 + \delta$ (compact) dimensions by analyzing the behavior of the VPF of the photon. We computed first the imaginary part of the VPF by using unitarity arguments, and found that it rapidly reaches the value obtained in a non-compact theory (only a few modes are necessary). We also showed that it grows as $(s/M_c^2)^{\delta/2}$, exhibiting clearly the non-renormalizability of theories in extra dimensions. To obtain the full VPF, one can use an appropriately subtracted dispersion relation. Instead, we use the full quantum effective field theory, with the expectation, suggested by the calculation of the imaginary part of the VPF, that the bad UV behavior of the theory is captured by the behavior of the uncompactified theory. To check this idea, we computed the VPF in the uncompactified theory, regularized by dimensional regularization ($\delta \rightarrow \delta - \epsilon$). We found that, after analytical continuation, the one loop VPF is finite, and proportional to Q^δ for an odd number of dimensions, and has a simple pole, proportional to Q^δ , for an even number of dimensions. This result can be understood easily, because there are no possible Lorentz and gauge invariant operators in the Lagrangian able to absorb a term like Q^δ for odd δ . For δ even it shows that higher dimension operators are needed to regularize the theory. As a check we also recovered the imaginary part of the VPF in the limit of infinite compactification radius.

For comparison with other approaches, we also obtained the VPF in the case that a hard cutoff is used to regularize it. We found that the pieces that do not depend on the cutoff are exactly the same as those obtained by dimensional regularization, while the cutoff dependent pieces are arbitrary, and can be changed at will by changing the cutoff procedure.

Next we computed the VPF in the compactified theory, and showed that it can be separated into a UV and an IR

finite contribution and the VPF calculated in the uncompactified theory; as was shown previously, the latter can be controlled using dimensional regularization. The finite part is more complicated, but can be computed numerically for any number of dimensions. Also, some analytical approximations were obtained for the low and the high energy limits ($Q \ll M_c$ and $Q \gg M_c$ respectively). Adding these two pieces, together with the counterterms coming from higher dimension operators, we obtain a finite expression for an effective charge which can be extrapolated continuously from $Q \ll M_c$ to $Q \gg M_c$; however, its value does depend on higher dimension operator couplings.

Decoupling of all KK modes in this effective charge is smooth and physically meaningful, and the low energy logarithmic running is recovered. We use this effective charge to connect the low energy couplings [i.e. $\alpha_{\text{eff}}(m_Z)$] with the coupling of the theory including all KK modes, regularized by dimensional regularization. We find that this matching involves only the standard logarithmic running from m_Z to the compactification scale M_c . In particular, no power corrections appear in this matching. However, if cutoffs are used to regularize the VPF in the non-compact space, one does find power corrections, exactly as expected from naive dimensional analysis. In the EFT language one could interpret these corrections as an additional matching between the ef-

fective D dimensional field theory and some more complete theory. The question is how reliably can this matching be estimated without knowing the complete theory. By adding to our theory an additional fermion with $M_f \gg M_c$, and integrating it out, we argue that power corrections cannot be computed without knowing the details of the complete theory, in which the D dimensional theory is embedded. Some examples in which this matching can, in principle, be computed are some 5D GUT's and string models [8,14,20,57–59], and the recently proposed de-constructed extra dimensions [18,60–63]. For the question of unification of couplings this result seems rather negative, at least when compared with standard grand unified theories, where gauge coupling unification can be tested without knowing their details. Alternatively, one can approach this result from a more optimistic point of view, and regard the requirement of low-energy unification of couplings as a stringent constraint on the possible extra-dimensional extensions of the SM.

ACKNOWLEDGMENTS

This work has been funded by the Spanish MCyT under the Grants BFM2002-00568 and FPA2002-00612; and by the OCyT of the “Generalitat Valenciana” under the Grant GV01-94.

-
- [1] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Rev. D **59**, 086004 (1999).
 - [2] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B **429**, 263 (1998).
 - [3] I. Antoniadis, Phys. Lett. B **246**, 377 (1990).
 - [4] I. Antoniadis and K. Benakli, Phys. Lett. B **326**, 69 (1994).
 - [5] C. P. Bachas, J. High Energy Phys. **11**, 023 (1998).
 - [6] K. R. Dienes, E. Dudas, and T. Gherghetta, Phys. Lett. B **436**, 55 (1998).
 - [7] K. R. Dienes, E. Dudas, and T. Gherghetta, Nucl. Phys. **B537**, 47 (1999).
 - [8] D. Ghilencea and G. G. Ross, Phys. Lett. B **442**, 165 (1998).
 - [9] C. D. Carone, Phys. Lett. B **454**, 70 (1999).
 - [10] A. Delgado and M. Quiros, Nucl. Phys. **B559**, 235 (1999).
 - [11] E. G. Floratos and G. K. Leontaris, Phys. Lett. B **465**, 95 (1999).
 - [12] P. H. Frampton and A. Rasin, Phys. Lett. B **460**, 313 (1999).
 - [13] Z. Kakushadze, Nucl. Phys. **B548**, 205 (1999).
 - [14] Z. Kakushadze and T. R. Taylor, Nucl. Phys. **B562**, 78 (1999).
 - [15] A. Perez-Lorenzana and R. N. Mohapatra, Nucl. Phys. **B559**, 255 (1999).
 - [16] D. Dumitru and S. Nandi, Phys. Rev. D **62**, 046006 (2000).
 - [17] M. Masip, Phys. Rev. D **62**, 065011 (2000).
 - [18] N. Arkani-Hamed, A.G. Cohen, and H. Georgi, “Accelerated Unification,” hep-th/0108089.
 - [19] Z. Berezhiani, I. Gogoladze, and A. Kobakhidze, Phys. Lett. B **522**, 107 (2001).
 - [20] A. Hebecker and A. Westphal, “Power-Like Threshold Corrections to Gauge Unification in Extra Dimensions,” hep-ph/0212175.
 - [21] J. Kubo, H. Terao, and G. Zoupanos, Nucl. Phys. **B574**, 495 (2000).
 - [22] R. Contino, L. Pilo, R. Rattazzi, and E. Trincherini, Nucl. Phys. **B622**, 227 (2002).
 - [23] G. Rodrigo and A. Santamaria, Phys. Lett. B **313**, 441 (1993).
 - [24] H. Georgi, Annu. Rev. Nucl. Part. Sci. **43**, 209 (1993).
 - [25] K. G. Wilson and J. B. Kogut, Phys. Rep. **12**, 75 (1974).
 - [26] S. Weinberg, Physica A **96**, 327 (1979).
 - [27] S. Weinberg, Phys. Lett. **91B**, 51 (1980).
 - [28] L. J. Hall, Nucl. Phys. **B178**, 75 (1981).
 - [29] H. Leutwyler, Ann. Phys. (N.Y.) **235**, 165 (1994).
 - [30] A. V. Manohar, “Effective field theories,” hep-ph/9606222.
 - [31] A. Pich, “Effective Field Theory,” hep-ph/9806303.
 - [32] E. Witten, Nucl. Phys. **B122**, 109 (1977).
 - [33] H. Georgi, H. R. Quinn, and S. Weinberg, Phys. Rev. Lett. **33**, 451 (1974).
 - [34] J. Callan, G. Curtis, S. R. Coleman, J. Wess, and B. Zumino, Phys. Rev. **177**, 2247 (1969).
 - [35] S. R. Coleman, J. Wess, and B. Zumino, Phys. Rev. **177**, 2239 (1969).
 - [36] S. Weinberg, Phys. Rev. **166**, 1568 (1968).
 - [37] J. Gasser and H. Leutwyler, Ann. Phys. (N.Y.) **158**, 142 (1984).
 - [38] A. Pich, Rep. Prog. Phys. **58**, 563 (1995).
 - [39] G. Ecker, Prog. Part. Nucl. Phys. **35**, 1 (1995).
 - [40] C. P. Burgess and D. London, Phys. Rev. Lett. **69**, 3428 (1992).
 - [41] C. P. Burgess and D. London, Phys. Rev. D **48**, 4337 (1993).
 - [42] M. Quiros, J. Phys. G **27**, 2497 (2001).
 - [43] S. Hamidi and C. Vafa, Nucl. Phys. **B279**, 465 (1987).

- [44] L. J. Dixon, D. Friedan, E. J. Martinec, and S. H. Shenker, Nucl. Phys. **B282**, 13 (1987).
- [45] D. Binosi and J. Papavassiliou, “The QCD Effective Charge to All Orders,” hep-ph/0209016.
- [46] W. Bernreuther and W. Wetzel, Nucl. Phys. **B197**, 228 (1982).
- [47] G. Rodrigo, A. Pich, and A. Santamaria, Phys. Lett. B **424**, 367 (1998).
- [48] S. J. Brodsky, M. S. Gill, M. Melles, and J. Rathsmann, Phys. Rev. D **58**, 116006 (1998).
- [49] T. Appelquist, H.-C. Cheng, and B. A. Dobrescu, Phys. Rev. D **64**, 035002 (2001).
- [50] A. Pomarol and M. Quiros, Phys. Lett. B **438**, 255 (1998).
- [51] A. Delgado, A. Pomarol, and M. Quiros, J. High Energy Phys. **01**, 030 (2000).
- [52] J. Papavassiliou and A. Santamaria, Phys. Rev. D **63**, 125014 (2001).
- [53] A. Muck, A. Pilaftsis, and R. Ruckl, Phys. Rev. D **65**, 085037 (2002).
- [54] T. R. Taylor and G. Veneziano, Phys. Lett. B **212**, 147 (1988).
- [55] C. Itzykson and J. B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- [56] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Addison-Wesley, Reading, MA, 1995).
- [57] K. Choi, H. D. Kim, and Y. W. Kim, J. High Energy Phys. **11**, 033 (2002).
- [58] F. Paccetti Correia, M. G. Schmidt, and Z. Tavatkiladze, “Precursors From $S^{(1)}/Z_2 \times Z_2'$ Orbifold GUTs,” hep-ph/0302038.
- [59] F. Paccetti Correia, M. G. Schmidt, and Z. Tavatkiladze, Nucl. Phys. **B649**, 39 (2003).
- [60] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, Phys. Rev. Lett. **86**, 4757 (2001).
- [61] H.-C. Cheng, C. T. Hill, S. Pokorski, and J. Wang, Phys. Rev. D **64**, 065007 (2001).
- [62] S. Pokorski, J. Phys. G **27**, 2479 (2001).
- [63] P. H. Chankowski, A. Falkowski, and S. Pokorski, J. High Energy Phys. **08**, 003 (2002).