

On slow light as a black hole analogue

W. G. Unruh* and R. Schützhold†

Canadian Institute for Advanced Research Cosmology Program, Department of Physics and Astronomy, University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

(Received 7 March 2003; published 3 July 2003)

Although slow light (electromagnetically induced transparency) would seem an ideal medium in which to institute a “dumb hole” (black hole analogue), it suffers from a number of problems. We show that the high phase velocity in the slow light regime ensures that the system cannot be used as an analogue displaying Hawking radiation. Even though an appropriately designed slow-light setup may simulate classical features of black holes—such as horizon, mode mixing, “Bogoliubov” coefficients, etc.—it does not reproduce the related quantum effects.

DOI: 10.1103/PhysRevD.68.024008

PACS number(s): 04.70.Dy, 04.60.-m, 04.80.-y, 42.50.Gy

I. INTRODUCTION

The astonishing ability to slow light to speeds of a few meters per second has been a striking development in quantum optics, see e.g. [1]. The idea to use matter systems as analogues [2] to the (yet unobserved) Hawking effect [3] for black holes¹ has raised the possibility of experimentally testing certain assumptions which enter into those calculations, see e.g. [5]. The dependence of those analogs on the detection of sound waves however causes problems, as the detection technology for light is much more developed than for sound, and finding an optical analog to black holes [6–9] could make the experimental detection of the analog for Hawking radiation easier, cf. [10].

Recently Leonhardt [6,8] has suggested that slow light systems could be used to create such an analog, but that approach has been criticized by one of us [10]. In this paper, we look in more detail at the use of slow light in such an analog, and try to understand in what sense slow light could be used to create an analog for black holes, and why, despite that analog, it will not create the thermal radiation characteristic of the Hawking process.

II. DESCRIPTION OF THE SET-UP

In order to generate slow light, one first chooses an atom with a convenient set of atomic transitions, cf. [1,11]. In particular, a system is chosen with two long lived meta-stable or stable states, and with one state which is coupled to these two states via dipole electromagnetic transitions (Λ -system). Let us call the two lower meta-stable states $|a\rangle$ and $|b\rangle$. The third higher energy state is $|c\rangle$. The two states $|a\rangle$ and $|b\rangle$ are assumed to have energy $-\omega_a$, $-\omega_b$, and $|c\rangle$ has energy zero and decay constant $\Gamma > 0$. (That is, this

higher energy state is assumed to have decay channels other than electromagnetic radiation to the $|a\rangle$ and $|b\rangle$ states.)

The electromagnetic field, which we will assume has a fixed polarization, will be represented by the vector potential A where $E = \partial_t A$ (temporal gauge).

A. Effective Lagrangian

The effective Lagrangian for this system can be written as ($\hbar = c = 1$ throughout)

$$L = \int dx \mathcal{L}^A + \sum_j (L_j^\psi + L_j^{A\psi}), \quad (1)$$

with the usual term governing the dynamics of the electromagnetic field

$$\mathcal{L}^A = \frac{1}{2}[E^2 - B^2] = \frac{1}{2}[(\partial_t A)^2 - (\partial_x A)^2], \quad (2)$$

and the Lagrangian of the atomic states

$$L_j^\psi = i(\psi_{aj}^* \partial_t \psi_{aj} + \psi_{bj}^* \partial_t \psi_{bj} + \psi_{cj}^* \partial_t \psi_{cj}) + \omega_a \psi_{aj}^* \psi_{aj} + \omega_b \psi_{bj}^* \psi_{bj} + i\Gamma \psi_{cj}^* \psi_{cj}, \quad (3)$$

as well as the interaction term in the dipole approximation

$$L_j^{A\psi} = E(x_j)(\epsilon_a \psi_{cj}^* \psi_{aj} + \epsilon_b \psi_{cj}^* \psi_{bj}) + \text{H.c.}, \quad (4)$$

where x_j is the location of the j -th atom. Here the $\psi_{\dots j}$ are the amplitudes for the j -th particle being in the corresponding state $|\text{atom } j\rangle = \psi_{aj}|a\rangle + \psi_{bj}|b\rangle + \psi_{cj}|c\rangle$ and ϵ_a , ϵ_b are the associated dipole transition amplitudes.

In contrast to the usual setup, i.e., a strong control beam and a weak (perpendicular) probe beam, let us assume that there is a strong background counter-propagating electromagnetic field

$$A_0(t, x) = \Omega \left(\frac{\cos \theta}{\epsilon_a \omega_a} e^{i\omega_a(t-x)} + \frac{\sin \theta}{\epsilon_b \omega_b} e^{i\omega_b(t+x)} \right) + \text{H.c.}, \quad (5)$$

i.e., at the resonant frequencies of the two transitions. The mixing angle θ controls the relative strength of the left- and

*Email address: unruh@physics.ubc.ca

†Email address: schuetz@physics.ubc.ca

¹It should be mentioned here that the analogy between the propagation of light in gravitational fields on the one hand and within dielectric and permeable media (moving or at rest) on the other hand in general has been noticed much earlier, see, e.g., [4], i.e., even before the Hawking effect has been found.

right-moving beam and Ω denotes the averaged Rabi frequency.² For a single beam ($\theta=0$ or $\theta=\pi/2$) Ω reduces to the exact Rabi frequency of that beam. The fact that the phase velocity is unity (i.e., the light speed) prefigures the fact that the effective dielectric constant of the atoms is unity at these transition frequencies when the atoms are in the so called “dark state,” cf. [1,11].

In the following we shall assume that we can and are making the rotating wave approximation. One solution, the only (up to an overall phase) nondecaying solution, for the atoms is

$$\begin{aligned}\psi_{aj}^0 &= +e^{i\omega_a(t-x_j)}\sin\theta, \\ \psi_{bj}^0 &= -e^{i\omega_b(t+x_j)}\cos\theta, \\ \psi_{cj}^0 &= 0.\end{aligned}\quad (6)$$

Since the Rabi oscillations between the states $|a\rangle$ and $|c\rangle$ interfere destructively with those between the states $|b\rangle$ and $|c\rangle$ (leading to a vanishing occupation of $|c\rangle$), this solution is called a dark state (no spontaneous emission).

B. Linearization

Let us redefine our electromagnetic field such that

$$\begin{aligned}A(t,x) &= \left(\Omega \frac{\cos\theta}{\epsilon_a\omega_a} + \Phi_a(t,x) \right) e^{-i\omega_a(t-x)} \\ &+ \left(\Omega \frac{\sin\theta}{\epsilon_b\omega_b} + \Phi_b(t,x) \right) e^{-i\omega_b(t+x)} + \text{H.c.},\end{aligned}\quad (7)$$

where we are going to assume that both Φ_b and Φ_a are slowly varying functions of time and space (i.e., beat fluctuations).

Furthermore, let us define

$$\begin{aligned}\psi_{aj} &= (\Psi_{aj} + \sin\theta)e^{i\omega_a(t-x_j)}, \\ \psi_{bj} &= (\Psi_{bj} - \cos\theta)e^{i\omega_b(t+x_j)}, \\ \psi_{cj} &= \Psi_{cj},\end{aligned}\quad (8)$$

where the new variables Ψ are also assumed to be slowly varying.

Substituting into the Lagrangian, retaining only the second order terms³ in the Ψ , Φ_b , Φ_a , using the rotating wave approximation, and neglecting time derivatives of Φ_b and Φ_a with respect to ω_a and ω_b we get the effective (approximated) Lagrangian for the beat fluctuations

$$\mathcal{L}^A \approx 2i\omega_a\Phi_a^*(\partial_t + \partial_x)\Phi_a + 2i\omega_b\Phi_b^*(\partial_t - \partial_x)\Phi_b, \quad (9)$$

²Note that Ω is often defined differently, i.e., with an additional factor of two.

³The zeroth-order contributions decouple and the first-order terms vanish after an integration by parts, since the background fields solve the equations of motion.

and the atomic states

$$\begin{aligned}L_j^\Psi &\approx i(\Psi_{aj}^*\partial_t\Psi_{aj} + \Psi_{bj}^*\partial_t\Psi_{bj} \\ &+ \Psi_{cj}^*\partial_t\Psi_{cj} + \Gamma\Psi_{cj}^*\Psi_{cj}) - i\Omega(\Psi_{cj}^*\Psi_{aj}\cos\theta \\ &+ \Psi_{cj}^*\Psi_{bj}\sin\theta - \text{H.c.}),\end{aligned}\quad (10)$$

as well as the interaction

$$\begin{aligned}L_j^{A\Psi} &\approx -i\omega_a\epsilon_a\sin\theta\Phi_a(x_j)\Psi_{cj}^* \\ &+ i\omega_b\epsilon_b\cos\theta\Phi_b(x_j)\Psi_{cj}^* + \text{H.c.}\end{aligned}\quad (11)$$

III. EQUATIONS OF MOTION

The equations of motion for the particle amplitudes can be derived from the effective Lagrangian

$$\begin{aligned}\partial_t\Psi_{aj} &= -\Omega\cos\theta\Psi_{cj}, \\ \partial_t\Psi_{bj} &= -\Omega\sin\theta\Psi_{cj}, \\ \partial_t\Psi_{cj} &= \Omega(\cos\theta\Psi_{aj} + \sin\theta\Psi_{bj}) - \Gamma\Psi_{cj} \\ &+ \omega_a\epsilon_a\sin\theta\Phi_a(x_j) - \omega_b\epsilon_b\cos\theta\Phi_b(x_j),\end{aligned}\quad (12)$$

and the equation of motion for the fields Φ_a and Φ_b are

$$\begin{aligned}2(\partial_t + \partial_x)\Phi_a &= -\epsilon_a\sin\theta\sum_j\Psi_{cj}\delta(x-x_j), \\ 2(\partial_t - \partial_x)\Phi_b &= +\epsilon_b\cos\theta\sum_j\Psi_{cj}\delta(x-x_j).\end{aligned}\quad (13)$$

Assuming that the particles are sufficiently closely spaced so that there are many particles in a space of the order of a wavelength of the field, the sum over j can be replaced by the density of the particles

$$\begin{aligned}2(\partial_t + \partial_x)\Phi_a &= -\rho(x)\epsilon_a\sin\theta\Psi_c(x), \\ 2(\partial_t - \partial_x)\Phi_b &= +\rho(x)\epsilon_b\cos\theta\Psi_c(x).\end{aligned}\quad (14)$$

A. Effective dispersion relation

Assuming harmonic space-time dependence $e^{-i\omega t + i\kappa x}$ of all of the variables, we can solve the equations of motion for the atomic amplitudes (12)

$$\begin{aligned}\Psi_{cj}(\omega) &= [\epsilon_a\omega_a\sin\theta\Phi_a(\omega, x_j) - \epsilon_b\omega_b\cos\theta\Phi_b(\omega, x_j)] \\ &\times \frac{i\omega}{\omega^2 - \Omega^2 + i\Gamma\omega},\end{aligned}\quad (15)$$

and inserting this result into Eq. (14) we finally obtain the dispersion relation

$$[\omega + X(\omega) - \kappa][\omega + Y(\omega) + \kappa] = X(\omega)Y(\omega), \quad (16)$$

where

$$X(\omega) = \frac{\omega}{2} \frac{\rho \omega_a \epsilon_a^2 \sin^2 \theta}{\Omega^2 - \omega^2 - i\Gamma \omega},$$

$$Y(\omega) = \frac{\omega}{2} \frac{\rho \omega_b \epsilon_b^2 \cos^2 \theta}{\Omega^2 - \omega^2 - i\Gamma \omega}. \quad (17)$$

B. Adiabatic regime

For small ω and κ , the dispersion relation derived above turns out to be linear, i.e., $\omega \propto \kappa$. Let us specify the required conditions. As already mentioned above, Eq. (14) is valid for wavelengths which are much larger than the interatomic distance Δx (typically a few hundreds of nanometers) only

$$\kappa \ll \frac{1}{\Delta x}. \quad (18)$$

In addition, the manipulations of the previous section (rotating wave approximation) are based on the assumption that the fields Φ_b and Φ_a are slowly varying, i.e., $\omega \ll \omega_a, \omega_b$. However, since the Rabi frequency Ω is supposed to be much smaller than the atomic transition energies ω_a, ω_b and the decay rate is assumed to be small $\Gamma < \Omega$, the knee frequency $\tilde{\Omega}$ of the above dispersion relation yields the relevant frequency cutoff

$$\omega \ll \min \left\{ \Omega, \omega_a, \omega_b, \frac{\Omega^2}{\Gamma} \right\} = \Omega. \quad (19)$$

In this limit, i.e., in the adiabatic regime, Eq. (12) can be solved via

$$\Psi_c = \frac{\omega_a \epsilon_a \sin \theta}{\Omega^2} \Phi_a - \frac{\omega_b \epsilon_b \cos \theta}{\Omega^2} \Phi_b. \quad (20)$$

Rescaling the fields via

$$\tilde{\Phi}_a = \omega_a \epsilon_a \sin \theta \Phi_a,$$

$$\tilde{\Phi}_b = \omega_b \epsilon_b \cos \theta \Phi_b, \quad (21)$$

Eqs. (13) and (14) become

$$(\partial_t + \partial_x) \tilde{\Phi}_a = - \frac{\rho \omega_a \epsilon_a^2 \sin^2 \theta}{2\Omega^2} (\partial_t \tilde{\Phi}_a - \partial_t \tilde{\Phi}_b),$$

$$(\partial_t - \partial_x) \tilde{\Phi}_b = + \frac{\rho \omega_b \epsilon_b^2 \cos^2 \theta}{2\Omega^2} (\partial_t \tilde{\Phi}_a - \partial_t \tilde{\Phi}_b). \quad (22)$$

In order to cast these two first-order differential equations into the usual second-order form, let us choose θ such that⁴

⁴Otherwise one would obtain a velocity-like term even for a medium at rest, cf. Sec. V below. However, this term alone cannot generate an effective horizon.

$$\frac{\rho \omega_a \epsilon_a^2 \sin^2 \theta}{2\Omega^2} = \frac{\rho \omega_b \epsilon_b^2 \cos^2 \theta}{2\Omega^2} = \aleph, \quad (23)$$

where the dimensionless quantity \aleph describes the slow-down of the waves and can be very large $\aleph \gg 1$. In terms of the fields

$$\Phi_{\pm} = \tilde{\Phi}_a \pm \tilde{\Phi}_b, \quad (24)$$

we can indeed combine the two first-order equalities above into one second-order equation

$$\left(\frac{\partial^2}{\partial t^2} - \frac{\partial}{\partial x} \frac{1}{1+2\aleph} \frac{\partial}{\partial x} \right) \Phi_{\pm} = 0. \quad (25)$$

Obviously, small background fields, i.e., small Rabi frequencies Ω , may generate a drastic slow-down $\aleph \gg 1$.

Note, however, that the above wave equation differs from the equation of motion describing a slow-light pulse in the usual setup—i.e., a strong control beam and a weak (perpendicular) probe beam, cf. [1,11]

$$([1 + \aleph] \partial_t \pm \partial_x) \Phi = 0. \quad (26)$$

Hence the slow-down in Eq. (25) $v_{\text{group}} = v_{\text{phase}} = 1/\sqrt{1+2\aleph}$ of the design proposed in the present article is not as extreme as that of the usual setup $v_{\text{group}} = 1/(1+\aleph) \neq v_{\text{phase}} \approx 1$, but still substantial.

IV. EFFECTIVE GEOMETRY

So far we considered a static medium at rest with a possibly position-dependent $\aleph = \aleph(x)$. Now we allow for a space-time varying variable $\aleph = \aleph(t, x)$, where the medium is still at rest. A change of \aleph can be generated by varying ρ , i.e., by adiabatically adding or removing atoms. The other parameters in Eq. (23) remain constant—a time-dependent Ω , for example, would generate additional source terms and thereby invalidate the background solution.

Furthermore, we shall assume $\omega_a = \omega_b$ as well as $\epsilon_a = \epsilon_b$ (which is a reasonable approximation) and hence $\theta = \pi/4$ for the sake of simplicity and absorb these quantities by rescaling the fields Φ_{\pm} .

A. Effective action

Introducing the abbreviation $\Psi = (\Psi_a, \Psi_b, \Psi_c)^T$ the linearized Lagrangian governing the dynamics of the Ψ -fields in Eqs. (10) and (11) can be cast into the following form:

$$\mathcal{A}^{\Psi} = \int d^2x (i\Psi^{\dagger} \cdot \dot{\Psi} + \Psi^{\dagger} \cdot \mathbf{M} \cdot \Psi + [\Psi^{\dagger} \cdot \mathbf{N}] \Phi_{-} + [\mathbf{N}^{\dagger} \cdot \Psi] \Phi_{*}), \quad (27)$$

with \mathbf{M} denoting a (self-adjoint) 3×3 matrix and \mathbf{N} a three-component vector as determined by Eqs. (10) and (11). In terms of the differential operator defined via $\hat{\mathbf{D}} = i\partial_t + \mathbf{M}$ and its formal inverse $\hat{\mathbf{D}}^{-1}$ we may complete the square

$$\mathcal{A}^\Psi = \int d^2x (\tilde{\Psi}^\dagger \cdot \hat{\mathbf{D}} \cdot \tilde{\Psi} - \Phi_-^* N^\dagger \cdot \hat{\mathbf{D}}^{-1} \cdot N \Phi_-), \quad (28)$$

with

$$\tilde{\Psi} = \Psi + \hat{\mathbf{D}}^{-1} \cdot N \Phi_- . \quad (29)$$

Assuming that the quantum state of the Ψ -fields is adequately described by the path-integral with the usual (regular) measure $\mathfrak{D}\Psi$ we are now able to integrate out (i.e., eliminate) those degrees of freedom explicitly arriving at an effective action for the Φ -fields alone

$$\exp\{i\mathcal{A}_{\text{eff}}\} = \frac{1}{Z_\Psi} \int \mathfrak{D}\Psi \exp\{i(\mathcal{A}^\Phi + \mathcal{A}^\Psi)\}. \quad (30)$$

As demonstrated in Eq. (28), the above path-integral is Gaussian ($\mathfrak{D}\Psi = \mathfrak{D}\tilde{\Psi}$) and the associated Jacobi determinant is independent of Φ . Hence we obtain

$$\mathcal{A}_{\text{eff}} = \mathcal{A}^\Phi - \int d^2x \Phi_-^* N^\dagger \cdot \hat{\mathbf{D}}^{-1} \cdot N \Phi_- . \quad (31)$$

As usual, the inverse differential operator $\hat{\mathbf{D}}^{-1}$ causes the effective action to be nonlocal (in time)—but in the adiabatic limit $\omega \ll \Omega$, $\aleph \omega \ll \omega_{a,b}$, $\kappa \ll 1/\Delta x$, and $\aleph \kappa \ll \omega_{a,b}$ the low-energy effective action is local $i\aleph \Phi_-^* \Phi_-$. An easy way to reproduce this result is to remember the original equation of motion

$$\hat{\mathbf{D}} \cdot \Psi + N \Phi_- = 0 \rightsquigarrow \Psi = -\hat{\mathbf{D}}^{-1} \cdot N \Phi_- , \quad (32)$$

and its solution in the adiabatic limit as given by Eq. (20). Together with Eq. (9) we finally arrive at

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & \frac{i}{2} (\Phi_+^* \dot{\Phi}_+ + [1 + 2\aleph] \Phi_-^* \dot{\Phi}_- + \Phi_+^* \dot{\Phi}'_- + \Phi_-^* \dot{\Phi}'_+) \\ & + \text{H.c.} \end{aligned} \quad (33)$$

Strictly speaking, one obtains an effective action for each atom

$$\mathcal{A}_{\text{eff}}^j \propto i \int dt \Phi_-^*(t, x^j) \frac{d\Phi_-(t, x^j)}{dt} + \text{H.c.}, \quad (34)$$

where the total effective action incorporates the sum over all atoms. With the assumption that the atoms are sufficiently closely spaced, cf. Eq. (18), and moving in a direction perpendicular to the beam (e.g., in the y -direction) only, we recover Eq. (33).

An alternative method for effectively changing the density ρ is to cause transitions between the states $|a\rangle$ and $|b\rangle$ and further states $|d\rangle$ and $|e\rangle$, which do not couple to the electromagnetic field Φ_\pm under consideration. The dynamics of these additional states is governed by the Lagrangian

$$\begin{aligned} \mathcal{L}_{\text{add}} = & i\psi_d^* \dot{\psi}_d + i\psi_e^* \dot{\psi}_e + \omega_d \psi_d^* \psi_d + \omega_e \psi_e^* \psi_e \\ & + (i\zeta \psi_d^* \psi_a + i\zeta \psi_e^* \psi_b + \text{H.c.}), \end{aligned} \quad (35)$$

where $i\zeta$ denotes the space-time dependent transition amplitude. (This particular parametrization will be more convenient later on.)

If the amplitude (population) of the states $|d\rangle$ and $|e\rangle$ is large $\psi_{d,e} \gg \psi_{a,b}$ and the transition weak $\zeta \ll 1$, we may neglect the back-reaction ($\psi_{a,b} \rightarrow \psi_{d,e}$) as well as the associated (quantum) fluctuations and describe the process by a classical external source for $\psi_{a,b}$.

Furthermore, assuming $\omega_a = \omega_d$ and $\omega_b = \omega_e$ as well as

$$\begin{aligned} \psi_d &= + e^{i\omega_a(t-x)} \sin \theta + \mathcal{O}(\zeta^2), \\ \psi_e &= - e^{i\omega_b(t+x)} \cos \theta + \mathcal{O}(\zeta^2), \end{aligned} \quad (36)$$

the background solution in Eq. (6) acquires an overall prefactor

$$\begin{aligned} \psi_a^0 &= + \zeta(t, x) e^{i\omega_a(t-x)} \sin \theta, \\ \psi_b^0 &= - \zeta(t, x) e^{i\omega_b(t+x)} \cos \theta, \\ \psi_c^0 &= 0. \end{aligned} \quad (37)$$

This scale factor $\zeta(t, x)$ enters the subsequent formulas and effectively changes the density of the contributing atoms $\rho_{\text{eff}} = \zeta^2 \rho$. In particular, the wave equation (22) gets modified via

$$\partial_t \Phi_- + \partial_x \Phi_+ + 2\aleph \zeta \partial_t (\zeta \Phi_-) = 0, \quad (38)$$

which is exactly the same equation as derived from the effective action in Eq. (33) with $\aleph \rightarrow \zeta^2 \aleph$.

B. Effective spinor-representation

According to Eq. (33) the total effective action for the beat fluctuations Φ_\pm of the electromagnetic field can be written as

$$\begin{aligned} \mathcal{A} = & \frac{i}{2} \int d^2x [(1 + 2\aleph)(\Phi_-^* \partial_t \Phi_- - [\partial_t \Phi_-^*] \Phi_-) + (\Phi_+^* \partial_t \Phi_+ \\ & - [\partial_t \Phi_+^*] \Phi_+) + (\Phi_+^* \partial_x \Phi_- - [\partial_x \Phi_+^*] \Phi_-) \\ & + (\Phi_-^* \partial_x \Phi_+ - [\partial_x \Phi_-^*] \Phi_+)]. \end{aligned} \quad (39)$$

Introducing the effective two-component spinor $\boldsymbol{\psi}$ (not to be confused with the atomic amplitudes $\psi_{a,b,c}$)

$$\boldsymbol{\psi} = \begin{pmatrix} \sqrt{1 + 2\aleph} \Phi_- \\ \Phi_+ \end{pmatrix}, \quad (40)$$

this action can be rewritten as

$$\begin{aligned} \mathcal{A} = & \frac{i}{2} \int \frac{d^2x}{\sqrt{1 + 2\aleph}} \left[\sqrt{1 + 2\aleph} (\boldsymbol{\psi}^\dagger \partial_t \boldsymbol{\psi} - [\partial_t \boldsymbol{\psi}^\dagger] \boldsymbol{\psi}) \right. \\ & \left. + (\boldsymbol{\psi}^\dagger \sigma_x \partial_x \boldsymbol{\psi} - [\partial_x \boldsymbol{\psi}^\dagger] \sigma_x \boldsymbol{\psi}) - \frac{\partial_x \aleph}{1 + 2\aleph} \boldsymbol{\psi}^\dagger i \sigma_y \boldsymbol{\psi} \right], \end{aligned} \quad (41)$$

with $\sigma_x, \sigma_y, \sigma_z$ being the Pauli (spin) matrices obeying $\sigma_x^2 = \sigma_y^2 = \sigma_z^2 = I$.

But this exactly corresponds to the expression for a (1+1)-dimensional Dirac field ψ

$$A = \int d^2x \sqrt{-g} \left[\frac{i}{2} (\bar{\psi} \gamma^\mu \nabla_\mu \psi - [\nabla_\mu \bar{\psi}] \gamma^\mu \psi) - m \bar{\psi} \psi \right], \quad (42)$$

if we define the Dirac γ -matrices via

$$\begin{aligned} \gamma^0 &= \sqrt{1 + 2\aleph} \sigma_y, \\ \gamma^1 &= -i \sigma_z, \end{aligned} \quad (43)$$

and, accordingly, the Dirac adjoint ($\bar{\psi} \psi, \bar{\psi} \gamma^\mu \psi \in \mathbb{R}$)

$$\bar{\psi} = \psi^\dagger \sigma_y, \quad (44)$$

as well as introduce the effective mass

$$m = -\frac{1}{2} \frac{1}{1 + 2\aleph} \frac{\partial \aleph}{\partial x}. \quad (45)$$

The effective metric is given by $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$

$$ds^2 = \frac{dt^2}{1 + 2\aleph} - dx^2, \quad (46)$$

and displays the expected slow-down.

For deriving the identity of Eqs. (39) and (42) we need the properties of the spin connection Γ_μ (Fock-Ivanenko coefficient) which enters into the spin derivative [remember $\partial_\mu (\bar{\psi} \psi) = (\nabla_\mu \bar{\psi}) \psi + \bar{\psi} \nabla_\mu \psi$]

$$\nabla_\mu \psi = \partial_\mu \psi + \Gamma_\mu \psi, \quad \nabla_\mu \bar{\psi} = \partial_\mu \bar{\psi} - \bar{\psi} \Gamma_\mu, \quad (47)$$

and is defined by $[\nabla_\mu (\bar{\psi} \gamma^\nu \psi) = (\nabla_\mu \bar{\psi}) \gamma^\nu \psi + \bar{\psi} \gamma^\nu \nabla_\mu \psi]$

$$\partial_\mu \gamma^\nu + \Gamma_{\rho\mu}^\nu \gamma^\rho = [\gamma^\nu, \Gamma_\mu], \quad (48)$$

with $\Gamma_{\rho\mu}^\nu$ being the Christoffel symbol. In our (1+1)-dimensional representation, the l.h.s. is a linear combination of σ_y and σ_z , cf. Eq. (43), and, therefore, the spin connection Γ_μ has to be proportional to σ_x . As a result we obtain the relation

$$\{\Gamma_\mu, \gamma_\nu\} = 0, \quad (49)$$

and thus confirm the identity of Eqs. (39) and (42)

$$\begin{aligned} 2i \mathcal{L}_{m=0} &= [\nabla_\mu \bar{\psi}] \gamma^\mu \psi - \bar{\psi} \gamma^\mu \nabla_\mu \psi \\ &= [\partial_\mu \bar{\psi}] \gamma^\mu \psi - \bar{\psi} \gamma^\mu \partial_\mu \psi - \bar{\psi} \{\Gamma_\mu, \gamma^\mu\} \psi \\ &= [\partial_\mu \bar{\psi}] \gamma^\mu \psi - \bar{\psi} \gamma^\mu \partial_\mu \psi. \end{aligned} \quad (50)$$

Finally, if we were to choose (over some finite region, since $\aleph > 0$)

$$1 + 2\aleph(t, x) = f(t) e^{-4mx}, \quad (51)$$

the effective mass m in Eq. (45) would be constant (which, however, is not necessary for the introduction of an effective geometry) and the analogy to the (1+1)-dimensional massive Dirac field complete

$$(i \gamma^\mu \nabla_\mu - m) \psi = 0. \quad (52)$$

C. Effective energy

The energy-momentum tensor of a Dirac field reads

$$\begin{aligned} T_{\mu\nu} &= \frac{i}{2} (\bar{\psi} \gamma_{(\mu} \nabla_{\nu)} \psi - [\nabla_{(\mu} \bar{\psi}] \gamma_{\nu)} \psi) \\ &= \frac{i}{2} (\bar{\psi} \gamma_{(\mu} \partial_{\nu)} \psi - [\partial_{(\mu} \bar{\psi}] \gamma_{\nu)} \psi), \end{aligned} \quad (53)$$

where the second equality sign holds in general only in (1+1) dimensions (in analogy to the simplifications above). For an arbitrarily space-time dependent \aleph , however, there is no energy or momentum conservation law associated to this tensor. But assuming time-translation symmetry as described by the Killing vector $\xi = \partial/\partial t$ we may construct a conserved energy via

$$E = \int d\Sigma_\mu T^{\mu\nu} \xi_\nu = \int dx \sqrt{-g} T_0^0, \quad (54)$$

which, for the Dirac field in Eq. (53) and the (static) metric in Eq. (46), reads

$$E = \int dx \frac{i}{2} (\psi^\dagger \dot{\psi} - \dot{\psi}^\dagger \psi). \quad (55)$$

On the classical level, this quantity is (even in flat space-time) not positive definite (as is well-known). For quantum fields the situation can be different. Imposing fermionic (i.e., anti) commutation relations the energy operator is—after renormalization of the zero-point energy and definition of the vacuum state as the filled Dirac sea—indeed non-negative (again in flat space-time). However, the fields $\psi = (\sqrt{1 + 2\aleph} \Phi_-, \Phi_+)^T$ do not obey fermionic but bosonic statistics (as one would expect, cf. Sec. VI below) and, therefore, the effective energy possesses negative parts.

This fact is not surprising in the context of the electromagnetic field, since one has the huge background field with which these perturbations can exchange energy. However, since in the laboratory frame, the background metric is stationary, the energy is a conserved quantity, and the potential instability of the negative energy will not be triggered.

D. Inner product

Since the (classical) equation of motion (52) can be described by means of an effective metric in Eqs. (46) and (60) below, we can introduce a conserved inner product for two solutions of the wave equation ψ_1 and ψ_2 . As usual, the inner product of the Dirac field can be derived by means of the Noether theorem associated to the global $U(1)$ -symmetry $\psi \rightarrow e^{i\varphi} \psi$ and reads

$$\begin{aligned}
(\psi_1|\psi_2) &= \int d\Sigma_\mu \bar{\psi}_1 \gamma^\mu \psi_2 \\
&= \int dx \sqrt{-g} \bar{\psi}_1 \gamma^0 \psi_2 \\
&= \int dx \psi_1^\dagger \psi_2.
\end{aligned} \tag{56}$$

In contrast to the energy in Eq. (55), this quantity is non-negative on the classical level (as well as for quantum fields with bosonic statistics). If we were to impose fermionic commutation relations, the above pseudo-norm would equal the difference of the number of particles and antiparticles and hence not be positive definite. But for bosonic statistics it is non-negative.

Note that, for a scalar field, the situation is completely different since in that case, the inner product is not positive definite: $(F^*|F^*) = -(F|F)$.

V. BLACK HOLE ANALOGUE

After introducing the notion of the effective geometry we can now design an analogue of a black or white hole. To this end we move the medium with a constant velocity v in order to be able to correct (tune) the background beam according to the resulting Doppler shift. Again the background solution, i.e., Ω and θ , should be homogeneous if we want to avoid additional source terms for the linearized fields. (In the reference frame of the moving atoms, an inhomogeneous background becomes time-dependent and thereby also causes a deviation from the dark state.)

The only parameter left for influencing the effective geometry is the density ρ . In order to arrive at a stationary effective metric, the density profile in the laboratory frame should be time-independent $\rho = \rho(x)$. In the rest frame of the fluid, this requirement implies a space-time dependence of $\rho = \rho(x - vt)$. At a first glance, such a scenario seems to be inconsistent with a constant velocity v , but one could arrange a flow profile such as $\mathbf{v} = v e_x - v y \rho' e_y / \rho$ which, for a light beam at $y=0$, reproduces these properties. As already mentioned at the end of Sec. IV A, an alternative possibility is to cause transitions $|d\rangle, |e\rangle \rightarrow |a\rangle, |b\rangle$.

Since we are still working with nonrelativistic velocities $v \ll 1$, the rest frame of the medium and the laboratory are related by a Galilei transformation

$$\frac{\partial}{\partial t} \rightarrow \frac{\partial}{\partial t} + v \frac{\partial}{\partial x}, \quad \frac{\partial}{\partial x} \rightarrow \frac{\partial}{\partial x}. \tag{57}$$

Having derived a covariant, i.e., coordinate-independent, representation of the effective action in Eq. (42), this transformation is completely equivalent to a corresponding change of the effective geometry

$$\gamma^0 \rightarrow \gamma^0, \quad \gamma^1 \rightarrow \gamma^1 + v \gamma^0. \tag{58}$$

The effective metric is then given by the well-known Painlevé-Gullstrand-Lemaître form [12]

$$g_{\text{eff}}^{\mu\nu} = (1 + 2\aleph) \begin{pmatrix} 1 & v \\ v & v^2 - 1/(1 + 2\aleph) \end{pmatrix}. \tag{59}$$

The inverse metric simply reads

$$g_{\text{eff}}^{\mu\nu} = \begin{pmatrix} 1/(1 + 2\aleph) - v^2 & v \\ v & -1 \end{pmatrix}. \tag{60}$$

Obviously, a horizon ($g_{00}^{\text{eff}} = 0$) occurs for $v^2 = 1/(1 + 2\aleph)$, which could be a relatively low velocity and perhaps experimentally accessible.

A. Negative effective energy

For stationary (in the laboratory frame) parameters $\aleph = \aleph(x)$ and $v = \text{const}$ one may construct a conserved energy (Noether theorem) of the beat fluctuations ψ via Eq. (54). Since for a moving medium, the effective metric in Eq. (60) has off-diagonal elements, the resulting expression is more complicated than in Eq. (55). For the sake of convenience, we adopt the geometric-optics approximation $\omega, \kappa \gg \aleph'/\aleph$ and obtain

$$E \approx \int dx \psi^\dagger \frac{\omega + (1 + \sigma_x v \sqrt{1 + 2\aleph})(\omega + v \kappa)}{2} \psi, \tag{61}$$

and, after diagonalization and normalization ($\psi|\psi\rangle = 1$), the solutions for the effective energy assume the following form:

$$E_{\text{eff}} \approx \frac{1}{2} [\omega + (1 \pm v \sqrt{1 + 2\aleph})(\omega + v \kappa)]. \tag{62}$$

Using the approximate ($\omega, \kappa \gg \aleph'/\aleph \approx 0$) equations of motion, we obtain

$$E_{\text{eff}} \approx \omega + v \kappa, \tag{63}$$

which is the frequency as measured in the rest frame of the fluid.

We observe that even the branch $\omega > 0$ of the dispersion relation which would correspond to a positive energy in flat space-time (with $v = 0$) can have a negative effective energy beyond the horizon $v > 1/\sqrt{1 + 2\aleph}$ since $\omega > 0$ does not imply $\omega + v \kappa > 0$ anymore. This purely classical phenomenon—i.e., that the energy (density) measured at infinity can become negative beyond the ergo-sphere $g_{00} = 0$ —occurs for real black holes as well and can be considered as the underlying reason for the mechanism of super-radiance, etc.

Of course, the total energy of the system as derived by the total action in Eq. (1) is always positive. The modes with a negative effective energy (pseudo- or quasienergy, cf. [13]) possess a total energy which is smaller than that of the background. In this regard a (classical) mixing of positive and negative (effective) energy modes is possible.

B. “Bogoliubov” coefficients

If the effective metric possesses a horizon, one would expect the usual mixing of positive and negative energy solutions as governed by the “Bogoliubov” coefficients α_E and β_E defined via

$$\psi_E^{\text{out}} = \alpha_E \psi_+^{\text{in}} + \beta_E \psi_-^{\text{in}}, \quad (64)$$

where ψ_E^{out} denotes the outgoing “would-be” Hawking particle and E its (positive) energy according to Eq. (54); whereas ψ_+^{in} and ψ_-^{in} are incoming (trans-Planckian, cf. [17]) modes whose energy (i.e., frequency) in the fluid’s rest frame is positive (ψ_+^{in}) or negative (ψ_-^{in}), respectively. As we shall see in the next section, the commutation relations are such that Eq. (64) describes a mixing of annihilation operators only and hence does not imply any particle creation. (Hence the quotation marks in “Bogoliubov” coefficients and the notion of a “would-be” Hawking particle.)

These wave-packets are normalized via

$$(\psi_+^{\text{in}} | \psi_+^{\text{in}}) = (\psi_-^{\text{in}} | \psi_-^{\text{in}}) = (\psi_E^{\text{out}} | \psi_E^{\text{out}}) = 1, \quad (65)$$

and orthogonal

$$(\psi_+^{\text{in}} | \psi_-^{\text{in}}) = 0. \quad (66)$$

Owing to the positivity of the inner product in Sec. IV D the completeness relation has a plus sign instead of a minus as for the scalar field, i.e.,

$$|\alpha_E|^2 + |\beta_E|^2 = 1. \quad (67)$$

Consequently we obtain the Fermi-Dirac factor for the scattering (“Bogoliubov”) coefficients (instead of the Bose-Einstein distribution for scalar fields)

$$|\beta_E| = e^{\pi E / \kappa_s} |\alpha_E| \rightsquigarrow |\beta_E|^2 = \frac{1}{e^{2\pi E / \kappa_s} + 1}. \quad (68)$$

However, it should be emphasized here that this mode mixing is *a priori* a purely classical phenomenon and independent of the quantum features (commutation relations)—the fields $\hat{\psi}$ do not obey Fermi-Dirac statistics, see the next section. Only if the quantum commutation relations assigned a physically reasonable particle interpretation to the modes ψ_E —as it is the case for a truly fermionic Dirac quantum field, for example, but not for the fields $\hat{\Phi}_\pm$ (see below)—could one infer the (quantum) Hawking radiation.

The surface gravity of the effective horizon at $v^2 = 1/(1 + 2\aleph) = c_{\text{slow}}^2$ depends on the rate of change of the velocity of light in the laboratory frame across the horizon

$$\begin{aligned} \kappa_s &= \left| \frac{\partial(|v| - c_{\text{slow}})}{\partial x} \right|_{\text{horizon}} \\ &= \left| \frac{\partial c_{\text{slow}}}{\partial x} \right|_{\text{horizon}} \\ &= \frac{1}{\sqrt{(1 + 2\aleph)^3}} \left| \frac{\partial \aleph}{\partial x} \right|_{\text{horizon}}. \end{aligned} \quad (69)$$

By comparison with Eq. (45), we observe that κ_s is of the same order of magnitude as the rest energy induced by the effective mass [remember the homogeneous Dirac equation ($i\gamma^0 \partial_t - m$) $\psi = 0$]

$$\omega_m = \frac{m}{\sqrt{1 + 2\aleph}} = -\frac{1}{2} \frac{1}{\sqrt{(1 + 2\aleph)^3}} \frac{\partial \aleph}{\partial x}. \quad (70)$$

As a result, the relevant mode-mixing effects—i.e., the “Bogoliubov” β -coefficients—are not strongly suppressed by the effective mass.

VI. COMMUTATION RELATIONS

Having derived an effective metric which may exhibit a horizon, one is immediately led to the question of whether the system under consideration could be used to simulate the Hawking effect. As it will turn out, the answer is “no”—since the Hawking effect is a quantum effect, it is not sufficient to consider the wave equation, one also has to check the commutation relations which generate the zero-point fluctuations (the source of the Hawking radiation) according to the Heisenberg uncertainty principle. For convenience we shall transform back into the rest frame of the medium and assume a constant \aleph for the calculations in this section.

A. Commutators

Obviously the effective action derived above is intrinsically different from the one of a charged scalar field, for example. To make the difference more explicit let us consider the effective (adiabatic limit) commutation relations following from Eq. (33).

For any given time t_0 , the equal-time commutation relations of the fields $\hat{\Phi}_\pm$ vanish. Since the equations of motion do not mix $\hat{\Phi}_\pm$ with $\hat{\Phi}_\pm^\dagger$, this remains true for all times

$$[\hat{\Phi}_\pm(t, x), \hat{\Phi}_\pm(t', x')] = [\hat{\Phi}_\pm^\dagger(t, x), \hat{\Phi}_\pm^\dagger(t', x')] = 0. \quad (71)$$

According to Eq. (33) the canonical conjugated momenta are $i\Phi_+^*$ and $i[1 + 2\aleph]\Phi_-^*$, respectively, and hence we obtain

$$[\hat{\Phi}_+(t, x), \hat{\Phi}_+^\dagger(t, x')] = \delta(x - x'), \quad (72)$$

and

$$[\hat{\Phi}_-(t, x), \hat{\Phi}_-^\dagger(t, x')] = \frac{\delta(x - x')}{1 + 2\aleph}. \quad (73)$$

The remaining (equal-time) commutators vanish

$$[\hat{\Phi}_+^\dagger(t, x), \hat{\Phi}_-(t, x')] = [\hat{\Phi}_+(t, x), \hat{\Phi}_-^\dagger(t, x')] = 0, \quad (74)$$

and the commutation relations for the time-derivatives of the fields can be inferred from the equations of motion $\hat{\Phi}_+ + \hat{\Phi}'_- = 0$ and $(1 + 2\aleph)\hat{\Phi}_- + \hat{\Phi}'_+ = 0$.

Remembering the definition of the effective two-component spinor in Eq. (40) the above relations can be cast into the compact form

$$[\hat{\psi}_A(t, x), \hat{\psi}_B(t', x')] = [\hat{\psi}_A^\dagger(t, x), \hat{\psi}_B^\dagger(t', x')] = 0, \quad (75)$$

as well as

$$[\hat{\psi}_A(t,x), \hat{\psi}_B^\dagger(t,x')] = \delta_{AB} \delta(x-x'). \quad (76)$$

Since the beat fluctuation of the electromagnetic field (coupled to the medium) does not obey the Pauli exclusion principle, one cannot fill the Dirac sea consisting of all negative (effective) energy (in flat space-time) states and thereby define a new vacuum state—as it is possible for fermionic quantum fields.

B. Comparison with other fields

Let us compare the above commutation relations with those of a [(1+1) dimensional] Schrödinger field ψ

$$[\hat{\psi}(t,x), \hat{\psi}(t',x')] = [\hat{\psi}^\dagger(t,x), \hat{\psi}^\dagger(t',x')] = 0, \quad (77)$$

as well as

$$[\hat{\psi}(t,x), \hat{\psi}^\dagger(t,x')] = \delta(x-x'), \quad (78)$$

on the one hand and with the commutators of a [(1+1)-dimensional] charged scalar field ϕ

$$\begin{aligned} [\hat{\phi}(t,x), \hat{\phi}(t',x')] &= [\hat{\phi}^\dagger(t,x), \hat{\phi}^\dagger(t',x')] = 0, \\ [\hat{\phi}(t,x), \hat{\phi}^\dagger(t,x')] &= [\hat{\phi}^\dagger(t,x), \hat{\phi}(t,x')] = 0, \end{aligned} \quad (79)$$

as well as

$$[\hat{\phi}(t,x), \partial_t \hat{\phi}^\dagger(t,x')] = i \delta(x-x'), \quad (80)$$

on the other hand. In the latter case (charged scalar field ϕ), the equation of motion can mix positive and negative frequencies and thereby lead to particle production—whereas in the former situation (Schrödinger field ψ), the number of particles is conserved. This difference becomes more evident when one decomposes the fields into real (self-adjoint) and imaginary (anti-self-adjoint) parts. For ψ , the independent canonical conjugated variables are $\Re\psi$ and $\Im\psi$ —whereas for ϕ , they are $\Re\phi$ and $\Re\dot{\phi}$ (as well as $\Im\phi$ and $\Im\dot{\phi}$).

Obviously, the commutation relations of the fields Φ_\pm are clearly inconsistent with those of a charged scalar field ϕ and show more similarity to the (bosonic) Schrödinger field. Therefore, the system under consideration cannot serve as a true analog for the quantum effects in the presence of a black hole horizon—such as Hawking radiation—although it reproduces all classical phenomena.

Since the fields $\hat{\Phi}_\pm$ describe fluctuations of the electromagnetic field, it is also clear that they do not obey the fermionic (anti)commutation rules

$$\{\hat{\psi}_A(t,x), \hat{\psi}_B(t,x')\} = \{\hat{\psi}_A^\dagger(t,x), \hat{\psi}_B^\dagger(t,x')\} = 0, \quad (81)$$

as well as

$$\{\hat{\psi}_A(t,x), \hat{\psi}_B^\dagger(t,x')\} = \delta_{AB} \delta(x-x'). \quad (82)$$

An effective Dirac field satisfying bosonic commutation relations might seem rather strange in view of the spin-

statistics theorem. Indeed, one key ingredient needed for the derivation of this theorem, the spectral condition (which is one of the Wightman axioms), is not satisfied in our case, since the effective energy can become negative owing to the huge total energy of the background field, see also Sec. IV C.

C. Particle creation

In order to answer the question of whether there is any particle creation at all in the described slow-light system, one has to clarify the notion of (quasi)particles to be created (or not) and to specify the corresponding (in/out) vacuum state.

For example, an appropriate initial state $|\text{in}\rangle$, which is a coherent state in terms of the fundamental creation and annihilation operators of the electromagnetic field, could be chosen such that it is annihilated⁵ by all fields $\hat{\Phi}_\pm$,

$$\forall x : \hat{\Phi}_+(t_{\text{in}}, x)|\text{in}\rangle = \hat{\Phi}_-(t_{\text{in}}, x)|\text{in}\rangle = 0. \quad (83)$$

This is possible because (in the rotating wave approximation) the fields $\hat{\Phi}_\pm$ are composed only of positive frequency parts of the electromagnetic field \hat{A} , cf. Eq. (7), i.e., they are annihilators displaced by c -numbers describing the background field (coherent transformation).

In this case the initial (vacuum) state is annihilated by the fields $\hat{\Phi}_\pm$ at all times

$$\forall t, x : \hat{\Phi}_+(t, x)|\text{in}\rangle = \hat{\Phi}_-(t, x)|\text{in}\rangle = 0, \quad (84)$$

as the time-evolution does not mix $\hat{\Phi}_\pm$ with $\hat{\Phi}_\pm^\dagger$, and there is no particle creation.

For another initial (vacuum) state (e.g., a squeezed state) and a different particle concept,

$$f[\hat{\Phi}_\pm, \hat{\Phi}_\pm^\dagger]|\text{in}'\rangle = 0, \quad (85)$$

however, some effects of (quasi)particle creation might occur. These phenomena could be tested by sending in a (multi-mode) squeezed state and comparing the number of photons per mode in the in- and out-states.

Another possible source for (quasi)particle creation is the finite life-time of the atomic state $|c\rangle$ as represented by the effective decay rate Γ . Realistically, this decay corresponds to some spontaneous emission process generated by the quantum fluctuations of the electromagnetic field, for example. Consistent with the fluctuation-dissipation theorem this coupling also introduces (quantum) noise, which is not included in our treatment and could possibly lead to particle creation. However, this is clearly a pure trans-Planckian effect and cannot be interpreted as Hawking radiation.

⁵Note that this state is therefore annihilated by the (normal-ordered) effective energy in Eq. (55), which is, however, not positive definite (no ground state). Consequently, it cannot be the equivalent of the Israel-Hartle-Hawking [14] state, in which the Hawking radiation is somewhat hidden by the fact that there is no net energy flux.




FIG. 1. One branch of the dispersion relation of the Φ -field in Eq. (86). Frequency ω and wave-number κ are plotted in units of the Rabi frequency Ω for $\aleph=10$ and $\Gamma/\Omega=1/10$. These values (of order one) are but illustrative and chosen in order to resolve the characteristic features in one figure—realistically the orders of magnitude are different. The imaginary part describes the absorption and does not change significantly in the limit $\Gamma \downarrow 0$. For very large as well as for very small ω the medium becomes transparent. The steep slope within the transparency window $\omega \ll \Omega$ corresponds to the reduced propagation velocity—whereas the effect of the medium for large ω is negligible. As one can observe, the anomalous frequency solutions $\omega > \Omega$ are separated from the normal ones $\omega < \Omega$ by a large region of absorption.

VII. DISPERSION RELATION

Although slow light cannot be used to simulate the Hawking effect it can reproduce various classical effects associated to horizons,⁶ such as mode mixing and the associated Bogoliubov coefficients, see Sec. V B. In view of the red- or blue-shift near the horizon deviations from the linear dispersion relation have to be taken into account, cf. [17]. With the choice in Eq. (23) the dispersion relation in Sec. III A simplifies because of $X(\omega)=Y(\omega)$, and we obtain for a medium at rest, cf. Figs. 1 and 2

$$\kappa = \pm \omega \sqrt{1 + 2\aleph \frac{\Omega^2}{\Omega^2 - \omega^2 - i\Gamma\omega}}. \quad (86)$$

We observe two major differences between the dispersion relation above and that for the sonic black hole analogs, for example in Bose-Einstein condensates (see [18] and Sec. IX A) with

⁶Other systems which are potentially capable of simulating those classical effects with present-day technology are discussed in Refs. [15] and [16].




FIG. 2. The real part of the dispersion relation in Fig. 1 as ω vs κ with the same values. One can easily recognize that the first deviation from the linear dispersion relation at $\omega \ll \Omega$ is “subluminal”—although it becomes finally “superluminal” for $\omega \gg \Omega$. The solutions with an anomalous (negative or even infinite) group velocity lie completely in the absorptive region, cf. Fig. 1.

$$\omega^2 = c_{\text{sound}}^2 \kappa^2 (1 + \xi^2 \kappa^2), \quad (87)$$

where ξ denotes the so-called healing length and provides a wave-number cutoff, cf. Fig. 3.

The sonic black hole analogs generate a deviation from the linear dispersion relation via the spatial dependence (κ) and, consequently, for each value of the wave-number κ there exist two possible solutions for the frequency ($\pm \omega$ for a medium at rest). In contrast, for the black hole analogs based on slow light the deviation is mainly⁷ caused by the (nonlocal) temporal dependence. (This remains true for all dielectric/optical black hole analogs, cf. [7,9].) As a result, one has two values of κ for each value of ω , but can have more than two solutions for ω for some values of κ . Even though these anomalous solutions for ω are separated from the normal ones by a relatively large region of absorption, it would be interesting to see under which circumstances this peculiar behavior may give raise to additional effects (such as mode mixing, etc.).

Another major difference between the dispersion relations (86) and (87) is that the sonic dispersion relation (87) is “superluminal”/supersonic for large wave-numbers $v_{\text{group}} = d\omega/d\kappa > c_{\text{sound}}$ for $\xi\kappa \ll 1$ whereas the slow-light dispersion relation (86) is “subluminal” $v_{\text{group}} = d\omega/d\kappa < 1/\sqrt{1+2\aleph}$ within the transparency window, say $|\omega|$

⁷Of course, the finite interatomic distance results in a deviation from the linear dispersion relation too, but the cutoff given by the Rabi frequency is usually reached earlier.




FIG. 3. One branch of the dispersion relation of (zero) sound waves in Bose-Einstein condensates at rest, cf. Eq. (87), in arbitrary units. If the condensate is moving the various κ -solutions for a given frequency ω in the laboratory frame can be found by the points of intersection with straight lines as determined by Eq. (88). For a subsonic velocity $v < c_{\text{sound}}$, there is only one solution, denoted by $s+$, which has a small wave-number and a positive pseudo-norm, i.e., a positive $\omega_{\text{fluid's rest-frame}}$ (assuming $\omega_{\text{lab-frame}} > 0$). For supersonic velocities, on the other hand, i.e., beyond the horizon, there are three possible solutions—one with a small wave-number and a negative pseudo-norm ($s-$) as well as two others with large wave-numbers and positive ($I+$) and negative ($I-$) pseudo-norm, respectively. The mixing between these modes at the horizon generates the Hawking radiation ($s+$).

$< \Omega/2$, but $|\omega| \notin \Omega$. For very large frequencies $\omega \gg \Omega$ one recovers the speed of light in vacuum $\omega = \kappa$ —although this limit is totally outside the region of applicability of our approximations.

VIII. PROBLEMS OF SLOW LIGHT

The direct (naive) way to use the most common setup (see Figs. 4 and 5) for slow-light experiments—i.e., a strong control beam and a weak (perpendicular) probe beam—in order to build a black hole analog goes along with a number of (somewhat related) difficulties listed below. Whereas the first three obstacles can be avoided by the arrangement proposed in this article, the fourth one persists—indicating that this system is a classical, but not a quantum analogue of a black hole.

A. Frequency window

Light pulses (of the probe beam) are only slowed down drastically—or may propagate at all—in an extremely narrow frequency window in the optical or near-optical regime. But the frequency of the particles constituting the Hawking




FIG. 4. One branch of the dispersion relation of a slow-light pulse (in the usual setup) $\kappa^2 = \omega^2 [1 + \wp(\omega + \omega_0) - \wp(\omega - \omega_0)]$ where $\wp(\omega) = 2\aleph(\Omega^2/\omega_0)\omega/(\omega^2 - \Omega^2 + i\Gamma\omega)$, see e.g., [1,10], in units of the Rabi frequency Ω for $\omega_0/\Omega = 20$, $\Gamma/\Omega = 1/2$, and $\aleph = 5$. Again, these unrealistic values have been chosen in order to illustrate the characteristic features. For more realistic values the peaks would be more pronounced, the transparency windows $|\omega \pm \omega_0| \ll \Omega$ narrower, and the slope inside them steeper, etc., but the main structure remains. For $|\omega \pm \omega_0| \gg \Omega$ the influence of the medium is negligible. Within the transparency windows $|\omega \pm \omega_0| \ll \Omega$, the steep slope indicates a reduced group velocity and the solutions with an anomalous group velocity $|\omega \pm \omega_0| = \mathcal{O}(\Omega)$ lie inside the absorptive regions.

radiation cannot be much larger than the surface gravity (e.g., the gradient of the fluid's velocity) which makes an experimental verification in this way very unlikely.

B. Doppler shift

In a stationary medium, the frequency as measured in the laboratory frame is conserved—but the frequency in the atom's rest frame changes as soon as the velocity of the medium (Doppler shift) or the wave-number (redshift) varies (which necessarily happens near the horizon). Hence the beam will leave the narrow frequency window—which is generated by the (moving) atoms—in general.

C. Group and phase velocity

Since the group and the phase velocity of the probe beam are extremely different $v_{\text{group}} \ll v_{\text{phase}} \approx 1$, it is not possible to describe its dynamics by an effective local wave equation resembling a scalar field in a curved space-time.

D. Positive/negative frequency-mixing

In order to obtain particle creation, one has to have a mixing of positive and negative frequencies, or, more accurately, positive and negative pseudo-norm (as induced by the inner product, cf. Sec. V B) solutions. In a stationary flowing




FIG. 5. The real part of the dispersion relation in Fig. 4 as ω vs κ with the same values. The additional line demonstrates the slope corresponding to a motion of the medium with the reduced group velocity as in Fig. 3. Obviously, there can be no mixing of positive and negative pseudo-norms via the usual mechanism sketched below Fig. 3 in this case. Even though the peaks can be much higher for small Γ and thereby could possibly intersect with the straight line, the resulting solutions would lie completely in the region of strong absorption (cf. Fig. 4) and therefore do certainly not model Hawking radiation.

medium (as used for the black hole analogs), this can occur by tilting the dispersion relation due to the Doppler effect caused by the velocity of the medium

$$\omega_{\text{lab-frame}} = \omega_{\text{fluid's rest-frame}} + v_{\text{medium}} \kappa. \quad (88)$$

As soon as the velocity on the medium exceeds $|\omega/\kappa|$, i.e., the phase velocity, a mixing of positive and negative frequencies (in the fluid's rest frame) becomes possible, cf. [10]. However, since the phase velocity of the slow-light pulse is basically the same as in vacuum, this mechanism does not work in this situation and, consequently, there is no particle creation.

IX. COMPARISON WITH OTHER SYSTEMS

One of the main points of the present article is the observation that an appropriate wave equation and the resulting effective geometry of a black hole analog is *not* enough for predicting Hawking radiation. Although all the classical effects can be reproduced in such a situation, the adequate simulation of the *quantum* effects requires the correct commutation relations as well.

In view of this observation one might wonder whether this is actually the case for the currently discussed (e.g., sonic/acoustic and dielectric/optical) black hole analogs. In the following we shall deal with this question for two repre-

sentative examples, for which the commutation relations can be derived easily.

A. Bose-Einstein condensates

The dynamics of Bose-Einstein condensates are to a very good approximation described by the Gross-Pitaevskii equation

$$i\dot{\psi} = \left(-\frac{\nabla^2}{2m} + V(\mathbf{r}) + \eta^2 |\psi|^2 \right) \psi, \quad (89)$$

where ψ denotes the mean-field amplitude, m the mass of the bosons, V an external (trapping) potential, and η is the scattering parameter governing the two-body repulsion of the constituents. Inserting the eikonal ansatz (Madelung representation),

$$\psi = \sqrt{\varrho} e^{iS}, \quad (90)$$

and introducing the (mean-field) velocity $\mathbf{v} = \nabla S/m$, one obtains the equation of continuity $\dot{\varrho} + \nabla(\varrho \mathbf{v})$ and the equivalent of the Bernoulli or the Hamilton-Jacobi equation

$$\dot{S} + V + \eta^2 \varrho + \frac{(\nabla S)^2}{2m} = \frac{1}{2m} \frac{\nabla^2 \sqrt{\varrho}}{\sqrt{\varrho}}. \quad (91)$$

Within the Thomas-Fermi approximation, one neglects the quantum potential, i.e., the term on the l.h.s., and hence recovers the usual equations of fluid dynamics, see also [18]. The linearization around a given (stationary) background profile ϱ_0 and $S_0 \rightarrow \mathbf{v}_0$ yields the well-known wave equation

$$(\partial_t + \nabla \cdot \mathbf{v}_0)(\partial_t + \mathbf{v}_0 \cdot \nabla) \delta S = \frac{\eta^2}{m} \nabla \varrho_0 \nabla \delta S. \quad (92)$$

The commutation relations of δS , which we are interested in, can be derived from the commutator of the fundamental fields

$$[\hat{\psi}(t, \mathbf{r}), \hat{\psi}^\dagger(t, \mathbf{r}')] = \delta^3(\mathbf{r} - \mathbf{r}'). \quad (93)$$

Inserting the linearization of $\hat{\psi} = \sqrt{\hat{\varrho}} \exp(i\hat{S})$ around a classical background via $\hat{\varrho} = \varrho_0 + \delta\hat{\varrho}$ and $\hat{S} = S_0 + \delta\hat{S}$ we obtain (note that $\hat{\varrho} = \hat{\varrho}^\dagger$ and $\hat{S} = \hat{S}^\dagger$)

$$[\delta\hat{\varrho}(t, \mathbf{r}), \delta\hat{S}(t, \mathbf{r}')] = i\delta^3(\mathbf{r} - \mathbf{r}'). \quad (94)$$

The relation between $\delta\hat{\varrho}$ and $\delta\hat{S}$ follows from Eq. (91) in the Thomas-Fermi approximation

$$\delta\hat{\varrho} = -\frac{1}{\eta^2} (\partial_t + \mathbf{v}_0 \cdot \nabla) \delta\hat{S}. \quad (95)$$

Hence $\delta\hat{\varrho}$ is indeed the (negative) canonical conjugated momentum to $\delta\hat{S}$ —provided that one inserts the constant factor η^2 correctly into the (effective) action—and the commutation relations are equivalent (within the used approximation) to those of a quantum field in a curved space-time.

B. Nondispersive dielectric media

As another example we study nondispersive and linear dielectric media, see e.g. [7]. For a medium at rest the fundamental Lagrangian describing the electromagnetic field, the dynamics of the medium ($\mathcal{L}[\mathbf{P}]$), as well as their mutual interaction ($\mathbf{E} \cdot \mathbf{P}$) is given by

$$\mathcal{L} = \frac{1}{2}(\mathbf{E}^2 - \mathbf{B}^2) + \mathbf{E} \cdot \mathbf{P} + \mathcal{L}[\mathbf{P}]. \quad (96)$$

Accordingly, using the temporal gauge and introducing the vector potential via $\mathbf{E} = -\partial_t \mathbf{A}$ and $\mathbf{B} = -\nabla \times \mathbf{A}$, the canonical momentum is just the electric displacement

$$\mathbf{\Pi} = \mathbf{D} = \mathbf{E} + \mathbf{P}. \quad (97)$$

Performing basically the same steps as in Sec. IV A we may integrate out the degrees of freedom associated to the medium \mathbf{P} and thereby arrive at an effective (low-energy) action for the (macroscopic) electromagnetic field alone, cf. [7]. But, in contrast to the highly resonant behavior of \mathbf{P} in slow-light systems, nondispersive media respond adiabatically with a constant susceptibility $\chi = \varepsilon - 1$, i.e., $\mathbf{P} = \chi \mathbf{E}$ and thus $\mathbf{\Pi} = \mathbf{D} = \varepsilon \mathbf{E}$, to the external field (at sufficiently low frequencies), cf. [7].

If the (nondispersive) medium is moving with the velocity $\boldsymbol{\beta}$ the electric and magnetic fields get mixed and one obtains

$$\mathbf{\Pi} = \mathbf{D} = \varepsilon \mathbf{E} + (\varepsilon - 1) \mathbf{B} \times \boldsymbol{\beta} + \mathcal{O}(\boldsymbol{\beta}^2). \quad (98)$$

Again, the commutation relations fit to an effective-metric description—which is not completely surprising because the effective action has the same form as in curved space-times, cf. [7].

X. DISCUSSION

Let us summarize: The naive application of slow light (i.e., the most common setup) in order to create a black hole analog goes along with several problems, cf. Sec. VIII. With the scenario proposed in this article, the problems associated with the classical wave equation can be solved and it is—at least in principle—possible to create a (classical) black hole analog for the Φ field. At low wave-number, the corresponding dispersion relation represents a quadratic relation between κ and ω , and can thus be written in terms of an effective metric. If the fluid is in motion, this low wave-number equation can be changed into a black hole type wave equation.

However, this classical black hole analog does *not* reproduce the expected quantum effects—such as Hawking radiation.⁸ In order to simulate the Hawking effect, it is not sufficient to design a system with an equivalent effective

equation of motion—the commutation relations have to match as well. This is indeed the case for the sonic black hole analogs in Bose-Einstein condensates and nondispersive dielectric black hole analogs—but for sound waves in more complicated systems, for example, it is not immediately obvious.

Nevertheless, in the scenario described in this article, the field Φ governing the beat fluctuations of an electromagnetic background field obey the same equation of motion as in the presence of a horizon and hence can be used to model several classical effects associated with black holes—for example the mode mixing at the horizon as described by the “Bogoliubov” coefficients, see Sec. V B. One way of measuring the “Bogoliubov” coefficients could be to send in a “classical” pulse above the background—i.e., a particular coherent state in terms of the fundamental electromagnetic field—and compare it with the outgoing pulse. As another (more fancy) possibility one might think of a multimode squeezed state—which in some sense simulates the vacuum fluctuations which are transformed into quasiparticles by the mode mixing.

However, one should bear in mind that, as the wave-packets propagate away from the horizon and get strongly blueshifted, they eventually reach the regime where the concept of the effective geometry breaks down and effects like dispersion, nonlocality (in time) of the effective action, and, finally, absorption become relevant. For a reasonably clean interpretation, therefore, one should investigate the scattering of the wave-packets not too far away from the horizon.

A. Miles instability

Another interesting classical effect is related to the negative parts of the energy in Eq. (62). Since a conserved positive definite energy functional of the linearized perturbations would demonstrate linear stability, the negative contribution in Eq. (62) can be interpreted as an indicator for a potential instability (e.g., super-radiance)—provided a suitable coupling between positive and negative (effective) energy modes.

As an example, let us assume that the “superluminally” flowing $v > 1/\sqrt{1+2\aleph}$ slow-light medium interacts with the environment in the laboratory frame via a friction term such as $\Gamma \partial_t \Phi$ (with possible spatial derivatives). For small ω and κ the resulting dissipation alters the dispersion relation via

$$(\omega + v\kappa)^2 = \frac{\kappa^2}{1+2\aleph} - i\omega\Gamma(\kappa), \quad (99)$$

with the potentially κ -dependent (additional spatial derivatives) friction term $\Gamma(\kappa)$ describing the interaction of the Φ -field with the environment at rest.

For small Γ the imaginary part of the solutions for the frequency ω (assuming a real wave-number $\kappa \in \mathbb{R}$) reads

$$\Im(\omega) = -\frac{\Gamma(\kappa)}{2} (1 \pm v\sqrt{1+2\aleph}). \quad (100)$$

⁸This conclusion applies in the same way to the scenario proposed in Ref. [8], where the Schwarzschild metric is simulated by a medium at rest with the horizon corresponding to a singularity in the effective refractive index. Such static analogs of the Schwarzschild geometry (see also [19]) have further problems.

Consequently, beyond the horizon $v > 1/\sqrt{1+2\aleph}$ one of the allowed frequency solutions acquires a positive imaginary part and thus the dissipation (interaction with the environment) generates an instability. Note that the relative velocity $v > 1/\sqrt{1+2\aleph}$ between the slow-light medium and the environment (at rest) is crucial since a friction term like $\Gamma(\partial_t + v\partial_x)\Phi \rightarrow i(\omega + v\kappa)\Gamma$ would of course not lead to any instability.

This instability is somewhat analogous to the Miles instability [20] generating surface waves in water by wind blowing over it. In Ref. [16], this phenomenon is called thermodynamic instability since it occurs when the free energy of the medium acquires negative parts in the frame of the environment.

B. Outlook

Apart from the aforementioned experiments there are many more conceivable tests one could perform with the proposed classical black hole analog based on slow light. A more drastic way of investigating the interior structure of the

sample (than the mere comparison of the in- and out-states) could be to freeze the dark state by completely switching off the background field and take a “snap-shot” of the state of the atoms by illuminating them with strong laser beams with frequencies corresponding to certain atomic transitions and measuring the absorption.

Furthermore it would be interesting to investigate the influence of the anomalous frequency solutions of the dispersion relation generated by the nonlocal temporal dependence (cf. Sec. VII), for example, on additional mode-mixing. This question is relevant for more general (nondispersive) dielectric black hole analogs and might also lead to some insight into the trans-Planckian problem.

ACKNOWLEDGMENTS

The authors acknowledge comments from R. Parentani and M. Scully. This work was supported by the Alexander von Humboldt Foundation, the Canadian Institute for Advanced Research, and the Natural Science and Engineering Research Council of Canada.

-
- [1] L.V. Hau *et al.*, Nature (London) **397**, 594 (1999); M.M. Kash *et al.*, Phys. Rev. Lett. **82**, 5229 (1999); D. Budker *et al.*, *ibid.* **83**, 1767 (1999); O. Kocharovskaya *et al.*, *ibid.* **86**, 628 (2001); D.F. Phillips *et al.*, *ibid.* **86**, 783 (2001); A.V. Turukhin *et al.*, *ibid.* **88**, 023602 (2002); see also M.O. Scully and M.S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, England, 1997).
- [2] W.G. Unruh, Phys. Rev. Lett. **46**, 1351 (1981).
- [3] S.W. Hawking, Nature (London) **248**, 30 (1974); Commun. Math. Phys. **43**, 199 (1975).
- [4] W. Gordon, Ann. Phys. (Leipzig) **72**, 421 (1923); J.E. Tamm, J. Russ. Phys. Chem. Soc. **56(2-3)**, 284 (1924); G.V. Skrotskii, Sov. Phys. Dokl. **2**, 226 (1957); N.L. Balazs, Phys. Rev. **110**, 236 (1958); J. Plebanski, *ibid.* **118**, 1396 (1960); F. Winterberg, Nuovo Cimento Soc. Ital. Fis., B **53**, 264 (1968); F. de Felice, Gen. Relativ. Gravit. **2**, 347 (1971); L.D. Landau and E.M. Lifschitz, *The Classical Theory of Fields*, Course of Theoretical Physics Vol. 2 (Pergamon, London, 1971); W. Schleich and M.O. Scully, in *New Trends in Atomic Physics*, edited by R. Stora and G. Grynberg, Proc. Les Houches (Session 38) 1982 (North Holland, Amsterdam, 1984), p. 995; V.V. Zheleznyakov, V.V. Kocharovskiy, and V.I. Kocharovskiy, Sov. Phys. Usp. **32**, 835 (1989), and references therein.
- [5] M. Visser, Class. Quantum Grav. **15**, 1767 (1998); M. Visser, C. Barcelo, and S. Liberati, Gen. Relativ. Gravit. **34**, 1719 (2002); H.C. Rosu, Int. J. Mod. Phys. D **3**, 545 (1994); Gravitation Cosmol. **7**, 1 (2001).
- [6] U. Leonhardt and P. Piwnicki, Phys. Rev. Lett. **84**, 822 (2000); see also M. Visser, *ibid.* **85**, 5252 (2000); and U. Leonhardt and P. Piwnicki, *ibid.* **85**, 5253 (2000).
- [7] R. Schützhold, G. Plunien, and G. Soff, Phys. Rev. Lett. **88**, 061101 (2002); see also I. Brevik and G. Halmes, Phys. Rev. D **65**, 024005 (2002).
- [8] U. Leonhardt, Nature (London) **415**, 406 (2002); Phys. Rev. A **65**, 043818 (2002).
- [9] M. Novello, S. Perez Bergliaffa, J. Salim, V. De Lorenci, and R. Klippert, Class. Quantum Grav. **20**, 859 (2003); V.A. De Lorenci, R. Klippert, and Y.N. Obukhov, in “Aguas de Lindoia 2002, Physics of particles and fields,” gr-qc/0210104; V.A. De Lorenci and R. Klippert, Phys. Rev. D **65**, 064027 (2002); M. Novello, Int. J. Mod. Phys. A **17**, 4187 (2002); M. Novello and J.M. Salim, Phys. Rev. D **63**, 083511 (2001); M. Novello, V.A. De Lorenci, J.M. Salim, and R. Klippert, *ibid.* **61**, 045001 (2000); V.A. De Lorenci, R. Klippert, M. Novello, and J.M. Salim, Phys. Lett. B **482**, 134 (2000); F. Baldovin, M. Novello, S.E. Perez Bergliaffa, and J.M. Salim, Class. Quantum Grav. **17**, 3265 (2000).
- [10] W. G. Unruh, in *Artificial Black Holes*, edited by M. Novello, M. Visser, and G. Volovik (World Scientific, Singapore, 2002).
- [11] U. Leonhardt, in *Artificial Black Holes*, edited by M. Novello, M. Visser, and G. Volovik (World Scientific, Singapore, 2002).
- [12] P. Painlevé, C. R. Hebd. Seances Acad. Sci. **173**, 677 (1921); A. Gullstrand, Ark. Mat., Astron. Fys. **16**, 1 (1922); G. Lemaître, Ann. Soc. Sci. (Bruxelles) **53**, 51 (1933).
- [13] M. Stone, Phys. Rev. E **62**, 1341 (2000).
- [14] W. Israel, Phys. Lett. **57A**, 107 (1976); J.B. Hartle and S.W. Hawking, Phys. Rev. D **13**, 2188 (1976).
- [15] R. Schützhold and W.G. Unruh, Phys. Rev. D **66**, 044019 (2002).
- [16] G.E. Volovik, Pis'ma Zh. Éksp. Teor. Fiz. **75**, 691 (2002) [JETP Lett. **75**, 418 (2002)]; **76**, 296 (2002) [**76**, 240 (2002)]; see also G.E. Volovik, *Universe in a Helium Droplet* (Oxford University Press, Oxford, 2003) and D.A. Abanin, Pis'ma Zh. Eksp. Teor. Fiz. **77**, 222 (2003) [JETP Lett. **77**, 191 (2003)].
- [17] T. Jacobson, Phys. Rev. D **44**, 1731 (1991); **48**, 728 (1993); **53**, 7082 (1996); Prog. Theor. Phys. Suppl. **136**, 1 (1999); W.G. Unruh, Phys. Rev. D **51**, 2827 (1995); R. Brout, S. Massar, R. Parentani, and P. Spindel, *ibid.* **52**, 4559 (1995); S. Corley and T. Jacobson, *ibid.* **54**, 1568 (1996); S. Corley, *ibid.* **55**, 6155

- (1997); **57**, 6280 (1998); B. Reznik, *ibid.* **55**, 2152 (1997).
- [18] L.J. Garay, J.R. Anglin, J.I. Cirac, and P. Zoller, *Phys. Rev. Lett.* **85**, 4643 (2000); *Phys. Rev. A* **63**, 023611 (2001); C. Barcelo, S. Liberati, and M. Visser, *Class. Quantum Grav.* **18**, 1137 (2001); L.J. Garay, *Int. J. Theor. Phys.* **41**, 2073 (2002).
- [19] B. Reznik, *Phys. Rev. D* **62**, 044044 (2000); G. Chapline *et al.*, *Philos. Mag. B* **81**, 235 (2001); R.B. Laughlin, *Int. J. Mod. Phys. A* **18**, 831 (2003); see also *On the Static Analogs of the Schwarzschild Geometry*, in preparation.
- [20] J.W. Miles, *J. Fluid Mech.* **3**, 185 (1957); see also G.E. Vekstein, *Am. J. Phys.* **66**, 886 (1998).