

Using untagged $B^0 \rightarrow DK_S$ to determine γ M. Gronau,¹ Y. Grossman,^{1,2,3} N. Shuhmaher,¹ A. Soffer,⁴ and J. Zupan^{1,5}¹*Department of Physics, Technion–Israel Institute of Technology, Technion City, 32000 Haifa, Israel*²*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309, USA*³*Santa Cruz Institute for Particle Physics, University of California, Santa Cruz, California 95064, USA*⁴*Department of Physics, Colorado State University, Fort Collins, Colorado 80523, USA*⁵*J. Stefan Institute, Jamova 39, P.O. Box 3000, 1001 Ljubljana, Slovenia*

(Received 12 February 2004; published 2 June 2004)

It is shown that the weak phase $\gamma \equiv \arg(-V_{ud}V_{ub}^*V_{cb}V_{cd}^*)$ can be determined using only untagged decays $B^0/\bar{B}^0 \rightarrow DK_S$. In order to reduce the uncertainty in γ , we suggest combining information from $B^\pm \rightarrow DK^\pm$ and from untagged B^0 decays, where the D meson is observed in common decay modes. Theoretical assumptions, which may further reduce the statistical error, are also discussed.

DOI: 10.1103/PhysRevD.69.113003

PACS number(s): 12.15.Hh, 11.30.Er, 13.25.Hw

I. INTRODUCTION

CP violation measured in $B \rightarrow J/\psi K_S$ [1] is interpreted in terms of the phase $\beta \equiv \arg(-V_{tb}V_{td}^*V_{cb}V_{cb}^*)$ in a way which is practically free of theoretical uncertainties, providing an important test of the Kobayashi-Maskawa mechanism [2]. On the other hand, the current interpretation of CP asymmetry measurements in $B \rightarrow \pi^+\pi^-$ [3] in terms of the phase $\alpha \equiv \arg(-V_{td}V_{ub}^*V_{ub}V_{ud}^*)$ involves an uncertainty in the ratio of penguin-to-tree amplitudes [4]. A theoretically clean method [5,6] for measuring the phase $\gamma \equiv \arg(-V_{ud}V_{ub}^*V_{cb}V_{cd}^*)$ involves interference between tree amplitudes $\bar{b} \rightarrow \bar{c}u\bar{s}$ and $\bar{b} \rightarrow \bar{u}c\bar{s}$, governing $B \rightarrow \bar{D}^0 X_s$ and $B \rightarrow D^0 X_s$, where the \bar{D}^0 and D^0 decay to a common hadronic state, and $X_s = K, K^*, K\pi, \dots$ is a strangeness-1 state.

Originally, this idea for measuring γ was proposed for charged B decays [5] and for time-dependent neutral B decays [6] of the type $B \rightarrow D_{CP}K$. Here positive (negative) CP eigenstates, such as K^+K^- ($K_S\pi^0$), identify equal admixtures of D^0 and \bar{D}^0 states with equal (opposite) signs. A variety of other D decay final states can be used as well, leading to several variants of the original method [7,8]. Every hadronic state accessible at the tree level to D^0 decay is also accessible at the tree level to \bar{D}^0 decay with varying levels of Cabibbo suppression. Flavor states, $K^-\pi^+$ and $K^*\pi^+$, which are Cabibbo-favored in D^0 decays, are doubly Cabibbo-suppressed in \bar{D}^0 decays. Flavorless states, such as K^*+K^- and K^+K^* , are produced in singly Cabibbo-suppressed decays of both D^0 and \bar{D}^0 . Recently, it was shown that a model-independent extraction of γ is also possible by considering $B^\pm \rightarrow DK^\pm$ with subsequent multibody D decay, such as $D \rightarrow K_S\pi^+\pi^-$ [9,10]. Other variants make use of the decays $B^\pm \rightarrow D^*K^\pm, D^* \rightarrow D\pi^0$, the self-tagged decay mode $B^0 \rightarrow DK^*0, K^*0 \rightarrow K^+\pi^-$ [11], multibody B decays of the type $B \rightarrow DK\pi$ [12] and combinations of these processes [13].

First results for $B^\pm \rightarrow D_{CP}K^\pm$ were presented recently by the Belle [14] and BaBar [15] Collaborations. These studies were based on several tens of events in each experiment and demonstrate the potential of a larger data sample in provid-

ing useful constraints on γ [16]. The Belle Collaboration has also presented a preliminary analysis of about 100 events of the type $B^\pm \rightarrow DK^\pm, D \rightarrow K_S\pi^+\pi^-$ [17], from which constraints on γ were obtained. The main difficulty of each of these methods is that each decay mode by itself has a very low rate. Reaching high sensitivity in near future measurements of γ requires combining several relevant B and D decay modes.

Measurements relevant to studying γ have so far focused on charged B decays, $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow DK^*0$ [18]. At first glance, neutral B decays seem to be less promising for two reasons: they have much smaller rates and require B^0 flavor tagging. Let us discuss these two points one at a time:

(i) The processes $B^0 \rightarrow DK^0$ and $B^0 \rightarrow DK^*0$ are expected to be color-suppressed [11], implying that their rates are about an order of magnitude below the rates of corresponding charged B decays. However, the crucial factor determining the sensitivity of a measurement of γ is not the decay rate itself. Rather, the sensitivity is governed by the magnitude of the smaller of the two interfering amplitudes in a given process. Since the smaller amplitudes in B^+ and B^0 decays are both color-suppressed, they are expected to be of comparable magnitudes. Therefore, this by itself is not a limiting factor for neutral B decays.

(ii) Only time-integrated rates have so far been measured in $B^0 \rightarrow DK^0$, combining rates for $B^0 \rightarrow \bar{D}^0K^0$ and $\bar{B}^0 \rightarrow \bar{D}^0\bar{K}^0$ [19]. It was shown in [6,20,21] that a determination of γ is possible from time-dependent measurements, for which one must tag the flavor of the initial B^0 . The effective flavor tagging efficiency at B factories is about 30% (and much smaller at hadron machines), resulting in a doubling of the statistical error relative to the perfect-tag case. As we show below, γ can be determined using untagged data alone. This makes use of events that cannot be tagged or even events that are mis-tagged, regaining a significant part of the sensitivity lost due to the low effective tagging efficiency.

In the present paper we investigate what can be learned from untagged decays $B^0/\bar{B}^0 \rightarrow DK_S$, where the D meson is observed in several decay modes. A potential lower bound on $|\cos\gamma|$ from untagged decays, in which D mesons are observed in CP eigenstates, was noted by Fleischer [22]. We

will go beyond this bound by showing that γ can actually be completely determined in the range $0 < \gamma < \pi$, using only untagged decays. In practice, it is useful to combine information from untagged neutral B decays with information from charged B decays, since the observables related to D decays are common to both cases. This provides an overconstrained information, permitting a more accurate determination of γ than when using B^+ decays alone.

The plan of the paper is as follows. In Sec. II we introduce parallel notation for charged and neutral $B \rightarrow DK$ decays, discussing briefly the relative magnitudes of decay amplitudes in these processes. Section III studies two-body and quasi-two-body D decays, distinguishing between several classes of decay modes. We show that γ can be determined from untagged neutral B decays alone and derive an explicit expression for $\tan^2 \gamma$ in terms of measurable rates. Multibody D decays in $B^0 \rightarrow DK_S$ are studied in Sec. IV. In Sec. V we discuss a way of reducing the number of hadronic parameters by assuming isospin symmetry and by neglecting an annihilation contribution in $B \rightarrow DK$. Finally, Sec. VI concludes. We also add an appendix studying the time dependence in $B^0(t) \rightarrow (K_S \pi^+ \pi^-)_D K_S$.

II. AMPLITUDES IN $B \rightarrow DK$ DECAYS

We define decay amplitudes of $B \rightarrow DK$ for charged B mesons,

$$\begin{aligned} A(B^+ \rightarrow \bar{D}^0 K^+) &\equiv A_c, \\ A(B^+ \rightarrow D^0 K^+) &\equiv A_c r_c e^{i(\delta_c + \gamma)}, \end{aligned} \quad (1)$$

and for neutral B mesons,

$$\begin{aligned} A(B^0 \rightarrow \bar{D}^0 K_S) &\equiv A_n, \\ A(B^0 \rightarrow D^0 K_S) &\equiv A_n r_n e^{i(\delta_n + \gamma)}. \end{aligned} \quad (2)$$

By convention, $A_i \geq 0$, $r_i \geq 0$, and $0 \leq \delta_i \leq 2\pi$ ($i = c, n$). Amplitudes for the CP conjugated decays have the same expressions, but the phase γ occurs with an opposite sign.

Let us discuss briefly the relevant ratios of amplitudes. The amplitude A_c is a combination of color-allowed and color-suppressed contributions, while the amplitude A_n is purely color-suppressed. The ratio A_n/A_c may be estimated in two ways, leading to comparable values. We mention in each case the required approximation:

(i) Measurements of $B^+ \rightarrow \bar{D}^0 K^+$ [14,15,23] and $B^0 \rightarrow \bar{D}^0 K^0$ [19] imply

$$\frac{A_n}{A_c} \simeq \sqrt{\frac{\Gamma(B^0/\bar{B}^0 \rightarrow \bar{D}^0 K_S)}{\Gamma(B^+ \rightarrow \bar{D}^0 K^+)}} = 0.25 \pm 0.07. \quad (3)$$

In addition to a term A_n^2 , the untagged rate in the numerator includes also a smaller term $A_n^2 r_n^2$ which we neglect.

(ii) Using flavor SU(3) [24], one may relate A_n/A_c to a corresponding ratio measured in $B \rightarrow \bar{D} \pi$ [25]:

$$\frac{A_n}{A_c} \simeq \sqrt{\frac{\Gamma(B^0 \rightarrow \bar{D}^0 \pi^0)}{\Gamma(B^+ \rightarrow \bar{D}^0 \pi^+)}} = 0.24 \pm 0.01. \quad (4)$$

This relation is affected by SU(3) breaking corrections and by a small exchange amplitude in $B^0 \rightarrow \bar{D}^0 \pi^0$ [26,27]. SU(3) breaking corrections, which are common in $A(B^+ \rightarrow \bar{D}^0 K^+)/A(B^+ \rightarrow \bar{D}^0 \pi^+)$ and $A(B^0 \rightarrow \bar{D}^0 K_S)/A(B^0 \rightarrow \bar{D}^0 \pi^0)$, cancel in Eq. (4) and do not affect this estimate.

It is more difficult to obtain reliable estimates for r_c and r_n . The two parameters are expected to be smaller than one since they contain a Cabibbo-Kobayashi-Maskawa (CKM) factor $|V_{ub}V_{cs}/V_{cb}V_{us}| \simeq 0.4$. The parameter r_c involves also an unknown color-suppression factor in $\bar{b} \rightarrow \bar{u}c\bar{s}$, while r_n involves the ratio of color-suppression factors in $\bar{b} \rightarrow \bar{u}c\bar{s}$ and $\bar{b} \rightarrow \bar{c}u\bar{s}$. Since the dynamics of $B \rightarrow \bar{D}K$ decays (caused by $\bar{b} \rightarrow \bar{c}u\bar{s}$) and $B \rightarrow DK$ (caused by $\bar{b} \rightarrow \bar{u}c\bar{s}$) are different, color suppression may be different in the two cases. This introduces large uncertainties in r_c and r_n [28].

It is easier to justify an approximate relation between the magnitudes of the two color-suppressed amplitudes $A_c r_c$ and $A_n r_n$. Noting that the two processes $B^+ \rightarrow D^0 K^+$ and $B^0 \rightarrow D^0 K^0$ differ only by the flavor of the spectator quark, one expects

$$A_c r_c \simeq \sqrt{2} A_n r_n. \quad (5)$$

In Sec. V we will discuss the approximation involved in this relation and a way of testing it experimentally. This approximate equality implies that the sensitivity to γ is comparable in charged and neutral B decays, since the sensitivity in each case is governed by the smaller of the two interfering amplitudes. This point provides a major motivation for our study.

III. TWO-BODY AND QUASI-TWO-BODY D DECAYS

Considering decays of \bar{D}^0 and D^0 into a generic two-body or quasi-two-body hadronic state f_D and its CP conjugate \bar{f}_D , we denote the corresponding amplitudes by

$$\begin{aligned} A(\bar{D}^0 \rightarrow f_D) &= A(D^0 \rightarrow \bar{f}_D) \equiv A_f, \\ A(D^0 \rightarrow f_D) &= A(\bar{D}^0 \rightarrow \bar{f}_D) \equiv A_f r_f e^{i\delta_f}, \end{aligned} \quad (6)$$

where by convention $A_f \geq 0$, $r_f \geq 0$, and $0 \leq \delta_f \leq 2\pi$. Here and below we set the weak phase in D decays to zero and neglect D^0 - \bar{D}^0 mixing. The effects of D^0 - \bar{D}^0 mixing can be included as in [29], but are not further discussed here.

Using this notation, one finds expressions for decay rates in charged B decays:

$$\Gamma(B^+ \rightarrow f_D K^+) = A_c^2 A_f^2 [1 + r_c^2 r_f^2 + 2r_c r_f \cos(\delta_c + \delta_f + \gamma)],$$

$$\Gamma(B^- \rightarrow \bar{f}_D K^-) = A_c^2 A_f^2 [1 + r_c^2 r_f^2 + 2r_c r_f \cos(\delta_c + \delta_f - \gamma)],$$

$$\Gamma(B^+ \rightarrow \bar{f}_D K^+) = A_c^2 A_f^2 [r_c^2 + r_f^2 + 2r_c r_f \cos(\delta_c - \delta_f + \gamma)],$$

$$\Gamma(B^- \rightarrow f_D K^-) = A_c^2 A_f^2 [r_c^2 + r_f^2 + 2r_c r_f \cos(\delta_c - \delta_f - \gamma)]. \quad (7)$$

Combining B^+ and B^- decay rates for states involving a common D decay mode, f_D or \bar{f}_D , one finds

$$\begin{aligned} \langle \Gamma(B \rightarrow f_D K_c) \rangle &\equiv \Gamma(B^+ \rightarrow f_D K^+) + \Gamma(B^- \rightarrow f_D K^-) \\ &= A_c^2 A_f^2 [(1+r_c^2)(1+r_f^2) \\ &\quad + 4r_c r_f \cos(\delta_f + \gamma) \cos \delta_c], \\ \langle \Gamma(B \rightarrow \bar{f}_D K_c) \rangle &\equiv \Gamma(B^+ \rightarrow \bar{f}_D K^+) + \Gamma(B^- \rightarrow \bar{f}_D K^-) \\ &= A_c^2 A_f^2 [(1+r_c^2)(1+r_f^2) \\ &\quad + 4r_c r_f \cos(\delta_f - \gamma) \cos \delta_c]. \end{aligned} \quad (8)$$

Studying neutral B decays, one finds similar expressions for untagged decay rates [20]:

$$\begin{aligned} \langle \Gamma(B \rightarrow f_D K_n) \rangle &\equiv \Gamma(B^0 \rightarrow f_D K_S) + \Gamma(\bar{B}^0 \rightarrow f_D K_S) \\ &= A_n^2 A_f^2 [(1+r_n^2)(1+r_f^2) \\ &\quad + 4r_n r_f \cos(\delta_f + \gamma) \cos \delta_n], \\ \langle \Gamma(B \rightarrow \bar{f}_D K_n) \rangle &\equiv \Gamma(B^0 \rightarrow \bar{f}_D K_S) + \Gamma(\bar{B}^0 \rightarrow \bar{f}_D K_S) \\ &= A_n^2 A_f^2 [(1+r_n^2)(1+r_f^2) \\ &\quad + 4r_n r_f \cos(\delta_f - \gamma) \cos \delta_n]. \end{aligned} \quad (9)$$

Individual time-dependent decay rates for $B^0(t) \rightarrow f_D K_S$, $\bar{B}^0(t) \rightarrow f_D K_S$ and their CP conjugates are given in [20], and include more information than the untagged rates. These, however, will not be needed in the following.

The decay rates in Eqs. (8) and (9) display a dependence on two types of quantities. Amplitudes and strong phases in $B \rightarrow DK$ (A_i, r_i, δ_i ; $i=c, n$), which in general obtain different values in charged and neutral B decays, and the corresponding quantities in $D^0/\bar{D}^0 \rightarrow f_D$, (A_f, r_f, δ_f), which are common to both B^+ and B^0 decays. We will refer to these quantities as B and D decay parameters, respectively. In the following we will assume that the D decay quantities A_f and r_f have been measured and are known. They can be obtained through branching ratio measurements in an independent sample of neutral D mesons, flavor-tagged through their production in the decay $D^{*+} \rightarrow D^0 \pi^+$ [30]. In quasi-two-body D decays the three parameters A_f, r_f and δ_f can be determined simultaneously through a complete Dalitz plot analysis. Although the phases δ_f can in principle be measured [10,31–33], we will treat them as unknown, unless indicated otherwise.

Note that the combined B^\pm rates in Eqs. (8) and those in the untagged B^0 decays, Eqs. (9), depend in each case only on two combinations of B decay parameters:

$$X_i \equiv A_i^2 (1+r_i^2), \quad Y_i \equiv 2A_i^2 r_i \cos \delta_i, \quad i=c, n, \quad (10)$$

which obey

$$|Y_i| \leq X_i. \quad (11)$$

We see that the individual branching ratios for $B^0 \rightarrow \bar{D}^0 K_S$ and $B^0 \rightarrow D^0 K_S$, proportional to A_n^2 and $A_n^2 r_n^2$, respectively, cannot be measured from untagged decays alone.

The three D decay parameters A_f, r_f and δ_f in Eqs. (6) depend, of course, on the final state f_D . One may distinguish between three cases for which we give examples:

- (i) $f_D = CP$ state (e.g., $f_{CP+} = K^+ K^-$, $f_{CP-} = K_S \pi^0$), for which $r_{CP\pm} = 1$, $\cos \delta_{CP\pm} = \pm 1$.
- (ii) $f_D =$ flavorless (e.g., $K^{*+} K^-$), for which $r_f = \mathcal{O}(1)$ but generally $r_f \neq 1$, $\delta_f =$ unknown.
- (iii) $f_D =$ flavor state (e.g., $K^+ \pi^-$), for which $r_f \approx \tan^2 \theta_C$ [30], $\delta_f =$ unknown, where θ_C is the Cabibbo angle.

Using Eqs. (9) and the observation (10), it is simple to show that γ may be determined solely from untagged B^0 decays. Consider N different non- CP neutral D decay modes f_D^k ($k=1, \dots, N$) together with their CP conjugates \bar{f}_D^k , as the final states in the $B^0 \rightarrow DK_S$ decay chain. The unknown variables are γ , X_n , Y_n and N strong phases δ_f^k . Equations (9), which provide $2N$ measurables for $N+3$ unknowns, are solvable for $N \geq 3$. That is, γ may be determined from untagged $B^0 \rightarrow DK_S$ decay rates, where D^0 is observed in at least three different non- CP decay modes and their CP conjugates. This argument may be generalized to include other untagged B^0 decays, such as $B^0 \rightarrow D^* K_S (D^* \rightarrow D^0 \pi^0)$. Assuming M different B^0 decay modes of this kind, each of which introduces a pair of unknowns X_n^j and Y_n^j ($j=1, \dots, M$), one has $2MN$ measurables for $2M+N+1$ unknowns. For $M \geq 2$ this set of equations is solvable for $N \geq 2$. Namely, two non- CP decay modes of D^0 are sufficient for determining γ from untagged $B^0 \rightarrow DK_S$ and $B^0 \rightarrow D^* K_S$.

For a CP eigenstate the strong phase $\delta_{f(CP)}$ is either 0 or π . In this case the two equations in Eqs. (9) become identical and provide a single measurable. Choosing the decay modes f_D^k to be (i) an even- CP state, (ii) an odd- CP state, and (iii) a single non- CP eigenstate and its CP conjugate (involving an unknown phase δ_f), one can solve the four equations for γ , X_n , Y_n and δ_f . For this case we now derive an explicit expression for $\tan^2 \gamma$ in terms of measurable rates. The derivation holds for both charged and neutral B decays.

Using Eqs. (8) and (9), one has

$$\langle \Gamma(B \rightarrow f_{CP\pm} K_i) \rangle = 2A_{CP\pm}^2 [X_i \pm Y_i \cos \gamma] \quad (i=c, n), \quad (12)$$

where the two signs on the right-hand side correspond to positive and negative CP eigenstates. Adding and subtracting rates for even- CP and odd- CP eigenmodes, one finds

$$\Sigma_{CP}^i \equiv \frac{\langle \Gamma(B \rightarrow f_{CP+} K_i) \rangle}{2A_{CP+}^2} + \frac{\langle \Gamma(B \rightarrow f_{CP-} K_i) \rangle}{2A_{CP-}^2} = 2X_i, \quad (13)$$

$$\begin{aligned}\Delta_{CP}^i &\equiv \frac{\langle \Gamma(B \rightarrow f_{CP+} K_i) \rangle}{2A_{CP+}^2} - \frac{\langle \Gamma(B \rightarrow f_{CP-} K_i) \rangle}{2A_{CP-}^2} \\ &= 2Y_i \cos \gamma \quad (i=c,n).\end{aligned}\quad (14)$$

These definitions apply in practice to sums over individual CP states. Equation (13) provides the most direct way to determine X_c and X_n . We define CP -conserving rate asymmetries between even and odd CP states:

$$\mathcal{A}_{CP}^i \equiv \frac{\Delta_{CP}^i}{\Sigma_{CP}^i} \quad (i=c,n).\quad (15)$$

Using Eq. (11), one obtains two potential inequalities for $\cos \gamma$ in terms of these ratios:

$$|\cos \gamma| > |\mathcal{A}_{CP}^i| \quad (i=c,n).\quad (16)$$

This inequality holds separately for charged and neutral B decays. The inequality for neutral B decays was noted in [22].

As mentioned above, in order to determine γ from untagged neutral B decays, one needs in addition to the two rates for CP eigenstates given in Eq. (12) ($i=n$) two rate measurements for a non- CP state f_D and its CP conjugate \bar{f}_D . These rates are given by the two equations in Eqs. (9). We denote the sum and difference of these rates and of the corresponding rates in charged B decays by

$$\begin{aligned}\Sigma_f^i &\equiv \frac{\langle \Gamma(B \rightarrow \bar{f}_D K_i) \rangle + \langle \Gamma(B \rightarrow f_D K_i) \rangle}{A_f^2(1+r_f^2)}, \\ \Delta_f^i &\equiv \frac{\langle \Gamma(B \rightarrow \bar{f}_D K_i) \rangle - \langle \Gamma(B \rightarrow f_D K_i) \rangle}{A_f^2(1+r_f^2)} \\ &(i=c,n),\end{aligned}\quad (17)$$

which imply CP -violating asymmetries:

$$\mathcal{A}_f^i \equiv \Delta_f^i / \Sigma_f^i \quad (i=c,n).\quad (18)$$

It is then straightforward to show that $\tan^2 \gamma$ is proportional to $(\mathcal{A}_f^i)^2$ and is given by

$$\begin{aligned}\tan^2 \gamma &= \frac{(\Delta_f^i)^2}{\rho_f^2 (\Delta_{CP}^i)^2 - (\Sigma_{CP}^i - \Sigma_f^i)^2} \\ &= \frac{(\mathcal{A}_f^i)^2 (\Sigma_f^i / \Sigma_{CP}^i)^2}{\rho_f^2 (\mathcal{A}_{CP}^i)^2 - (1 - \Sigma_f^i / \Sigma_{CP}^i)^2} \quad (i=c,n),\end{aligned}\quad (19)$$

where

$$\rho_f \equiv \frac{2r_f}{1+r_f^2}.\quad (20)$$

This result applies to both charged and neutral B mesons. We stress that when determining γ we do not rely on separating the two terms A_i^2 and $A_i^2 r_i^2$ contributing to X_i .

While the possibility of measuring γ from untagged B^0 decays alone is interesting, the most efficient way to determine the weak phase would be to combine information from decays of charged B decays and untagged neutral B decays. The derivation we have just presented applies also to charged B decays alone. One needs to measure only the combined B^\pm rates given in Eqs. (8), without needing to separate the small $A_c^2 r_c^2$ term. Thus, Eq. (19) also gives $\tan^2 \gamma$ in terms of decay rates for combined $B^\pm \rightarrow DK^\pm$ events, where D mesons are observed in decays into an even- CP eigenstate, an odd- CP eigenstate, and a non- CP flavorless state or a flavor state. We see that, in principle, a determination of γ does not require measuring CP asymmetries in $B^\pm \rightarrow f_{CP} K^\pm$. These asymmetry measurements, for even- CP and odd- CP states [14,15], provide additional useful information. Using all these measurements together with rate measurements for untagged neutral B decays will lead to a more accurate determination of γ than when using only charged B mesons.

Equation (19) displays an explicit dependence of γ on rate measurements defined in Eqs. (13), (14) and (17). We see that $\tan \gamma$ is proportional to the CP asymmetries $\mathcal{A}_f^i \sim r_f \cos \delta_i \sin \gamma$ ($i=c,n$), indicating that the sensitivity for measuring γ increases with r_f . The sensitivity depends also on the value of δ_f . In the extreme case that δ_f vanishes (mod π) the two rates in Eqs. (8) [and in Eqs. (9)] become equal and γ cannot be extracted. For the two-body flavor state $f=K^+\pi^-$, the phase δ_f vanishes in the SU(3) symmetry limit. However, SU(3) breaking effects are known to be large in D decays. Consequently, sizable values of δ_f have been calculated for this final state in several models [34]. This phase can be measured at a charm factory [31,32]. There are experimental indications for small phases in two cases of quasi-two-body states, $f=K^{*+}\pi^-$, where $\delta_f=(12 \pm 3)^\circ$ (mod π) was measured [17], and $f=\rho^+\pi^-$ where $\delta_f=(4 \pm 3 \pm 4)^\circ$ was measured [35]. These results were obtained by studying the Dalitz plot of $D^0 \rightarrow K_S \pi^+ \pi^-$ and $D^0 \rightarrow \pi^+ \pi^- \pi^0$, respectively. No measurement exists for δ_f in $D^0 \rightarrow K^{*+} K^-$, which can be measured by studying the Dalitz plot of $D^0 \rightarrow K^+ K^- \pi^0$ [33].

As we note in the next section, a complete Dalitz plot analysis of three body D^0 decays involves other strong phases, which are large in regions where two resonances overlap or when resonances interfere with non-resonant contributions. These phases will be shown to be useful when studying γ in $B \rightarrow DK$, where the D meson is observed in a three-body final state.

IV. $B \rightarrow DK_S$ OBSERVED IN MULTI-BODY D DECAYS

The study of untagged $B^0 \rightarrow DK_S$ presented in the previous section for D mesons decaying in two-body modes may be extended to multibody decays. To be specific, we focus on the case of the three-body D decay,

$$D \rightarrow K_S \pi^- \pi^+, \quad (21)$$

following the discussion of $B^\pm \rightarrow DK^\pm$ in [9]. In order to make our point, we start with a model-independent approach. We also explain how modeling the amplitude for D^0

$\rightarrow K_S \pi^- \pi^+$ in terms of a sum of a given set of intermediate resonances [36], as done recently in [17], may help in reducing the experimental error in γ .

We denote the amplitude for $D^0 \rightarrow K_S \pi^- \pi^+$ at a given point in the Dalitz plot by

$$A(D^0 \rightarrow K_S(p_1) \pi^-(p_2) \pi^+(p_3)) \equiv A(s_{12}, s_{13}) e^{i\delta(s_{12}, s_{13})}, \quad (22)$$

where $s_{ij} \equiv (p_i + p_j)^2$. As in two body decays, we use the convention $A(s_{12}, s_{13}) \geq 0$ and $0 \leq \delta(s_{12}, s_{13}) \leq 2\pi$. Assuming that CP is conserved in this decay, one has

$$\begin{aligned} A(\bar{D}^0 \rightarrow K_S(p_1) \pi^-(p_2) \pi^+(p_3)) \\ = A(D^0 \rightarrow K_S(p_1) \pi^-(p_3) \pi^+(p_2)) \\ \equiv A(s_{13}, s_{12}) e^{i\delta(s_{13}, s_{12})}. \end{aligned} \quad (23)$$

That is, the (complex) decay amplitude for \bar{D}^0 at a given point (s_{12}, s_{13}) in the Dalitz plot equals the decay amplitude for D^0 at a point (s_{13}, s_{12}) obtained by reflection across a symmetry axis corresponding to exchanging the momenta of the two pions.

The density of events in the D decay Dalitz plot for untagged $B \rightarrow (K_S \pi^- \pi^+)_D K_S$ is obtained using Eqs. (2), (22) and (23) (or from the time dependence in the Appendix),

$$\begin{aligned} \frac{d^2\Gamma}{ds_{12} ds_{13}} (B^0/\bar{B}^0 \rightarrow [K_S(p_1) \pi^-(p_2) \pi^+(p_3)]_D K_S) \\ = A_n^2 [A^2(s_{12}, s_{13}) + A^2(s_{13}, s_{12})] (1 + r_n^2) \\ + 4r_n A(s_{12}, s_{13}) A(s_{13}, s_{12}) \\ \times \cos[\delta(s_{12}, s_{23}) - \delta(s_{13}, s_{12}) + \gamma] \cos \delta_n. \end{aligned} \quad (24)$$

This is in complete analogy with the first of Eqs. (9). The second equation, describing the density at the point of reflection across the symmetry axis, involves an opposite sign for γ . Integrating Eq. (24) over an area (a bin) i lying below the symmetry axis and over a corresponding symmetry-reflected area \bar{i} lying above the symmetry axis, one has¹

$$\begin{aligned} \langle \Gamma_i \rangle &\equiv \int_i d\Gamma(B^0/\bar{B}^0 \rightarrow [K_S \pi^- \pi^+]_D K_S) \\ &= X_n(T_i + T_{\bar{i}}) + 2Y_n[c_i \cos \gamma - s_i \sin \gamma], \\ \langle \Gamma_{\bar{i}} \rangle &\equiv \int_{\bar{i}} d\Gamma(B^0/\bar{B}^0 \rightarrow [K_S \pi^- \pi^+]_D K_S) \\ &= X_n(T_i + T_{\bar{i}}) + 2Y_n[c_i \cos \gamma + s_i \sin \gamma], \end{aligned} \quad (25)$$

where we define

¹The method outlined here applies to any multibody D^0 decay with the set of equations (25) unchanged. If the i th bin is in the phase space of a final state f_D , then the \bar{i} th bin is a CP transformed bin in the phase space of the corresponding \bar{f}_D state.

$$T_i \equiv \int_i ds_{12} ds_{13} A^2(s_{13}, s_{23}),$$

$$\begin{aligned} c_i &\equiv \int_i ds_{12} ds_{13} A(s_{12}, s_{13}) A(s_{13}, s_{12}) \\ &\times \cos[\delta(s_{12}, s_{23}) - \delta(s_{13}, s_{12})], \end{aligned}$$

$$\begin{aligned} s_i &\equiv \int_i ds_{12} ds_{13} A(s_{12}, s_{13}) A(s_{13}, s_{12}) \sin[\delta(s_{12}, s_{23}) \\ &- \delta(s_{13}, s_{12})]. \end{aligned} \quad (26)$$

The partial rates T_i in D decays may be measured using flavor-tagged D^0 decays and are assumed to be known. The other D decay variables, c_i and s_i , which in principle can be measured model independently at a charm factory (up to a sign ambiguity in s_i), will nonetheless be taken as unknown. Consider k different bins i lying below the symmetry axis, each contributing two unknowns c_i and s_i . Together with X_n , Y_n and γ , there are $2k + 3$ unknowns. Equations (25), which provide $2k$ measurables ($\langle \Gamma_i \rangle$ and $\langle \Gamma_{\bar{i}} \rangle$), are therefore unsolvable.

The situation changes when one measures another neutral B decay of this type, e.g. the sequence $B^0 \rightarrow D^* K_S$, $D^* \rightarrow D^0 \pi^0$, $D^0 \rightarrow K_S \pi^+ \pi^-$, which introduces a pair of new variables analogous to X_n , Y_n . In this case one has $4k$ measurables for $2k + 5$ unknowns, a solution for which requires $k \geq 3$. That is, γ may be determined by measuring partial rates in the two untagged neutral B decay modes, $B \rightarrow DK_S$ and $B \rightarrow D^* K_S$, for at least three pairs of Dalitz plot bins in $D \rightarrow K_S \pi^+ \pi^-$.

A more powerful approach is to combine information from all the untagged neutral B decays and charged B decays, using both multibody and two-body D decays. For instance, in analogy with Eqs. (7), the decays $B^\pm \rightarrow (K_S \pi^+ \pi^-)_D K^\pm$ provide four measurables for each bin [9], instead of the two in Eqs. (25). Combining charged and untagged neutral $B \rightarrow DK$ decays, where $D \rightarrow K_S \pi^+ \pi^-$, yields $6k$ measurables for $2k + 6$ unknowns, $c_i, s_i, A_c, r_c, \delta_c, X_n, Y_n$ and γ . Therefore, two pairs of bins provide an overconstrained system of equations for determining γ .

The two equations (25) are not mutually independent when $s_i = 0$, in analogy with the singularity noted in Eq. (19) when $\delta_f = 0 \pmod{\pi}$. However, the relevant strong phase differences which determine s_i are large at least in some regions of the Dalitz plot of $D^0 \rightarrow K_S \pi^+ \pi^-$. Consider, for instance, the two overlapping regions of a vertical band describing the Cabibbo-allowed mode $K^{*-} \pi^+$ and a horizontal band describing the doubly Cabibbo-suppressed mode $K^{*+} \pi^-$ with a diagonal band representing the Cabibbo-allowed mode $\rho^0 \bar{K}^0$. The local strong phases which determine s_i for these two regions are the phase differences between amplitudes describing the sum of the $K^{*-} \pi^+$ and $\rho^0 \bar{K}^0$ contributions and the sum of the $K^{*+} \pi^-$ and $\rho^0 \bar{K}^0$ contributions. These phases are large and vary a lot over the

overlapping regions because of the two largely different $K^*\pi$ contributions in the two amplitudes.

One can reduce the number of unknowns appearing in the determination of γ , if the unknowns coming from the D decay, c_i and s_i , appearing in Eqs. (25), are determined independently. This can be done by assuming a Breit-Wigner (BW) form for the intermediate resonances contributing to this decay [9]. The parameters of the model describing the D decay amplitude can then be fitted to data of tagged D decays, which are abundant at B factories. The observables in Eqs. (25) now depend only on three unknowns, X_n , Y_n and γ .

It is hard to quantify the theoretical error introduced by assuming a BW form. One way to proceed is to change the number of resonances and see how the sensitivity changes. This is only a partial determination of the error. Another source of error is the accuracy of the BW assumption. This can be determined by the goodness of the fit to the tagged D decays or by using a different model for resonances, such as a K-matrix model for wide resonances [37]. A rough estimate of the theoretical error caused by assuming a superposition of BW amplitudes is about 10° [17]. Further studies are required in order to evaluate possible contributions of non-BW terms in the D decay amplitude and their effect on determining γ . As mentioned, this model dependence can be avoided by measuring the parameters c_i and s_i at a charm factory.

V. USING ISOSPIN AND NEGLECTING AN ANNIHILATION AMPLITUDE

As already mentioned, the most powerful approach for improving the determination of γ is to combine charged B decays with the information from untagged neutral B decays. Adding untagged neutral B decays to a sample of charged B decays with the same D decay final states introduces only two unknown parameters, X_n and Y_n . Here we discuss an approximation which may be used to reduce the number of parameters further. This introduces a theoretical error in γ . Yet it is worthwhile considering such an approximation as long as this error is smaller than the statistical error.

We recall two isospin relations [38], one for $\bar{b} \rightarrow \bar{c}u\bar{s}$ transitions,

$$A(B^0 \rightarrow D^- K^+) = A(B^+ \rightarrow \bar{D}^0 K^+) - A(B^0 \rightarrow \bar{D}^0 K^0), \quad (27)$$

and another for $\bar{b} \rightarrow \bar{u}c\bar{s}$ transitions,

$$A(B^0 \rightarrow D^0 K^0) = A(B^+ \rightarrow D^0 K^+) + A(B^+ \rightarrow D^+ K^0). \quad (28)$$

The amplitude of $B^+ \rightarrow D^+ K^0$ is pure annihilation, and is expected to be smaller than the other two amplitudes in the last relation [24]. The absence of rescattering effects, which may enhance this amplitude to a level comparable to the other two amplitudes in this relation, can be tested [39] by setting very stringent experimental bounds on the branching ratio for $B^+ \rightarrow D^+ K^0$. Neglecting $A(B^+ \rightarrow D^+ K^0)$, Eq. (28) reduces to [38]

$$A(B^0 \rightarrow D^0 K^0) = A(B^+ \rightarrow D^0 K^+). \quad (29)$$

Namely, the two color-suppressed amplitudes have equal magnitudes and equal strong phases. Note that the error due to isospin violation, at most a few percent, is likely to be much smaller than that involved in neglecting the annihilation amplitude.

Equations (27) and (29) may be used to simplify the determination of γ when combining B^\pm decays and untagged B^0 decays. The two equations imply

$$A_c^2 + 2A_n^2 - 2\sqrt{2}A_c A_n \cos(\delta_n - \delta_c) = \Gamma(B^0 \rightarrow D^- K^+),$$

$$\sqrt{2}A_n r_n = A_c r_c. \quad (30)$$

The right-hand side of the first equation, which involves a color-allowed process, has already been measured [40] and will be assumed to be given. Equations (30) reduce the six parameters describing charged and neutral $B \rightarrow DK$ decays, A_i, r_i, δ_i ($i = c, n$), to four independent ones. The measurable parameters in untagged $B^0 \rightarrow DK_S$ decays, $X_n \equiv A_n^2(1 + r_n^2)$ and $Y_n \equiv 2A_n^2 r_n \cos \delta_n$, can now be expressed in terms of the three B^+ decay parameters and a single B^0 decay parameter (δ_n , for instance). That is, under the above assumption, adding information from untagged neutral B^0 decays to studies of B^\pm decays involves a single new unknown parameter instead of two parameters. This is expected to reduce the statistical error in determining γ .

VI. CONCLUDING REMARKS

Before concluding let us make several general comments:

(i) Since the data set of $B^+ \rightarrow DK^+$ is larger than that of $B^0 \rightarrow DK_S$, hadronic parameters such as X_c are easier to measure than X_n . As noted, the approximate relation $\sqrt{2}A_n r_n \approx A_c r_c$ implies comparable sensitivities to γ in charged and neutral B decays. The sensitivity in B^0 decays is smaller by a factor of $\sqrt{2}$ since only K_S mesons are experimentally useful. Moreover, the detection efficiency for K_S is about a factor of 2 smaller than that for charged kaons. This is expected to reduce somewhat the effect of neutral B decays on determining γ .

(ii) Our study focused on $B \rightarrow DK$ decays. It can be extended to multibody B decays, including $B^+ \rightarrow D^* K^+$ and $B^0 \rightarrow D^* K_S$, where $D^* \rightarrow D\pi^0$, as well as to the self-tagged decays $B^+ \rightarrow D^{(*)} K^{*+}$ and $B^0 \rightarrow D^{(*)} K^{*0}$. This would add to the statistical power of the analysis since the parameters related to D decays are common to these processes and to $B \rightarrow DK$.

(iii) The method we discussed for B^0 decays can be applied also to B_s decays, replacing the K_S by ϕ, η', η . In that case the advantage of being able to use untagged data is greater, because the hadronic environment where B_s decays will be studied makes flavor tagging less efficient. We have neglected the width difference between the two neutral B meson states, which is a very good approximation for non-strange B mesons. In the case of B_s , the width difference is expected to be non-negligible and may be taken into account in a straightforward manner.

(iv) In our discussion we assumed that strong phases in D decays are unknown and need to be determined from the analysis simultaneously with γ . As we already mentioned, the strong phases in two-body and quasi-two-body D decays can be determined independently [10,31–33]. The strong phases in three-body decays may also be determined by assuming that the D decay amplitude is given as a sum of Breit-Wigner amplitudes. Knowledge of strong phases would imply, for instance, that fewer D decay modes are needed in order to determine γ from untagged decays alone. In practice, this implies that a combined fit of the data to fewer hadronic parameters will result in a smaller error in γ .

(v) The extraction of γ from $B^\pm \rightarrow DK^\pm$ involves a number of discrete ambiguities [5,41]. The ambiguities in untagged B^0 decays may be identified in Eqs. (9) which are invariant under

$$\begin{aligned} P_{\text{ex}} &\equiv \{\gamma \rightarrow \delta_f, \delta_f \rightarrow \gamma\}, \\ P_- &\equiv \{\gamma \rightarrow -\gamma, \delta_f \rightarrow -\delta_f\}, \\ P_\pi &\equiv \{\gamma \rightarrow \gamma + \pi, \delta_f \rightarrow \delta_f + \pi \text{ or } \delta_n \rightarrow \delta_n + \pi\}. \end{aligned} \quad (31)$$

Once several two-body D decay modes are combined or once a multibody D decay mode is used, the first two ambiguities may be resolved. The ambiguity P_{ex} is lifted, since δ_f is not expected to be the same for all two-body and quasi-two-body D decay modes, and is known to change over the Dalitz plot in three-body decays. Measuring the sign of s_i in Eqs. (25) through a fit to a sum of Breit-Wigner resonance functions would resolve the P_- ambiguity. This introduced essentially no model dependence, since one needs only the sign of s_i , which is easily determined in the vicinity of a BW resonance. Resolving P_- avoids the ambiguity $\gamma \rightarrow \pi - \gamma$, a combination of P_- and P_π , which is particularly problematic in view of the proximity of γ to $\pi/2$ [41]. The only remaining ambiguity is $\gamma \rightarrow \gamma + \pi$. This ambiguity is the least problematic, since the two corresponding values of γ are maximally separated.

To conclude, we have studied the information obtained from untagged neutral B decays of the type $B \rightarrow DK_S$ involving several D decay modes. We have shown that these measurements alone can, in principle, determine γ . Of course, B^0 tagging information, while limited, can only improve this determination. By combining information from untagged B^0 decays with that obtained in corresponding charged B decays one gains statistics, thereby permitting a more accurate determination of γ . While statistics are limited, one may neglect an annihilation amplitude in $B \rightarrow DK$, reducing by 1 the number of hadronic parameters and resulting in a smaller experimental error in γ . This introduces a theoretical error in γ which must be further studied.

ACKNOWLEDGMENTS

The work of Y.G. is supported in part by the Department of Energy under contracts DE-AC03-76SF00515 and DE-FG03-92ER40689. The work of A.S. is supported by the U.S. Department of Energy under contract DE-FG03-

93ER40788. The work of J.Z. is supported in part by EU grant HPRN-CT-2002-00277 and by the Ministry of Education, Science and Sport of the Republic of Slovenia.

APPENDIX: TIME-DEPENDENT $B \rightarrow DK_S$ WITH MULTIBODY D DECAYS

In this appendix we provide a formalism allowing the extraction of γ from time-dependent rates in $B^0 \rightarrow f_D K_S$ where f_D is a multibody final state. As we show, this not only serves the purpose of determining γ , but also helps resolve the current twofold ambiguity, $\beta \rightarrow \pi/2 - \beta$. For simplicity we take f_D to be the three-body final state $K_S \pi^+ \pi^-$ studied in Sec. IV.

Time-dependent partial rates, integrated over a bin i in the Dalitz plot of $D \rightarrow K_S \pi^+ \pi^-$ lying below the symmetry axis and over a corresponding symmetry-reflected bin \bar{i} above the axis, are readily calculated for initial B^0 and \bar{B}^0 states (a positive B meson bag parameter is assumed [42]):

$$\begin{aligned} \Gamma_i &\equiv \int_i d\Gamma(B^0(t) \rightarrow (K_S \pi^- \pi^+)_D K_S) \\ &= e^{-\Gamma_B t} A_n^2 \left\{ I_i^+ \cos^2 \left(\frac{\Delta m_B t}{2} \right) + I_i^- \sin^2 \left(\frac{\Delta m_B t}{2} \right) \right. \\ &\quad \left. + S_i \sin(\Delta m_B t) \right\}, \end{aligned} \quad (A1)$$

$$\begin{aligned} \bar{\Gamma}_i &\equiv \int_i d\Gamma(\bar{B}^0(t) \rightarrow (K_S \pi^- \pi^+)_D K_S) \\ &= e^{-\Gamma_B t} A_n^2 \left\{ I_i^- \cos^2 \left(\frac{\Delta m_B t}{2} \right) + I_i^+ \sin^2 \left(\frac{\Delta m_B t}{2} \right) \right. \\ &\quad \left. - S_i \sin(\Delta m_B t) \right\}, \end{aligned} \quad (A2)$$

$$\begin{aligned} \Gamma_{\bar{i}} &\equiv \int_{\bar{i}} d\Gamma(B^0(t) \rightarrow (K_S \pi^- \pi^+)_D K_S) \\ &= e^{-\Gamma_B t} A_n^2 \left\{ I_{\bar{i}}^+ \cos^2 \left(\frac{\Delta m_B t}{2} \right) + I_{\bar{i}}^- \sin^2 \left(\frac{\Delta m_B t}{2} \right) \right. \\ &\quad \left. + S_{\bar{i}} \sin(\Delta m_B t) \right\}, \end{aligned} \quad (A3)$$

$$\begin{aligned} \bar{\Gamma}_{\bar{i}} &\equiv \int_{\bar{i}} d\Gamma(\bar{B}^0(t) \rightarrow (K_S \pi^- \pi^+)_D K_S) \\ &= e^{-\Gamma_B t} A_n^2 \left\{ I_{\bar{i}}^- \cos^2 \left(\frac{\Delta m_B t}{2} \right) + I_{\bar{i}}^+ \sin^2 \left(\frac{\Delta m_B t}{2} \right) \right. \\ &\quad \left. - S_{\bar{i}} \sin(\Delta m_B t) \right\}. \end{aligned} \quad (A4)$$

The six observables determined from the time dependence, $I_i^\pm, I_{\bar{i}}^\pm, S_i$ and $S_{\bar{i}}$, are defined in terms of the quantities in Eqs. (26):

$$I_i^\pm \equiv T_{\bar{i}} + r_n^2 T_i + 2r_n [\cos(\gamma \pm \delta_n) c_i \mp \sin(\gamma \pm \delta_n) s_i], \quad (\text{A5})$$

$$S_i \equiv r_n T_{\bar{i}} \sin(2\beta + \gamma - \delta_n) + [\sin(2\beta) c_i - \cos(2\beta) s_i] \\ + r_n^2 [\sin(2\beta + 2\gamma) c_i + \cos(2\beta + 2\gamma) s_i] \\ + r_n T_i \sin(2\beta + \gamma + \delta_n). \quad (\text{A6})$$

Expressions for the observables $I_{\bar{i}}^\pm$ and $S_{\bar{i}}$ are obtained from Eqs. (A5) and (A6) by replacing $T_i \leftrightarrow T_{\bar{i}}$, $s_i \rightarrow -s_i$. Note that $I_{i,\bar{i}}^\pm$ correspond directly to the partial decay widths $\Gamma_{i,\bar{i}}^\pm$ for $B^\pm \rightarrow DK^\pm$ defined in [9].

Dividing the Dalitz plot into k pairs of bins, i and \bar{i} , the $6k$ observables permit an extraction of γ . There are $2k+4$ unknowns, $c_i, s_i, A_n, r_n, \delta_n, \gamma$ [$\sin(2\beta)$ is assumed to be known], so that the system is solvable for $k \geq 1$. Namely, in order to determine γ from time-dependent decay rates into $(K_S \pi^+ \pi^-)_D K_S$, it is sufficient to divide the D decay Dalitz plot into two bins, symmetric with respect to the symmetry axis.

The solution for γ involves a fourfold discrete ambiguity. Equations (A1)–(A4) are invariant under the following four independent discrete transformations:

$$P_\pi^\gamma \equiv \{\gamma \rightarrow \gamma + \pi, \delta_n \rightarrow \delta_n + \pi\},$$

$$P'_\pi \equiv \{\gamma \rightarrow \gamma + \pi, \beta \rightarrow \beta + \pi/2, c_i \rightarrow -c_i, s_i \rightarrow -s_i\},$$

$$P_\pi^\beta \equiv \{\beta \rightarrow \beta + \pi\},$$

$$P_- \equiv \{\gamma \rightarrow -\gamma, \beta \rightarrow \pi/2 - \beta, \delta_n \rightarrow -\delta_n, s_i \rightarrow -s_i\}. \quad (\text{A7})$$

The P'_π ambiguity can be resolved model independently either by using the sign of $\sin 2\beta$ or by measuring the sign of c_i at a $\Psi(3770)$ charm factory [9]. The P_- ambiguity can be resolved if one determines the sign of s_i by fitting the Dalitz plot to a sum of Breit-Wigner forms. (See Sec. VI.) Note that resolving the P_- ambiguity in this way leads to the determination of the sign of $\cos(2\beta)$ in an essentially model-independent way. It also determines the sign of γ or equivalently the sign of $\cos(2\alpha)$. Fixing the sign of $\cos(2\beta)$ is a consequence of knowing the sign of s_i in multibody decays. This is impossible in two-body D decays, where the sign of $\sin \delta_f$ cannot be determined. The remaining two ambiguities, P_π^γ and P_π^β , cannot be resolved without further theoretical input.

-
- [1] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002); Belle Collaboration, K. Abe *et al.*, Phys. Rev. D **66**, 071102 (2002).
- [2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] Belle Collaboration, K. Abe *et al.*, hep-ex/0401029; BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 281802 (2002); A. Jawahery, talk at the 21st International Symposium on Lepton and Photon Interactions at High Energies, Fermilab, Batavia, IL, 2003.
- [4] M. Gronau and J.L. Rosner, Phys. Rev. D **65**, 093012 (2002); **66**, 053003 (2002); **66**, 119901(E) (2002).
- [5] M. Gronau and D. Wyler, Phys. Lett. B **265**, 172 (1991); M. Gronau, Phys. Rev. D **58**, 037301 (1998).
- [6] M. Gronau and D. London, Phys. Lett. B **253**, 483 (1991).
- [7] D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997); Phys. Rev. D **63**, 036005 (2001).
- [8] Y. Grossman, Z. Ligeti, and A. Soffer, Phys. Rev. D **67**, 071301(R) (2003).
- [9] A. Giri, Y. Grossman, A. Soffer, and J. Zupan, Phys. Rev. D **68**, 054018 (2003).
- [10] D. Atwood and A. Soni, Phys. Rev. D **68**, 033003 (2003).
- [11] I. Dunietz, Phys. Lett. B **270**, 75 (1991).
- [12] D. Atwood, G. Eilam, M. Gronau, and A. Soni, Phys. Lett. B **341**, 372 (1995); R. Aleksan, T.C. Petersen, and A. Soffer, Phys. Rev. D **67**, 096002 (2003); M. Gronau, Phys. Lett. B **557**, 198 (2003).
- [13] D. Atwood and A. Soni, hep-ph/0312100.
- [14] Belle Collaboration, S.K. Swain *et al.*, Phys. Rev. D **68**, 051101 (2003).
- [15] BABAR Collaboration, B. Aubert *et al.*, hep-ex/0311032.
- [16] M. Gronau, Proceedings of Flavor Physics and CP Violation (FPCP 2003), Paris, France, 2003, hep-ph/0306308.
- [17] Belle Collaboration, K. Abe *et al.*, hep-ex/0308043.
- [18] BELLE Collaboration, K. Abe *et al.*, hep-ex/0307074; BABAR Collaboration, B. Aubert, Phys. Rev. D **69**, 051101 (2004).
- [19] Belle Collaboration, P. Krokovny *et al.*, Phys. Rev. Lett. **90**, 141802 (2003).
- [20] B. Kayser and D. London, Phys. Rev. D **61**, 116013 (2000).
- [21] D. Atwood and A. Soni, Phys. Rev. D **68**, 033009 (2003).
- [22] R. Fleischer, Phys. Lett. B **562**, 234 (2003); Nucl. Phys. **B659**, 321 (2003).
- [23] CLEO Collaboration, A. Bornheim *et al.*, Phys. Rev. D **68**, 052002 (2003).
- [24] M. Gronau, O.F. Hernandez, D. London, and J.L. Rosner, Phys. Rev. D **50**, 4529 (1994); **52**, 6356 (1995).
- [25] BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **88**, 052002 (2002); CLEO Collaboration, T.E. Coan *et al.*, *ibid.* **88**, 062001 (2002); BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. D **69**, 032004 (2004).
- [26] C.W. Chiang and J.L. Rosner, Phys. Rev. D **67**, 074013 (2003).
- [27] Belle Collaboration, P. Krokovny *et al.*, Phys. Rev. Lett. **89**, 231804 (2002); BABAR Collaboration, B. Aubert *et al.*, *ibid.* **90**, 181803 (2003).
- [28] After the submission of this paper an upper limit, $r_c < 0.22$ (90% C.L.), was reported by BABAR Collaboration, B. Aubert *et al.*, Report No. SLAC-PUB-10333, hep-ex/0402024.
- [29] J.P. Silva and A. Soffer, Phys. Rev. D **61**, 112001 (2000).
- [30] CLEO Collaboration, R. Godang *et al.*, Phys. Rev. Lett. **84**, 5038 (2000); FOCUS Collaboration, J.M. Link *et al.*, *ibid.* **86**, 2955 (2001); BABAR Collaboration, B. Aubert *et al.*, *ibid.* **91**, 171801 (2003).

- [31] A. Soffer, hep-ex/9801018.
- [32] M. Gronau, Y. Grossman, and J.L. Rosner, Phys. Lett. B **508**, 37 (2001).
- [33] J.L. Rosner and D.A. Suprun, Phys. Rev. D **68**, 054010 (2003).
- [34] See references in [32].
- [35] CLEO Collaboration, V.V. Frolov *et al.*, hep-ex/0306048.
- [36] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **308**, 435 (1993); E687 Collaboration, P.L. Frabetti *et al.*, *ibid.* **331**, 217 (1994); CLEO Collaboration, H. Muramatsu *et al.*, Phys. Rev. Lett. **89**, 251802 (2002).
- [37] FOCUS Collaboration, J.M. Link *et al.*, Phys. Lett. B **585**, 200 (2004).
- [38] M. Gronau and J.L. Rosner, Phys. Lett. B **439**, 171 (1998); J.H. Jang and P. Ko, Phys. Rev. D **58**, 111302 (1998); see also Z.z. Xing, *ibid.* **58**, 093005 (1998).
- [39] B. Blok, M. Gronau, and J.L. Rosner, Phys. Rev. Lett. **78**, 3999 (1997).
- [40] BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **87**, 111801 (2001).
- [41] A. Soffer, Phys. Rev. D **60**, 054032 (1999).
- [42] Y. Grossman, B. Kayser, and Y. Nir, Phys. Lett. B **415**, 90 (1997).