

---

**Errata**


---

**Scale Invariance and Spectral Forms for Conformal Stress Tensors**, Kimball A. Milton [Phys. Rev. D 7, 1120 (1973)]. Equation (3.7) defines  $v^2$  and  $w^2$ , not  $v$  and  $w$ . The sign of Eq. (5.27) is incorrect, which changes the sign of  $\Pi_1$  and  $\Pi_2$ . Consequently, the signs of Eqs. (5.45), (5.48), and of the integral in (5.52), should be reversed.  $\hat{\Lambda}$  is correct as written, but the anomaly, Eq. (5.64), should appear as

$$\mathcal{G} = - \left[ 4P^2 - \frac{(PQ)^2}{Q^2} \right] (\Pi - 3\Lambda)(Q=0).$$

Finally, at the top of p. 1131, the limit referred to is when  $Q=0$ , not  $Q^2=0$ .

**Massless, Euclidean Quantum Electrodynamics on the 5-Dimensional Unit Hypersphere**, Stephen L. Adler [Phys. Rev. D 6, 3445 (1972)]. 1. Page 3447, Table I. The normalizing factors for the 5-dimensional and Euclidean electron propagators should read  $(-1/\pi^2)$  and  $(-1/2\pi^2)$ , respectively, instead

of  $(-i/\pi^2)$  and  $(-i/2\pi^2)$  as given. (Corresponding changes should be made elsewhere in the paper.) 2. Page 3449, first column, third line following Eq. (44): Eq. (8a) should read Eq. (8).

**Symmetries and Nonsymmetries of the Relativistic Quark Model**, J. L. Rosner [Phys. Rev. D 6, 1781 (1972)]. Equation (C14), instead of

$$\langle K^0 \bar{d} | \bar{s} \rangle = 1/\sqrt{3},$$

should read

$$\langle K^0 \bar{d} | \bar{s} \rangle = (\frac{3}{8})^{1/2}.$$

This modifies all coefficients of  $a^{(3)}$  in Table II by a factor  $3/2\sqrt{2}$ , and all other expressions containing  $a^{(3)}$  accordingly.

The correct expression may be obtained by a direct tensor reduction of the product  $\bar{8} \otimes \bar{3}$ . I am grateful to H. Haut, C. Leroy, and J. van Parijs of the University of Louvain for pointing out the error to me.