

CP violation in the partial width asymmetries for $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ decaysS. Fajfer,^{1,2} T. N. Pham,³ and A. Prapotnik¹¹*J. Stefan Institute, Jamova 39, P. O. Box 3000, 1001 Ljubljana, Slovenia*²*Department of Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia*³*Centre de Physique Theorique, Centre National de la Recherche Scientifique, UMR, 7644, Ecole Polytechnique, 91128 Palaiseau Cedex, France*

(Received 7 May 2004; revised manuscript received 14 June 2004; published 31 August 2004)

We investigate the possibility of observing CP asymmetries in partial widths for the decays $B^- \rightarrow \pi^+ \pi^- K^-$ and $B^- \rightarrow K^+ K^- K^-$ produced by the interference of the nonresonant decay amplitude with resonant amplitudes. The resonant states that subsequently decay into $\pi^+ \pi^-$ and $K^+ K^-$ or $K^- \pi^+$ are charmonium $\bar{c}c$ states with $J^P = 0^+, 1^-, 1^+$ or the $\phi(1020)$ meson. We find that the largest partial width asymmetry comes from the χ_{c0} resonance, while the resonance $\psi(2S)$ gives a partial width asymmetry of the order 10%.

DOI: 10.1103/PhysRevD.70.034033

PACS number(s): 13.25.Hw, 12.15.Ji, 12.39.Hg

I. INTRODUCTION

The experimental data on B mesons decaying into three mesons accumulate [1–4], and a number of important questions on their decay dynamics and their relevance for the precise determination of the CP -violating phase γ should be answered [5–14]. Motivated by Belle and BaBar results on the B meson three-body decays [1–4], we continue with the study of CP -violating partial width asymmetry in the $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ and $B^\pm \rightarrow K^\pm K^+ K^-$ decay amplitudes.

Recently, we have studied a case of the partial width asymmetry resulting from the interference of the nonresonant $B^- \rightarrow M^+ M^- K^-$, $M = \pi, K$, and the resonant $B^- \rightarrow \chi_{c0} K^- \rightarrow M^+ M^- K^-$ decay amplitudes [5]. In both decay modes, the dominant contribution to the nonresonant amplitude comes from penguin operators. However, there is a small tree level contribution in which the weak CP -violating phase γ enters. The strong phase, which is necessary to obtain the CP -violating asymmetry, enters through the dispersive part of both nonresonant and resonant amplitudes.

It was pointed out by the authors of Refs. [9] and [15] that the dispersive part of the nonresonant amplitude exactly cancels the dispersive part of the resonant amplitude coming from the intermediate state, which is identical to the final state. Therefore, the partial width asymmetry for $B^\pm \rightarrow \mathcal{R} K^\pm \rightarrow M^+ M^- K^\pm$, $M = \pi, K$, will be proportional to the decay width of the resonant state \mathcal{R} to all channels excluding the $M^+ M^-$ state. It means that one would expect a large CP asymmetry for the two-meson invariant mass in the χ_{c0} mass region since the decay width of χ_{c0} is rather large and its branching ratio to $M^+ M^-$, $M = \pi, K$ is negligible. The amplitude for the χ_{c0} resonant decay mode was determined using the narrow width approximation [5,7] and the experimental results for the $B^- \rightarrow \chi_{c0} K^-$ and $\chi_{c0} \rightarrow M^+ M^-$ decay rates. The asymmetry was found to be about 20%. In the case of $B^- \rightarrow K^- M^+ M^-$ there are, however, additional important reasons why the partial width asymmetry can be sizable. In fact, if in the $B^- \rightarrow K^- M^+ M^-$ decays the partial widths coming from the nonresonant \mathcal{M}_{nr} and the resonant \mathcal{M}_r amplitude are of the same order of magnitude, as in our analysis

at the χ_{c0} resonance region [5], one obtains a significant CP -violating asymmetry. In the case of negligible nonresonant amplitude relative to the resonant amplitude (or vice versa) one would get a very small partial width asymmetry.

In this paper, we extend this analysis to the case of the CP -violating partial width asymmetry when the interference with the nonresonant amplitude occurs in the neighborhood of the resonance \mathcal{R} which is a charmonium $\bar{c}c$ state with $J^P = 0^+, 1^-, 1^+$ or a light vector and scalar meson. We will restrict our investigation only to those resonant states \mathcal{R} for which the decay $B^- \rightarrow \mathcal{R} M^-$, $M = K, \pi$ amplitude does not have two or more contributions with different weak phase, as from the experimental branching ratio we are able to extract only the absolute value of the amplitude. For example, in the case of $B^- \rightarrow \mathcal{R} K^-$ with $\mathcal{R} = \rho^0$ there is a penguin and a tree amplitude and one needs to know their relative sizes to constrain the partial width asymmetry. In this decay mode it has also been found that the naive factorization fails to describe the decay rate [16,17]. Therefore, we concentrate on the partial width asymmetry for the cases in which the relevant two-body amplitude can be completely extracted from the measured decay rates.

In the case of the $B^- \rightarrow K^- \pi^+ \pi^-$ partial width asymmetry, the intermediate resonant states of interest would be the light strange mesons $K^*(890)$, $K_1(1270)$, $K_1(1400)$, $K_0^*(1430)$, etc., in the decay chain $B^- \rightarrow \mathcal{R} \pi^- \rightarrow K^- \pi^+ \pi^-$ and the charmonium $\bar{c}c$ states in the decay chain $B^- \rightarrow \mathcal{R} K^- \rightarrow K^- \pi^+ \pi^-$. The B^- decays to these strange mesons in the final state occur as a pure penguin transition. Among all such decays only the rates for $\mathcal{R} = K^*(890)$ and $K_0^*(1430)$ were measured [3]. However, the $K^*(890)$ and $K_0^*(1430)$ mesons decay to $K^- \pi^+$ with the branching ratios close to 100%. In the case we consider it means that the partial decay width to the rest of the states is negligible and the corresponding CP -violating asymmetry vanishes. The relevant charmonium $\bar{c}c$ states in the decay chain $B^- \rightarrow \mathcal{R} K^- \rightarrow \pi^+ \pi^- K^-$ are produced by the $b \rightarrow \bar{c}cs$ transition. The resonant $B^- \rightarrow M^+ M^- K^-$ amplitude is obtained from the tree level contribution, which is proportional to the V_{cb}

and V_{cs} Cabibbo-Kobayashi-Maskawa (CKM) matrix elements, followed by the strong decay of the $\bar{c}c$ state into $\pi^+\pi^-$ or K^+K^- via the Okubo-Zweig-Iizuka (OZI) suppressed strong interaction. Apart from the already mentioned χ_{c0} state, this category includes also J/ψ , χ_{c1} , χ_{c2} , $\psi(2S)$, etc. We will consider contributions from all the above-mentioned states, even though the $B^- \rightarrow \chi_{c2}K^-$ and $\chi_{c1} \rightarrow M^-M^+$ branching ratios have not been measured yet. Nevertheless, we expect that the partial width asymmetry in this decay mode can be rather large. Although one would expect that the $b \rightarrow \bar{c}cs$ transition will give larger rates for the two-body decays than in the case of the $b \rightarrow \bar{u}us$ transition, the fact that the strong transition of the charmonium states is OZI suppressed makes the nonresonant and resonant partial width to be of the same size and this leads to a sizable CP -violating asymmetry.

In the case of the $B^- \rightarrow K^-K^+K^-$ decays with the two-meson invariant mass below the charmonium production threshold, the resonant contribution comes from the intermediate $\bar{s}s$ states. We consider only the CP asymmetry at the $\phi(1020)$ resonance and do not consider contributions from the scalar meson resonances due to the lack of knowledge on their structure.

In the analysis of the partial width CP asymmetry, one needs a knowledge of the nonresonant amplitudes. We compute the nonresonant decay amplitudes by using a model that combines the heavy quark effective theory and chiral Lagrangian, previously developed in Refs. [5–8]. This model assumes the naive factorization for the weak vertices. The fact that the factorization works reasonably well in the relevant two-body decay modes encourages us to apply it in the three-body decays we consider here. Even more, the experimental investigation of the nonresonant amplitudes done by the Belle Collaboration [3] indicates that one has to rely on a model when discussing the nonresonant background. In comparison with our previous investigation [5,6], we include now the contributions of $B^*(0^+)$ resonances.

In Sec. II we present the calculation and the results on the nonresonant $B^- \rightarrow K^-M^+M^-$, $M = \pi, K$ decay modes, while in Sec. III we analyze the partial width asymmetries. The summary of our results is given in Sec. IV.

II. NONRESONANT AMPLITUDES

The effective weak Hamiltonian relevant for the $B^\pm \rightarrow K^\pm M^+M^-$ decays and their CP conjugates after Fierz reordering of the quark fields and neglecting the contribution of the color octet operators is [16–21]

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \left(V_{us}^* V_{ub} (a_1 O_1 + a_2 O_2) + V_{cs}^* V_{cb} (a_{1c} O_{1c} + a_{2c} O_{2c}) - V_{ts}^* V_{tb} \sum_{i=3}^{10} a_i O_i \right), \quad (1)$$

The effective Wilson coefficients are denoted by a_i , and the operators O_i read

$$O_1 = (\bar{u}b)_{V-A} (\bar{s}u)_{V-A}, \quad O_2 = (\bar{u}u)_{V-A} (\bar{s}b)_{V-A}, \quad (2)$$

$$O_{1c} = (\bar{c}b)_{V-A} (\bar{s}c)_{V-A}, \quad O_{2c} = (\bar{c}c)_{V-A} (\bar{s}b)_{V-A}, \quad (3)$$

$$O_3 = \sum_{q=u,d,s} (\bar{q}q)_{V-A} (\bar{s}b)_{V-A},$$

$$O_4 = \sum_{q=u,d,s} (\bar{q}b)_{V-A} (\bar{s}q)_{V-A}, \quad (4)$$

$$O_5 = \sum_{q=u,d,s} (\bar{q}q)_{V+A} (\bar{s}b)_{V-A},$$

$$O_6 = -2 \sum_{q=u,d,s} [\bar{q}(1-\gamma_5)b][\bar{s}(1+\gamma_5)q], \quad (5)$$

$$O_7 = \sum_{q=u,d,s} \frac{3}{2} e_q (\bar{q}q)_{V+A} (\bar{s}b)_{V-A},$$

$$O_8 = -2 \sum_{q=u,d,s} \frac{3}{2} e_q [\bar{q}(1-\gamma_5)b][\bar{s}(1+\gamma_5)q], \quad (6)$$

$$O_9 = \sum_{q=u,d,s} \frac{3}{2} e_q (\bar{q}q)_{V-A} (\bar{s}b)_{V-A},$$

$$O_{10} = \sum_{q=u,d,s} \frac{3}{2} e_q (\bar{q}b)_{V-A} (\bar{s}q)_{V-A}, \quad (7)$$

where $(\bar{q}_1 q_2)_{V\pm A}$ stands for $\bar{q}_1 \gamma^\mu (1 \pm \gamma_5) q_2$. Here O_1 and O_2 are the tree level operators, $O_3 - O_6$ are the QCD penguin operators and $O_7 - O_{10}$ are the electromagnetic penguin operators. From Ref. [21] we take $a_1 = 1.05$, $a_2 = 0.07$, $a_4 = -0.043 - 0.016i$, and $a_6 = -0.054 - 0.016i$. The values of the other Wilson coefficients are at least one order of magnitude smaller and therefore we can safely neglect them.

For the CKM matrix elements the Wolfenstein parametrization is used [$V_{ub} = A\lambda^3(\bar{\rho} - i\bar{\eta})$, $V_{us} = \lambda$, $V_{ts} = -A\lambda^2$, $V_{tb} = 1$], with $A = 0.8$, $\lambda = 0.228$, $\bar{\rho} = 0.118 - 0.273$ (the average value 0.222) and $\bar{\eta} = 0.305 - 0.393$ (the average value 0.339) [22]. The matrix elements of the four quark operators acting in O_i for the $B^- \rightarrow K^- \pi^+ \pi^-$ decay can be written using the factorization assumption as

$$\begin{aligned} & \langle \pi^+ \pi^- K^- | (\bar{s}b)(\bar{q}q) | B^- \rangle \\ & = \langle K^- | (\bar{s}b) | B^- \rangle \langle \pi^+ \pi^- | (\bar{q}q) | 0 \rangle, \end{aligned} \quad (8)$$

$$\begin{aligned} & \langle \pi^+ \pi^- K^- | (\bar{d}b)(\bar{s}d) | B^- \rangle \\ & = \langle \pi^- | (\bar{d}b) | B^- \rangle \langle K^- \pi^+ | (\bar{s}d) | 0 \rangle, \end{aligned} \quad (9)$$




FIG. 1. Feynman diagrams contributing to the nonresonant part of the amplitude.

$$\begin{aligned}
 & \langle \pi^+ \pi^- K^- | (\bar{u}b)(\bar{s}u) | B^- \rangle \\
 &= \langle \pi^+ \pi^- | (\bar{u}b) | B^- \rangle \langle K^- | (\bar{s}u) | 0 \rangle \\
 &+ \langle 0 | (\bar{u}b) | B^- \rangle \langle K^- \pi^+ \pi^- | (\bar{s}u) | 0 \rangle. \quad (10)
 \end{aligned}$$

In the above equations $(\bar{q}_i q_j)$ denotes the vector or axial-vector current or scalar or pseudoscalar density. By analyzing the matrix elements given above, one finds [5] that only the first term in (10) gives an important contribution to the nonresonant decay rate. Terms (8) and (9) contribute to the resonant part of the amplitude (through resonances which decay into $\pi^+ \pi^-$ or $K^- \pi^-$, respectively), while the annihilation term in (10) is found to be negligible as explained in Refs. [5]. In the matrix element of the O_6 operator, additional terms might arise, but they are either small or cancel among themselves [5].

The $B^- \rightarrow K^- K^+ K^-$ amplitude can be factorized in the same way by replacing π^\pm with K^\pm in (8)–(10). However, in this case, the contribution coming from $B^- \rightarrow \rho^0 K^- \rightarrow K^- K^+ K^-$ [Eq. (8)] is part of the nonresonant amplitude, since the ρ^0 mass is below the $K^- K^+$ threshold. Nevertheless, we find this contribution to be small due to the suppression of the ρ^0 propagator in the high-energy regions and due to the smallness of its Wilson coefficients (a_2 and a_9) and will therefore neglect it. The same argument holds if the ρ meson is replaced by similar resonances (σ , etc.).

Next, we proceed with the determination of $\mathcal{A}_\pi = \langle \pi^+(p_2) \pi^-(p_1) K^-(p_3) | O_1 | \bar{B}^- \rangle$ and $\mathcal{A}_K = \langle K^+(p_2) K^-(p_1) K^-(p_3) | O_1 | \bar{B}^- \rangle$. The approach used in the calculation of these matrix elements was already explained in Refs. [5,23–25]. Here, we follow the same method, but add the contributions of the B_0^* scalar meson

resonances. We introduce the $b\bar{q}$ states ($q = u, d, s$), with the $J^P = 1^+, 0^+$ assignment incorporated in the field S [26]:

$$S = \frac{1}{2} (1 + v_\alpha \gamma^\alpha) [D_1^\mu \gamma_\mu \gamma_5 - D_0], \quad (11)$$

which then interacts with the $b\bar{q}$ $J^P = 1^-, 0^-$ multiplet (H) and the pseudo-Goldstone mesons by the means of the Lagrangian:

$$\mathcal{L}_s = ih \text{Tr}(S_b \gamma_\mu \gamma_5 \mathcal{A}_{ba}^\mu \bar{H}_a), \quad (12)$$

where $\mathcal{A}^\mu = \frac{1}{2} (\xi^\dagger \partial^\mu \xi - \xi \partial^\mu \xi^\dagger)$ with the light pseudoscalar fields in ξ . The weak current is given by

$$j_\mu^S = i \frac{F^+}{2} \text{Tr}[\gamma^\mu (1 - \gamma_5) S_b \xi_{ba}^\dagger]. \quad (13)$$

The parameter $h = -0.6 \pm 0.2$ is taken from the recent study of the $D_s(0^+)$ state [27], while for the scalar meson decay constant we use $F^+ = 0.46 \text{ GeV}^{3/2}$ [26].

The matrix element \mathcal{A}_π can be written as

$$\begin{aligned}
 \mathcal{A}_\pi = & -f_3 \left[m_3^2 r^{\text{nr}} + \frac{1}{2} (m_B^2 - m_3^2 - s) w_+^{\text{nr}} + \frac{1}{2} (s + 2t - m_B^2 \right. \\
 & \left. - 2m_2^2 - m_3^2) w_-^{\text{nr}} \right], \quad (14)
 \end{aligned}$$

where the form factors w_+^{nr} , w_-^{nr} , and r^{nr} are determined by calculating contributions coming from the Feynman diagrams in Fig. 1:

$$w_+^{\text{nr}} = -\frac{g}{f_1 f_2} \frac{f_B m_B^{3/2} m_B^{1/2}}{t - m_{B^*}^2} \left(1 - \frac{m_B^2 - m_1^2 - t}{2m_{B^*}^2} \right) + \frac{f_B}{2f_1 f_2} - \frac{\sqrt{m_B} \alpha_2}{2f_1 f_2 m_B^2} (2t + s - m_B^2 - m_3^2 - 2m_1^2) + \frac{F^+ h \sqrt{m_B}}{2f_1 m_B^2} \frac{m_B^2 - t}{t - m_{B_0^*}^2}, \quad (15)$$

$$w_-^{\text{nr}} = \frac{g}{f_1 f_2} \frac{f_B m_B^{3/2} m_B^{1/2}}{t - m_{B^*}^2} \left(1 + \frac{m_B^2 - m_1^2 - t}{2m_{B^*}^2} \right) + \frac{\sqrt{m_B} \alpha_1}{f_1 f_2} + \frac{F^+ h \sqrt{m_B}}{2f_1 m_B^2} \frac{m_B^2 - t}{t - m_{B_0^*}^2}, \quad (16)$$




FIG. 2. Dalitz plots for the nonresonant $B^- \rightarrow K^- \pi^- \pi^+$ (left) and $B^- \rightarrow K^- K^- K^+$ (right) decay modes.

$$\begin{aligned}
 r^{\text{nr}} = & -\frac{f_B}{2f_1 f_2 (m_3^2 - m_B^2)} (2t + s - m_B^2 - m_3^2 - 2m_1^2) + \frac{f_B}{2f_1 f_2} + \frac{g f_B}{f_1 f_2 (t - m_{B^*}^2)} (m_B^2 - m_1^2 - t) - \frac{\sqrt{m_B} \alpha_2}{2f_1 f_2 m_B^2} (2t + s - m_B^2 - m_3^2 - 2m_1^2) \\
 & - \frac{4g^2 f_B m_B m_{B^*}}{f_1 f_2 (m_3^2 - m_B^2) (t - m_{B^*}^2)} \left(\frac{s - m_1^2 - m_2^2}{2} - \frac{(t + m_2^2 - m_3^2)(m_B^2 - m_1^2 - t)}{4m_{B^*}^2} \right) + \frac{F^+ h \sqrt{m_B}}{f_1 m_B^2} \frac{m_B^2 - t}{t - m_{B_0^*}^2} \\
 & + \frac{F^+ h^2 \sqrt{m_B}}{f_1 f_2 m_B^3} \frac{(m_B^2 - t)(t - m_3^2)}{(t - m_{B_0^*}^2)(m_3^2 - m_{B_0^*}^2)}. \tag{17}
 \end{aligned}$$

We used Mandelstam's variables $s = (p_B - p_3)^2$ and $t = (p_B - p_1)^2$. Indices 1, 2, and 3 correspond to π^- , π^+ , and K^- , respectively ($f_1 = f_2 = f_\pi$, $f_3 = f_K$, $m_1 = m_2 = m_\pi$, $m_3 = m_K$). The masses m_B , m_{B^*} , and $m_{B_0^*}$ correspond to the B^- , $B^{0*}(1^-)$, and $B^{0*}(0^+)$ mesons, (1^-) denoting vector and (0^+) scalar states. The rest of parameters are taken to be $f_\pi = 0.132$ GeV, $f_K = 0.16$ GeV, $f_B = 0.175$ GeV, $f_{B_s} = 1.16f_B$, $\alpha_1 = 0.16$ GeV $^{1/2}$, $\alpha_2 = 0.15$ GeV $^{1/2}$ as in Ref. [5]. For the strong coupling g we use $g = 0.56$ according to the measurement of Ref. [28]. Note that in Ref. [5] there are misprints in Eq. (16): the sign in front of α_2 is reversed, as well as the overall sign in (22).

The matrix element of O_4 has the same structure as the matrix element of O_1 , while for determining the matrix element of O_6 we follow the approach described in Ref. [5].

Using the expressions (18)–(20) of Ref. [5], we find that its contribution is proportional to the matrix element of O_1 or O_4 with the proportionality factor $k_6 = -2\mathcal{B}f_\pi^2/m_b f_K^2$.

The matrix element $\langle K^+(p_2)K^-(p_1)K^-(p_3)|O_1|\bar{B}^- \rangle$ is calculated in the same way. The expression for \mathcal{A}_K and its form factors can be derived from the Eqs. (14)–(17), adding the additional contribution obtained by interchanging s and t and by taking $f_1 = f_2 = f_3 = f_K$, $m_1 = m_2 = m_3 = m_K$. In the propagators the B meson masses are replaced by the B_s mass.

Now, the nonresonant amplitudes for $B^- \rightarrow M^+ M^- K^-$ can be written as

$$\mathcal{M}_{\text{nr}} = \frac{G_F}{\sqrt{2}} \mathcal{A}_{K,\pi} [V_{ub} V_{us}^* a_1 - V_{tb} V_{ts}^* (a_4 + k_6 a_6)], \tag{18}$$




FIG. 3. Diagrams presenting the nonresonant (left) and the resonant (right) contributions to dispersive part of the amplitude in the phase space region of the P_2 and P_3 invariant mass close to the \mathcal{R} mass. Blob in the left diagram presents the nonresonant weak decay mode (see Fig. 1).

TABLE I. The decay $B^- \rightarrow \mathcal{R}K^-$ width and the branching ratios for $\mathcal{R} \rightarrow M^+ M^-$.

R	$B^- \rightarrow \mathcal{R}K^-$	$\mathcal{R} \rightarrow \pi^+ \pi^-$	$\mathcal{R} \rightarrow K^+ K^-$
ϕ	$(7.9 \pm 2.0) \times 10^{-6}$		$(49.2 \pm 0.7)\%$
J/ψ	$(1.01 \pm 0.05) \times 10^{-3}$	$(1.47 \pm 0.23) \times 10^{-4}$	$(2.37 \pm 0.31) \times 10^{-4}$
χ_{c0}	$(6.5 \pm 1.1) \times 10^{-4}$	$(5.0 \pm 0.7) \times 10^{-3}$	$(5.9 \pm 0.9) \times 10^{-3}$
χ_{c1}	$(6.0 \pm 2.4) \times 10^{-4}$	$< 2.1 \times 10^{-3}$	$< 2.1 \times 10^{-3}$
$\psi(2S)$	$(6.6 \pm 0.6) \times 10^{-4}$	$(8 \pm 5) \times 10^{-5}$	$(1.0 \pm 0.7) \times 10^{-4}$

with $\mathcal{A}_{\pi,K}$ defined in Eqs. (14)–(17), while for $B^+ \rightarrow M^+ M^- K^+$ we have

$$\bar{\mathcal{M}}_{\text{nr}} = \frac{G_F}{\sqrt{2}} \mathcal{A}_{K,\pi} [V_{ub}^* V_{us} a_1 - V_{tb}^* V_{ts} (a_4 + k_6 a_6)]. \quad (19)$$

Using the above expressions, we obtain the following branching ratios:

$$\begin{aligned} \text{BR}(B^\pm \rightarrow K^\pm \pi^+ \pi^-)_{\text{nr}} &= 14 \times 10^{-6}, \\ \text{BR}(B^\pm \rightarrow K^\pm K^+ K^-)_{\text{nr}} &= 9.0 \times 10^{-6}, \end{aligned} \quad (20)$$

where $\text{BR}(B^\pm \rightarrow K^\pm M^+ M^-)_{\text{nr}}$ stands for the CP -averaged rates for $B^- \rightarrow M^- M^+ K^-$ and $B^+ \rightarrow M^- M^+ K^+$ ($[\text{BR}(B^- \rightarrow K^- M^+ M^-) + \text{BR}(B^+ \rightarrow K^+ M^+ M^-)]/2$). In Ref. [5] it was found that due to the imaginary part of the a_4 and a_6 Wilson coefficients, we can have a large CP asymmetry between the nonresonant $B^+ \rightarrow M^- M^+ K^+$ and $B^- \rightarrow M^- M^+ K^-$ amplitudes. The size of this asymmetry depends on the $\bar{\rho}$ and $\bar{\eta}$ CKM parameters and is rather large (60% in the case of $B^\pm \rightarrow \pi^- \pi^+ K^\pm$ and 40% in a case of $B^\pm \rightarrow K^- K^+ K^\pm$ decay mode). The largest error in $\text{BR}(B^\pm \rightarrow K^\pm M^+ M^-)_{\text{nr}}$, due to the model parameters, comes from the uncertainty in the CKM weak phase γ , the decay constants, and the coupling g . For example, by taking two times smaller g , the rate $\text{BR}(B^\pm \rightarrow K^\pm \pi^+ \pi^-)_{\text{nr}}$ decreases by 40% and $\text{BR}(B^\pm \rightarrow K^\pm \pi^+ \pi^-)_{\text{nr}}$ by 30%. Varying $\bar{\rho}$ between 0.118 and 0.273 and $\bar{\eta}$ between 0.305 and 0.393 gives $\text{BR}(B^\pm \rightarrow K^\pm \pi^+ \pi^-)_{\text{nr}} = (6.2 - 12.6) \times 10^{-6}$ and $\text{BR}(B^\pm \rightarrow K^\pm K^+ K^-)_{\text{nr}} = (11 - 17) \times 10^{-6}$. The uncertainty in the branching ratios coming from the B decay constants is not larger than 10%.

The Dalitz plots for $B^- \rightarrow K^- M^+ M^-$ ($M = \pi, K$) decays, are given in Fig. 2 ($g = 0.56$). We can see that the nonresonant $B^- \rightarrow K^- K^+ K^-$ decay amplitude is rather flat, while in the case of $B^- \rightarrow \pi^- \pi^+ K^-$, an increase at low K and π momenta phase space region is evident. The inclusion of the scalar states $B^*(0^+)$ does not give a significant contribution to the decay rate, increasing it by a few percent in both decay modes.

Recently B factories [1,3,4] obtained some insight into the nonresonant contribution to the $B^- \rightarrow K^- M^+ M^-$ decay widths. The preliminary results of the Belle Collaboration are [1,3] $\text{BR}(B^\pm \rightarrow K^\pm \pi^+ \pi^-)_{\text{nr,exp}} = (14 \pm 6) \times 10^{-6}$ and $\text{BR}(B^\pm \rightarrow K^\pm K^+ K^-)_{\text{nr,exp}} = (22.5 \pm 4.9) \times 10^{-6}$, while the

BaBar Collaboration still has only the upper limit¹ [1,4]. The inclusion of the nonresonant contribution in the $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ Dalitz plot analysis [3] was motivated by the obvious deficit of the data in the low $K^+ \pi^-$ invariant mass phase space region (see Fig. 11, first row of Ref. [3]). They used a rather simple fit [see Eq. (11) [3]] for the nonresonant amplitude. Nevertheless, as pointed out by Fry [1], this contribution is not yet well understood and more studies of this problem are expected. Calculated ranges for the branching ratios within our model $\text{BR}(B^\pm \rightarrow K^\pm \pi^+ \pi^-)_{\text{nr}} = (11 - 17) \times 10^{-6}$ and $\text{BR}(B^\pm \rightarrow K^\pm K^+ K^-)_{\text{nr}} = (6.2 - 12.6) \times 10^{-6}$ agree with the Belle Collaboration results within two standard deviations. Unfortunately, the experimental statistics is still too low to compare the distributions of the differential decay rate of the model and the experiment. It is interesting that our model predicts a rather small differential decay width distribution in the region of the low $\pi^+ K^-$ invariant mass. In order to describe data given in Fig. 11 of Ref. [3] it seems that one needs such behavior of the nonresonant amplitude.

In addition the results of Ref. [3] indicate the existence of the broad structures in the experimental data at $\sqrt{s} \approx 1.3$ GeV in the $K^+ \pi^+ \pi^-$ final state and at $\sqrt{s} \approx 1.5$ GeV in the $K^+ K^+ K^-$ final state. Although one explanation is that light scalar resonances might be responsible for this effect [3], we suggest that these increases might be induced by the nonresonant effects also, as can be seen in the presented Dalitz plots (Fig. 2).

III. PARTIAL WIDTH ASYMMETRY

For the resonances in the s channel, the partial decay width Γ_p for $B^- \rightarrow M \bar{M} K^-$, $M = \pi^+, K^+$, which contains both the nonresonant and resonant contributions, is obtained by integrating the amplitude from $s_{\min} = (m_R - 2\Gamma_R)^2$ to $s_{\max} = (m_R + 2\Gamma_R)^2$:

$$\Gamma_p = \frac{1}{(2\pi)^3} \frac{1}{32m_B^3} \int_{s_{\min}}^{s_{\max}} ds \int_{t_{\min}(s)}^{t_{\max}(s)} dt |\mathcal{M}_{\text{nr}} + \mathcal{M}_r|^2. \quad (21)$$

Similarly, the partial decay width $\Gamma_{\bar{p}}$ for $B^+ \rightarrow M \bar{M} K^+$, $M = \pi^+, K^+$ is defined in the same way. The CP -violating asymmetry is then

¹After completing this paper, we learned that the BaBar Collaboration published their new results [29].

$$A_p = \frac{|\Gamma_p - \Gamma_p^-|}{|\Gamma_p + \Gamma_p^-|}. \quad (22)$$

It is important to notice that at the phase space region where the invariant mass of M^+M^- approaches the mass of the \mathcal{R} resonant state, M^+M^- can rescatter through that resonance as it is visualized in Fig. 3 (left figure). If $\text{BR}(\mathcal{R} \rightarrow M^+M^-)$ is large, this can lead to a significant absorptive amplitude, and it contributes to the partial decay width asymmetry. As mentioned in the Introduction and explained in the Appendix, such a contribution is exactly canceled by the absorptive part of a resonant decay, where the resonance rescatters through the intermediate states equal to final states [Fig. 3 (right figure)]. This implies that one has to include the

TABLE II. The parameters used in our numerical calculations.

V	K_V	$g_{V\pi\pi}$	$g_{VKK}(g_{VK\pi})$
ϕ	2.26×10^{-9}		6.34
J/ψ	1.41×10^{-7}	2.76×10^{-4}	4.37×10^{-3}
$\psi(2S)$	2.04×10^{-7}	1.41×10^{-3}	.166
χ_{c1}	1.68×10^{-7}	< 0.0126	< 0.0134

factor $[1 - \text{BR}(\mathcal{R} \rightarrow M^+M^-)]$ in the equation for the partial decay asymmetry.

In the calculation of the $\Gamma_p - \Gamma_p^-$, by taking $V_{ub} = |V_{ub}|e^{i\gamma} \simeq A\lambda^3(\bar{\rho} - i\bar{\eta})$, we derive

$$\begin{aligned} \Gamma_p - \Gamma_p^- = \sin \gamma \frac{4m_R \Gamma_R [1 - \text{BR}(\mathcal{R} \rightarrow M^+M^-)]}{(2\pi)^3 32m_B^3} \int_{s_{\min}}^{s_{\max}} ds \int_{t_{\min}(s)}^{t_{\max}(s)} dt \frac{G_F}{\sqrt{2}} |V_{ub}| |V_{us}^*| a_1 \langle K\pi\pi | O_1 | B \rangle_{\text{nr}} \\ \times |\mathcal{M}(B^- \rightarrow \mathcal{R}K^-)| \frac{1}{(s - m_R^2)^2 + (m_R \Gamma_R)^2} |\mathcal{M}(\mathcal{R} \rightarrow \pi^- \pi^+)|, \end{aligned} \quad (23)$$

while the $\Gamma_p + \Gamma_p^-$ is given by

$$\begin{aligned} \Gamma_p + \Gamma_p^- = 2 \frac{1}{(2\pi)^3 32m_B^3} \int_{s_{\min}}^{s_{\max}} ds \int_{t_{\min}(s)}^{t_{\max}(s)} dt \left\{ |\mathcal{M}_{\text{nr}}|^2 \right. \\ \left. + \left| \mathcal{M}(B^- \rightarrow \mathcal{R}K^-) \frac{1}{s - m_R^2 + im_R \Gamma_R} \right. \right. \\ \left. \left. \times \mathcal{M}(\mathcal{R} \rightarrow \pi^- \pi^+) \right|^2 \right\}. \end{aligned} \quad (24)$$

The $B^- \rightarrow \mathcal{R}K^- \rightarrow K^- M^+ M^-$ amplitudes are obtained from the experimental data [29] and the measured branching ratios for $B^- \rightarrow \mathcal{R}K^-$ and $R \rightarrow M^+ M^-$ are given in Table I.

For the scalar resonance exchange (χ_0 in our case) in the $B^- \rightarrow SK^- \rightarrow M^+ M^- K^-$ decay, we have

$$\begin{aligned} \mathcal{M}(B^- \rightarrow SP_1(q_1) \rightarrow P_1(q_1)P_2(q_2)P_3(q_3)) \\ = \mathcal{M}(B^- \rightarrow SP_1(q_1)) \frac{1}{m_{23}^2 - m_S^2 + i\Gamma_S m_S} \\ \times \mathcal{M}(S \rightarrow P_2(q_2)P_3(q_3)), \end{aligned} \quad (25)$$

where $m_{23}^2 = (q_2 + q_3)^2$, while m_S and Γ_S are the mass and the decay width of the scalar resonance, respectively. We find $\mathcal{M}(B^- \rightarrow \chi_{c0} K^-) = 3.34 \times 10^{-7}$ GeV, $\mathcal{M}(\chi_{c0} \rightarrow \pi^+ \pi^-) = 0.118$ GeV, and $\mathcal{M}(\chi_{c0} \rightarrow K^+ K^-) = 0.132$ GeV.

The amplitude for the B^- decay into light vector and pseudoscalar resonance and the amplitude for the vector meson decay into two pseudoscalar states are given by

$$\begin{aligned} \mathcal{M}(B^- \rightarrow V(\varepsilon)P_1(q_1)) = K_V q_1 \cdot \varepsilon^*, \\ \mathcal{M}(V \rightarrow P_2(q_2)P_3(q_3)) = \frac{g_{VP_2 P_3}}{\sqrt{2}} (q_2 - q_3) \cdot \varepsilon. \end{aligned} \quad (26)$$

TABLE III. The partial width asymmetry for $B^- \rightarrow K^- \pi^+ \pi^-$, calculated with $g=0.56$ and given $\bar{\rho}$ and $\bar{\eta}$. $A_p(\chi_{c1})$ is obtained by taking the upper bound for g_{VMM} .

	$\bar{\rho}=0.118$ $\bar{\eta}=0.305$	$\bar{\rho}=0.118$ $\bar{\eta}=0.393$	$\bar{\rho}=0.273$ $\bar{\eta}=0.305$	$\bar{\rho}=0.273$ $\bar{\eta}=0.393$	$\bar{\rho}=0.222$ $\bar{\eta}=0.339$
$A_p(\psi(2S))$	10.2%	13.0%	10.3%	13.1%	11.3%
$A_p(J/\psi)$	0.8%	1.1%	0.8%	1.1%	0.9%
$A_p(\chi_{c1})$	3.5%	4.5%	3.5%	4.5%	3.9%
$A_p(\chi_{c0})$	17.3%	21.8%	17.6%	22.1%	19.3%

TABLE IV. The partial width asymmetry for $B^- \rightarrow K^- \pi^+ \pi^-$, calculated with $g=0.27$ and given $\bar{\rho}$ and $\bar{\eta}$. $A_p(\chi_{c1})$ is obtained by taking the upper bound for g_{VMM} .

	$\bar{\rho}=0.118$ $\bar{\eta}=0.305$	$\bar{\rho}=0.118$ $\bar{\eta}=0.393$	$\bar{\rho}=0.273$ $\bar{\eta}=0.305$	$\bar{\rho}=0.273$ $\bar{\eta}=0.393$	$\bar{\rho}=0.222$ $\bar{\eta}=0.339$
$A_p(\psi(2S))$	13.5%	17.3%	13.7%	17.3%	15.1%
$A_p(J/\psi)$	1.2%	1.6%	1.2%	1.6%	1.4%
$A_p(\chi_{c1})$	5.0%	6.4%	5.0%	6.5%	5.6%
$A_p(\chi_{c0})$	12.8%	16.1%	12.9%	16.3%	14.2%

The amplitude for the three-body resonant decay for this case is

$$\begin{aligned} \mathcal{M}(B^- \rightarrow V(\varepsilon)P_1(q_1) \rightarrow P_1(q_1)P_2(q_2)P_3(q_3)) \\ = \frac{K_V g_{VP_2P_3}}{\sqrt{2}} \\ \times \frac{-q_1 \cdot (q_2 - q_3) + [q_1 \cdot (q_2 + q_3)(m_2^2 - m_3^2)]/m_V^2}{m_{23}^2 - m_V^2 + i\Gamma_V m_V}, \end{aligned} \quad (27)$$

where $m_{23}^2 = (q_2 + q_3)^2$, while m_1 , m_2 , m_3 , and m_V are the masses of particles P_1 , P_2 , P_3 , and V , respectively, and Γ_V is the width of the vector resonance. Using the above formulas, we find the expression for the resonance exchange in the s channel:

$$\begin{aligned} \mathcal{M}(B^- \rightarrow VK^- \rightarrow K^- M^+ M^-) \\ = \frac{K_V g_{VMM}}{2\sqrt{2}} \frac{m_B^2 + 2m_M^2 + m_K^2 - 2t - s}{s - m_V^2 + i\Gamma_V m_V}, \end{aligned} \quad (28)$$

where M stands for K or π . In the case of the $K^- K^+ K^\pm$ mode the contributions coming from the s and t channels are completely symmetric. Values of K_V and $g_{VP_1P_2}$ are given in Table II.

The results for the asymmetries are presented in Tables III–VI. Tables III and V contain the asymmetries for $g=0.56$. The off-shell mass effects might reduce this coupling as mentioned in Ref. [5], and therefore we present the partial width asymmetries for $g=0.27$ (Tables IV and VI). We calculate asymmetries for the ranges $\bar{\rho}=0.118-0.273$ (the av-

erage value 0.222) and $\bar{\eta}=0.305-0.393$ (the average value 0.339) as in Ref. [22]. The subtraction of $\text{BR}(\mathcal{R} \rightarrow M^+ M^-)$ in Eq. (23) makes a sizable effect in the case of the $B^- \rightarrow K^- \phi \rightarrow M^+ M^- K^-$ asymmetry, but it is negligible in the case of partial width asymmetry in the neighborhood of charmonium resonances. Then we can draw the following conclusions: In the case of $B^- \rightarrow K^- \pi^+ \pi^-$, all partial width asymmetries are not very large. The largest asymmetry was found in the case of the χ_{c0} resonance and then in the case of the $\psi(2S)$. The partial width asymmetry $A_p(\chi_{c1})$ is calculated by taking the upper bounds for $\chi_{c1} M^+ M^-$ coupling. All these asymmetries are rather stable on the variations of g . In the case of $B^- \rightarrow K^- K^+ K^-$ the situation is different. Calculated partial width asymmetries except the $A_p(\chi_{c0})$ are smaller than in the case of $B^- \rightarrow K^- \pi^+ \pi^-$. They depend more on the variations of the g coupling. The only relatively sizable partial width asymmetry in addition to $A_p(\chi_{c0})$ is $A_p(\psi(2S))$. We have also estimated the partial width asymmetry for the $B^- \rightarrow \chi_{c2} K^-$ channel, by assuming the $B_{\chi_{c2}K}$ coupling to be of the same size as for the vector (scalar) mesons and we found it negligible.

IV. SUMMARY

In this paper we have investigated the partial width asymmetry for the $B^- \rightarrow M^+ M^- K^-$, $M = \pi, K$ decays which results from the interference of nonresonant and resonant amplitudes. First, we have calculated the nonresonant branching ratios and found that the model we use gives the decay rates in the reasonable agreement with the Belle Collaboration results [3]. Comparing the Dalitz plots for the nonresonant decay modes obtained from our model with the experimental data [3], we find that our model reproduces the data quite well. The inclusion of the B_0^* scalar meson is rather insig-

TABLE V. The partial width asymmetry for $B^- \rightarrow K^- K^+ K^-$, calculated with $g=0.56$ and given $\bar{\rho}$ and $\bar{\eta}$. $A_p(\chi_{c1})$ is obtained by taking the upper bound for g_{VMM} .

	$\bar{\rho}=0.118$ $\bar{\eta}=0.305$	$\bar{\rho}=0.118$ $\bar{\eta}=0.393$	$\bar{\rho}=0.273$ $\bar{\eta}=0.305$	$\bar{\rho}=0.273$ $\bar{\eta}=0.393$	$\bar{\rho}=0.222$ $\bar{\eta}=0.339$
$A_p(\phi)$	0.3%	0.3%	0.3%	0.3%	0.3%
$A_p(\psi(2S))$	3.1%	3.8%	3.0%	3.7%	3.3%
$A_p(J/\psi)$	0.03%	0.04%	0.03%	0.04%	0.03%
$A_p(\chi_{c1})$	0.5%	0.7%	0.5%	0.3%	0.6%
$A_p(\chi_{c0})$	28.8%	35%	27.6%	33.8%	30.6%

TABLE VI. The partial width asymmetry for $B^- \rightarrow K^- K^+ K^-$, calculated with $g=0.27$ and given $\bar{\rho}$ and $\bar{\eta}$. $A_p(\chi_{c1})$ is obtained by taking the upper bound for g_{VMM} .

	$\bar{\rho}=0.118$ $\bar{\eta}=0.305$	$\bar{\rho}=0.118$ $\bar{\eta}=0.393$	$\bar{\rho}=0.273$ $\bar{\eta}=0.305$	$\bar{\rho}=0.273$ $\bar{\eta}=0.393$	$\bar{\rho}=0.222$ $\bar{\eta}=0.339$
$A_p(\phi)$	0.3%	0.3%	0.3%	0.3%	0.3%
$A_p(\psi(2S))$	8.1%	10.1%	7.9%	9.8%	8.8%
$A_p(J/\psi)$	0.55%	0.71%	0.55%	0.71%	0.61%
$A_p(\chi_{c1})$	3.0%	3.8%	3.0%	3.8%	3.3%
$A_p(\chi_{c0})$	23.1%	28.7%	22.5%	28.0%	25%

nificant, contributing only by a few percent to the rate.

We then consider the partial width asymmetries for a few resonant decay modes for which the amplitude does not contain the weak phase γ . In the case of $B^- \rightarrow \pi^+ \pi^- K^-$ the largest partial width asymmetry arises from the interference of the nonresonant amplitude with the resonant amplitude coming from the χ_{c0} and $\psi(2S)$ states. In the case of $B^- \rightarrow K^- K^+ K^-$ the largest partial width asymmetry comes from the χ_{c0} scalar resonance and is about 10% in the case of $\psi(2S)$ state.

ACKNOWLEDGMENTS

We thank our colleagues B. Golob and P. Križan for stimulating discussions on experimental aspects of this investigation and D. Bečirević for very useful comments. The research of S.F. and A.P. was supported in part by the Ministry of Education, Science and Sport of the Republic of Slovenia.

APPENDIX

Following the approach of Ref. [15], the total amplitude contributing to the partial decay width Γ_p can be written as a sum of the resonant and nonresonant contributions as defined in Eq. (21) in the following form:

$$\mathcal{M} = \mathcal{M}_{nr} + \mathcal{M}_r = T e^{-i\gamma} + P + R, \quad (\text{A1})$$

where T is the nonresonant tree contribution, P the nonresonant penguin contribution and R the resonant contribution to the amplitude. The partial width asymmetry defined in Eq. (22) is proportional to

$$A_p \propto \sin \gamma \text{Im}[T(P^* + R^*)], \quad (\text{A2})$$

where $\text{Im}(A)$ stands for the imaginary part of A [similarly $\text{Re}(A)$ stands for the real part of A]. If we neglect the small imaginary part of the penguin Wilson coefficients, T and P will have the same strong phase. This implies that the only contribution to the partial decay asymmetry will come from the interference of the tree nonresonant and the resonant amplitude. One can write

$$A_p \propto \text{Im}(T)\text{Re}(R) - \text{Im}(R)\text{Re}(T). \quad (\text{A3})$$

The imaginary part of T is given by the absorptive part of the left diagram on the Fig. 3. Using Cutkosky's rules, its contribution can be written as

$$\text{Im}(T)\text{Re}(R) = \frac{(2\pi)^4}{2} \int \text{Re}(T)\text{Re}(R)\text{Re}(S)d\Phi, \quad (\text{A4})$$

where the integration is taken over the $P_2 P_3$ phase space. Here S denotes the strong rescattering amplitude of $P_2 P_3$ through the resonance \mathcal{R} visualized in Fig. 4. Similarly, the imaginary part of R is given by the absorptive part of the right diagram on the Fig. 3, where now the sum of all possible intermediate states into which \mathcal{R} decays should be taken into account. One can separate this contribution into the part with P_2 and P_3 as an intermediate state $[\text{Im}(R)_{P_{2,3}}]$ and the part with all other intermediate states $(\text{Im}(R)')$. Again with the use of Cutkosky's rules, one obtains

$$\text{Im}(R_{P_{2,3}})\text{Re}(T) = \frac{(2\pi)^4}{2} \int \text{Re}(T)\text{Re}(R)\text{Re}(S)d\Phi. \quad (\text{A5})$$

The right-hand sides of (A4) and (A5) are equal and therefore these two contributions to (A3) cancel among themselves and we have

$$A_p \sim \text{Im}(R)'\text{Re}(T) \sim \Gamma_R [1 - \text{BR}(\mathcal{R} \rightarrow P_2 P_3)]. \quad (\text{A6})$$

This cancellation is obviously a result of the unitarity and it maintains the equality of the total decay widths for the meson and the antimeson as required by CPT theorem [30]. That was already noticed in Refs. [9] and [15], where the more general proof is presented.




FIG. 4. Rescattering of the P_2 and P_3 states through the resonance \mathcal{R} .

- [1] J. R. Fry, talk given at Lepton Photon Conference, Fermilab, 2003.
- [2] Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **88**, 031802 (2002).
- [3] Belle Collaboration, K. Abe *et al.*, Belle-CONF-0338.
- [4] B. Aubert *et al.*, Phys. Rev. Lett. **91**, 051801 (2003).
- [5] S. Fajfer, R.J. Oakes, and T.N. Pham, Phys. Lett. B **539**, 67 (2002).
- [6] B. Bajc, S. Fajfer, R.J. Oakes, T.N. Pham, and S. Prelovšek, Phys. Lett. B **447**, 313 (1999).
- [7] N.G. Deshpande, G. Eilam, X.G. He, and J. Trampetić, Phys. Rev. D **52**, 5354 (1995).
- [8] H.Y. Cheng and K.C. Yang, Phys. Rev. D **66**, 054015 (2002).
- [9] G. Eilam, M. Gronau, and R.R. Mendel, Phys. Rev. Lett. **74**, 4984 (1995).
- [10] S. Fajfer, R.J. Oakes, and T.N. Pham, Phys. Rev. D **60**, 054029 (1999).
- [11] I. Bediaga, R.E. Blanco, C. Göbel, and R. Mendez-Galain, Phys. Rev. Lett. **81**, 4067 (1998).
- [12] R.E. Blanco, C. Göbel, and R. Mendez-Galain, Phys. Rev. Lett. **86**, 2720 (2001).
- [13] R. Enomoto, Y. Okada, and Y. Shimizu, Phys. Rev. B **433**, 109 (1998).
- [14] M. Hazumi, Phys. Lett. B **583**, 285 (2004).
- [15] H. Simma, G. Eliam, and D. Wyler, Nucl. Phys. **B352**, 367 (1991).
- [16] A. Ali, G. Kramer, and C.D. Lü, Phys. Rev. D **59**, 014005 (1999).
- [17] A. Ali, G. Kramer, and C.D. Lü, Phys. Rev. D **58**, 094009 (1998).
- [18] A. Ali and C. Greub, Phys. Rev. D **57**, 2996 (1998).
- [19] N.G. Deshpande and X.G. He, Phys. Lett. B **336**, 471 (1994); N.G. Deshpande, X.G. He, W.S. Hou, and S. Pakvasa, Phys. Rev. Lett. **82**, 2240 (1999).
- [20] C. Isola and T.N. Pham, Phys. Rev. D **62**, 094002 (2000).
- [21] T.E. Browder, A. Datta, X.G. He, and S. Pakvasa, Phys. Rev. D **57**, 6829 (1998).
- [22] Data taken from www.ckmfitter.in2p3.fr; A. Hocker *et al.*, Eur. Phys. J. C **21**, 225 (2001).
- [23] B. Bajc, S. Fajfer, R.J. Oakes, and T.N. Pham, Phys. Rev. D **58**, 054009 (1998).
- [24] B. Bajc, S. Fajfer, R.J. Oakes, and S. Prelovsek, Phys. Rev. D **56**, 7207 (1997).
- [25] S. Fajfer, R.J. Oakes, and T.N. Pham, Phys. Rev. D **60**, 054029 (1999).
- [26] R. Casalbuoni *et al.*, Phys. Rep. **281**, 145 (1997).
- [27] P. Colangelo and F. De Fazio, Phys. Lett. B **570**, 180 (2003).
- [28] CLEO Collaboration, T.E. Coan *et al.*, hep-ex/0102007; paper submitted to the Fourth International Conference on B Physics and CP Violation (BCP4), Ise-Shima, Japan.
- [29] BaBar Collaboration, B. Aubert *et al.*, hep-ex/0408032. Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
- [30] J-M. Gerard and W-s. Hou, Phys. Rev. Lett. **62**, 855 (1989); L. Wolfenstein, Phys. Rev. D **43**, 151 (1991); D. Atwood and A. Soni, *ibid.* **58**, 036005 (1998).