

**Vector fluctuations from multidimensional curvature bounces**

Massimo Giovannini

*Department of Physics, Theory Division, CERN, 1211 Geneva 23, Switzerland*

(Received 14 July 2004; published 11 November 2004)

It is argued that in the case of a (four-dimensional) smooth transition across a (dilaton-driven) curvature bounce growing vector fluctuations evolve continuously into decaying solutions at later times in the post-bounce epoch. Analytical examples of this observation are given. In the case of multi-dimensional bouncing models the situation is different, since the system of differential equations describing the vector modes of the geometry has a richer structure. The amplification of the vector modes of the geometry is specifically investigated in a regular five-dimensional bouncing curvature model where scale factors of the external and internal manifolds evolve at a dual rate. Vector fluctuations, in this case, can be copiously produced and are continuous across the bounce. The relevance of these results is critically illustrated.

DOI: 10.1103/PhysRevD.70.103509

PACS numbers: 98.80.Cq, 04.50.+h, 04.62.+v, 11.25.Mj

**I. FORMULATION OF THE PROBLEM**

It is known that in conventional inflationary models vector fluctuations of the metric are not amplified by the pumping action of the gravitational field [1]. On the contrary, both scalar and tensor modes of the geometry are copiously generated in the transition from a quasi-de Sitter stage of expansion to a radiation-dominated evolution.

Vector modes of the geometry may have interesting phenomenological implications. For instance, as argued long ago by Wassermann [2] (see also [3]) the existence of large-scale vorticity (sometimes connected with a large-scale magnetic field) may have relevant implications for the origin of global rotation of spiral galaxies. The possible existence of primordial vorticity has been the subject of various investigations. Harrison [4] (see also [5] for a complementary perspective) noticed that if vorticity is present over sufficiently large length scales in a plasma of ions and electrons, magnetic fields can be induced over scales comparable with the typical scale of the vorticity. The dynamical origin of primordial vorticity has been, since then, a subject of debate. In [6] it was suggested that cosmic strings with small scale structure may be responsible for the dynamical origin of vorticity. The dynamical friction between cosmic strings and matter may also be responsible for further sources of vorticity [7].

In spite of the fact that the considerations quoted so far may indeed produce large-scale vorticity, it is useful to take another approach and consider the possibility that, thanks to the pumping action of the gravitational field, the vector fluctuations of the geometry will be amplified, providing, at some later epoch, the initial condition for the evolution of the rotational fluctuations of the fluid. The motivation of the present investigation is to discuss the evolution of vector modes of the geometry in the case of a bouncing background. As is known, the ten-dimensional degrees of freedom of a four-dimensional metric can be decomposed, with respect to three-

dimensional rotations, into four scalar modes, two tensor modes (described by a divergenceless and traceless rank-two symmetric tensor in three dimensions), and four vector degrees of freedom (described by two divergenceless vectors in three dimensions). Out of these ten degrees of freedom, four can be fixed by a choice of coordinate system. While the tensor modes are invariant under infinitesimal gauge transformations, the scalar and vector modes are not.

While this work was in progress, Battefeld and Brandenberger [8] argued that it is not unlikely that vector modes of the geometry may be amplified in the case of four-dimensional curvature bounces as the ones provided by cyclic/ekpyrotic scenarios [9] or in the context of the prebig bang scenario [10]. The authors of Ref. [8] worked in a four-dimensional conformally flat Friedmann-Robertson-Walker (FRW) model and discussed the dynamics in the context of an Einsteinian theory of gravity. In the case of a fluid with stiff equation of state, Ref. [8] finds a growing velocity field and the authors correctly point out that the specific fate of such evolution is rather sensitive to the detailed dynamics of the bounce.

There are, in principle, different bouncing dynamics that can be analyzed. The simplest case is the one of a four-dimensional background in the presence of fluid sources together with a dilatonic background. In this case, different examples reported in this paper seem to suggest that prior to the bounce vector fluctuations of the geometry may increase. However, the same specific examples also suggest that the growing solutions turn into decaying modes after the bounce. In the multidimensional case the situation becomes increasingly interesting. There, more scalar and vector modes arise and their quantum-mechanical fluctuations can be amplified.

For the sake of completeness, even if the scalar modes of the geometry are not the subject of the present investigation, we want to mention that curvature bounces have been the subject of various debates concerning the fate of

the scalar fluctuations of the geometry (see [11] for a list of references of this matter).

The plan of the present paper will be the following. In Sec. II the evolution of vector modes of the geometry will be analyzed in the context of four-dimensional curvature bounces. In the string frame description, it will be shown that the vector modes present prior to the curvature bounce match continuously with decreasing vector modes after the bounce. Specific analytical examples will be presented in the presence of fluid sources. In Sec. III the specific case of multidimensional curvature bounces will be discussed. The evolution of the vector fluctuations of a five-dimensional curvature bounce will be derived in fully gauge-invariant terms. In Sec. IV the spectrum of the vector modes will be derived and it will be shown that vector modes can be amplified in the five-dimensional model of curvature bounce. Section V contains the concluding remarks with a critical summary of the main suggestions stemming from the present study. Finally, the appendix contains useful technical results needed for the derivation of the evolution equations of vector modes in more than four dimensions.

## II. FOUR-DIMENSIONAL CURVATURE BOUNCES

In a conformally flat FRW background and assuming the dilaton field is homogeneous

$$ds^2 = G_{\alpha\beta} dx^\alpha dx^\beta = a^2(\eta)[d\eta^2 - d\vec{x}^2], \quad \varphi = \varphi(\eta), \quad (2.1)$$

the vector modes of the geometry are parametrized by two vectors,  $Q_i$  and  $W_i$ , which are divergenceless, i.e.,  $\partial_i Q^i = \partial_i W^i = 0$ <sup>1</sup>:

$$\begin{aligned} \delta G_{0i} &= -a^2(\eta)Q_i, \\ \delta G_{ij} &= 2a^2(\eta)\partial_{(i}W_{j)} \equiv a^2(\eta)(\partial_i W_j + \partial_j W_i). \end{aligned} \quad (2.2)$$

From Eq. (2.2), the components Ricci tensor can be computed to first-order in the fluctuations of the metric:

$$\begin{aligned} \delta R_{0i} &= (\mathcal{H}' + 2\mathcal{H}^2)Q_i - \frac{1}{2}\nabla^2[Q_i + W'_i], \\ \delta R_{ij} &= -\partial_{(i}Q'_{j)} - \partial_{(i}W''_{j)} - 2(\mathcal{H}' + 2\mathcal{H}^2)\partial_{(i}W_{j)} \\ &\quad - 2\mathcal{H}[\partial_{(i}W'_{j)} + \partial_{(i}Q_{j)}], \end{aligned} \quad (2.3)$$

where  $\mathcal{H} = (\ln a)'$  and the prime denotes a derivation with respect to the conformal time coordinate  $\eta$ .

Under infinitesimal diffeomorphisms  $Q_i$  and  $W_i$  transform as

<sup>1</sup>In the present investigation the attention will be confined to spatially flat manifolds. However, this parametrization of the vector modes can be extended to the case of nonflat spatial geometry by replacing, when needed, the ordinary derivatives with covariant derivatives with respect to the spatial part of the geometry.

$$\tilde{W}_i = W_i + \zeta_i, \quad \tilde{Q}_i = Q_i - \zeta'_i, \quad (2.4)$$

where  $\partial_i \zeta^i = 0$  is the gauge function. Hence  $V_i$  is invariant under infinitesimal coordinate transformations. This observation also implies that the condition  $W_i = 0$  fixes the gauge freedom completely and is analogous, in this sense, to the conformally Newtonian gauge often employed in the analysis of scalar perturbations<sup>2</sup>. This gauge choice would have led to the evolution equations for  $Q_i$ , coinciding, in such a gauge, with the evolution of the gauge-invariant vector  $V_i$ . We will go back to the gauge issue during the discussion of vector fluctuations in the multidimensional case.

The aim will now be to show that in a simple model of low-curvature transition, the growing vector mode will match to a decreasing solution in the post-bounce epoch. As was shown in [10], a smooth transition at low curvatures can be achieved if the dilaton potential is a local function of  $\bar{\varphi}$ . The generally covariant action describing this model is given, in  $D$  space-time dimensions by

$$\begin{aligned} S &= -\frac{1}{\lambda_s^{D-2}} \int d^{D-1}x \sqrt{|G|} e^{-\varphi} [R + G^{\alpha\beta} \nabla_\alpha \varphi \nabla_\beta \varphi \\ &\quad + V(\bar{\varphi})] + S_m, \end{aligned} \quad (2.5)$$

where  $S_m$  represents the contribution of effective (fluid) sources and where

$$\begin{aligned} V &= V(e^{-\bar{\varphi}}), \\ e^{-\bar{\varphi}(x)} &= \frac{1}{\lambda_s^{D-1}} \int d^D w \sqrt{|G(w)|} e^{-\varphi(w)} \\ &\quad \times \sqrt{G^{\alpha\beta} \partial_\alpha \varphi(w) \partial_\beta \varphi(w)} \delta[\varphi(x) - \varphi(w)]. \end{aligned} \quad (2.6)$$

The variation of the action (2.5) with respect to  $G_{\mu\nu}$  and  $\varphi$  leads to the following equations:

$$\begin{aligned} \mathcal{G}_{\mu\nu} + \nabla_\mu \nabla_\nu \varphi + \frac{1}{2} G_{\mu\nu} \left[ G^{\alpha\beta} \nabla_\alpha \varphi \nabla_\beta \varphi \right. \\ \left. - 2G^{\alpha\beta} \nabla_\alpha \nabla_\beta \varphi - V \right], \\ - \frac{e^{-\varphi}}{2} \sqrt{G^{\alpha\beta} \partial_\alpha \varphi \partial_\beta \varphi} \gamma_{\mu\nu} I_1 = \lambda_s^{D-2} e^\varphi T_{\mu\nu}, \end{aligned} \quad (2.7)$$

<sup>2</sup>Notice that, in the scalar case, the longitudinal gauge condition (imposed on the gauge parameters preserving the scalar nature of the fluctuation) eliminates the off-diagonal entries of the perturbed metric. In the case of vector modes the situation is, in a sense, opposite since the gauge choice fixing the coordinate system completely keeps only the off-diagonal elements of the perturbed metric tensor.

$$R + 2G^{\alpha\beta}\nabla_\alpha\nabla_\beta\varphi - G^{\alpha\beta}\partial_\alpha\varphi\partial_\beta\varphi + V - \frac{\partial V}{\partial\varphi} + e^{-\varphi}\frac{\hat{\nabla}^2\varphi}{\sqrt{G^{\alpha\beta}\partial_\alpha\varphi\partial_\beta\varphi}}I_1 - e^{-\varphi}V'I_2 = 0, \quad (2.8)$$

where<sup>3</sup>

$$\hat{\nabla}^2 = \gamma^{\mu\nu}\partial_\mu\partial_\nu, \quad G_\mu^\nu = R_\mu^\nu - \frac{1}{2}\delta_\mu^\nu R, \quad (2.9)$$

and

$$\gamma_{\mu\nu} = G_{\mu\nu} - \frac{\partial_\mu\varphi\partial_\nu\varphi}{\sqrt{G^{\alpha\beta}\partial_\alpha\varphi\partial_\beta\varphi}} \quad (2.10)$$

is the induced metric. In Eqs. (2.7) and (2.8) the following integrals

$$I_1 = \frac{1}{\lambda_s^{D-1}} \int d^D w \sqrt{|G(w)|} V'(e^{-\varphi(w)}) \delta[\varphi(x) - \varphi(w)],$$

$$I_2 = \frac{1}{\lambda_s^{D-1}} \int d^D w \sqrt{|G(w)|} \times \sqrt{G^{\alpha\beta}\partial_\alpha\varphi(w)\partial_\beta\varphi(w)} \delta'[\varphi(x) - \varphi(w)], \quad (2.11)$$

have also been defined.

By combining Eqs. (2.7) and (2.8) in order to eliminate the Ricci scalar we obtain the following equation

$$R_{\mu\nu} + \nabla_\mu\nabla_\nu\varphi - \frac{1}{2}G_{\mu\nu}\left[\frac{\partial V}{\partial\varphi} + e^{-\varphi}V'I_2\right] + \frac{1}{2}e^{-\varphi}\left[G_{\mu\nu}\frac{\hat{\nabla}^2\varphi}{\sqrt{G^{\alpha\beta}\partial_\alpha\varphi\partial_\beta\varphi}} - \gamma_{\mu\nu}\sqrt{G^{\alpha\beta}\partial_\alpha\varphi\partial_\beta\varphi}\right]I_1 = \lambda_s^{D-2}e^\varphi T_{\mu\nu}. \quad (2.12)$$

In the case of a homogeneous dilaton and for a conformally flat metric of FRW type Eqs. (2.7) and (2.12) lead, in  $D = 4$  and in units  $2\lambda_s^2 = 1$ , to the following system of equations:

$$\dot{\bar{\varphi}}^2 - 3H^2 - V = e^{\bar{\varphi}}\bar{\rho}, \quad (2.13)$$

$$\dot{H} = \dot{\bar{\varphi}}H + \frac{1}{2}e^{\bar{\varphi}}\bar{p}, \quad (2.14)$$

$$2\ddot{\bar{\varphi}} - \dot{\bar{\varphi}}^2 - dH^2 + V - \frac{\partial V}{\partial\varphi} = 0, \quad (2.15)$$

<sup>3</sup>In Eq. (2.8) as well as in Eqs. (2.11) and (2.12) the prime denotes a derivation with respect to the argument of the given functional and *not* the derivative with respect to the conformal time coordinate. The two notations cannot be confused since the latter is only employed when dealing with the explicit form of the equations on a given background.

$$\dot{\bar{\rho}} + dH\bar{\rho} = 0, \quad (2.16)$$

where the stress tensor of the fluid sources

$$T_\mu^\nu = (p + \rho)u_\mu u^\nu - p\delta_\mu^\nu \quad (2.17)$$

has been assumed and where the overdot denotes a derivation with respect to the cosmic time coordinate. In Eqs. (2.13)–(2.15) the energy and pressure densities have been rescaled as

$$\bar{\rho} = a^3\rho, \quad \bar{p} = a^3p. \quad (2.18)$$

For instance, in the case  $\bar{\rho} = \bar{p} = 0$ , Eqs. (2.13)–(2.16) admit the solution [10]

$$V(\bar{\varphi}) = -V_0 e^{4\bar{\varphi}}, \quad (2.19)$$

$$a(t) = a_0 \left[ \tau + \sqrt{\tau^2 + 1} \right]^{1/\sqrt{3}}, \quad (2.20)$$

$$\bar{\varphi} = -\frac{1}{2} \log(1 + \tau^2) + \varphi_0, \quad (2.21)$$

where

$$\tau = \frac{t}{t_0}, \quad t_0 = \frac{e^{-2\varphi_0}}{\sqrt{V_0}}. \quad (2.22)$$

The scale  $t_0$  fixes the width of the curvature bounce<sup>4</sup>.

Consider the case  $\bar{p} = 0$ . In this case various solutions are possible depending upon the shape of the potential. In this case Eq. (2.14) can be immediately integrated and an exact regular solution of the remaining equations is

$$V = -V_0 e^{\bar{\varphi}} - V_1 e^{4\bar{\varphi}}, \quad (2.23)$$

with

$$H = \frac{1}{\sqrt{3}} \frac{t_0}{\sqrt{t^2 + t_0^2}}, \quad \bar{\varphi} = \varphi_0 - \frac{1}{2} \log \left[ \frac{t^2}{t_0^2} + 1 \right], \quad \rho = \bar{\rho}_0 a^{-3}, \quad (2.24)$$

subject to the conditions:

$$\bar{\rho}_0 = V_0, \quad V_1 e^{2\varphi_0} = \frac{1}{t_0^2}. \quad (2.25)$$

Another solution, which is effective in illustrating some aspects of the present analysis, may be obtained if the scale factor and the energy density of the fluid are both *constant*. In this toy example, Eqs. (2.13)–(2.16) may be

<sup>4</sup>The possibility of expressing the evolution of the background geometry in terms of the rescaled cosmic time  $\tau$  will also prove useful in the numerical discussion of the spectrum of vector fluctuations in multidimensional models.

integrated easily and the result is

$$a = 1, \quad \rho = \rho_0, \quad p = 0, \quad V = -V_0 e^{2\bar{\varphi}},$$

$$e^{\varphi} = \frac{e^{\varphi_0}}{1 + (t/t_0)^2}, \quad e^{\varphi_0} \rho_0 = \frac{4}{t_0^2} = V_0 e^{2\varphi_0}. \quad (2.26)$$

Finally, consider the case of a radiation fluid, i.e.,  $3\bar{p} = \bar{\rho}$  and the potential is  $V = -V_0 e^{2\bar{\varphi}}$ . In this case Eqs. (2.14) and (2.15) can be integrated leading to the expressions

$$H = \frac{1}{3} \frac{e^{\bar{\varphi}}}{L} (x + x_1), \quad \dot{\bar{\varphi}} = -\frac{e^{\bar{\varphi}}}{L} (x + x_0), \quad (2.27)$$

where

$$\frac{dx}{dt} = \frac{L}{2} \bar{\rho}, \quad (2.28)$$

having defined the integration constants  $x_1, x_0$ . By fixing  $x_1 = 3x_0, x_0 = L$  the explicit solution can be written in terms of the variable  $x$  as

$$a(x) = \sqrt{x^2 + L^2} e^{3 \arctan(x/L)}, \quad \frac{d \ln a}{dx} = \frac{x + 3L}{x^2 + L^2},$$

$$\frac{d\bar{\varphi}}{dx} = -\frac{3(x + L)}{x^2 + L^2}, \quad \frac{dx}{dt} = \frac{\rho_0 L}{2} \frac{1}{a(x)}. \quad (2.29)$$

The shifted dilaton is

$$\bar{\varphi} = \varphi_0 - 3 \arctan(x/L) - \frac{3}{2} \ln(x^2 + L^2). \quad (2.30)$$

Looking at the constraint, we should also bear in mind that

$$e^{\bar{\varphi}_0} = \frac{3}{2} \bar{\rho}_0 L^2. \quad (2.31)$$

We will now analyze the evolution of the vector modes of the geometry in the case  $D = 4$  and in the gauge  $W_i = 0$ . The evolution equations of the vector modes can be derived by perturbing to first-order Eq. (2.7) and by using the explicit form of the background equations which we write in the conformal time parametrization

$$\bar{\varphi}'^2 = 3\mathcal{H}^2 + V a^2 + e^{\bar{\varphi}} a^2 \bar{\rho}, \quad (2.32)$$

$$\mathcal{H}' - \mathcal{H}^2 = \mathcal{H} \bar{\varphi}' + \frac{a^2 e^{\bar{\varphi}}}{2} \bar{p}, \quad (2.33)$$

$$\bar{\varphi}'^2 - 2\bar{\varphi}'' + 2\mathcal{H} \bar{\varphi}' + 3\mathcal{H}^2 + \frac{\partial V}{\partial \bar{\varphi}} a^2 - V a^2 = 0, \quad (2.34)$$

$$\bar{\rho}' + 3\mathcal{H} \bar{\rho} = 0. \quad (2.35)$$

Combining the above equations

$$4\mathcal{H}^2 + 2\mathcal{H}' - 2\bar{\varphi}'' + \frac{\partial V}{\partial \bar{\varphi}} a^2 = e^{\bar{\varphi}} a^2 (\bar{p} - \bar{\rho}). \quad (2.36)$$

The combination (2.36) is rather useful in simplifying the evolution equations of the fluctuations which become

$$\nabla^2 Q_i = -a^2 e^{\varphi} (p + \rho) \mathcal{V}_i, \quad (2.37)$$

$$Q_i' = (\bar{\varphi}' + \mathcal{H}) Q_i, \quad (2.38)$$

where  $\mathcal{V}_i$  is the fluctuation of the velocity field and where we used the explicit perturbed form of the energy-momentum tensor

$$\delta T_i^0 = (p + \rho) \mathcal{V}_i, \quad u^0 = \frac{1}{a}, \quad \delta u_i = a \mathcal{V}_i. \quad (2.39)$$

Equation (2.37) follows from the perturbation of the (0i) component of Eq. (2.7) in the gauge  $W_i = 0$  and using the explicit expressions reported in Eqs. (2.2) and (2.3).

Equations (2.19)–(2.21) can be used in order to obtain the explicit solution for the vector modes associated with the velocity field. From Eq. (2.38) we have

$$Q_i = c_i a e^{\bar{\varphi}}, \quad (2.40)$$

where, as before,  $c_i$  is an integration constant carrying vector indices. Equation (2.37) can be written, in Fourier space and bearing in mind the definitions of  $\bar{\rho}$  and  $\bar{p}$ , as

$$\mathcal{V}_i = \frac{k^2}{a^2 (\bar{\rho} + \bar{p})} e^{-\bar{\varphi}} Q_i. \quad (2.41)$$

Combining Eqs. (2.40) and (2.41) the following relation can be obtained

$$\mathcal{V}_i = \frac{c_i k^2}{a(\bar{\rho} + \bar{p})}. \quad (2.42)$$

Consider now, to begin with, the solution reported in Eq. (2.24). In this case, from Eq. (2.40), it can be easily obtained

$$Q_i(t) = c_i t_0 e^{\varphi_0} \frac{(t + \sqrt{t^2 + t_0^2})^{1/\sqrt{3}}}{\sqrt{t^2 + t_0^2}}. \quad (2.43)$$

Notice that for  $t \ll 0$ ,  $Q_i$  grows as

$$Q_i \sim |\eta|^{-1} \sim |t|^{-\frac{\sqrt{3}+1}{\sqrt{3}}}. \quad (2.44)$$

However, for  $t \gg 0$  this possibly dangerous growing mode evolves towards a decreasing solution going as  $Q_i \sim t^{(1-\sqrt{3})/\sqrt{3}}$ . From Eq. (2.42) the velocity field is found to be

$$\mathcal{V}_i(t) = \frac{c_i k^2}{\bar{\rho}_0} (t + \sqrt{t^2 + t_0^2})^{-1/\sqrt{3}}. \quad (2.45)$$

Again, in the string frame metric  $\mathcal{V}_i$  is always decreasing and regular.

In the case of the solution given in Eq. (2.26), Eqs. (2.37) and (2.38) lead to

$$Q_i = c_i \frac{e^{\varphi_0} a_0}{1 + \tau^2}, \quad (2.46)$$

$$\mathcal{V}_i = \frac{c_i k^2}{a_0 \rho_0}, \quad (2.47)$$

where, as previously defined,  $\tau$  denotes the rescaled cosmic time coordinate. As in the previous example (see Eq. (2.43))  $Q_i$  increases before the bounce and decreases later on.

Finally, in the case of the fluid solution illustrated in Eqs. (2.29)–(2.30) Eqs. (2.40) and (2.42) imply

$$Q_i = \frac{e^{\varphi_0} c_i}{x^2 + L^2}, \quad (2.48)$$

$$\mathcal{V}_i = \frac{c_i k^2}{\rho_0}, \quad (2.49)$$

where, as previously introduced,  $x$  is related to the cosmic time coordinate by a monotonic relation specified by the last relation of Eq. (2.29). Notice that  $Q_i$  exhibits, qualitatively, the same behavior already discussed in Eqs. (2.44) and (2.46).

In this paper the description of the vector modes has been conducted in the string frame. However, the connection to the Einstein frame is rather simple in the case of vector modes of the geometry. Since

$$\delta G_{\mu\nu}^E = e^{-\varphi} \delta G_{\mu\nu}^s, \quad \delta T_{\mu\nu}^E = e^{2\varphi} \delta T_{\mu\nu}^s, \quad (2.50)$$

then

$$Q_i^E = Q_i^s = Q_i, \quad W_i^E = W_i^s = W_i, \quad \mathcal{V}_i^E = \mathcal{V}_i^s = \mathcal{V}_i, \quad (2.51)$$

and

$$a_E = e^{-\varphi/2} a_s, \quad \mathcal{H}_E = \mathcal{H}_s - \frac{\varphi'}{2}, \quad \rho_E = e^{2\varphi} \rho_s, \quad p_E = e^{2\varphi} p_s, \quad \varphi_E = \varphi_s = \varphi. \quad (2.52)$$

In Eqs. (2.50)–(2.52) the labels “E” and “s” refer, respectively, to Einstein and string frame quantities.

Recalling that in the transformation from string to Einstein frame the conformal time coordinate does not change (i.e.,  $\eta_s = \eta_E = \eta$ ), the above relations imply

$$Q_i' = -2\mathcal{H}_E Q_i$$

$$k^2 Q_i = \frac{a_E^2}{2} (p_E + \rho_E) \mathcal{V}_i. \quad (2.53)$$

### III. MULTIDIMENSIONAL CURVATURE BOUNCES

The analysis of gauge-invariant fluctuations in multidimensional and/or anisotropic cosmological models has been discussed, through the years, in different contexts. One of the first attempts to adapt the original Bardeen [12] formalism to a multidimensional framework was discussed in [13] with the aim of performing a general discussion of the main features of cosmological fluctuations in Kaluza-Klein models [14]. Of related interest are

some works discussing the theory of cosmological fluctuations in Bianchi models [15,16]. In [17], a perturbative framework for the gauge-invariant discussion of multidimensional pre-Big Bang models was introduced.

The problem of gauge-invariant fluctuations in models with extradimensions received new momentum in the context of brane models, and different approaches (see, for instance [18–20] and references therein) were developed in order to study the possible localization of modes of various spin of both gravitational and nongravitational origin. In [21] vector modes of the geometry were analyzed in the case of a six-dimensional Abelian vortex leading to the localization of gravitational interaction (see also [22] for a more general discussion of codimension-two braneworlds and of their vector excitations).

It is useful to notice, preliminarily, that the action given in Eq. (2.5) also admits classical solutions in the presence of internal (contracting) dimensions. In this section, the vector fluctuations of multidimensional cosmological models will be studied. The line element will be written, in this case, as

$$ds^2 = G_{\mu\nu} dx^\nu dx^\mu = [dt^2 - a^2(t) g_{ij} dx^i dx^j] - b^2(t) g_{AB} dy^A dy^B, \quad (3.1)$$

where  $G_{\mu\nu}$  is the  $D$ -dimensional metric;  $g_{ij}(x)$  and  $g_{AB}(y)$  are the metric tensors of two maximally symmetric Euclidian manifolds. The indices  $\mu, \nu$  run over the full  $D = d + n + 1$ -dimensional space-time; the  $d$ -expanding dimensions are typically three and the  $n$ -contracting dimensions (labeled by capital Latin letters) will be, for a while, taken to be generic but the case  $n = 1$  will be of particular interest in the following part of the investigation.

From Eq. (3.1) in the particular case  $g_{ij} = \delta_{ij}$  and  $g_{AB} = \delta_{AB}$ , Eqs. (2.7) and (2.8) can be written in explicit terms by introducing the expansion rates  $H = \dot{a}/a$  and  $F = \dot{b}/b$  (the dot denotes a derivation with respect to the cosmic time coordinate):

$$\ddot{\varphi}^2 - dH^2 - nF^2 - V = 0, \quad (3.2)$$

$$\dot{H} = H\dot{\varphi}, \quad (3.3)$$

$$\dot{F} = F\dot{\varphi}, \quad (3.4)$$

$$2\ddot{\varphi} - \dot{\varphi}^2 - dH^2 - nF^2 + V - \frac{\partial V}{\partial \varphi} = 0. \quad (3.5)$$

Equations (3.2)–(3.4) come, respectively, from the  $(0, 0)$ ,  $(i, j)$ ,  $(a, b)$  components of Eq. (2.12), while Eq. (3.5) can be derived from Eq. (2.8).

In the case of  $V = -V_0 e^{4\varphi}$ , Eqs. (3.2)–(3.5) are solved by the following ansatz:

$$a(t) = (\tau + \sqrt{\tau^2 + 1})^{\frac{1}{\sqrt{d+n}}}, \quad b(t) = (\tau + \sqrt{\tau^2 + 1})^{-\frac{1}{\sqrt{d+n}}}, \quad (3.6)$$

$$\bar{\varphi} = \varphi_0 - \frac{1}{2} \ln[1 + \tau^2], \quad \tau = \frac{t}{t_0}. \quad (3.7)$$

The solution given in Eqs. (3.6) and (3.7) describes the expansion of the  $d$  dimensions and the contraction (at a dual rate) of the  $n$  internal dimensions.

Recalling the relation between the cosmic time coordinate  $t$  and the conformal time coordinate  $\eta$ , i.e.,  $a(\eta)d\eta = dt$ , Eqs. (3.2)–(3.5) can be written as

$$\bar{\varphi}'^2 - d\mathcal{H}^2 - n\mathcal{F}^2 - Va^2 = 0, \quad (3.8)$$

$$\mathcal{H}' = \mathcal{H}^2 + \mathcal{H}\bar{\varphi}', \quad (3.9)$$

$$\mathcal{F}' = \mathcal{H}\mathcal{F} + \mathcal{F}\bar{\varphi}', \quad (3.10)$$

$$\bar{\varphi}'^2 - 2\bar{\varphi}'' + 2\mathcal{H}\bar{\varphi}' + d\mathcal{H}^2 + n\mathcal{F}^2 + \frac{\partial V}{\partial \bar{\varphi}}a^2 - Va^2 = 0. \quad (3.11)$$

where, in full analogy with the four-dimensional case, the rates of expansion in conformal time have been defined:

$$\mathcal{H} = (\ln a)', \quad \mathcal{F} = (\ln b)', \quad \iota = \frac{\partial}{\partial \eta}. \quad (3.12)$$

A particularly interesting case is the one where only one extradimension is present (i.e.,  $n = 1$ ). In this case, setting  $n = 1$  and  $d = 3$  in Eq. (3.6) the scale factors are

$$a(t) = \frac{1}{b(t)} = \sqrt{\tau + \sqrt{\tau^2 + 1}}. \quad (3.13)$$

From Eq. (3.13), it can be easily deduced that the scale factor  $a(t)$  smoothly interpolates between  $a(t) \sim t^{-1/2}$  for  $t \rightarrow -\infty$  and  $a(t) \sim t^{1/2}$  for  $t \rightarrow +\infty$ . Moreover,  $b(t)$  will contract like  $t^{1/2}$  for  $t \ll -t_0$  and will contract like  $t^{-1/2}$  for  $t \gg t_0$ . Both the scale factors and the curvature invariants are regular in the origin and for  $|t| \rightarrow \infty$ . Finally, a pleasant feature of the case  $d = 3$  and  $n = 1$  is that in the limit  $t \rightarrow +\infty$  the usual radiation-dominated evolution is correctly reproduced. In Fig. 1 the time-evolution of the scale factors  $a(\tau)$  and  $b(\tau)$  and of the associated rates, i.e.,  $H(\tau)$  and  $F(\tau)$ , is illustrated. This model describes the decoupling of internal and external dimensions.

In the following part of the present section, the vector modes arising in the model of Eq. (3.13) will be specifically analyzed. The equations of motion will be derived in a gauge-independent way. The fluctuations of a five-dimensional line element of the type of Eq. (3.1) are, in general, described by 15 degrees of freedom which can be formally decomposed in terms of their transformation properties with respect to three-dimensional rotations as

$$\begin{aligned} \delta G_{00} &= 2a^2\phi, & \delta G_{0i} &= -a^2\partial_i P - a^2Q_i, & \delta G_{0y} &= -abC, \\ \delta G_{ij} &= 2a^2\psi\delta_{ij} - 2a^2\partial_i\partial_j E + a^2h_{ij} \\ &\quad + a^2(\partial_i W_j + \partial_j W_i), & \delta G_{iy} &= -ab\partial_i D \\ &\quad - abH_i, & \delta G_{yy} &= 2b^2\xi, \end{aligned} \quad (3.14)$$




FIG. 1 (color online). The time-evolution of the scale factors  $a(\tau)$  (solid line) and  $b(\tau)$  (dashed line) introduced in Eq. (3.13). The time-evolution of the associated rates of expansion and contraction is also reported.

where

$$\partial_j h_i^j = 0, \quad h_i^i = 0, \quad (3.15)$$

$$\partial_i W^i = 0, \quad \partial_i Q^i = 0, \quad \partial_i H^i = 0. \quad (3.16)$$

According to (3.15)  $h_{ij}$  carries 2 degrees of freedom since it is symmetric in the two indices. Equation (3.16) tells us that  $W_i$ ,  $Q_i$  and  $H_i$  carry overall 6 degrees of freedom. The remaining degrees of freedom are seven scalars, which will not be specifically discussed since the attention of the present analysis is mainly centered on the vector modes.

The vectors  $W_i$ ,  $Q_i$  and  $H_i$  obey a set of differential equations, which we ought to derive now. Notice, preliminarily, that in terms of the divergenceless gauge parameter  $\zeta_i$  the vectors  $W_i$ ,  $Q_i$  and  $H_i$  transform, for infinitesimal diffeomorphisms preserving the vector nature of the fluctuation, as

$$\begin{aligned} \tilde{Q}_i &= Q_i - \zeta'_i, & \tilde{W}_i &= W_i + \zeta_i, \\ \tilde{H}_i &= H_i - \frac{a}{b} \partial_y \zeta_i, \end{aligned} \quad (3.17)$$

As in the four-dimensional case, gauge-invariant fluctuations can be defined, namely

$$V_i = Q_i + W'_i, \quad (3.18)$$

$$Z_i = H_i + \frac{a}{b} \partial_y W_i. \quad (3.19)$$

Notice that (3.18) corresponds to the gauge-invariant fluctuation of the four-dimensional case.

In order to derive the coupled evolution of  $V_i$  and  $Z_i$ , Eq. (2.7) shall be linearized to first-order in  $Q_i$ ,  $W_i$  and  $H_i$ . By inserting the definitions (3.18) and (3.19) into the obtained equations, the system will only contain the relevant gauge-invariant combinations, i.e.,  $V_i$  and  $Z_i$ .

From Eq. (2.7), the general form of the various components of the perturbed equation can be written, in a compact form, as

$$a^2 \delta \mathcal{G}_i^j - \frac{\varphi'}{2} [\partial_i Q^j + \partial^j Q_i] - \frac{\varphi'}{2} [\partial_i W^{j'} + \partial^j W'_i] = 0, \quad (3.20)$$

$$a^2 \delta \mathcal{G}_i^y + \frac{a}{2b} \varphi' [H'_i + (\mathcal{F} - \mathcal{H}) H_i] - \frac{\varphi'}{2} \frac{a^2}{b^2} \partial_y Q_i = 0, \quad (3.21)$$

$$a^2 \delta \mathcal{G}_0^i - Q^i (\varphi'' - 2\mathcal{H} \varphi') + \frac{a^2}{2} \frac{\partial V}{\partial \varphi} Q^i = 0. \quad (3.22)$$

In order to derive Eqs. (3.20)–(3.22) the perturbed connections reported in the appendix (see Eq. (A3)) have been inserted in the linearized form of Eq. (2.7), bearing in mind that the fluctuation of the induced metric (2.10)

implies

$$\delta \gamma_0^i = Q^i. \quad (3.23)$$

From Eqs. (3.20)–(3.22), using Eqs. (A7)–(A9) reported in the appendix, the following expressions are obtained:

$$\begin{aligned} & \frac{1}{2} [(\partial_i Q^j + \partial^j Q_i)' - (\bar{\varphi}' + \mathcal{H})(\partial_i Q^j + \partial^j Q_i)] \\ & - \frac{a}{2b} \partial_y [\partial_i H^j + \partial^j H_i] + \frac{1}{2} [(\partial_i W^j + \partial^j W_i)'' \\ & + (\bar{\varphi}' + \mathcal{H})(\partial_i W^j + \partial^j W_i)'] - \frac{a^2}{2b^2} \partial_y [\partial_i W^j \\ & + \partial^j W_i] = 0, \end{aligned} \quad (3.24)$$

$$\begin{aligned} Q_i \left[ \bar{\varphi}^2 + 3\mathcal{H}^2 + \mathcal{F}^2 - 2\bar{\varphi}'' + 2\mathcal{H}\bar{\varphi}' - Va^2 + \frac{\partial V}{\partial \bar{\varphi}} a^2 \right] \\ + \frac{1}{2} \nabla^2 Q_i + \frac{1}{2} \nabla^2 W'_i + \frac{a^2}{2b^2} \nabla_y^2 Q_i - \frac{a}{2b} \partial_y [H'_i + (\mathcal{F} \\ - \mathcal{H}) H_i] = 0, \end{aligned} \quad (3.25)$$

$$\begin{aligned} & \frac{1}{2} \partial_y [Q'_i + (\mathcal{H} - 2\mathcal{F} - \bar{\varphi}') Q_i] + \frac{1}{2} \partial_y \nabla^2 W_i \\ & - \frac{b}{2a} [H''_i - (\bar{\varphi}' + \mathcal{H}) H'_i - \nabla^2 H_i - (\mathcal{H} - \mathcal{F})^2 H_i \\ & - \nabla^2 H_i] = 0. \end{aligned} \quad (3.26)$$

Recalling the definitions of the gauge-invariant fluctuations given in Eqs. (3.18) and (3.19), Eq. (3.24) becomes

$$\begin{aligned} & - \frac{a}{2b} \partial_y [\partial_i Z^j + \partial^j Z_i] + \frac{1}{2} [(\partial_i V^j + \partial^j V_i)' \\ & - (\bar{\varphi}' + \mathcal{H})(\partial_i V^j + \partial^j V_i)] = 0, \end{aligned} \quad (3.27)$$

By virtue of Eq. (3.11), the first term of Eq. (3.25) vanishes. The use of the gauge-invariant fluctuations  $V_i$  and  $Z_i$  into Eq. (3.25) then leads to

$$- \frac{a}{2b} \partial_y [Z'_i + (\mathcal{F} - \mathcal{H}) Z_i] + \frac{a^2}{2b^2} \nabla_y^2 V_i + \frac{1}{2} \nabla^2 V_i = 0. \quad (3.28)$$

Finally, inserting Eqs. (3.18) and (3.19) into Eq. (3.26), the last gauge-invariant evolution equation will be: the (iy) equation

$$\begin{aligned} & \frac{1}{2} \partial_y [V'_i + (\mathcal{H} - 2\mathcal{F} - \bar{\varphi}') V_i] - \frac{b}{2a} [Z''_i - (\bar{\varphi}' + \mathcal{H}) Z'_i \\ & - (\mathcal{H} - \mathcal{F})^2 Z_i - \nabla^2 Z_i] = 0. \end{aligned} \quad (3.29)$$

Equations (3.27)–(3.29) can also be derived with gauge-dependent methods, as can be expected from general considerations. For instance, fixing the gauge in such away that  $W_i = 0$  the set of Eqs. (3.27)–(3.29) can be

exactly reproduced, as was checked by explicit calculation.

#### IV. SPECTRUM OF VECTOR MODES

For the purposes of this investigation, it is interesting to consider the spectrum of the zero modes with respect to the internal dimensions. This is achieved by setting to zero all the terms containing one (or more) derivatives with respect to  $y$ . Thus, from Eqs. (3.28)–(3.29) the following system:

$$\nabla^2 Q_i = 0, \quad (4.1)$$

$$Q'_i = (\mathcal{H} + \bar{\varphi}') Q_i, \quad (4.2)$$

$$H''_i - (\mathcal{H} + \bar{\varphi}') H'_i - (\mathcal{H} - \mathcal{F})^2 H_i - \nabla^2 H_i = 0, \quad (4.3)$$

is obtained. The evolution of  $H_i$  and  $Q_i$  is, in this approximation, decoupled. Furthermore, it can be noticed that the evolution of  $Q_i$  is the one already discussed in the four-dimensional case, with the minor difference that, in the present case, the background evolution differs from the four-dimensional case. From Eq. (4.2) we then have

$$Q_i(t) = c_i \frac{\sqrt{\tau(\tau^2 + 1) + (\tau^2 + 1)^{3/2}}}{\tau^2 + 1}, \quad (4.4)$$

which also implies that  $Q_i \sim t^{-1/2}$  for  $t \gg t_0$ . Hence, even if the time dependence is different from the four-dimensional case, it is still true that the  $Q_i$  decreases at large (positive) times even if the solution is growing just before the bounce.

Let us now concentrate on the evolution of  $H_i$  whose equation can be written, in Fourier space and for a single vector polarization as<sup>5</sup>

$$u'' + [k^2 - \mathcal{U}(\eta)]u = 0, \quad (4.5)$$

$$\mathcal{U}(\eta) = (\mathcal{H} - \mathcal{F})^2 - \left( \frac{e^{-\bar{\varphi}/2}}{\sqrt{a}} \right)'' \sqrt{a} e^{\bar{\varphi}/2}, \quad (4.6)$$

where  $H_i = \sqrt{a} e^{\bar{\varphi}/2} u_i$ . This equation will be numerically solved in a moment for different values of  $k$ . However, in order to cross-check the numerical results it is useful to anticipate the analytical estimate. For this purpose a useful observation is that, for times that are sufficiently far from the bounce, the asymptotic solutions are an excellent approximation for the background evolution.

With this logic in mind, for  $t \ll -t_0$  the solution given by Eqs. (3.6) and (3.7) can be approximated, in conformal time, by

$$a(\eta) \sim (-\eta)^{-1/3}, \quad b(\eta) \sim (-\eta)^{1/3}, \quad \bar{\varphi} \sim -\frac{2}{3} \ln(-\eta). \quad (4.7)$$

For  $t \gg t_0$ , the solution (3.6) and (3.7) behaves, in conformal time, as

$$a(\eta) \sim \eta, \quad b(\eta) \sim \frac{1}{\eta}, \quad \bar{\varphi} \sim -2 \ln \eta. \quad (4.8)$$

Therefore, in the two opposite limits, Eq. (4.5) is given, respectively, by

$$u''_i + \left[ k^2 - \frac{4\nu^2 - 1}{4\eta^2} \right] u_i = 0, \quad \eta < -\eta_1, \quad (4.9)$$

$$u''_i + \left[ k^2 - \frac{4\mu^2 - 1}{4\eta^2} \right] u_i = 0, \quad \eta > \eta_1, \quad (4.10)$$

where  $\eta_1 \sim \eta_0$  is a time scale comparable with the size of the bounce.

Inserting Eqs. (4.7) and (4.8) into Eq. (4.6) the indices  $\mu$  and  $\nu$  are determined to be  $\nu = 2/3$  and  $\mu = 2$ . Equations (4.9) and (4.10) can be solved exactly in the two limits. Now, imposing quantum-mechanical initial conditions for  $\eta \rightarrow -\infty$ ,

$$u_-(\eta) = \frac{\sqrt{-\pi\eta}}{2} e^{i\frac{\pi}{2}(\nu+1/2)} H_\nu^{(1)}(-k\eta), \quad \eta < -\eta_0, \quad (4.11)$$

the solution for  $\eta \rightarrow +\infty$  can be written in terms of the mixing coefficients  $c_\pm(k)$

$$u_+(\eta) = \frac{\sqrt{\pi\eta}}{2} e^{i\frac{\pi}{2}(\mu+1/2)} [c_-(k) H_\mu^{(1)}(k\eta) + c_+(k) e^{-i\pi(\mu+1/2)} H_\mu^{(2)}(k\eta)], \quad \eta > \eta_0, \quad (4.12)$$

where  $H_\alpha^{(1)}(z) = H_\alpha^{(2)*}(z) = J_\alpha(z) + iY_\alpha(z)$  are the usual Hankel functions of argument  $z$  and index  $\alpha$  [23]. Notice that the phases in Eqs. (4.11) and (4.12) have been carefully selected in such a way as to recover mode functions with canonical quantum-mechanical normalization in the limits  $\eta \rightarrow \pm\infty$ .

The two solutions (4.11) and (4.12) can be continuously matched across the bounce, i.e., imposing  $u_-(-\eta_1) = u_+(\eta_1)$  and similarly for the first derivative. The mixing coefficients turn out to be

$$c_+(k) = i \frac{\pi}{4} x_1 e^{i\pi(\nu+\mu+1)/2} \left[ -\frac{\nu + \mu + 1}{x_1} H_\mu^{(1)}(x_1) H_\nu^{(1)}(x_1) + H_\mu^{(1)}(x_1) H_{\nu+1}^{(1)}(x_1) + H_{\mu+1}^{(1)}(x_1) H_\nu^{(1)}(x_1) \right], \quad (4.13)$$

<sup>5</sup>Notice that in the following the index denoting the Fourier mode will be omitted in order to avoid confusion with the vector polarization.



$$c_-(k) = i \frac{\pi}{4} x_1 e^{i\pi(\nu-\mu)/2} \left[ -\frac{\nu + \mu + 1}{x_1} H_\mu^{(2)}(x_1) H_\nu^{(1)}(x_1) + H_\mu^{(2)}(x_1) H_{\nu+1}^{(1)}(x_1) + H_{\mu+1}^{(2)}(x_1) H_\nu^{(1)}(x_1) \right], \quad (4.14)$$

satisfying the exact Wronskian normalization condition  $|c_+(k)|^2 - |c_-(k)|^2 = 1$ . In the small argument limit, i.e.,  $k\eta_1 \sim k\eta_0 \ll 1$  the leading term in Eq. (4.14) leads to

$$c_-(k) \simeq \frac{i2^{\mu+\nu}}{4\pi} e^{i\pi(\nu-\mu)/2} x_1^{-\mu-\nu} (\nu + \mu - 1) \Gamma(\mu) \Gamma(\nu), \quad (4.15)$$

which implies, after inserting the specific values of  $\mu$  and  $\nu$  that the number of produced vector quanta is given, for each polarization, by

$$|c_-(k)|^2 \sim |k/k_1|^{-2\gamma}, \quad \gamma = (\mu + \nu) \sim 2.6, \quad (4.16)$$

implying, in turn, that the energy density per logarithmic frequency interval, i.e.,  $k^4 |c_-(k)|^2$ , increases at large distance-scales (small  $k$ ) as  $k^{-1.12}$ .

The same estimate will now be obtained through a totally different procedure, where the evolution equation for  $H_i$  is solved directly using the exact background solutions. Since the background solutions have a simple form in cosmic time, it is more practical to solve the evolution equation of the fluctuations directly in cosmic time. This will not affect the determination of the mixing

coefficients, which are independent of the specific parametrization of the time coordinate.

Equation (4.3) becomes, in cosmic time,

$$\ddot{H}_i - \dot{\bar{\varphi}} \dot{H}_i + [\omega^2 - (H - F)^2] H_i = 0, \quad (4.17)$$

where  $\omega(t) = k/a(t)$  is the physical momentum. By defining now the rescaled variable  $v_i = e^{-\bar{\varphi}/2} H_i$ , Eq. (4.17) becomes:

$$\ddot{v}_i + \left[ \omega^2 + \frac{\ddot{\bar{\varphi}}}{2} - \frac{\dot{\bar{\varphi}}^2}{2} - (H - F)^2 \right] v_i = 0. \quad (4.18)$$

Inserting Eqs. (3.6) and (3.7) into Eq. (4.18), the following simple equation for a single vector polarization can be obtained:

$$\frac{d^2 v}{d\tau^2} + \left[ \frac{\kappa^2}{\tau \sqrt{\tau^2 + 1}} - \frac{3(\tau^2 + 2)}{4(\tau^2 + 1)^2} \right] v = 0, \quad (4.19)$$

where  $\tau = t/t_0$  is the usual rescaled cosmic time and  $\kappa = kt_0$ .

Notice that the variable  $v$  defined in Eq. (4.19) is simply related to the variable  $u$  defined in Eq. (4.5):

$$H_i = \sqrt{a} e^{\bar{\varphi}/2} u_i = e^{\bar{\varphi}/2} v_i. \quad (4.20)$$

Since  $v = \sqrt{a} u$ , the quantum-mechanical initial conditions will be, in terms of  $v$

$$v(t) = \frac{1}{\sqrt{2\omega}} e^{-i \int \omega dt}, \quad (4.21)$$




FIG. 2 (color online). The numerical calculation of the mixing coefficients for different values of  $\kappa = kt_0$ , the comoving momentum rescaled through the size of the bounce. The dashed line is Eq. (4.25) whose right-hand side has been numerically computed.

for  $t \rightarrow -\infty$ . For  $t \rightarrow +\infty$  the solution can again be written in terms of the mixing coefficients  $c_{\pm}$  as

$$v(t) = \frac{1}{\sqrt{2\omega}} \left[ c_+(k) e^{-i \int \omega dt} + c_-(k) e^{i \int \omega dt} \right]. \quad (4.22)$$

Equation (4.19) was solved for different values of  $k$  (or  $\kappa = kt_0$ , the dimensionless comoving momentum). Quantum-mechanical initial conditions are given when the given mode is inside the horizon at the initial time of integration. From the numerical solution for the real and imaginary parts of  $v$  and  $\dot{v}$ , i.e.,

$$v = f + iq, \quad \dot{v} = g + ip, \quad (4.23)$$

it is then easy to obtain the mixing coefficients after the relevant mode reentered during radiation, i.e.,

$$|c_+(k)|^2 + |c_-(k)|^2 = \frac{1}{\omega} \left[ \omega^2 (f^2 + q^2) + \left( g - \frac{H}{2} f \right)^2 + \left( p - \frac{H}{2} q \right)^2 \right], \quad (4.24)$$

$$|c_+(k)|^2 - |c_-(k)|^2 = 2(qg - fp). \quad (4.25)$$

The results of this calculation are reported in Fig. 2. Recall that, according to Eq. (4.25),  $|c_+(k)|^2 - |c_-(k)|^2 = 1$  since  $|c_+(k)|^2 - |c_-(k)|^2$  is proportional to the Wronskian of the solution. From Fig. 2, it is apparent that for an increase of 1 order of magnitude in  $\kappa$ ,  $|c_-(k)|^2$  increases by 5.1 orders of magnitude, exactly as predicted in Eq. (4.16).

## V. CONCLUDING REMARKS

In the present paper the evolution of vector modes of the geometry has been discussed in the case of bouncing cosmologies both in four and in higher dimensions. The main suggestions following from a series of detailed examples can be summarized as follows:

- (i) in the four-dimensional case, as also noted in [8], growing vector modes may be present in the regime preceding the curvature bounce;
- (ii) however, in the analytical models described in the present paper, the growing mode matches with a decaying mode after the curvature bounce, so that no further dangerous divergences are introduced;
- (iii) in higher dimensions, more vector modes are expected and their spectrum was computed in a simple five-dimensional model of curvature bounce, where the compactification of the internal dimension is achieved dynamically;
- (iv) the typical spectra of vector modes obtained in the specific (but fully consistent) example are red after the modes reenter the horizon.

The most problematic aspect of this analysis is represented by the possible occurrence of red spectra, which may be rather dangerous in models based on curvature bounces.

## APPENDIX: VECTOR MODES OF THE GEOMETRY

The three divergenceless vectors appearing in five dimensions are defined as

$$\begin{aligned} \delta G_{0i} &= -a^2 Q_i, & \delta G_{ij} &= a^2 (\partial_i W_j + \partial_j W_i), \\ \delta G_{iy} &= -ab H_i \end{aligned} \quad (A1)$$

and

$$\begin{aligned} \delta G^{0i} &= -\frac{Q^i}{a^2}, & \delta G^{ij} &= -\frac{1}{a^2} (\partial^i W^j + \partial^j W^i), \\ \delta G^{iy} &= \frac{H^i}{ab}. \end{aligned} \quad (A2)$$

Notice that Eqs. (A1) and (A2) are general in the sense that no specific gauge choice has been imposed. From Eqs. (A1) and (A2)

$$\begin{aligned} \delta \Gamma_{i0}^0 &= \mathcal{H} Q_i, & \delta \Gamma_{ij}^0 &= -\frac{1}{2} (\partial_i Q_j + \partial_j Q_i) - \mathcal{H} (\partial_i W_j + \partial_j W_i) - \frac{1}{2} (\partial_i W_j' + \partial_j W_i'), & \delta \Gamma_{00}^i &= Q^{i'} + \mathcal{H} Q^i, \\ \delta \Gamma_{i0}^j &= \frac{1}{2} (\partial_i Q^j - \partial^j Q_i) - \frac{1}{2} (\partial^j W_i' + \partial_i W^{j'}), & \delta \Gamma_{yy}^i &= \frac{b}{a} \partial_y H^i - \frac{b^2}{a^2} \mathcal{F} Q^i, \\ \delta \Gamma_{yi}^0 &= -\frac{1}{2} \partial_y Q_i + \frac{b}{2a} [H_i' + (\mathcal{H} + \mathcal{F}) H_i], & \delta \Gamma_{i0}^y &= \frac{a}{2b} [H_i' + (\mathcal{F} - \mathcal{H}) H_i] - \frac{a^2}{2b^2} \partial_y Q_i, \\ \delta \Gamma_{0y}^i &= \frac{1}{2} \partial_y Q^i + \frac{b}{2a} [H^{i'} + (\mathcal{H} - \mathcal{F}) H^i], & \delta \Gamma_{ij}^y &= \frac{a}{2b} (\partial_i H_j + \partial_j H_i) + \frac{a^2}{2b^2} (\partial_i W_j + \partial_j W_i), \\ \delta \Gamma_{ij}^k &= -\mathcal{H} Q^k \delta_{ij} + \frac{1}{2} \partial^k (\partial_i W_j + \partial_j W_i) - \frac{1}{2} \partial_j (\partial^k W_i + \partial_i W^k) - \frac{1}{2} \partial_i (\partial_j W^k + \partial^k W_j), \\ \delta \Gamma_{iy}^j &= \frac{b}{2a} (\partial_i H^j - \partial^j H_i) - \frac{1}{2} \partial_y (\partial^j W_i + \partial_i W^j). \end{aligned} \quad (A3)$$

From the expressions of the Christoffel connections, the first-order fluctuations of the Ricci tensors can be determined:

$$\delta R_{0i} = [\mathcal{H}' + 2\mathcal{H}^2 + \mathcal{H}\mathcal{F}]Q^i - \frac{a^2}{2b^2}\nabla_y^2 Q_i - \frac{1}{2}\nabla^2 Q_i - \frac{1}{2}\nabla^2 W_i' + \frac{a}{2b}\partial_y[H_i' + (\mathcal{F} - \mathcal{H})H_i], \quad (\text{A4})$$

$$\begin{aligned} \delta R_{ij} = & \frac{a}{2b}\partial_y[\partial_i H_j + \partial_j H_i] - \frac{1}{2}\{(\partial_i Q_j + \partial_j Q_i)' \\ & + (2\mathcal{H} + \mathcal{F})(\partial_i Q_j + \partial_j Q_i)\} - \frac{1}{2}(\partial_i W_j'' + \partial_j W_i'') \\ & - \frac{1}{2}(2\mathcal{H} + \mathcal{F})(\partial_i W_j' + \partial_j W_i') + \frac{a^2}{2b^2}\partial_y(\partial_i W_j \\ & + \partial_j W_i) - \frac{1}{2}(\partial_i W_j + \partial_j W_i)(2\mathcal{H}' + 4\mathcal{H}^2 \\ & + 2\mathcal{H}\mathcal{F}), \end{aligned} \quad (\text{A5})$$

$$\begin{aligned} \delta R_{iy} = & -\frac{1}{2}\partial_y[Q_i' + (4\mathcal{H} - \mathcal{F})Q_i] - \frac{1}{2}\partial_y\nabla^2 W_i \\ & + \frac{b}{2a}\{H_i'' + (2\mathcal{H} + \mathcal{F})H_i' + (\mathcal{H}' + \mathcal{F}' + \mathcal{H}^2 \\ & + 5\mathcal{H}\mathcal{F})H_i - \nabla^2 H_i\}. \end{aligned} \quad (\text{A6})$$

In Eqs. (A4)–(A6),  $\nabla_y^2$  denotes the Laplacian of the internal dimensions. The perturbed Ricci tensors with mixed indices are also useful for a swifter calculation of the perturbed equations:

$$\begin{aligned} \delta R_0^i = & \frac{Q^i}{a^2}[2\mathcal{H}' + \mathcal{F}' - 2\mathcal{H}^2 - 2\mathcal{H}\mathcal{F} + \mathcal{F}^2] \\ & + \frac{1}{2b^2}\nabla_y^2 Q^i + \frac{1}{2a^2}\nabla^2 Q^i + \frac{1}{2a^2}\nabla^2 W_i \\ & - \frac{1}{2ab}\partial_y[H^{ii'} + (\mathcal{F} - \mathcal{H})H^{i'}], \end{aligned} \quad (\text{A7})$$

$$\begin{aligned} \delta R_i^j = & -\frac{1}{2ab}\partial_y[\partial_i H^j - \partial^j H_i] + \frac{1}{2a^2}\{(\partial_i Q^j + \partial^j Q_i)' \\ & + (2\mathcal{H} + \mathcal{F})(\partial_i Q^j + \partial^j Q_i)\} + \frac{1}{2a^2}\{(\partial_i W^j \\ & + \partial^j W_i)'' + (2\mathcal{H} + \mathcal{F})(\partial_i W^j + \partial^j W_i)'\} \\ & - \frac{1}{2b^2}\partial_y[\partial_i W^j + \partial^j W_i], \end{aligned} \quad (\text{A8})$$

$$\begin{aligned} \delta R_i^y = & \frac{1}{2b^2}\partial_y[Q_i' + (4\mathcal{H} - \mathcal{F})Q_i] + \frac{1}{2b^2}\partial_y\nabla^2 W_i \\ & - \frac{1}{2ab}\{H_i'' + (2\mathcal{H} + \mathcal{F})H_i' + (\mathcal{F}' - \mathcal{H}' - 3\mathcal{H}^2 \\ & + 3\mathcal{H}\mathcal{F})H_i - \nabla^2 H_i\}. \end{aligned} \quad (\text{A9})$$

Clearly, since vector fluctuations are parametrized by divergenceless vectors in three dimensions, their contribution to the first-order fluctuation of the Ricci scalar vanishes. Hence, recalling the definition of the Einstein tensor  $\mathcal{G}_\mu^\nu$ ,

$$\delta\mathcal{G}_0^i = \delta R_0^i, \quad \delta\mathcal{G}_i^j = \delta R_i^j, \quad \delta\mathcal{G}_i^y = \delta R_i^y. \quad (\text{A10})$$

- 
- [1] L. P. Grishchuk, Phys. Rev. D **48**, 5581 (1993).  
[2] I. Wasserman, Astrophys. J. **224**, 337 (1978).  
[3] P. Coles, Comments Astrophys. **16**, 45 (1992).  
[4] E. Harrison, Mon. Not. R. Astron. Soc. **147**, 279 (1970).  
[5] I. Mishustin and A. Ruzmaikin, Sov. Phys. JETP **34**, 223 (1972).  
[6] T. Vachaspati and A. Vilenkin, Phys. Rev. Lett. **67**, 1057 (1991).  
[7] P. Avelino and P. Shellard, Phys. Rev. D **51**, 5946 (1995).  
[8] T. J. Battefeld and R. Brandenberger, hep-th/0406180.  
[9] J. Khouri, B. A. Ovrut, P. J. Steinhardt, and N. Turok, Phys. Rev. D **66**, 046005 (2002); **65**, 086007 (2002); N. Turok and P. J. Steinhardt, hep-th/0403020.  
[10] M. Gasperini, M. Giovannini, and G. Veneziano, Phys. Lett. B **569**, 113 (2003); M. Gasperini, M. Giovannini, and G. Veneziano, Nucl. Phys. B **694**, 206 (2004); M. Giovannini, Classical Quantum Gravity **21**, 4209 (2004).  
[11] D. H. Lyth, Phys. Lett. B **526**, 172 (2002); *ibid.* **524**, 1 (2002); R. Brandenberger and F. Finelli, J. High Energy Phys. **11** (2001) 054; Phys. Rev. D **65**, 103520 (2002); F. Finelli, J. Cosmol. Astropart. Phys. **0310** (2003) 011; S. Tsujikawa, R. Brandenberger, and F. Finelli, Phys. Rev. D **66**, 083513 (2002); S. Gratton, J. Khoury, P. J. Steinhardt, and N. Turok, Phys. Rev. D **69**, 103505 (2004); C. Cartier, hep-th/0401036; L. E. Allen and D. Wands, Phys. Rev. D **70**, 063515 (2004); R. Kallosh, L. Kofman, and A. D. Linde, Phys. Rev. D **64**, 123523 (2001); J. Martin and P. Peter, Phys. Rev. D **68**, 103517 (2003); **69**, 107301 (2004); P. Peter and N. Pinto-Neto, Phys. Rev. D **66**, 063509 (2002); J. Martin *et al.*, Phys. Rev. D **65**, 123513 (2002); J. C. Hwang and H. Noh, Phys. Rev. D **65**, 124010 (2002); Phys. Lett. B **545**, 207 (2002); J. C. Hwang, Phys. Rev. D **65**, 063514 (2002); C. Cartier, R. Durrer, and E. J. Copeland, Phys. Rev. D **67**, 103515 (2003); C. Gordon and N. Turok, Phys. Rev. D **67**, 123508 (2003).  
[12] J. M. Bardeen, Phys. Rev. D **22**, 1882 (1980).  
[13] R. Abbott, B. Bednarz, and S. D. Ellis, Phys. Rev. D **33**, 2147 (1986).  
[14] D. Sahdev, Phys. Rev. D **30**, 2495 (1984); **39**, 3155 (1989); R. Abbott, S. M. Barr, and S. D. Ellis, Phys. Rev. D **30**, 720 (1984); E. W. Kolb, D. Lindley, and D. Seckel, Phys. Rev. D **30**, 1205 (1984).

- [15] K. Tomita and M. Den, Phys. Rev. D **34**, 3570 (1986).
- [16] B. L. Hu, Phys. Rev. D **18**, 969 (1978).
- [17] M. Giovannini, Phys. Rev. D **55**, 595 (1997).
- [18] C. van de Bruck, M. Dorca, R. H. Brandenberger, and A. Lukas, Phys. Rev. D **62**, 123515 (2000).
- [19] E.W. Kolb, G. Servant, and T.M.P. Tait, J. Cosmol. Astropart. Phys. 0307 (2003) 008.
- [20] M. Giovannini, Phys. Rev. D **64**, 064023 (2001); **64**, 124004 (2001).
- [21] M. Giovannini, Phys. Rev. D **66**, 044016 (2002); **65**, 124019 (2002); M. Giovannini, J.V. Le Be, and S. Riederer, Classical Quantum Gravity **19**, 3357 (2002).
- [22] J. M. Cline, J. Descheneau, M. Giovannini, and J. Vinet, J. High Energy Phys. 0306 (2003) 048.
- [23] A. Erdelyi, W. Magnus, F. Obhehttinger, and F. Tricomi, *Higher Transcendental Functions* (McGraw-Hill, New York, 1953).