

Realization of minimal supergravity

M. Ibe, Izawa K.-I., and T. Yanagida

Department of Physics, University of Tokyo, Tokyo 113-0033, Japan

(Received 18 October 2004; published 8 February 2005)

Minimal supergravity mediation of supersymmetry breaking has attracted much attention due to its simplicity, which leads to its predictive power. We consider how Nature possibly realizes minimal supergravity through inflationary selection of the theory. Minimality is impressively consistent with the present observational bounds and it might be tested with the aid of low-energy soft parameters obtained in future experiments.

DOI: 10.1103/PhysRevD.71.035005

PACS numbers: 12.60.Jv

I. INTRODUCTION

The minimal supergravity (mSUGRA) [1] is a very interesting framework, since it has definite predictions on low-energy physics, which are well consistent with the present observations. In particular, the absence of flavor-changing neutral currents (FCNC's) in the present experimental precisions is one of the predictions of mSUGRA. If mSUGRA is (approximately¹) realized in truth, we wonder the reason of its selection by Nature, since it seems no more symmetric than nonminimal supergravity in the presence of a nontrivial superpotential, as is the case for the standard model of elementary particles.

In this paper, we point out that supergravity effective theory with a large cutoff scale M_* ($> M_G$) may be chosen by inflationary selection of background vacuum structures [2], which implies a specific type of mSUGRA theory. Here, M_G is the reduced Planck-scale, $M_G \simeq 2.4 \times 10^{18}$ GeV. The large cutoff suggests relatively small gaugino masses, which in turn indicate masses of squarks as large as a few TeV. In this parameter region, the mass of the lightest Higgs boson is easily raised up to the current experimental limit. In spite of the large stop mass, we may naturally obtain the breaking scale of electroweak symmetry at $\mathcal{O}(100)$ GeV [3,4] due to renormalization group (RG) focus point behavior [5] of a supersymmetry (SUSY) breaking soft mass of a Higgs boson.

The large-cutoff theories might be realized in various corners of theory moduli space, or (string [6,7]) landscape. We do not specify concrete construction of such theories but simply assume their presence. The task in this paper is not to achieve constructive realization of the large-cutoff but to seek a plausible way to select it among vast possibilities on the landscape.²

The rest of the paper goes as follows. In the next section, we consider possible inflationary selection of minimality in supergravity. In section III, we specify plausible boundary conditions on gravity mediation of SUSY breaking. In section IV, low-energy phenomenology is investigated by

means of RG analysis. Section V is devoted to conclusions and discussion.

II. POSSIBLE INFLATIONARY SELECTION

We are led by the following question: what is expected beyond the standard model³ as a typical structure of the natural laws? We here consider inflationary selection of background vacuum structures [2] and dwell on mediocrity principle, which may prefer flatter inflaton potential [7,9].⁴

For concreteness of presentation, let us adopt a simplest case of supergravity inflation model [10,11] as an example. Namely, we consider a single-superfield model for slow-roll inflation. In terms of a single chiral superfield ϕ , an inflaton φ can be provided by $\sqrt{2}$ times the real part of its lowest component. We adopt a natural superpotential⁵

$$W = v^2 \phi - \frac{y}{n+1} \phi^{n+1} \quad (1)$$

and a generic Kähler potential

$$K = |\phi|^2 + \frac{\kappa}{4} |\phi|^4 + \dots, \quad (2)$$

where $v^2, y, \kappa > 0$ and the ellipsis denotes higher-order terms, which may be disregarded. Here and henceforth in this section, we have taken the unit with a cutoff scale M_* equal to one. Note that the small scale v^2 can be generated dynamically [10,12].

The potential for the lowest component ϕ is given in supergravity by

$$V = \exp\left(\frac{K}{M_G^2}\right) \left\{ \left(\frac{\partial^2 K}{\partial \phi \partial \phi^\dagger} \right)^{-1} |DW|^2 - 3 \left| \frac{W}{M_G} \right|^2 \right\}, \quad (3)$$

where we have defined

³We suppose the standard model as a prerequisite with its presently measured values of the couplings.

⁴This is a terminology adopted by Vilenkin [9]: “We are one of the infinite number of civilizations living in thermalized regions of the metauniverse. Although it may be tempting to believe that our civilization is very special, the history of cosmology demonstrates that the assumption of being average is often a fruitful hypothesis.”

⁵This form is protected by nonrenormalization or R symmetry.

¹We use the term mSUGRA in an approximate sense and do not mean the strictly minimal Kähler potential in this paper.

²See the conclusion in Ref. [8].

$$DW \equiv \frac{\partial W}{\partial \phi} + \frac{\partial K}{\partial \phi} \frac{W}{M_G^2}. \quad (4)$$

Thus, the potential of the real part φ is approximately given by

$$V(\varphi) = v^4 - \frac{\kappa}{2} v^4 \varphi^2 - \frac{\lambda}{n!} \varphi^n \quad (5)$$

for $n \geq 3$ and $0 < \lambda, v^2, \varphi \ll 1$ with $\lambda/n! \equiv yv^2/2^{(n/2)-1}$. The parameters n, v , and λ are potentially under control by symmetry. Let us fix them hereafter, for simplicity of argument.

We adopt slow-roll approximation [13]. The slow-roll inflationary regime is prescribed by the condition

$$\epsilon(\varphi) = \frac{1}{2} \left(M_G \frac{V'(\varphi)}{V(\varphi)} \right)^2 \leq 1, \quad |\eta(\varphi)| \leq 1, \quad (6)$$

where

$$\eta(\varphi) = M_G^2 \frac{V''(\varphi)}{V(\varphi)}. \quad (7)$$

For the potential Eq. (5), we obtain

$$\begin{aligned} \frac{\epsilon}{M_G^2} &\simeq \frac{1}{2} \left(\frac{-\kappa v^4 \varphi - \frac{\lambda}{(n-1)!} \varphi^{n-1}}{v^4} \right)^2 \\ &= \frac{\varphi^2}{2} \left(\kappa + \frac{\lambda v^{-4}}{(n-1)!} \varphi^{n-2} \right)^2, \\ \frac{\eta}{M_G^2} &\simeq \frac{-\kappa v^4 - \frac{\lambda}{(n-2)!} \varphi^{n-2}}{v^4} = -\kappa - \frac{\lambda v^{-4}}{(n-2)!} \varphi^{n-2}, \end{aligned} \quad (8)$$

as slow-roll parameters.⁶ These parameters, ϵ and $|\eta|$, characterize the flatness of the inflaton potential.⁷ From a viewpoint of mediocrity principle, some tuning for flatter inflaton potential may be favored. For flatter potential, total

⁶Thus the slow-roll condition Eq. (6) is satisfied for $\varphi \leq \varphi_f$ where

$$\varphi_f^{n-2} \simeq \frac{(n-2)!(1 - \kappa M_G^2)}{\lambda v^{-4} M_G^2},$$

which provides the value φ_f of the inflaton field at the end of inflation. An initial value φ_i of the inflaton field amounts to the corresponding number N of total e -folding as [11]

$$\begin{aligned} N &= \int_{\varphi_f}^{\varphi_i} d\varphi M_G^{-2} \frac{V(\varphi)}{V'(\varphi)} \\ &\simeq \frac{1}{(n-2)\kappa M_G^2} \ln \left[\frac{1 - \kappa M_G^2}{1 + (n-2)\kappa M_G^2} \left(1 + \frac{(n-1)!\kappa}{\lambda v^{-4} \varphi_i^{n-2}} \right) \right]. \end{aligned}$$

⁷The spectral index n_s for fluctuations of the microwave background radiation is given by [14,15],

$$n_s = 1 - 6\epsilon + 2\eta \simeq 1 - 2\kappa \times \left(\frac{M_G}{M_*} \right)^2.$$

The present observation $n_s = 0.97 \pm 0.03$ [16] implies $(M_G/M_*)^2 \leq 3 \times 10^{-2}$ for $|\kappa| = 1$ which corresponds to the lower bound on M_* as $M_* \geq 6 \times M_G$.

amount of inflation becomes larger and inflation lasts longer, even possibly turns out to be eternal, to result in larger volume of habitable universe.

Small parameters ϵ and $|\eta|$ are achieved by tuning two apparent factors for flatter inflaton potential. One is obviously the coupling κ , which is small for small ϵ and $|\eta|$. The other is the reduced Planck scale M_G , which is also small for small ϵ and $|\eta|$, provided radiative corrections due to gravitational interaction controlled by M_G are loop suppressed and affect the effective coupling κ by at most order unity for $M_G \simeq M_*/4\pi$. This latter case is assumed in the following discussion,⁸ which corresponds to mSUGRA with a large cutoff M_* compared to the gravitational scale M_G . It leads to a particular pattern of effective Lagrangian parameters, whose details will be given in the following sections.

In the remainder of this section, let us further see possible implications of the large-cutoff $M_* \simeq 4\pi M_G$ on inflationary selection of background vacua. For that purpose, we consider multiple succession of inflations with each inflationary stage naturally preparing the initial conditions for the next stage [19]. The background vacua with multiple inflations seem to constitute remarkable ingredients in inflationary selection, based on which we seek a typical structure of the natural laws.

The point is that the tuning of the scale M_G might simultaneously realize successive inflations which are favorable according to mediocrity principle. This is to be contrasted to multiple tunings of each coupling (κ in the above example) corresponding to each inflaton potential.

In the theory (moduli) space, the background vacua (i.e., particular theories) with small Kähler couplings and/or small gravitational scale may induce inflation and be realized in Nature, which is at the heart of the inflationary selection.

III. PLAUSIBLE MINIMALITY

Motivated by the discussion in the previous section, we assume a large-cutoff M_* at an input scale Q_0 , below which we adopt the RG equations of the minimal supersymmetric standard model. This hypothesis leads to mSUGRA theory, since the large-cutoff suppresses higher dimensional operators in the Kähler potential. Before we explore low-energy implications of our large-cutoff hypothesis in the next section, let us set more detailed RG boundary conditions at the input scale⁹ around M_G for gravity mediation of SUSY-breaking.

⁸This seems possible in view of potential quantization [17] of Newton's constant in supergravity, though the size of the radiative corrections may be sensitive to the structure of ultraviolet physics [18] and beyond the scope of effective theory approach.

⁹We utilize the input scale $Q_0 = M_G$ or $Q_0 = M_{\text{GUT}}$ on occasion postulating that the ultraviolet contributions of RG above the so-called GUT scale $M_{\text{GUT}} \simeq 2 \times 10^{16}$ GeV are not significant.

As usual, the minimal Kähler potential generates a universal soft SUSY-breaking mass for chiral multiplets $m_{\text{scalar}} = m_0$. The universal scalar mass m_0 results in the universality of the scalar masses of squarks and sleptons in the first two generations providing a solution to the FCNC problem.

This minimality will be tested in the next generation accelerator experiments by examining spectra of the squarks and sleptons for $m_0 < \mathcal{O}(10)$ TeV. Hence we restrict our attention to this range in this paper. In order to attain sizable gaugino masses $M_i (i = 1, 2, 3)$, we adopt a singlet chiral superfield Z (Polonyi field) [20] with its F term as the dominant SUSY-breaking source in the hidden sector of gravity mediation. Note that the SUSY-breaking scale can be generated dynamically without its cosmological problem [12,14]. Then we obtain $M_i = \mathcal{O}(M_G/M_*)m_{3/2}$ through the F term of

$$\frac{f_i}{M_*} Z \mathcal{W}_{i\alpha} \mathcal{W}_i^\alpha, \quad (9)$$

where $\mathcal{W}_{i\alpha}$ and $f_i (i = 1, 2, 3)$ denote field strength chiral superfields for gauge multiplets and their order one coefficients, respectively, and Z has a SUSY-breaking F term as $F_Z \simeq \sqrt{3}m_{3/2}M_G$. Furthermore, the universal scalar mass m_0 can be expressed in terms of the gravitino mass $m_{3/2}$ as $m_0 = m_{3/2}$. Namely, for $M_* \simeq 4\pi M_G$ advocated in the previous section, the spectrum of the supersymmetric standard model (SSM) particles has a hierarchical structure, $m_0 \gg |M_i|$. For $m_0 = m_{3/2} < \mathcal{O}(1)$ TeV, gaugino masses of $\mathcal{O}(M_G/M_*)m_{3/2}$ would be too small and thus we are led to adopt an mSUGRA boundary condition $m_0 = \mathcal{O}(1 - 10)$ TeV.

In addition to m_0 and M_i , the Polonyi field Z in mSUGRA also determines so-called A parameters of SUSY-breaking. The A parameter for each (scalar)³ coupling is proportional to a universal A_0 parameter and the corresponding Yukawa coupling constant for the minimal Kähler potential. Since A_0 is proportional to the vacuum expectation value (VEV) of the Polonyi field Z , that is, $A_0 \simeq (\langle Z^* \rangle / M_G)m_{3/2}$, the assumption $|A_0| \lesssim m_0$ is not so implausible within a well-controlled expansion on Z/M_G in supergravity effective theory.¹⁰

So far, we have concentrated on the SUSY-breaking parameters in mSUGRA. By virtue of the Polonyi field, we also naturally obtain the supersymmetric Higgs mixing parameter μ of the electroweak order: the μ term can be provided through the Giudice-Masiero (GM) term [21] in the Kähler potential

$$K \supset \frac{\mathcal{O}(1)}{M_*} Z^* H_u H_d, \quad (10)$$

¹⁰The potential in supergravity has the e^{K/M_G^2} factor, which forces $|\langle Z \rangle| \lesssim M_G$ provided $K \simeq |Z|^2$. Note that such a range of the A_0 parameter is adequate to avoid color symmetry breaking.

which relates μ to the SUSY-breaking parameters, where $H_{u,d}$ denote the up-type and down-type Higgs superfields. Then the parameter μ is expected to be of the same order of the gaugino masses: $\mu \sim M_i \simeq \mathcal{O}(M_G/M_*)m_{3/2}$. This GM term yields a specific form of the so-called B term,¹¹ which is given explicitly in section IV B.

IV. LOW-ENERGY PHENOMENOLOGY

In the previous section, we have proposed specific mSUGRA boundary conditions: $m_0 = \mathcal{O}(1 - 10)$ TeV with a hierarchical structure¹² $m_0 \gg |M_i|, |\mu|$. It is non-trivial that the present scenario admits the electroweak symmetry breaking at the correct energy scale, in particular, since masses of squarks, sleptons, and Higgs bosons are very large at the input scale Q_0 . Fortunately, the mSUGRA boundary conditions turn out to be consistent with the present observational constraints on the electroweak physics. In this section, we show numerical analyses which indicate this consistency. For the sake of explanatory convenience, let us consider the case $\tan\beta \lesssim 10$ in the followings, though numerical estimates include results on the case with smaller $\tan\beta$ (see Fig. 5). Here, $\tan\beta \equiv v_u/v_d$ is the ratio of the two VEV's of the neutral Higgs fields, $v_{u,d} \equiv \langle H_{u,d}^0 \rangle$.

A. Consistency of the electroweak symmetry breaking

With large $\tan\beta$, the scale of electroweak symmetry breaking tends to be controlled by the value of a SUSY-breaking soft mass parameter $m_{H_u}^2$ (see Eq. (13) below) at a relevant scale for the electroweak physics, to be called the electroweak scale.

In mSUGRA, the running value of the parameter $m_{H_u}^2$ is related to the original parameters m_0, M_i , and A_0 by

$$m_{H_u}^2 = am_0^2 + b^i |M_i|^2 + c |A_0|^2 + d^i \text{Re}(M_i A_0^*), \quad (11)$$

where the coefficients $a - d^i$ are scale-dependent functions of dimensionless gauge and Yukawa coupling constants.¹³ As discussed in Refs. [3–5], the coefficients a and c are of order $10^{-2} - 10^{-1}$ at the electroweak scale for the pole mass of the top quark $m_t \simeq 170 - 180$ GeV, which results in the RG focus point behavior.

The smallness of the coefficient a comes from a cancellation between $m_{H_u}^2(Q_0) = m_0^2$ at the input scale Q_0 and RG contributions at the renormalization scale Q :

¹¹Our convention for the Higgs mass parameters is given in the Appendix.

¹²This hierarchy is natural in the large-cutoff theory with $M_* \simeq 4\pi M_G$, whereas it requires at least $\mathcal{O}(M_G^2/M_*^2)$ tuning (implemented by some flavor symmetry) in the ordinary mSUGRA theory with M_G as a cutoff scale, even if we presuppose minimality to put aside the corresponding tuning.

¹³Approximate analytical expressions for the coefficients $a - d^i$ can be found in Ref. [22], for instance.




FIG. 1 (color online). The m_0 dependence of $m_{H_u}^2$ at the stop mass scale $Q = Q_{\bar{t}}$ for $\text{sgn}(\mu) = +1$ and for the top quark pole mass (a) $m_t = 174$ GeV or (b) $m_t = 178$ GeV. The solid lines correspond to $M_{1/2} = 200$ GeV and $A_0 = 0$ GeV, the dashed lines $M_{1/2} = 300$ GeV and $A_0 = 0$ GeV, the dash-dot-dotted lines $M_{1/2} = 200$ GeV and $A_0 = 0.5m_0$, and the dash-dotted lines $M_{1/2} = 200$ GeV and $A_0 = m_0$. The three lines of each type correspond to $\tan\beta = 10, 20,$ and 30 , respectively, from below.

$$\begin{aligned} \delta a(Q)m_0^2 &\simeq -\frac{1}{2}\bar{m}_0^2 \cdot (1 - e^{L(Q)}) \simeq -\frac{1}{3}\bar{m}_0^2; \\ \bar{m}_0^2 &\equiv (m_{H_u}^2 + m_{Q_3}^2 + m_{U_3}^2)(Q_0) = 3m_0^2, \\ L(Q) &= \frac{1}{16\pi^2} \int_{\ln Q_0}^{\ln Q} 12|y_t|^2(Q')d \ln Q', \end{aligned} \quad (12)$$

where the subscripts Q_3 and U_3 represent an $SU(2)_L$ doublet quark and a singlet up-type quark, respectively, in the third family, and y_t denotes the top Yukawa coupling constant.¹⁴ In the first equation above, we use the fact $e^{L(Q)} \simeq 1/3$ at the electroweak scale for $m_t \simeq 170$ GeV.

On the other hand, the smallness of the coefficient c can be traced to the RG evolutions of (scalar)3 coupling constants $A_i \cdot y_i (i = t, b, \tau)$. As the renormalization scale is lowered from the input scale Q_0 , the A_i 's become small exponentially from A_0 and the contributions from the A terms to the RG evolution of $m_{H_u}^2$ become small. Thus, the A_0 dependence of $m_{H_u}^2$ at the electroweak scale (i.e. the coefficient c) is relatively small.

As a result, $m_{H_u}^2$ is suppressed compared to m_0 at the electroweak scale for $|M_i| \ll m_0$, $|A_0| < m_0$ and becomes of the same order of the gaugino masses.

In Fig. 1, we show the m_0 dependence of $m_{H_u}^2$ at a typical stop mass scale $Q = Q_{\bar{t}} = (m_{\bar{t}_1}m_{\bar{t}_2})^{1/2}$. In this computation, we impose the boundary condition $m_{\text{scalar}}^2 = m_0^2$ at the GUT scale (i.e. $Q_0 = M_{\text{GUT}}$),¹⁵ and take a universal gaugino mass $M_i = M_{1/2}$, for simplicity. As expected, the value of $m_{H_u}^2$ is much suppressed compared to the corresponding m_0^2 for the case of $|M_i| \ll m_0$ and $|A_0| < m_0$. We have plotted it for two central values of the observed top

¹⁴Here, we assume that the bottom Yukawa coupling constant y_b is negligible compared to y_t . Even if $y_b \simeq y_t$, the m_0 insensitivity of $m_{H_u}^2$ is valid, although the RG contribution from the bottom-type squark becomes important [5].

¹⁵We assume that the change of the input scale from M_{GUT} to M_G does not disturb the hierarchy $|m_{H_u}^2(Q_{\bar{t}})| \ll m_0^2$. For instance, this is the case in the grand unification scenario, since $\mathbf{10} \supset (Q_3, U_3)$ and $\mathbf{5} \supset (H_u)$ have the same RG trajectory between the M_G and M_{GUT} in the limit of M_i vanishing.

quark pole mass m_t : one is extracted from the Particle Data Group [16] $m_t = 174.3 \pm 5.1$ GeV and the other from the recent CDF and D0 results [23] $m_t = 178.0 \pm 4.3$ GeV.

In the SSM, the parameter μ is related to the Z^0 boson mass m_{Z^0} by minimizing the effective Higgs potential, and it can be expressed at the tree-level as

$$\begin{aligned} \frac{1}{2}m_{Z^0}^2 &= \frac{m_{H_d}^2 - m_{H_u}^2 \tan^2\beta}{\tan^2\beta - 1} - |\mu|^2 \\ &\simeq \frac{m_{H_d}^2}{\tan^2\beta} - m_{H_u}^2 - |\mu|^2. \end{aligned} \quad (13)$$

Thus, $m_{H_u}^2$ of the order of the gaugino masses manages to generate the electroweak symmetry breaking at the correct energy scale, or $m_{Z^0} = 91.2$ GeV, with the μ parameter naturally implied by the GM term (10). Here, the parameters in Eq. (13) are regarded to be values at the electroweak scale, while the RG evolution of μ from the electroweak scale to the input scale is negligible in order estimation (see Eq. (A2) in the Appendix).

In Fig. 2, we show admissible values of the μ parameter which yield the observed value of $m_{Z^0} = 91.2$ GeV. To determine the value of μ , we have used the ISAJET 7.69 code [24], which takes into account the one-loop corrections to the effective Higgs potential and the two-loop RG evolutions of parameters.¹⁶ The minimization of the effective Higgs potential is also performed at the typical stop mass scale $Q = Q_{\bar{t}}$. We see that the value of μ should be much suppressed compared to m_0 for $|M_i| \ll m_0$, $|A_0| < m_0$ and hence it is consistent with our large-cutoff hypothesis.

Let us comment on the falling-off behavior of the allowed μ in the very large m_0 region in Fig. 2. This behavior stems from a large cancellation between the suppressed m_0^2 contribution and the remaining ones to the value of the μ parameter. In such a region, the μ becomes very small due to the cancellation.

¹⁶The discrepancy of the value of the μ parameter among computational codes is discussed in Ref. [25].




FIG. 2 (color online). The m_0 dependence of μ at the stop mass scale $Q = Q_{\bar{t}}$ for $\text{sgn}(\mu) = +1$ and for the top quark pole mass (a) $m_t = 174$ GeV or (b) $m_t = 178$ GeV. The notations for the lines are the same as in Fig. 1, except for the corresponding values of $\tan\beta$ reversed in order: $\tan\beta = 10, 20$, and 30 from above.

From a cosmological point of view, the parameter regions with tiny μ may provide a natural explanation for the observed dark matter density [26], since the lightest neutralino is a bino-Higgsino mixture for such regions, and its relic abundance is in a cosmologically interesting range.

B. Consistency of the tree-level B term

We have assumed the GM term as the origin of the μ parameter of the electroweak order. Then, the SUSY-breaking Higgs mixing parameter B_0 is related to the A_0 parameter at the input scale. For the term (10), the tree-level relation is given by [21]

$$B_0 = B_0^{GM} \equiv \frac{2A_0 - 3m_{3/2}}{A_0 - 3m_{3/2}} m_{3/2}. \quad (14)$$

In this subsection, we examine how this condition is satisfied in the electroweak physics.¹⁷

To study the matching condition Eq. (14), we take the following procedure: We first fix sampling values of $[m_0, M_i, A_0, \text{sgn}(\mu), \tan\beta]$, and determine the required values of μ and B that reproduce $m_{Z^0} = 91.2$ GeV. In addition to Eq. (13), we have a relation

$$B\mu = \frac{\sin 2\beta}{2} (m_{H_u}^2 + m_{H_d}^2 + 2|\mu|^2), \quad (15)$$

which is also obtained by minimizing the tree-level effective Higgs potential. By means of Eqs. (13) and (15),¹⁸ we can obtain μ and B at the electroweak scale for the given mSUGRA parameters. Then, from the value of B at the electroweak scale, we compute the B parameter at the input scale (i.e. B_0) and compare it with B_0^{GM} in Eq. (14) to see whether or not the condition Eq. (14) is satisfied.

¹⁷A similar analysis is performed in Ref. [27] for small m_0 regions.

¹⁸More precisely, their one-loop corrections are taken into account in the numerical analysis below.

We find from Eq. (15) that the required value of B at the electroweak scale is given by

$$B \sim \frac{m_0^2}{\mu \tan\beta}, \quad (16)$$

since $m_{H_d}^2 \sim m_0^2 \gg |m_{H_u}^2|, |\mu|^2$ and $\sin 2\beta/2 \simeq 1/\tan\beta$ for $\tan\beta \gtrsim 10$.

In Fig. 3, we show numerical results on the value of B at the stop mass scale $Q = Q_{\bar{t}}$ as a function of m_0 . By comparing it with the μ parameter in Fig. 2, we find that the value of B becomes very large when the value of μ becomes very small, as expected from Eq. (16).

On the other hand, from the relation Eq. (14), we see $B_0^{GM} \sim m_0$ for $|A_0| < m_0 = m_{3/2}$. Thus Eq. (16) implies that the condition Eq. (14) can be satisfied for $\mu \sim m_0/\tan\beta \ll m_0$, since the RG evolution of B from the electroweak scale to the input scale is not significant: the RG equation of B is controlled by relatively small M_i and A parameter contributions (see Eq. (A3) in the Appendix), and thus $B \sim B_0$. As a result, we find that the condition Eq. (14) is satisfied in a certain parameter region.

In Figs. 4 and 5, we plot the contours of the values of B_0 at the input scale $Q_0 = M_{\text{GUT}}$ on the $(m_0, M_{1/2} \geq 0)$ plane with the fixed values of $A_0 = 0$, $\text{sign}(\mu) = +1$, and $\tan\beta = 20, 10, 5, 4$ as demonstrations. In the figures, we also show the value of μ at the stop mass scale $Q = Q_{\bar{t}}$.

Before examining the contours of B_0 , let us first understand the behavior of $\mu = \text{constant}$ lines in the figures. Generic behavior of the $\mu = \text{constant}$ lines may be seen from the panels (b,d) in Fig. 4 and (a) in Fig. 5. The lines are elliptic in small m_0 regions (elliptic domains) and hyperbolic in large m_0 regions (hyperbolic domains). They are parabolic in between (parabolic domains), where the value of μ is relatively insensitive to the variation of m_0 with $M_{1/2}$ fixed. In this perspective, the panels (a,c) in Fig. 4 belong to hyperbolic domains (continued from parabolic domains), where the $\mu = \text{constant}$ lines are hyperbolae (in the large $\tan\beta$ case in Ref. [4]); and the panels (b,c,d) in Fig. 5 belong to elliptic domains (continued on parabolic domains), where the $\mu = \text{constant}$ lines are el-




FIG. 3 (color online). The m_0 dependence of B at the stop mass scale $Q = Q_i$ for $\text{sgn}(\mu) = +1$ and for the top quark pole mass (a) $m_t = 174$ GeV or (b) $m_t = 178$ GeV. The notations for the lines are the same as in Fig. 1.




FIG. 4 (color online). Contour plots for the required values of $B = B_0$ at the input scale $Q_0 = M_{\text{GUT}} \approx 2 \times 10^{16}$ GeV in the $(m_0, M_{1/2})$ plane for the top quark pole mass (a,c) $m_t = 174$ GeV or (b,d) $m_t = 178$ GeV and for (a,b) $\tan\beta = 20$ or (c,d) $\tan\beta = 10$. In all the panels, we have fixed $A_0 = 0$, $\text{sgn}(\mu) = +1$, and the solid lines correspond to $B_0 = 0.2m_0, 0.5m_0, m_0$, respectively, from the left to the right. The GM term implies the parameter regions on the $B_0 = B_0^{\text{GM}} = m_0$ lines (see Eq. (15)). The gray shaded regions correspond to the parameters where the required value of μ is smaller than 700 GeV, 500 GeV, and 300 GeV, respectively, from above. The dark shaded regions at the bottom-left represent the Higgs mass bound 114 GeV[29], and the black shaded regions are excluded by the chargino mass limit, $m_{\chi^\pm} \geq 104$ GeV [30].




FIG. 5 (color online). Contour plots for the required values of $B = B_0$ at the input scale $Q_0 = M_{\text{GUT}} \approx 2 \times 10^{16}$ GeV in the $(m_0, M_{1/2})$ plane for the top quark pole mass (a,c) $m_t = 174$ GeV or (b,d) $m_t = 178$ GeV and for (a,b) $\tan\beta = 5$ or (c,d) $\tan\beta = 4$. In all the panels, we have fixed $A_0 = 0$, $\text{sgn}(\mu) = +1$, and the solid lines correspond to B_0 fixed as indicated. The GM term implies the parameter regions on the $B_0 = B_0^{\text{GM}} = m_0$ lines (see Eq. (15)). The shaded regions correspond to the parameters where the required value of μ is smaller than 1200 GeV, 900 GeV, and 600 GeV from above in the panel (a); smaller than 2000 GeV, 1500 GeV, and 1000 GeV from the right in the panels (b,c); and smaller than 3000 GeV, 2000 GeV, and 1000 GeV from the right in the panel (d). The dark shaded regions at the bottom-left represent the Higgs mass bound 114 GeV [29], and the black shaded regions are excluded by the chargino mass limit, $m_{\chi^\pm} \geq 104$ GeV [30].

lipes (in the small $\tan\beta$ case in Ref. [4]). We note that such behavior may be obtained from Eqs. (11)–(13), with the stop mass as the renormalization scale¹⁹ $Q = Q_t \sim m_0$.

Now, by inspection of $B_0 = B_0^{\text{GM}} = m_0$ lines due to the GM term (10) for $A_0 = 0$ (see Eq. (15)) in the figures, it is apparent that the large-cutoff theories with $m_0 \gg |M_{1/2}|, |\mu|$ correspond to those lines along the hyperbolic domains in Fig. 4 with moderate $\tan\beta$. In particular, the lower ends of the $B_0 = m_0$ lines are in small $M_{1/2}$ regions

with m_0 of a few TeV, which may be within the reach of accelerator experiments in the near future.

More generally, the figures imply that, in the large-cutoff theory, the tree-level relation²⁰ of the GM term (10) in mSUGRA can be satisfied for various parameters with $m_0 \gg |M_i|, |\mu|$, which are consistent with the present observational bounds. Hence we conclude that the GM term works very well with our large-cutoff hypothesis.

¹⁹In contrast, the RG focus point behavior manifests itself under the m_0 independent choice of the renormalization scale Q with the focus point given by $a(Q) = 0$ in Eq. (11) [5].

²⁰This tree-level relation may suffer from possible corrections of order M_G/M_* at the input scale, which, we hope, is to be compared with future experimental results.

V. CONCLUSIONS AND DISCUSSION

Gravity-mediated supersymmetry breaking and primordial inflation are expected to open windows into the Planck-scale physics through observations on superpartner spectra [1] and on temperature fluctuations of cosmic microwave background radiation [13]. In this paper, we have discussed realization of the large-cutoff theory in supergravity, which is possibly selected through inflationary dynamics. This large-cutoff hypothesis implies an mSUGRA spectrum with a large hierarchy between the universal scalar mass and the gaugino masses, $m_0 \gg |M_i|$. Very encouragingly, despite of relatively large masses of scalar particles, the electroweak symmetry breaking can occur at the correct energy scale with $m_0 \gg |\mu|$ in phenomenologically viable parameter regions.

In the large-cutoff hypothesis, the absence of the FCNC process is automatic, since all of the corresponding higher dimensional operators in the Kähler potential are suppressed by the large cutoff M_* . In addition, with the current chargino mass bound, the hierarchical spectrum $m_0 \gg |M_i|, |\mu|$ predicts heavy sfermions at a few TeV, and hence the CP problem in the SSM is ameliorated. In most of the parameter region ($m_0 \gg |M_i|, |\mu|$) we are interested in, the lightest supersymmetric particle is a neutralino, which

is a good candidate for the dark matter (see also the remark at the end of section IVA).

Finally, let us comment on the origin of matters in the universe in the present scenario. The hierarchical spectrum implies that the mass of the gravitino is of the order of a few TeV. In this case, the primordial abundance of the gravitino should be suppressed not to disturb the Big-Bang Nucleosynthesis, which implies the reheating temperature $T_R \lesssim 10^{6-7}$ GeV [28]. Hence, the baryon asymmetry must be provided at the corresponding low temperatures, $T \lesssim 10^{6-7}$ GeV.

ACKNOWLEDGMENTS

The authors wish to thank Y. Sinbara for valuable discussion. This work is partially supported by Grand-in-Aid Scientific Research (s) 14102004.

APPENDIX: NOTATION FOR THE HIGGS POTENTIAL

In this appendix, we list our convention for the Higgs mass parameters. We adopt the following form of the effective Higgs potential at the tree-level:

$$V = (|\mu|^2 + m_{H_u}^2)(|H_u^0|^2 + |H_u^+|^2) + (|\mu|^2 + m_{H_d}^2)(|H_d^0|^2 + |H_d^+|^2) + \{B\mu(H_u^+ H_d^- - H_u^0 H_d^0) + c.c.\} + \frac{1}{2}g^2|H_u^+ H_d^{0*} + H_u^0 H_d^{-*}|^2 + \frac{1}{8}(g^2 + g'^2)(|H_u^0|^2 + |H_u^+|^2 - |H_d^0|^2 - |H_d^+|^2)^2, \quad (\text{A1})$$

where the superscript (0, +, -) of each field denotes its electric charge, and g and g' denote the $SU(2)_W$ and $U(1)_Y$ gauge coupling constants, respectively. With this convention, the RG equations for the μ and B parameters are given at the one-loop level by

$$\frac{d}{d \ln Q} \mu = \frac{\mu}{16\pi^2} (3|y_t|^2 + 3|y_b|^2 + |y_\tau|^2 - 3g^2 - g'^2), \quad (\text{A2})$$

$$\frac{d}{d \ln Q} B = \frac{1}{16\pi^2} (6A_t|y_t|^2 + 6A_b|y_b|^2 + 2A_\tau|y_\tau|^2 + 6g^2 M_2 + 2g'^2 M_1), \quad (\text{A3})$$

where $A_i \cdot y_i$ ($i = t, b, \tau$) denotes the (scalar)³ coupling constant that is the supersymmetric counterpart of the Yukawa coupling y_i for each flavor t, b , or τ .

-
- [1] For reviews, H. P. Nilles, Phys. Rep. **110**, 1 (1984); D. J. H. Chung, L. L. Everett, G. L. Kane, S. F. King, J. Lykken, and L.-T. Wang, hep-ph/0312378.
 - [2] Izawa K.-I. and T. Yanagida, Prog. Theor. Phys. **101**, 171 (1999); **103**, 1077 (2000).
 - [3] R. Barbieri and G. F. Giudice, Nucl. Phys. **B306**, 63 (1988).
 - [4] K. L. Chan, U. Chattopadhyay, and P. Nath, Phys. Rev. D **58**, 096004 (1998).
 - [5] J. L. Feng, K. T. Matchev, and T. Moroi, Phys. Rev. Lett. **84**, 2322 (2000); Phys. Rev. D **61**, 075005 (2000).
 - [6] R. Bousso and J. Polchinski, J. High Energy Phys. **06** (2000) 06.
 - [7] See also L. Smolin, hep-th/0407213, and references therein.
 - [8] Izawa K.-I., Prog. Theor. Phys. **86**, 917 (1991).
 - [9] A. Vilenkin, Phys. Rev. Lett. **74**, 846 (1995).
 - [10] Izawa K.-I. and T. Yanagida, Phys. Lett. B **393**, 331 (1997).
 - [11] Izawa K.-I., Phys. Lett. B **576**, 1 (2003).
 - [12] Izawa K.-I. and T. Yanagida, Prog. Theor. Phys. **95**, 829 (1996); T. Hotta, Izawa K.-I., and T. Yanagida, Phys. Rev.

- D **55**, 415 (1997); Izawa K.-I., Y. Nomura, K. Tobe, and T. Yanagida, Phys. Rev. D **56**, 2886 (1997); Izawa K.-I., Prog. Theor. Phys. **99**, 157 (1998).
- [13] For reviews, D. H. Lyth and A. Riotto, Phys. Rep. **314**, 1 (1999); W. H. Kinney, astro-ph/0301448; S. F. King, Pramana **62**, 307 (2004).
- [14] Izawa K.-I. and T. Yanagida, Prog. Theor. Phys. **94**, 1105 (1995).
- [15] T. Asaka, K. Hamaguchi, M. Kawasaki, and T. Yanagida, Phys. Rev. D **61**, 083512 (2000).
- [16] S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [17] E. Witten and J. Bagger, Phys. Lett. B **115**, 202 (1982); M. K. Gaillard, Phys. Rev. D **58**, 105027 (1998).
- [18] M. K. Gaillard and V. Jain, Phys. Rev. D **49**, 1951 (1994); M. K. Gaillard, Phys. Lett. B **342**, 125 (1995); K. Choi, J. S. Lee, and C. Muñoz, Phys. Rev. Lett. **80**, 3686 (1998).
- [19] Izawa K.-I., M. Kawasaki, and T. Yanagida, Phys. Lett. B **411**, 249 (1997); Izawa K.-I., hep-ph/9710479.
- [20] See M. Dine and D. A. MacIntire, Phys. Rev. D **46**, 2594 (1992).
- [21] G. F. Giudice and A. Masiero, Phys. Lett. B **206**, 480 (1988).
- [22] S. Codoban and D. I. Kazakov, Eur. Phys. J. C **13**, 671 (2000).
- [23] P. Azzi *et al.*, hep-ex/0404010.
- [24] F. E. Paige, S. D. Protopescu, H. Baer, and X. Tata, hep-ph/0312045.
- [25] B. C. Allanach, S. Kraml, and W. Porod, J. High Energy Phys. **03** (2003) 16.
- [26] J. L. Feng, K. T. Matchev, and F. Wilczek, Phys. Lett. B **482**, 388 (2000).
- [27] J. R. Ellis, K. A. Olive, Y. Santoso, and V. C. Spanos, Phys. Rev. D **70**, 055005 (2004).
- [28] For recent developments, M. Kawasaki, K. Kohri, and T. Moroi, astro-ph/0408426.
- [29] G. Abbiendi *et al.*, Phys. Lett. B **565**, 61 (2003).
- [30] LEP2 SUSY Working Group, Combined LEP Chargino Results, up to 208 GeV for large m_0 , <http://lepsusy.web-cern.ch/lepsusy/www/inosmoriond01/charginospub.html>.