

Neutrino helicity asymmetries in leptogenesis

Luís Bento* and Francisco C. Santos†

Universidade de Lisboa, Faculdade de Ciências, Centro de Física Nuclear da Universidade de Lisboa, Avenida Prof. Gama Pinto 2, 1649-003 Lisboa, Portugal

(Received 10 November 2004; published 5 May 2005)

It is pointed out that the heavy singlet neutrinos characteristic of leptogenesis develop asymmetries in the abundances of the two helicity states as a result of the same mechanism that generates asymmetries in the standard lepton sector. Neutrinos and standard leptons interchange asymmetries in collisions with each other. It is shown that an appropriate quantum number, $B - L'$, combining baryon, lepton and neutrino asymmetries, is not violated as fast as the standard $B - L$. This suppresses the washout effects relevant for the derivation of the final baryon asymmetry. One presents detailed calculations for the period of neutrino thermal production in the framework of the singlet seesaw mechanism.

DOI: 10.1103/PhysRevD.71.096001

PACS numbers: 11.30.Fs, 14.60.St, 98.80.Cq

I. INTRODUCTION

Leptogenesis is an attractive way of generating the baryon number of the Universe [1]. The main idea put forward by Fukugita and Yanagida [2–4] is that a lepton number asymmetry can be produced in the decays of heavy singlet neutrinos into leptons and Higgs bosons and such an asymmetry is partially transferred to the baryon sector through electroweak sphaleron processes [5] that violate B and L but not $B - L$. The mechanism requires nonconservation of lepton number and CP provided by neutrino Majorana masses and complex Yukawa couplings. Both masses and couplings form the well known singlet seesaw model [6–10] of light neutrino masses and thus establish a close relationship between baryogenesis and low energy phenomenology [11–26]. This connection contributed for the present wide interest in leptogenesis.

The calculation of the final baryon asymmetry has been done in the literature with increasing levels of accuracy [14,15,21,25–30] but the main elements have remained the following. Singlet neutrino reactions are not symmetric under CP due to nontrivial complex Yukawa couplings in the neutrino mass eigenstate basis. Departure of neutrino densities from thermal equilibrium values are a necessary condition [31] to obtain net lepton asymmetry sources. This occurs if neutrinos are not produced in the inflaton decay but only gradually from active lepton and Higgs boson collisions. It occurs also to some extent when any of the neutrino species undergoes the transition to the respective nonrelativistic temperature epoch. Weak sphaleron processes transform a fraction of the generated $B - L$ asymmetry into baryon number. The final $B - L$ and baryon asymmetries depend on lepton number violating reactions whose net effect is to dissipate $B - L$. They include the reactions $\bar{\phi}\bar{\phi} \rightarrow l_i l_j$, $\bar{\phi}\bar{l}_j \rightarrow \phi l_i$, $\bar{l}_i l_j \rightarrow \phi\phi$, that violate lepton number by two units, but also processes that violate standard lepton number by one unit [3] such as

top quark and electroweak gauge boson scatterings like $\bar{t}q_t \rightarrow N_a l_i$, $\bar{\phi}W \rightarrow N_a l_i$ and $\bar{\phi}B \rightarrow N_a l_i$. Neutrino number densities and the set of standard lepton, quark and Higgs boson number asymmetries obey a system of coupled Boltzmann equations that is necessary to integrate to derive the final (present) $B - L$ asymmetry.

And here comes the main point of this paper. This system is incomplete because, contrary to what has ever been assumed, the two helicity states of singlet Majorana neutrinos do not have exactly the same abundances. We make the case that leptogenesis mechanisms naturally generate asymmetries in the two neutrino helicity state abundances, in the same way as the standard lepton asymmetries, and that the neutrino helicity asymmetries play a role in the system of Boltzmann equations that govern the evolution and transport of particle asymmetries, lepton number, in particular, and therefore contribute to the determination of the final $B - L$ asymmetry as a function of the fundamental parameters of the theory. We present explicit calculations in the framework of the singlet seesaw mechanism.

In next section we review some of the properties of this model that are relevant for leptogenesis in relation with the light neutrino mass spectra [32–34] as indicated by solar [35–41], atmospheric [42–44] and terrestrial neutrino experiments [45,46]. In Sec. III we introduce the concept of neutrino helicity asymmetries and discuss their relevance for the transport of lepton number. An appropriate quantum number $B - L'$ is proposed to replace the usual $B - L$ difference. In Sec. IV we calculate explicitly the lepton and neutrino asymmetries generated during the phase of singlet neutrino thermal production. This was presented in a brief fashion in a meeting [47]. Here we give a complete account of the work and improve the integrations over phase space by including Pauli blocking and lepton thermal mass effects in the numerical calculations. The Higgs boson thermal mass had already been taken into account. In Sec. V we study the washout processes and evaluate the damping rate of $B - L'$. We compare with the traditional treatment without neutrino helicity asymmetries and take

*Electronic address: lbento@cii.fc.ul.pt

†Electronic address: fsantos@cii.fc.ul.pt

the appropriate lessons. The main results are summarized in the final section.

II. SEESAW MODEL

The singlet seesaw mechanism [6] adds to the standard model singlet (left-handed) neutrinos, N_a , with heavy Majorana masses and Yukawa couplings with the standard lepton and Higgs doublets, l_i and ϕ , of the form

$$h_{ia} l_i N_a \phi + \frac{1}{2} M_a N_a N_a + \text{H.C.} \quad (1)$$

Spontaneous breaking of $SU(2) \times U(1)$ yields the light neutrino mass matrix ($v = \langle \phi^0 \rangle$)

$$m_{ij} = -(hM^{-1}h^T)_{ij}v^2. \quad (2)$$

The proper decay rate of N_a into leptons and Higgs is given by

$$\Gamma_{N_a}^0 = \frac{(h^\dagger h)_{aa}}{8\pi} M_a, \quad (3)$$

if one ignores thermal effects. Delayed decay occurs when the ratio to the Hubble expansion rate H at the temperature $T = M_a$,

$$K_a = \frac{\Gamma_{N_a}^0}{H(T=M_a)}, \quad (4)$$

is small. In the radiation era, $H = 1.66g_*^{1/2}T^2/M_P$, where g_* denotes the number of relativistic degrees of freedom, 107.5 in the standard model. It is enough to compare the sum

$$K = \sum K_a = (10^3 \text{ eV}^{-1}) \text{Tr}[hM^{-1}h^\dagger]v^2 \quad (5)$$

with the light neutrino mass scale $\text{Tr}[m]$ to conclude that the delayed decay condition, $K_a < 1$, is in general in conflict with the atmospheric neutrino mass gap [42], $\Delta m^2 \approx 2.5 \cdot 10^{-3} \text{ eV}^2$, which implies $K > 50$. Strictly speaking, the delayed decay scenario only requires that the lightest of the heavy neutrinos satisfies $K_a < 1$. But that is not the most natural picture, in particular, if light neutrinos are quasidegenerate. In Sec. IV we will assume that all parameters K_a are large, of the order of 50 or more, which has the effect that singlet neutrinos enter in thermal equilibrium at relativistic temperatures $T_a \gg M_a$.

III. NEUTRINO HELICITY ASYMMETRIES

Sterile neutrino Yukawa couplings $l_i N_a \phi$ conserve total lepton number, L_T , assigned as $L = -1$ for left-handed neutrino fields, N_a , and $L = +1$ for the right-handed neutrino conjugate fields \bar{N}_a . Neutrino Majorana masses break lepton number but it is clear that in the ultrarelativistic limit the masses are negligible and neutrinos and antineutrinos may have unequal abundances and symmetric

chemical potentials like any other particles. In that case neutrinos carry lepton number that can be exchanged between them and standard leptons in collisions mediated by Yukawa interactions. The Majorana masses do not change this completely. One faces a similar problem at defining lepton number of Majorana mass solar or atmospheric neutrinos, or lepton asymmetries of neutrinos at the time of Big Bang Nucleosynthesis [48]. The main difference is that they deal with standard active neutrinos and not necessarily sterile neutrinos. Free massive neutrino states are solutions of the Dirac equation that can be discriminated as spin eigenstates and, in particular, as helicity eigenstates. The $\pm 1/2$ helicity eigenstates spinors satisfy the relation

$$u\bar{u} = \frac{1}{2} \left(1 \pm \gamma_5 \frac{E - M\gamma^0}{p} \right) (\not{p} + M), \quad (6)$$

where E is the neutrino energy, p the momentum and M the mass. When neutrinos are ultrarelativistic, helicity and chiral states are almost identical and their lepton number is maximal (± 1). For an arbitrary neutrino state the lepton number current density is evaluated as the expectation value

$$J_L^\mu = \langle \bar{N} \gamma^\mu \gamma_5 N \rangle = \frac{1}{2} \langle \bar{\chi} \gamma^\mu \gamma_5 \chi \rangle, \quad (7)$$

where N is either a left-handed chiral field N_a or its conjugate \bar{N}_a , and $\chi = N_a + \bar{N}_a$ is a Majorana field (for the quantization of Majorana mass fields see Refs. [49–51]). As a result a positive (negative) helicity eigenstate carries a well defined average lepton number equal to the neutrino speed $v = p/E$ ($-v$):

$$L = \frac{u^\dagger \gamma_5 u}{u^\dagger u} = \pm v. \quad (8)$$

The lepton number vanishes in the nonrelativistic limit due to the Majorana nature of the neutrino mass.

One may observe that in contrast to lepton number, helicity is not invariant under Lorentz transformations. However, in an isotropic Universe the comoving thermal bath frame is a privileged frame where isotropy enforces the spin density matrix to be diagonal in the helicity basis. That means that each of the neutrino flavors N_a can be divided in two populations of opposite helicities and well defined distribution functions f_a^\pm . The total lepton number carried by each neutrino species is equal to

$$L_a = V \int \frac{d^3 p_a}{(2\pi)^3} (f_a^+ - f_a^-) v_a, \quad (9)$$

where v_a is the neutrino speed and V the spatial volume. It will prove convenient to work instead with the helicity asymmetries i.e., the differences between positive and negative helicity neutrino abundances,

$$\Lambda_a = V \int \frac{d^3 p_a}{(2\pi)^3} (f_a^+ - f_a^-). \quad (10)$$

Sterile neutrinos have been deprived of lepton number in the leptogenesis literature because they have Majorana masses (one exception is the oscillation mechanism of Ref. [52]). But contrary to the lepton number assignment, that is to an extent arbitrary, it is a unambiguous fact that neutrinos develop helicity asymmetries as a result of collisions with standard leptons and Higgs and also directly from the same leptogenesis processes that generate lepton asymmetries. That is shown explicitly in section IV C.

It is important to realize that neutrino helicity asymmetries affect the way lepton number is transported in collision processes and consequently the so-called lepton number washout effect. Take, for example, scatterings like $N_a l_i \leftrightarrow \bar{t} q_t$, $N_a \bar{l}_i \leftrightarrow t \bar{q}_t$, and crossed channels, that violate standard lepton number by one unit (t is the right-handed top quark and q_t the quark isodoublet linear combination that has a Yukawa coupling with it). The correct evaluation of the lepton number violation rate must take into account that the reaction rates depend on the neutrino helicity and the two helicity states have different abundances. Reactions with positive helicity neutrinos in initial or final states like $N_a^+ l_i$ are suppressed with respect to the opposite helicity states $N_a^- l_i$. The former are possible only because neutrinos have Majorana masses and their transition amplitudes are suppressed by the Lorentz contraction factors M_a/E . If neutrinos are in thermal equilibrium their helicity asymmetries can be parametrized with degeneracy parameters $\eta_a = \mu_a/T$ as any other fermions, η_a for positive helicity and $-\eta_a$ for negative helicity states, and the rate of standard lepton number violation contains terms like

$$\begin{aligned} \gamma_{N_a l_i \rightarrow \bar{t} q_t} - \gamma_{\bar{t} q_t \rightarrow N_a l_i} &= \gamma_+(\eta_\phi + \eta_{l_i} + \eta_a) \\ &+ \gamma_-(\eta_\phi + \eta_{l_i} - \eta_a), \end{aligned} \quad (11)$$

where γ_+ , γ_- stand for the average reaction rates of $N_a^+ l_i \leftrightarrow \bar{t} q_t$ and $N_a^- l_i \leftrightarrow t \bar{q}_t$ respectively. We have replaced η_t and η_{q_t} with the Higgs degeneracy parameter using the constraint $\eta_t - \eta_{q_t} = \eta_\phi$ enforced by the rapid top quark Yukawa interactions. Identical expressions apply to other single neutrino absorption and emission reactions, decays and inverse decays $N_a \leftrightarrow l_i \phi$ (at temperatures $T \lesssim M_a$ where the Higgs and lepton thermal masses are small enough), or $\bar{\phi} \leftrightarrow N_a l_i$ (at temperatures $T \gtrsim 2M_a$), and respective radiative processes with one electroweak gauge boson in the initial or final state. Notice that because γ_+ is smaller than γ_- the result depends on the neutrino chemical potentials. The two rates coincide with each other only when neutrinos are at rest. On the other hand one should not expect that the neutrino degeneracy parameters η_a are damped faster by the lepton number violating reactions than the combinations $\eta_\phi + \eta_{l_i}$, also damped by $\Delta L = 2$ collisions like $\phi l_i \leftrightarrow \bar{\phi} \bar{l}_j$.

Contrary to the partial lepton numbers of standard leptons, L_i , and neutrinos, L_a , the total lepton number

$$L_T = \sum_i L_i + \sum_a L_a \quad (12)$$

and the quantum number $B - L_T$ are conserved by neutrino-lepton Yukawa couplings. In addition, $B - L_T$ is conserved by sphalerons and is only violated by neutrino Majorana masses. For practical purposes, namely, application of the constraints imposed by *CPT* invariance and unitarity on the collision rates, it is more convenient to work with the helicity asymmetries Λ_a rather than the lepton numbers L_a because the states with definite distribution functions are helicity eigenstates, not (chiral) lepton number eigenstates. On the other hand, the quantum number

$$L' = \sum_i L_i + \sum_a \Lambda_a \quad (13)$$

has essentially the same interesting properties as the total lepton number L_T . The combination $B - L'$, also conserved by weak sphalerons, reduces to the standard model $B - L$ when the heavy neutrinos vanish from the Universe. It is conserved by neutrino Majorana masses but its violation by Yukawa couplings is suppressed by the neutrino masses i.e., mass over energy ratios.

Neutrino helicity asymmetries play a role in the transport of standard lepton number. We have shown for instance that processes that violate flavor quantum numbers L_i and Λ_a by one unit contribute to the violation rate of $L_i - B/3$ as follows (if all particles are in thermal equilibrium):

$$\begin{aligned} \frac{d(L_i - B/3)}{dt} &= -\gamma_{ia}^{(0)}(\eta_\phi + \eta_{l_i} - \eta_a) \\ &- \gamma_{ia}^{(2)}(\eta_\phi + \eta_{l_i} + \eta_a) + \dots \end{aligned} \quad (14)$$

where $\gamma_{ia}^{(0)}$ is the total rate of L' conserving reactions and $\gamma_{ia}^{(2)}$ the total rate of $\Delta L' = \pm 2$ reactions. In turn, the standard lepton asymmetries are transferred to the neutrino sector in collisions with them. The same processes as above contribute to the violation rates of helicity asymmetries Λ_a as

$$\begin{aligned} \frac{d\Lambda_a}{dt} &= \gamma_{ia}^{(0)}(\eta_\phi + \eta_{l_i} - \eta_a) \\ &- \gamma_{ia}^{(2)}(\eta_\phi + \eta_{l_i} + \eta_a) + \dots \end{aligned} \quad (15)$$

This proves that the neutrino helicity asymmetries cannot be assumed identically zero because the L' conserving reactions are faster than the L' violating ones. It is also clear that only the later contribute to the violation rate of $B - L'$. We will return to this point in Sec. V.

This discussion shows that in order to correctly evaluate the lepton number washout effects neutrino helicity asymmetries are an essential ingredient. They have been so far completely ignored in the leptogenesis literature. The calculation of the lepton number generated during the decay

phase of the lightest neutrino(s) and integration of the Boltzmann equations including neutrino helicity asymmetries is complicated by the fact that the neutrinos are neither purely nonrelativistic nor ultrarelativistic at temperatures close to their masses. In this paper we limit ourselves to the temperature range where neutrinos are ultrarelativistic which permits first order calculations on their mass over temperature ratios. In next section we calculate the $B - L'$ asymmetry generated during neutrino thermal production.

IV. GENERATION OF LEPTON ASYMMETRIES

A. Neutrino thermal production

Let us now examine the leptogenesis processes in detail during neutrino thermal production. We assume that the Universe is initially empty of singlet neutrinos, not produced in the inflaton decay but only thermally from standard leptons and Higgs. The dominant thermalizing reactions are identical to the ones of the charged lepton isosinglets [53], namely, top quark scattering processes like $q_i \bar{l}_R \rightarrow l_i N_a$, Higgs boson decay $\phi \rightarrow l_i N_a$ and related scattering processes with one additional electroweak gauge boson in the initial or final state. Higgs boson decays into leptons and neutrinos, first considered in Ref. [47] in the context of leptogenesis, are allowed as much as the decays into a lepton isodoublet and a charged lepton isosinglet, $\bar{\phi} \rightarrow l_i e_i$, because the Higgs has a significant thermal mass, $m_\phi = x_\phi T \sim 0.6T$, larger than the lepton thermal masses [54–56] as pointed out in Ref. [53].

Let n_a denote the average neutrino number densities per helicity degree of freedom and $Y_a = n_a V$ the abundances in a fixed comoving volume whose spatial volume V expands as $\dot{V} = 3HV$. For definiteness we assume that the reheating temperature is much higher than the neutrino masses so that neutrinos thermalize while they are ultrarelativistic. Then, the equilibrium densities and abundances are equal to $n_{\text{eq}} = 0.90T^3/\pi^2$ and $Y_{\text{eq}} = n_{\text{eq}}V$, respectively. Assuming for simplicity that the distribution functions scale with the equilibrium distribution functions f_a^{eq} as

$$f_a = \frac{n_a}{n_{\text{eq}}} f_a^{\text{eq}}, \quad (16)$$

and approximating the Pauli blocking factors $1 - f_a$ with $1 - f_a^{\text{eq}}$ in neutrino emission reactions, the neutrino abundances evolve as

$$\dot{Y}_a = \Gamma_a (Y_{\text{eq}} - Y_a). \quad (17)$$

Here we neglect reactions with two or more neutrino states that are of higher order in the Yukawa couplings.

Γ_a represent the neutrino collision frequencies. At temperatures much higher than the neutrino masses Γ_a scale with the temperature as follows:

$$\Gamma_a = \frac{\beta}{8\pi} (h^\dagger h)_{aa} T. \quad (18)$$

They are comparable with the $\bar{\phi} \rightarrow l_i N_a$ proper decay rate, equal to $\frac{1}{16\pi} (h^\dagger h)_{aa} m_\phi$ if one ignores thermal effects other than the Higgs boson mass. The coefficient β gets contributions from the Higgs decay, top quark and W, B gauge boson scatterings in analogy with the charged leptons case [25,53]. The relative weights of these reactions depend on the temperature because the couplings constants [57] and thermal mass factors [54–56,58] run with the energy scale [25]. Using $m_\phi = 0.6T$ for the Higgs mass and 10^9 GeV temperature scale coupling constants, namely, $\alpha_s \approx 1/26$, $\alpha_w \approx 1/38$, $\alpha' \approx 1/81$ for the strong, SU(2) and U(1)_Y gauge interactions, respectively, and $\lambda_t \approx 0.60$ for the top Yukawa coupling, one obtains $\beta \approx 1/7$, where 46% comes from the Higgs boson decay, 41% from W and B electroweak gauge boson scatterings, and 13% from right-handed top quark scatterings.

In the radiation era the Hubble expansion rate scales as $H = 1/2t \propto T^2$ and the assumed initially zero neutrino densities n_a converge exponentially to the equilibrium densities:

$$n_a = n_{\text{eq}} (1 - e^{-T_a/T}). \quad (19)$$

The relaxation temperatures T_a are given as

$$T_a = \frac{\Gamma_a T}{H} = \beta K_a M_a, \quad (20)$$

and the second identity establishes a relation with the parameters K_a of Eq. (3) that control the speed of neutrino decays when they become nonrelativistic and vanish from the Universe. The low energy neutrino data indicates that the parameters $K_a \gtrsim 50$ are large, see Eq. (5). As a result singlet neutrinos reach thermal equilibrium at temperatures $T_a \gtrsim 10M_a$ when they are still ultrarelativistic.

We have defined neutrino densities in the basis of Majorana mass eigenstates as usual. However, this is not valid for all temperature scales because neutrinos also get thermal masses from the interactions with the lepton-Higgs thermal bath. The chiral conserving thermal mass terms are [54,55]

$$m_{ab}^2 = \frac{1}{8} (h^\dagger h)_{ab} T^2. \quad (21)$$

At high enough temperatures the thermal masses dominate over the vacuum masses M_a and the neutrino Hamiltonian eigenstates are eigenstates of the matrix $(h^\dagger h)_{ab}$. The opposite happens when M_a are much larger than the thermal masses, and at temperatures where the two types of masses are comparable with each other the neutrinos undergo strong flavor oscillation processes. One can estimate the thermal over vacuum mass ratios using Eqs. (4) and (20):

$$\frac{m_{aa}^2}{M_a^2} = \pi K_a \frac{H}{M_a} \approx \frac{K_a^3 M_a}{10^{19} \text{ GeV}} \frac{T^2}{T_a^2}. \quad (22)$$

These have to be small at the relaxation temperatures T_a in order that the neutrino densities evolve as described in this section. For decay constants K_a of the order of 50 (100) this happens for neutrino masses M_a below 10^{12} (10^{11}) GeV. Then, the thermal masses can be neglected provided that the vacuum masses are not degenerate.

There is another point. The neutrino states produced in collisions are linear combinations of the mass eigenstates N_a of the form $h_{ia} N_a$. In a free path the mass eigenstate wave functions oscillate with unequal frequencies, $\sqrt{\mathbf{p}^2 + M_a^2}$, for the same linear momentum \mathbf{p} . They differ by $(M_a^2 - M_b^2)/2E$ for relativistic particles (energy $E \approx p$) and in a mean free path $1/\Gamma_a$ give rise to average phase differences equal to

$$\frac{M_a^2 - M_b^2}{2\Gamma_a} \langle E^{-1} \rangle \approx 5 \frac{M_a^2 - M_b^2}{m_{aa}^2}, \quad (23)$$

where we made use of Eqs. (20) and (22) with $\beta \approx 1/7$. This shows that as long as thermal masses are much smaller than the vacuum mass gaps the neutrino mass eigenstates develop large phase decoherence between collisions so that one can consider that the neutrino states with definite number densities are the vacuum mass eigenstates N_a . This is important because if, for example, the neutrino densities n_a were all equal to each other the leptogenesis sources would vanish in the ultrarelativistic regime at lowest order, as the results of next sections show.

B. Standard lepton asymmetries

Leptogenesis is dominated by the following CP asymmetric reactions [47]: Higgs decays into leptons and singlet neutrinos, inverse decays and scatterings of leptons off neutrinos. The CP asymmetries result from the diagrams of Fig. 1, more specifically, from the interference between




FIG. 1. Diagrams contributing to CP -asymmetries in decays, inverse decays and scatterings.

the tree level amplitude and the absorptive part of the one-loop amplitude of the Higgs boson decay (inverse decay) and from the interference between the t channel amplitude and the absorptive part of the s channel amplitude of neutrino-lepton scattering. The particle asymmetry sources vanish in thermal equilibrium but, as long as neutrinos stay rarefied scatterings and inverse decays do not match Higgs decays and particle asymmetries develop in the various lepton flavors, singlet neutrinos, and Higgs boson as well as enforced by hypercharge conservation.

The source terms (labeled with S) responsible for the generation of the standard family lepton numbers L_i are given in leading order by

$$(\dot{L}_i)_S = \sum_b \Delta\gamma(\bar{\phi} \rightarrow N_b l_i)_S - \Delta\gamma(N_b l_i \rightarrow \bar{\phi})_S + \sum_{abj} \Delta\gamma(N_a l_j \rightarrow N_b l_i)_S - \Delta\gamma(N_b l_i \rightarrow N_a l_j)_S, \quad (24)$$

where

$$\Delta\gamma(X \rightarrow Y) = \gamma(X \rightarrow Y) - \gamma(\bar{X} \rightarrow \bar{Y}) \quad (25)$$

denotes the difference between the rate of an arbitrary reaction $X \rightarrow Y$ integrated over a fixed comoving volume and the rate of the conjugate reaction $\bar{X} \rightarrow \bar{Y}$, where \bar{X} and \bar{Y} are CPT conjugate states of X and Y respectively (CPT transforms particles into antiparticles, reverses helicity but not momentum). The difference $\Delta\gamma(X \rightarrow Y)$ can be separated in two types of contributions, the transport terms proportional to particle antiparticle abundance asymmetries and the source terms responsible for primordial asymmetry generation. By definition the source terms exist in absence of particle asymmetries and in the $\Delta\gamma(X \rightarrow Y)_S$ expressions we assume that CPT conjugate states have exactly the same distribution functions.

CPT invariance and unitarity ensure that the total decay rates of any quantum state and its CPT conjugate state are equal to each other. This translates in constraints on the rate asymmetries:

$$\sum_Y \Delta\gamma(X \rightarrow Y)_S = 0. \quad (26)$$

In leading order, one gets:

$$\Delta\gamma(N_b l_i \rightarrow \bar{\phi})_S + \sum_{aj} \Delta\gamma(N_b l_i \rightarrow N_a l_j)_S = 0, \quad (27)$$

$$\sum_{bi} \Delta\gamma(\bar{\phi} \rightarrow N_b l_i)_S = 0. \quad (28)$$

Then, the L_i source terms of Eq. (24) can be written as,

$$(\dot{L}_i)_S = \sum_b \Delta\gamma(\bar{\phi} \rightarrow N_b l_i)_S + \sum_{abj} \Delta\gamma(N_a l_j \rightarrow N_b l_i)_S. \quad (29)$$

On the other hand, CPT invariance implies that any reaction $X \rightarrow Y$ has the same transition probability as $\bar{Y} \rightarrow \bar{X}$ where the bars indicate CPT conjugate states. As a result

the rate asymmetries $\Delta\gamma(X \rightarrow Y)$ obey the following constraints in thermal equilibrium assuming identical particle and antiparticle distribution functions (zero chemical potentials):

$$\Delta\gamma(X \rightarrow Y)_{\text{eq}} + \Delta\gamma(Y \rightarrow X)_{\text{eq}} = 0. \quad (30)$$

Here one employs the identity [59,60] satisfied by the thermal distribution functions of particles involved in two inverse reactions $X \rightarrow Y$ and $Y \rightarrow X$, schematically, $f_X(1 \pm f_Y) = f_Y(1 \pm f_X)$, where f_X (f_Y) stands for the product of initial state particle distribution functions and $1 \pm f_Y$ ($1 \pm f_X$) for the product of final state stimulated emission and/or Pauli blocking factors. Applying the above constraint on Eqs. (27) and (28) one obtains

$$\Delta\gamma(\bar{\phi} \rightarrow N_b l_i)_{\text{eq}} + \sum_{aj} \Delta\gamma(N_a l_j \rightarrow N_b l_i)_{\text{eq}} = 0, \quad (31)$$

$$\sum_{bi} \Delta\gamma(N_b l_i \rightarrow \bar{\phi})_{\text{eq}} = \sum_{abij} \Delta\gamma(N_a l_j \rightarrow N_b l_i)_{\text{eq}} = 0. \quad (32)$$

This ensures that Eq. (29) satisfies the well known property that the asymmetry source terms vanish in thermal equilibrium [59,60].

The diagrams of Fig. 1 lead to the following results:

$$\begin{aligned} \Delta\gamma(\bar{\phi} \rightarrow N_b l_i)_S &= \sum_{aj} J_{ijab} \int d\Phi f_\phi F_i F_j F_a F_b \\ &\times \frac{-4p'_i \cdot p'_j}{(p_a - p_i)^2 - m_\phi^2}, \end{aligned} \quad (33)$$

$$\begin{aligned} \Delta\gamma(N_a l_j \rightarrow N_b l_i)_S &= J_{ijab} \int d\Phi f_a f_j F_\phi F_i F_b \\ &\times \frac{4p'_i \cdot p'_j}{(p_a - p_i)^2 - m_\phi^2}, \end{aligned} \quad (34)$$

where f_α are particle distribution functions, F_α are fermion Pauli blocking factors $1 - f_\alpha$ or the stimulated emission factor $F_\phi = 1 + f_\phi$ and

$$\begin{aligned} d\Phi &= V \prod_\alpha \frac{d^3 p_\alpha}{(2\pi)^3 2E_\alpha} (2\pi)^8 \delta(p_a + p_j - p_\phi) \\ &\times \delta(p_b + p_i - p_\phi) \end{aligned} \quad (35)$$

is the phase space element running over all particles ϕ , l_i , l_j , N_a , N_b (V is the spatial volume). Standard leptons and Higgs boson are assumed to be in thermal equilibrium while singlet neutrinos have densities given by Eq. (19) and distribution functions proportional to the thermal distribution functions as in Eq. (16). The chemical potentials are equal to zero. The factors

$$J_{ijab} = \text{Im}\{h_{ia} h_{ja} h_{ib}^* h_{jb}^*\} M_a M_b \quad (36)$$

signal the necessary CP and lepton number violation

through Yukawa couplings and Majorana masses. J_{ijab} are antisymmetric under neutrino flavor exchange $a \leftrightarrow b$ and symmetric under lepton flavor exchange $i \leftrightarrow j$. The internal product $p'_i \cdot p'_j = p_i p_j - \mathbf{p}_i \cdot \mathbf{p}_j$ in the integrand functions refers to pseudo 4-vectors p'_i . The p'_i space components coincide with the 3-vector linear momentum \mathbf{p}_i and its time component is equal to the momentum absolute value, $p_i^0 = p_i$. The 4-vectors p'_i result from the spinor wave functions in a thermal bath [55]: $\sum u \bar{u} = \not{p}'_i$. In contrast, the time component of the 4-momentum p_i is the lepton energy, related with the lepton thermal mass as usual: $p_i^0 = (p_i^2 + m_i^2)^{1/2}$.

The generation rate of total standard lepton number $L = \sum L_i$ is obtained from Eq. (29). The constraints of Eqs. (27) and (28) indicate that only scatterings, or better, inverse decays, contribute as L sources:

$$(\dot{L})_S = \sum_{abij} \Delta\gamma(N_a l_j \rightarrow N_b l_i)_S = - \sum_{aj} \Delta\gamma(N_a l_j \rightarrow \bar{\phi})_S. \quad (37)$$

We evaluate the asymmetries including Pauli blocking effects and nonzero lepton thermal masses. However, we neglect the neutrino masses inside the integrals because we are calculating the leading order contributions at relativistic temperatures $T \gg M_a$. Neutrino masses appear in the constant factors J_{ijab} . All the dependence on the neutrino flavors goes in J_{ijab} and in the neutrino number densities n_a that are functions of the temperature and flavor dependent relaxation temperatures T_a . Notice that the following identity holds for the distribution functions in Eq. (34)

$$f_a f_j (1 + f_\phi) = \frac{n_a}{n_{\text{eq}}} f_\phi (1 - f_a^{\text{eq}}) (1 - f_j). \quad (38)$$

Under these conditions the generation rate of standard total lepton number is

$$(\dot{L})_S = - \frac{c}{(8\pi)^4} T^2 V \sum_{abij} \frac{n_a - n_{\text{eq}}}{n_{\text{eq}}} J_{ijab}, \quad (39)$$

with $c \sim 3$ ($c = 2.6$ for a Higgs thermal mass $m_\phi = 0.6T$ and lepton thermal square masses $m_l^2 = 0.036T^2$). It is clear that leptogenesis ceases when neutrinos reach thermal equilibrium abundances.

C. Neutrino asymmetries

In the above equations one sums over both N_a neutrino helicities. But as emphasized in this paper the leptogenesis processes generate also asymmetries in the abundances of the two helicity states N_a^+ and N_a^- . The helicity asymmetries are denoted as Λ_a and are defined with respect to the cosmological comoving frame. In the following the rate asymmetries like $\Delta\gamma(X \rightarrow N_a^+ Y)$ denote the difference between the rates of the CPT conjugate reactions $X \rightarrow N_a^+ Y$ and $\bar{X} \rightarrow N_a^- \bar{Y}$. Wherever the neutrino helicity does not appear explicitly a sum over helicities is assumed. The

leading order source terms are

$$\begin{aligned}
 (\dot{\Lambda}_b)_S = & \sum_i \{ \Delta\gamma(\bar{\phi} \rightarrow N_b^+ l_i) - \Delta\gamma(N_b^+ l_i \rightarrow \bar{\phi}) \\
 & + \Delta\gamma(\phi \rightarrow N_b^+ \bar{l}_i) - \Delta\gamma(N_b^+ \bar{l}_i \rightarrow \phi) \}_S \\
 & + \sum_{aij} \{ \Delta\gamma(N_a l_j \rightarrow N_b^+ l_i) - \Delta\gamma(N_b^+ l_i \rightarrow N_a l_j) \\
 & + \Delta\gamma(N_a \bar{l}_j \rightarrow N_b^+ \bar{l}_i) - \Delta\gamma(N_b^+ \bar{l}_i \rightarrow N_a \bar{l}_j) \}_S. \quad (40)
 \end{aligned}$$

When $a = b$ in the second summation one gets the correct factor of 2 for the helicity flip reactions $N_b^{\mp} \rightarrow N_b^{\pm}$.

The Λ_b source terms are subject to CPT invariance and unitarity conditions namely

$$\Delta\gamma(N_b^+ l_i \rightarrow \bar{\phi})_S + \sum_{aj} \Delta\gamma(N_b^+ l_i \rightarrow N_a l_j)_S = 0, \quad (41a)$$

$$\Delta\gamma(N_b^+ \bar{l}_i \rightarrow \phi)_S + \sum_{aj} \Delta\gamma(N_b^+ \bar{l}_i \rightarrow N_a \bar{l}_j)_S = 0. \quad (41b)$$

These constraints eliminate four of the source terms contained in Eq. (40):

$$\begin{aligned}
 (\dot{\Lambda}_b)_S = & \sum_i \Delta\gamma(\bar{\phi} \rightarrow N_b^+ l_i)_S + \Delta\gamma(\phi \rightarrow N_b^+ \bar{l}_i)_S \\
 & + \sum_{aij} \Delta\gamma(N_a l_j \rightarrow N_b^+ l_i)_S + \Delta\gamma(N_a \bar{l}_j \rightarrow N_b^+ \bar{l}_i)_S. \quad (42)
 \end{aligned}$$

From the CPT invariance condition under thermal equilibrium and zero chemical potentials, Eq. (30), one derives also that

$$\Delta\gamma(\bar{\phi} \rightarrow N_b^+ l_i)_{\text{eq}} + \sum_{aj} \Delta\gamma(N_a l_j \rightarrow N_b^+ l_i)_{\text{eq}} = 0, \quad (43a)$$

$$\Delta\gamma(\phi \rightarrow N_b^+ \bar{l}_i)_{\text{eq}} + \sum_{aj} \Delta\gamma(N_a \bar{l}_j \rightarrow N_b^+ \bar{l}_i)_{\text{eq}} = 0, \quad (43b)$$

which ensures that the Λ_b source terms vanish in thermal equilibrium.

The rate asymmetries in Eq. (29) contain by definition sums over neutrino helicity states and therefore relate to the ones of Eqs. (42) as

$$\begin{aligned}
 \Delta\gamma(\bar{\phi} \rightarrow N_b l_i) = & \Delta\gamma(\bar{\phi} \rightarrow N_b^+ l_i) \\
 & - \Delta\gamma(\phi \rightarrow N_b^+ \bar{l}_i), \quad (44a)
 \end{aligned}$$

$$\begin{aligned}
 \Delta\gamma(N_a l_j \rightarrow N_b l_i) = & \Delta\gamma(N_a l_j \rightarrow N_b^+ l_i) \\
 & - \Delta\gamma(N_a \bar{l}_j \rightarrow N_b^+ \bar{l}_i). \quad (44b)
 \end{aligned}$$

Replacing this in Eqs. (29) and (42) one obtains the leading source terms of the total 'lepton number' $L' = \sum L_i + \sum \Lambda_a$:

$$(\dot{L}')_S = 2 \sum_{bi} \Delta\gamma(\bar{\phi} \rightarrow N_b^+ l_i)_S + 2 \sum_{abij} \Delta\gamma(N_a l_j \rightarrow N_b^+ l_i)_S. \quad (45)$$

The asymmetries are calculated from the diagrams of Fig. 1. The N_b positive (negative) helicity states are speci-

fied with spinors satisfying Eq. (6). The results are the following, using the same notations as in Eqs. (33)–(35):

$$\begin{aligned}
 \Delta\gamma(\bar{\phi} \rightarrow N_b^+ l_i)_S = & \sum_{aj} J_{ijab} \int d\Phi f_{\phi} F_i F_j F_a F_b \\
 & \times \frac{-2\tau}{(p_a - p_i)^2 - m_{\phi}^2}, \quad (46)
 \end{aligned}$$

$$\begin{aligned}
 \Delta\gamma(N_a l_j \rightarrow N_b^+ l_i)_S = & J_{ijab} \int d\Phi f_a f_j F_{\phi} F_i F_b \\
 & \times \frac{2\tau}{(p_a - p_i)^2 - m_{\phi}^2}, \quad (47)
 \end{aligned}$$

with

$$\tau = p'_i \cdot p'_j + \frac{p_i}{p_b} p_b \cdot p'_j - \frac{p_j}{p_b} p_b \cdot p'_i. \quad (48)$$

Notice that the N_b helicity states as well as the absolute 3-momenta p_i, p_j, p_b in the expressions above are defined with respect to the cosmological comoving frame. As before we neglect the neutrino masses inside the phase space integrals. The integrand function τ is finite however, if one neglects lepton thermal masses and Pauli blocking factors the integration over the fermion angular variables yields a null result (recall that the distribution function factor in Eq. (47) obeys the relation (38)). This unexpected result means that in that approximation L' is not generated at all i.e., the total asymmetry, $\Lambda = \sum \Lambda_b$, generated in the neutrino sector cancels exactly the total lepton number generated in the standard lepton sector. But for nonzero lepton thermal masses the result is finite:

$$(\dot{L}')_S = \frac{c}{(8\pi)^4} T^2 V \sum_{abij} \frac{n_a - n_{\text{eq}}}{n_{\text{eq}}} J_{ijab}, \quad (49)$$

with $c \sim 1$ ($c = 1.2$ for Higgs thermal mass $m_{\phi} = 0.6T$ and lepton thermal square masses $m_l^2 = 0.036T^2$).

The comparison with Eq. (39) shows that leptogenesis generates a total neutrino helicity asymmetry Λ of opposite sign and larger than the standard lepton asymmetry L so that the source of $L' = L + \Lambda$ is of opposite sign to L . The neutrino and lepton asymmetries are not separately conserved. They are both violated by neutrino Yukawa couplings and the lepton number L is violated by weak sphalerons as well. These interactions cause the interchange of $B - L'$ between the singlet neutrino and standard lepton and quark sectors but conserve $B - L'$ in absence of neutrino Majorana masses. The neutrino masses are so responsible for the $B - L'$ dissipation. In next section we study this effect and establish the evolution of $B - L'$.

V. WASHOUT PROCESSES

Weak sphalerons and neutrino Yukawa couplings are the only ones that violate the standard partial lepton numbers

L_i . The processes like $\bar{\phi} \rightarrow N_a l_i$, $\bar{t} q_t \rightarrow N_a l_i$ violate the standard total lepton number $L = \sum L_i$ by one unit, but this does not mean that they dissipate the total lepton number that is generated in leptogenesis. Indeed, one has to distinguish between the two neutrino helicity states N_a^\pm . The reactions $\bar{\phi} \rightarrow N_a^- l_i$, $\bar{t} q_t \rightarrow N_a^- l_i$, $\bar{\phi} A \rightarrow N_a^- l_i$ (A is a gauge boson) $N_a^+ \bar{\phi} \rightarrow l_i A$, $N_a^+ A \rightarrow l_i \phi$, $N_a^+ \bar{t} \rightarrow \bar{q}_t l_i$, $N_a^+ q_t \rightarrow t l_i$ violate L by one unit but conserve the total 'lepton' number $L' = L + \Lambda$ because they decrease the total helicity $\Lambda = \sum \Lambda_a$ by one unit. On the contrary, the same processes with opposite helicity neutrino states violate L' by two units. They require nonzero neutrino Majorana masses and are suppressed at relativistic temperatures $T \gg M_a$. The L' conserving processes dominate the neutrino thermalization and enter in equilibrium with relaxation temperatures T_a given by Eq. (20). Below these temperatures the L' conserving processes are fast in comparison with the expansion rate and thus enforce constraints on the lepton and neutrino chemical potentials:

$$\eta_\phi + \eta_i - \eta_a = 0. \quad (50)$$

This implies that due to rapid flavor violation all lepton doublets l_i have equal degeneracy parameters $\eta_i = \eta_l$ and all positive (negative) helicity neutrinos N_a^+ (N_a^-) have equal degeneracy parameters $\eta_a = \eta_\phi + \eta_l$ ($-\eta_a$). As a result the partial asymmetries L_i and Λ_a are not directly related with the respective leptogenesis sources but are rather determined from the overall $B - L'$ asymmetry by the set of chemical potential constraints.

There are as much constraints as independent fast flavor changing reactions and this depends on the temperature scale [61,62]. In the standard model case the latest interactions to enter in equilibrium are the Yukawa couplings and their equilibrium temperatures depend on the particular right-handed quark or lepton isosinglet (for a detailed discussion see refs. [47,63]). In any case the constraints leave $B - L'$ as a free variable and determine the other quantum numbers, in particular L_i and Λ_a , as proportional to $B - L'$. In turn, $B - L'$ is completely determined by the leptogenesis sources on one hand and the dissipation (washout) processes on the other hand.

$B - L'$ is violated by two units in the $\Delta L = 1$, $\Delta \Lambda = 1$ reactions $\bar{\phi} \rightarrow N_a^+ l_i$, $\bar{t} q_t \rightarrow N_a^+ l_i$, $\bar{\phi} A \rightarrow N_a^+ l_i$, and crossed channels $N_a^- q_t \rightarrow t l_i$, ..., in the $\Delta L = 2$, $\Delta \Lambda = 0$ scatterings $\bar{\phi} \bar{\phi} \rightarrow l_i l_j$, $\bar{\phi} \bar{l}_j \rightarrow \phi l_i$, $\bar{l}_i \bar{l}_j \rightarrow \phi \phi$, and in the $\Delta L = 0$, $\Delta \Lambda = 2$ scatterings $l_i \bar{l}_j$, $\phi \bar{\phi} \rightarrow N_a^+ N_b^+$, $\phi N_b^- \rightarrow \phi N_a^+$, $l_j N_b^- \rightarrow l_i N_a^+$, $\bar{l}_j N_b^- \rightarrow \bar{l}_i N_a^+$, $N_a^- N_b^- \rightarrow l_i \bar{l}_j$, $\phi \bar{\phi}$. All these processes depend on the existence of neutrino Majorana masses. In the $\Delta L = 2$ case they contribute through the neutrino propagators and in the $\Delta L = 0$, 1 reactions they make possible that a neutrino be produced or annihilated with the 'wrong' helicity state i.e., helicity opposite to the chirality determined by the Yukawa couplings $l_i N_a \phi$. The left-handed (right-handed) chiral projection of a positive (negative) helicity state with energy E

goes as $M_a/2E$ in the relativistic limit. Thus, the L' violating processes are suppressed with respect to the respective L' conserving channels by a ratio going as $M_a^2/4E^2$. The thermal average of this ratio is about $M_a^2/16T^2$ for scatterings but in the case of $\bar{\phi} \rightarrow N_a l_i$ decays the center of mass energy, $m_\phi \sim 0.6T$, and average neutrino energy are much smaller. The $\bar{\phi} \rightarrow N_a^+ l_i$ branching fraction is given by

$$B_a^+ = \frac{\gamma(\bar{\phi} \rightarrow N_a^+ l_i)}{\gamma(\bar{\phi} \rightarrow N_a l_i)} \approx \frac{8M_a^2}{T^2}, \quad (51)$$

2 orders of magnitude higher than in the case of scatterings. As the $\Delta L = 1$ scatterings have total rates comparable with the Higgs decays, this means that the Higgs decays dominate the L' violating rates by 2 orders of magnitude over the scatterings. As far as $\Delta L = 0$ and $\Delta L = 2$ scatterings is concerned, $\phi \bar{\phi} \rightarrow N_a^+ N_b^+$, ..., $\bar{\phi} \bar{\phi} \rightarrow l_i l_j$, ..., they are also suppressed by the same 10^{-2} factor plus an extra factor of $|h_{ia}|^2$ with respect to the $\Delta L' = 2$ Higgs decays, because they get one more Yukawa coupling than the $\Delta L = 1$ scatterings.

From the discussion above one learns that the Higgs decays and inverse decays dominate the violation of $B - L'$. Hence, the evolution of $B - L'$ is well described by the equation

$$\dot{B} - \dot{L}' = -(\dot{L}')_S + 2 \sum_{ai} \{ \Delta \gamma(N_a^+ l_i \rightarrow \bar{\phi})_0 - \Delta \gamma(\bar{\phi} \rightarrow N_a^+ l_i)_0 \}, \quad (52)$$

where the label '0' indicates that the rates are calculated at tree level contrary to the leptogenesis source term $(\dot{L}')_S$ given in Eq. (45). At temperatures below T_a neutrinos have thermal equilibrium abundances and the transport terms depend on the chemical degeneracy parameters as follows:

$$\dot{B} - \dot{L}' = -(\dot{L}')_S + 4 \sum_{ai} \gamma_{\bar{\phi} \rightarrow N_a^+ l_i} (\eta_\phi + \eta_i + \eta_a). \quad (53)$$

Moreover, the L' conserving neutrino reactions are in equilibrium under T_a and the constraint of Eq. (50) applies, so that $\eta_i = \eta_l$ and $\eta_a = \eta_\phi + \eta_l$. The degeneracy parameters parametrize the particle antiparticle number asymmetries. For example, for a ultrarelativistic fermion f the asymmetry is given in leading order by $Y_f - Y_{\bar{f}} = 1.83 \eta_f Y_{\text{eq}}$ [60,64]. The constraints imposed by the standard model interactions leave one degeneracy parameter as free parameter [61,62]. In our case they lead to a relation of the form

$$B - L' = -r(\eta_\phi + \eta_l) Y_{\text{eq}}. \quad (54)$$

The precise value of the factor r depends on the set of chemical constraints and therefore on the temperature scale. It varies from $r = 12.8$ at temperatures above 10^{12} GeV, where the Yukawa couplings of the isosinglet right-handed b_R quark and τ_R lepton are not yet in equi-

librium, to $r = 18.6$ under 10^4 GeV, where all quark and lepton Yukawa couplings are in equilibrium.

The $\bar{\phi} \rightarrow N_a^+ l_i$ partial decay rate is a fraction of the $\bar{\phi} \rightarrow N_a l_i$ decay rate as indicated in Eq. (51). On the other hand, the Higgs decay rates scale with the first power of the temperature and can be related in the same way as in Eq. (20) with the Hubble expansion rate H :

$$\sum_i \gamma_{\bar{\phi} \rightarrow N_a l_i} = \frac{T'_a}{T} H Y_{\text{eq}}. \quad (55)$$

The temperatures T'_a are about one half smaller than the relaxation temperatures T_a because unlike the later T'_a do not receive contributions from scattering processes. Combining everything one rewrites Eq. (53) in the form,

$$\frac{d(B - L')}{dT^{-3}} = -\left(\frac{dT'}{dT^{-3}}\right)_s - T_\star^3 (B - L'), \quad (56)$$

$$T_\star^3 = \frac{8}{3r} \sum_a B_a^+ T_a^2 T'_a \sim \frac{2}{3} \sum_a M_a^2 T_a. \quad (57)$$

This shows that when leptogenesis processes are ineffective, at temperatures $T < T_a$, $B - L'$ decays exponentially with T^{-3} and damping constant equal to T_\star^3 . The same happens with the neutrino degeneracy parameters $\eta_a = \eta_\phi + \eta_l$, as determined by Eq. (54).

It is important to recognize the difference between T_\star and the relaxation temperatures T_a . At T_a neutrino reactions enter in thermal equilibrium and the lepton and neutrino partial quantum numbers $L_i - B/3$ and Λ_a begin to be rapidly violated. However, it does not mean that the standard lepton number L or $B - L'$ start to be washed out at T_a . It only means that the effective couplings $l_i N_a^- \phi$ enter in equilibrium and the constraints $\eta_\phi + \eta_i - \eta_a = 0$ are enforced. The complete set of chemical equilibrium constraints force $B - L'$ to distribute into B , $-L$ and $-\Lambda$ in similar proportions. On the other hand the effective couplings $l_i N_a^+ \phi$ are not yet in equilibrium because the temperatures T_a are much larger than the neutrino Majorana masses. These couplings enter in equilibrium later at the temperature T_\star . Below that temperature the constraints $\eta_\phi + \eta_i + \eta_a = 0$ should apply on top of the previous constraints $\eta_\phi + \eta_i - \eta_a = 0$. It means that the degeneracy parameters η_a , $\eta_\phi + \eta_i$, and $B - L'$ as well as all asymmetries proportional to $B - L'$ are strongly damped below the temperature T_\star . But not above T_\star . In fact, $B - L'$ is only marginally damped at the T_a temperature scale.

For decay constants K_a as large as 70, neutrinos enter in equilibrium at temperatures $T_a \approx \frac{1}{7} K_a M_a \sim 10 M_a$ but the $B - L'$ damping constant T_\star^3 is 2 orders of magnitude smaller than the T_a^3 scale: $T_\star^3 \sim \frac{1}{150} \sum_a T_a^3$. Moreover, the greater the decay constants K_a are the smaller is T_\star^3 in comparison with T_a^3 . On the other hand T_\star is quite close to the neutrino mass scale.

This gives an important lesson. If one ignores neutrino helicity asymmetries, as has ever been done in the literature, then one obtains that the temperatures T_a set the chemical potential constraints $\eta_i + \eta_\phi = 0$ and define the $B - L$ relaxation temperature scale. This is wrong. The correct constraints must include the neutrino degeneracy parameters η_a and are given by Eq. (50). Ignoring neutrino helicity asymmetries leads to an overestimate by 2 orders of magnitude of the damping rate of the lepton and quark asymmetries.

So far we have considered that singlet neutrinos have equilibrium temperatures $T_a \gtrsim 10 M_a$ larger than any of their masses M_a . Another scenario that has been often considered for simplicity is the hierarchical scenario [1–3,15,21,25,28] where one of the singlet neutrinos, N_1 , is much lighter than the others. Then, it is argued that any asymmetries generated at the heaviest neutrinos decaying phases are later washed out by the lightest neutrino $\Delta L = 1$ reactions before the temperature reaches the mass M_1 . As we have just shown this is not correct because it does not take into account the N_1 helicity asymmetry. One concludes from Eq. (57) that the quantum number $B - L'$ and all particle asymmetries related to it by chemical equilibrium constraints are damped with a relaxation temperature given by $T_\star^3 \approx \frac{2}{3} M_1^2 T_1$ (in cases where the $\Delta L = 2$ reactions mediated by off-shell neutrinos are not significant). T_\star is quite close to the mass M_1 which means that the asymmetries left after the heavy neutrino decays remain conserved at temperatures above the lightest neutrino mass.

Another possibility is that the lightest neutrino enters in equilibrium at a relaxation temperature T_1 below the other neutrinos masses. Then, the helicity asymmetry Λ_1 remains conserved when the heaviest neutrinos decay even if the standard lepton asymmetries are completely washed out by their $\Delta L = 1$ collisions or mediated $\Delta L = 2$ reactions. Later on, the neutrino asymmetry Λ_1 is converted into a standard lepton asymmetry at temperatures below T_1 . Preliminary numerical calculations [65] indicate that as much as 1% of the neutrino asymmetry may be converted into a final $B - L$ asymmetry even if there is no CP asymmetry associated with the lightest neutrino decay. This makes a radical contrast with the traditional scenario and will be further investigated.

VI. CONCLUSIONS

We studied an aspect of the leptogenesis mechanism that has ever been overlooked. We showed that as a rule the two helicity states of Majorana neutrinos do not have exactly the same abundances, contrary to what has been tacitly assumed. The helicity asymmetries defined as differences between the two helicity state abundances of each neutrino species can be parametrized with appropriate neutrino chemical potentials as any other particles asymmetries. The leptogenesis processes generate neutrino helicity

asymmetries of the same order of magnitude as the standard lepton asymmetries. This is quite natural because in the ultrarelativistic limit neutrino Majorana masses are negligible and the neutrino - lepton Yukawa couplings conserve a total lepton number assigned as +1 for right-handed neutrinos and -1 for left-handed neutrinos.

The neutrino helicity asymmetries participate in the system of Boltzmann equations that govern the evolution of particle asymmetries and do not decouple because for any particular reaction the two neutrino helicity states have distinct reaction rates. A reaction where an incoming left-handed lepton doublet goes with an incoming (outgoing) neutrino with positive (negative) helicity is suppressed with respect to the reaction where the same neutrino species is in the opposite negative (positive) helicity state. The former is possible only because the neutrino has a Majorana mass but its amplitude is suppressed by the Lorentz contraction factor. The two reaction rates coincide only when the neutrino is at rest. As a result the standard lepton and neutrino sectors interchange asymmetries with each other which affects significantly the transport and dissipation of lepton number.

It proves convenient to work with the quantum number $B - L'$ where L' is the sum of the total standard lepton number, L , and the total neutrino helicity asymmetry, Λ (equal to the difference between the total number of neutrinos with positive helicity and the total number of negative helicity neutrinos). $B - L'$ is conserved by sphalerons, its violation by Yukawa couplings is suppressed by the neutrino mass over temperature ratios, and reduces to the standard model $B - L$ number when the singlet neutrinos vanish from the Universe. The chemical potential constraints enforced by the fast reactions at any given moment set the asymmetries of particles in equilibrium as proportional to $B - L'$.

We made a detailed analysis of the period when neutrinos are ultrarelativistic which permits simplifying approximations. It shows that one has to distinguish between the temperature T_a at which a neutrino species N_a comes into equilibrium and the temperature scale where its reactions have a significant damping (washout) effect on $B - L'$ and all particle asymmetries proportional to $B - L'$ ($B - L$ in

particular). A neutrino species comes into equilibrium when its $\Delta L = 1$ reactions become fast in Hubble rate terms. In the existing literature all $\Delta L \neq 0$ reactions contribute to wash out the $B - L$ asymmetry. But this is not right because it does not differentiate between the neutrino helicity states. Some neutrino helicity configurations conserve L' . These channels dominate the rates and exist even in absence of neutrino Majorana masses. The other channels violate L' and contribute to wash out $B - L'$ but they are subdominant and are suppressed by the second power of neutrino Lorentz contraction factors. A consequence of this is that the contribution of $\Delta L = 1$ reactions to the $B - L'$ damping rate is 2 orders of magnitude smaller than expected if the neutrino helicity asymmetries are ignored.

In Sec. IV we studied the generation processes in the period of neutrino thermal production. It turns out that the $B - L'$ asymmetry generated directly in the neutrino helicity sector is of opposite sign and larger than the asymmetry generated in the standard lepton sector. We did not attempt to study the decaying phase of singlet neutrino(s). The calculations are more involved because the temperature is of the order of the neutrino mass(es). But one can identify one important difference with the traditional hierarchical scenario, where one of the singlet neutrinos is much lighter than the others: the asymmetries left when the heaviest neutrinos vanish from the Universe are not necessarily washed out by the fast $\Delta L = 1$ reactions of the lightest neutrino before the temperature reaches its mass scale. Moreover, even if the heaviest neutrino $\Delta L = 1$ collisions or $\Delta L = 2$ mediated reactions wash out the standard lepton and Higgs boson asymmetries, the helicity asymmetry carried by the lightest neutrino may survive if it enters in equilibrium at a relaxation temperature below the heaviest neutrino masses. The neutrino asymmetry is later converted into a standard global $B - L$ asymmetry at temperatures close to its mass. This has been supported by numerical calculations and will be reported elsewhere [65].

ACKNOWLEDGMENTS

This work was partially supported by the FCT grants CERN/FNU/43666/2001 and POCTI/FNU/43666/2002.

-
- [1] W. Buchmüller and M. Plümacher, *Int. J. Mod. Phys. A* **15**, 5047 (2000); *Phys. Rep.* **320**, 329 (1999).
 - [2] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174**, 45 (1986).
 - [3] M. A. Luty, *Phys. Rev. D* **45**, 455 (1992).
 - [4] L. Covi, E. Roulet, and F. Vissani, *Phys. Lett. B* **384**, 169 (1996); E. Roulet, L. Covi, and F. Vissani, *Phys. Lett. B* **424**, 101 (1998).
 - [5] V. Kuzmin, V. Rubakov, and M. Shaposhnikov, *Phys. Lett. B* **155**, 36 (1985).
 - [6] P. Minkowski, *Phys. Lett. B* **67**, 421 (1977).
 - [7] M. Gell-Mann, P. Ramond, and R. Slansky, in *Proceedings of the Workshop, Stony Brook, N.Y., 1979*, edited by P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979).
 - [8] T. Yanagida, in *Proceedings of the Workshop on Unified Theories and Baryon Number in the Universe, Tsukuba, Japan, 1979*, edited by A. Sawada and A. Sugamoto

- (unpublished); Prog. Theor. Phys. **64**, 1103 (1980).
- [9] S.L. Glashow, in *Quarks and Leptons, Cargèse, 1979*, edited M. Levy *et al.* (Plenum, New York, 1980).
- [10] R.N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
- [11] A.E. Nelson and S.M. Barr, Phys. Lett. B **246**, 141 (1990).
- [12] W. Fischler, G.F. Giudice, R.G. Leigh, and S. Paban, Phys. Lett. B **258**, 45 (1991).
- [13] W. Buchmüller and T. Yanagida, Phys. Lett. B **302**, 240 (1993).
- [14] A. Pilaftsis, Phys. Rev. D **56**, 5431 (1997); Int. J. Mod. Phys. A **14**, 1811 (1999).
- [15] M. Hirsch and S.F. King, Phys. Rev. D **64**, 113005 (2001).
- [16] W. Buchmüller and D. Wyler, Phys. Lett. B **521**, 291 (2001).
- [17] T. Endoh, S. Kaneko, S.K. Kang, T. Morozumi, and M. Tanimoto, Phys. Rev. Lett. **89**, 231601 (2002).
- [18] G.C. Branco, R. González Felipe, F.R. Joaquim, and M.N. Rebelo, Nucl. Phys. **B640**, 202 (2002).
- [19] S. Davidson and A. Ibarra, Phys. Lett. B **535**, 25 (2002); Nucl. Phys. **B648**, 345 (2003).
- [20] J. Ellis and M. Raidal, Nucl. Phys. **B643**, 229 (2002).
- [21] W. Buchmüller, P. Di Bari, and M. Plümacher, Nucl. Phys. **B643**, 367 (2002); **B665**, 445 (2003); Phys. Lett. B **547**, 128 (2002); Ann. Phys. (N.Y.) **315**, 305 (2005).
- [22] S. Davidson, J. High Energy Phys. 03 (2003) 037.
- [23] M.N. Rebelo, Phys. Rev. D **67**, 013008 (2003).
- [24] A. Ibarra and G.G. Ross, Phys. Lett. B **591**, 285 (2004).
- [25] G.F. Giudice, A. Notari, M. Raidal, A. Riotto, and A. Strumia, Nucl. Phys. **B685**, 89 (2004).
- [26] T. Hambye, Y. Lin, A. Notari, M. Papucci, and A. Strumia, Nucl. Phys. **B695**, 169 (2004).
- [27] M. Plümacher, Z. Phys. C **74**, 549 (1997).
- [28] M. Flanz and E.A. Paschos, Phys. Rev. D **58**, 113009 (1998).
- [29] R. Barbieri, P. Creminelli, A. Strumia, and N. Tetradis, Nucl. Phys. **B575**, 61 (2000).
- [30] A. Pilaftsis and T.E.J. Underwood, Nucl. Phys. **B692**, 303 (2004).
- [31] A.D. Sakharov, JETP Lett. **5**, 24 (1967).
- [32] J.N. Bahcall, M.C. Gonzalez-Garcia, and C. Pena-Garay, J. High Energy Phys. 08 (2004) 016.
- [33] M. Maltoni, T. Schwetz, M.A. Tortola, and J.W.F. Valle, New J. Phys. **6**, 122 (2004).
- [34] S. Goswami, A. Bandyopadhyay, and S. Choubey, Nucl. Phys. B Proc. Suppl. **143**, 121 (2005).
- [35] R. Davis, Prog. Part. Nucl. Phys. **32**, 13 (1994).
- [36] B.T. Cleveland *et al.*, Astrophys. J. **496**, 505 (1998).
- [37] GALLEX Collaboration, W. Hampel *et al.*, Phys. Lett. B **447**, 127 (1999).
- [38] GNO Collaboration, C. M. Cattadori, Nucl. Phys. B Proc. Suppl. **110**, 311 (2002).
- [39] SAGE Collaboration, J.N. Abdurashitov *et al.*, J. Exp. Theor. Phys. **95**, 181 (2002); [Zh. Eksp. Teor. Fiz. **122**, 211 (2002)].
- [40] Super-Kamiokande Collaboration, S. Fukuda *et al.*, Phys. Lett. B **539**, 179 (2002); Super-Kamiokande Collaboration, M.B. Smy *et al.*, Phys. Rev. D **69**, 011104 (2004).
- [41] SNO Collaboration, Q.R. Ahmad *et al.*, Phys. Rev. Lett. **89**, 011301 (2002); **89**, 011302 (2002); SNO Collaboration, S.N. Ahmed *et al.*, Phys. Rev. Lett. **92**, 181301 (2004).
- [42] Super-Kamiokande Collaboration, Y. Fukuda *et al.*, Phys. Rev. Lett. **81**, 1562 (1998); Super-Kamiokande Collaboration, Y. Ashie *et al.*, Phys. Rev. Lett. **93**, 101801 (2004).
- [43] MACRO Collaboration, M. Ambrosio *et al.*, Phys. Lett. B **566**, 35 (2003).
- [44] Soudan 2 Collaboration, M. Sanchez *et al.*, Phys. Rev. D **68**, 113004 (2003).
- [45] KamLAND Collaboration, K. Eguchi *et al.*, Phys. Rev. Lett. **90**, 021802 (2003).
- [46] K2K Collaboration, M.H. Ahn *et al.*, Phys. Rev. Lett. **90**, 041801 (2003).
- [47] L. Bento, in *Proceedings of the Fourth Tegernsee International Conference on Particle Physics Beyond the Standard Model, Tegernsee, Germany, 2003*, edited by H.-V. Klapdor Kleingrothaus (Springer, Berlin, 2004); hep-ph/0210274.
- [48] P. Langacker, G. Segrè, and S. Soni, Phys. Rev. D **26**, 3425 (1982); P. Langacker and G. Segrè, Phys. Rev. Lett. **49**, 1363 (1982).
- [49] K.M. Case, Phys. Rev. **107**, 307 (1957).
- [50] J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); **24**, 1883 (1981).
- [51] L. Bento and Z. Berezhiani, Phys. Rev. D **62**, 055003 (2000).
- [52] E. Kh. Akhmedov, V.A. Rubakov, and A. Yu. Smirnov, Phys. Rev. Lett. **81**, 1359 (1998).
- [53] J.M. Cline, K. Kainulainen, and K.A. Olive, Phys. Rev. Lett. **71**, 2372 (1993); Phys. Rev. D **49**, 6394 (1994); B. A. Campbell, S. Davidson, J. Ellis, and K.A. Olive, Phys. Lett. B **297**, 118 (1992).
- [54] V.V. Klimov, Sov. J. Nucl. Phys. **33**, 934 (1981); [Yad. Fiz. **33**, 1734 (1981)].
- [55] H.A. Weldon, Phys. Rev. D **26**, 2789 (1982); **40**, 2410 (1989).
- [56] S. Davidson, K. Kainulainen, and K.A. Olive, Phys. Lett. B **335**, 339 (1994).
- [57] C. Ford, D.R. Jones, P.W. Stephenson, and M.B. Einhorn, Nucl. Phys. **B395**, 17 (1993).
- [58] H.A. Weldon, Phys. Rev. D **26**, 1394 (1982).
- [59] S. Weinberg, Phys. Rev. Lett. **42**, 850 (1979).
- [60] E.W. Kolb and S. Wolfram, Nucl. Phys. **B172**, 224 (1980); **B195**, 542(E) (1982).
- [61] S. Yu. Khlebnikov and M.E. Shaposhnikov, Nucl. Phys. **B308**, 885 (1988); Phys. Lett. B **387**, 817 (1996).
- [62] J.A. Harvey and M.S. Turner, Phys. Rev. D **42**, 3344 (1990).
- [63] L. Bento, J. Cosmol. Astropart. Phys. 11 (2003) 002.
- [64] E.W. Kolb and M.S. Turner, *The Early Universe*, (Addison-Wesley, Reading, MA, 1990).
- [65] L. Bento and A. Cardoso (to be published).