

Generating perfect fluid spheres in general relativity

Petarpa Boonserm,^{*} Matt Visser,[†] and Silke Weinfurtner[‡]

School of Mathematics, Statistics, and Computer Science, Victoria University of Wellington, PO Box 600, Wellington, New Zealand
(Received 10 March 2005; published 28 June 2005)

Ever since Karl Schwarzschild's 1916 discovery of the spacetime geometry describing the interior of a particular idealized general relativistic star—a static spherically symmetric blob of fluid with position-independent density—the general relativity community has continued to devote considerable time and energy to understanding the general-relativistic static perfect fluid sphere. Over the last 90 years a tangle of specific perfect fluid spheres has been discovered, with most of these specific examples seemingly independent from each other. To bring some order to this collection, in this article we develop several new transformation theorems that map perfect fluid spheres into perfect fluid spheres. These transformation theorems sometimes lead to unexpected connections between previously known perfect fluid spheres, sometimes lead to new previously unknown perfect fluid spheres, and in general can be used to develop a systematic way of classifying the set of all perfect fluid spheres.

DOI: 10.1103/PhysRevD.71.124037

PACS numbers: 04.20.-q, 04.40.Dg, 95.30.Sf

I. INTRODUCTION

General relativistic perfect fluid spheres, or more precisely general relativistic static perfect fluid spheres, are interesting because they are first approximations to any attempt at building a realistic model for a general relativistic star [1–4]. The central idea is to start solely with spherical symmetry, which implies that in orthonormal components the stress energy tensor takes the form

$$T_{\hat{a}\hat{b}} = \begin{bmatrix} \rho & 0 & 0 & 0 \\ 0 & p_r & 0 & 0 \\ 0 & 0 & p_t & 0 \\ 0 & 0 & 0 & p_t \end{bmatrix} \quad (1)$$

and then use the perfect fluid constraint $p_r = p_t$, making the radial pressure equal to the transverse pressure. By using the Einstein equations, plus spherical symmetry, the equality $p_r = p_t$ for the pressures becomes the statement

$$G_{\hat{\theta}\hat{\theta}} = G_{\hat{r}\hat{r}} = G_{\hat{\phi}\hat{\phi}}. \quad (2)$$

In terms of the metric components, this leads to an ordinary differential equation (ODE), which then constrains the spacetime geometry, for *any* perfect fluid sphere.

Over the last 90 years, many “*ad hoc*” approaches to solving this differential equation have been explored, often by picking special coordinate systems, or making simple ansätze for one or another of the metric components [5–8]. (For recent overviews see [1–3].) The big change over the last several years has been the introduction of “algorithmic” techniques that permit one to generate large classes

perfect fluid spheres in a purely mechanical way [9–11]. In this article we will extend these algorithmic ideas, by proving several solution-generating theorems of varying levels of complexity. We shall then explore the formal properties of these solution-generating theorems (which generalize the notion of the Buchdahl transformation) and then will use these theorems to classify some of the previously known exact solutions, and additionally will generate several new previously unknown perfect fluid solutions.

II. SOLUTION-GENERATING THEOREMS

Start with some static spherically symmetric geometry in Schwarzschild (curvature) coordinates

$$ds^2 = -\zeta(r)^2 dt^2 + \frac{dr^2}{B(r)} + r^2 d\Omega^2 \quad (3)$$

and assume it represents a perfect fluid sphere. That is, $G_{\hat{\theta}\hat{\theta}} = G_{\hat{r}\hat{r}} = G_{\hat{\phi}\hat{\phi}}$. While $G_{\hat{\theta}\hat{\theta}} = G_{\hat{\phi}\hat{\phi}}$ is always fulfilled due to spherical symmetry, setting $G_{\hat{r}\hat{r}} = G_{\hat{\theta}\hat{\theta}}$ supplies us with an ODE

$$[r(\zeta)']B' + [2r^2\zeta'' - 2(r\zeta)']B + 2\zeta = 0, \quad (4)$$

which reduces the freedom to choose the two functions in Eq. (3) to one. This equation (4) is a first-order linear nonhomogeneous equation in $B(r)$. Thus—once you have chosen a $\zeta(r)$ —this equation can always be solved for $B(r)$. Solving for $B(r)$ in terms of $\zeta(r)$ is the basis of [10,11], (and is the basis for Theorem 1 below). On the other hand, we can also regroup this same equation as

$$2r^2B\zeta'' + (r^2B' - 2rB)\zeta' + (rB' - 2B + 2)\zeta = 0, \quad (5)$$

which is a linear homogeneous second-order ODE for $\zeta(r)$, which will become the basis for Theorem 2 below. The question we are going to answer in this section is, how to systematically “deform” the geometry (3) while still retaining the perfect fluid property. That is, suppose we start

^{*}Electronic address: Petarpa.Boonserm@mcs.vuw.ac.nz
URL: <http://www.mcs.vuw.ac.nz/research/people/Petarpa/name>
[†]Electronic address: matt.visser@vuw.ac.nz
URL: <http://www.mcs.vuw.ac.nz/~visser>
[‡]Electronic address: silke.weinfurtner@vuw.ac.nz
URL: <http://www.mcs.vuw.ac.nz/research/people/Silke/name>

with the specific geometry defined by

$$ds^2 = -\zeta_0(r)^2 dt^2 + \frac{dr^2}{B_0(r)} + r^2 d\Omega^2 \quad (6)$$

and assume it represents a perfect fluid sphere. We will show how to deform this solution by applying five different transformation theorems on $\{\zeta_0, B_0\}$, such that the outcome still presents a perfect fluid sphere. The outcome of this process will depend on one or more free parameters, and so automatically generates an entire family of perfect fluid spheres of which the original starting point is only one member. In addition, we analyze what happens if you apply these theorems more than once, iterating them in various ways.

A. Four theorems

The first theorem we present is a variant of a result first explicitly published in [11], though another limited variant of this result can also be found in [3]. We first rephrase the theorem of [11] in a slightly different formalism, and demonstrate an independent way of proving it. Using our proof it is easy to show that by applying Theorem 1 more than once no further solutions will be obtained, therefore the transformation in Theorem 1 is, (in a certain sense to be made precise below), “idempotent.”

Theorem 1.— Suppose $\{\zeta_0(r), B_0(r)\}$ represents a perfect fluid sphere. Define

$$\begin{aligned} \Delta_0(r) &= \left(\frac{\zeta_0(r)}{\zeta_0(r) + r\zeta_0'(r)} \right)^2 r^2 \\ &\times \exp \left\{ 2 \int \frac{\zeta_0'(r)}{\zeta_0(r)} \frac{\zeta_0(r) - r\zeta_0'(r)}{\zeta_0(r) + r\zeta_0'(r)} dr \right\}. \end{aligned} \quad (7)$$

Then for all λ , the geometry defined by holding $\zeta_0(r)$ fixed and setting

$$ds^2 = -\zeta_0(r)^2 dt^2 + \frac{dr^2}{B_0(r) + \lambda\Delta_0(r)} + r^2 d\Omega^2 \quad (8)$$

is also a perfect fluid sphere. That is, the mapping

$$\mathbf{T}_1(\lambda): \{\zeta_0, B_0\} \mapsto \{\zeta_0, B_0 + \lambda\Delta_0(\zeta_0)\} \quad (9)$$

takes perfect fluid spheres into perfect fluid spheres. Furthermore a second application of the transformation does not yield new information, \mathbf{T}_1 is “idempotent,” in the sense that

$$\begin{aligned} \mathbf{T}_1(\lambda_n) \circ \cdots \circ \mathbf{T}_1(\lambda_2) \circ \mathbf{T}_1(\lambda_1): \{\zeta_0, B_0\} \\ \mapsto \left\{ \zeta_0, B_0 + \left(\sum_{i=1}^n \lambda_i \right) \Delta_0(\zeta_0) \right\} \end{aligned} \quad (10)$$

We also note that \mathbf{T}_1 always has an inverse

$$[\mathbf{T}_1(\lambda)]^{-1} = \mathbf{T}_1(-\lambda). \quad (11)$$

Proof for Theorem 1.— Assume that $\{\zeta_0(r), B_0(r)\}$ is a solution for Eq. (4). Under what conditions does $\{\zeta_0(r), B_1(r)\}$ also satisfy Eq. (4)? Without loss of generality, we write

$$B_1(r) = B_0(r) + \lambda\Delta_0(r). \quad (12)$$

Equation (4) can now be used to determine $\Delta_0(r)$. That ordinary *inhomogeneous* first-order differential equation in B now simplifies to

$$[r(r\zeta_0)']\Delta_0' + [2r^2\zeta_0'' - 2(r\zeta_0)']\Delta_0 = 0, \quad (13)$$

which is an ordinary *homogeneous* first-order differential equation in Δ_0 . A straightforward calculation, including an integration by parts, leads to

$$\Delta_0(r) = \frac{r^2}{[r\zeta_0(r)]^2} \exp \left\{ \int \frac{4\zeta_0'(r)}{r\zeta_0(r)} dr \right\}. \quad (14)$$

Adding and subtracting $\pm 2(r\zeta_0'')/(\zeta_0(r\zeta_0'))$ to the argument under the integral leads to

$$\Delta_0 = \left(\frac{\zeta_0(r)}{\zeta_0(r) + r\zeta_0'(r)} \right)^2 r^2 \exp \left\{ 2 \int \frac{\zeta_0'(r)}{\zeta_0(r)} \frac{\zeta_0(r) - r\zeta_0'(r)}{\zeta_0(r) + r\zeta_0'(r)} dr \right\}, \quad (15)$$

as advertised. If we apply this transformation a second time we obtain no additional information. To see this, consider the sequence

$$\{\zeta_0, B_0\} \mapsto \{\zeta_0, B_1\} \mapsto \{\zeta_0, B_2\} \dots \quad (16)$$

But at the second step, since ζ_0 has not changed, we have $\Delta_1(r) = \Delta_0(r)$. More generally, at all subsequent steps, $\Delta_i(r) = \Delta_0(r)$. We can write this as

$$\prod_{i=1}^n \mathbf{T}_1(\lambda_i) = \mathbf{T}_1 \left(\sum_{i=1}^n \lambda_i \right). \quad (17)$$

or in the more suggestive form

$$\prod_{i=1}^n \mathbf{T}_1 \triangleq \mathbf{T}_1 \quad (18)$$

where the symbol \triangleq indicates “equality up to relabeling of the parameters.” That is, transformation \mathbf{T}_1 is idempotent up to relabeling of the parameters. (See Fig. 1.) \square

A version of Theorem 1 can also be found in [3]. Specifically, after several manipulations, changes of notation, and a change of coordinate system, the transformation exhibited in Eq. (16.11) of [3] can be converted into the transformation \mathbf{T}_1 of Theorem 1 above.

Applying Theorem 1 to a fixed $\{\zeta_0, B_0\}$ generates a one-dimensional space of perfect fluid spheres, which leads to the corollary below:

Corollary 1.— Let $\{\zeta_0, B_a\}$ and $\{\zeta_0, B_b\}$ both represent perfect fluid spheres, then for all p

$$\{\zeta_0, pB_a + (1-p)B_b\} \quad (19)$$




FIG. 1 (color online). The solid lines show $B(r)$ for 10 reapplications of Theorem 1 onto the Minkowski metric. The dashed line corresponds to a single application with the specific choice $\lambda_{\text{once}} = \sum \lambda_i$. It can be seen that 10 applications of Theorem 1 can be expressed by one application.

is also a perfect fluid sphere, furthermore all perfect fluid spheres for a fixed ζ_0 can be written in this form.

Proof.— The result is automatic once you note that for fixed ζ_0 the ODE for B is linear, (though not homogeneous, which is why the two coefficients p and $1 - p$ are chosen to add up to 1). \square

We defer extensive discussion of the application of this theorem and its corollary until Sec. III, where we use this and our other generating theorems as a basis for classifying perfect fluid spheres. At this stage we mention, only as a very simple example, that \mathbf{T}_1 applied to Minkowski space results in the geometry of the Einstein static universe.

Theorem 2.— Let $\{\zeta_0, B_0\}$ describe a perfect fluid sphere. Define

$$Z_0(r) = \sigma + \epsilon \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)}}. \quad (20)$$

Then for all σ and ϵ , the geometry defined by holding $B_0(r)$ fixed and setting

$$ds^2 = -\zeta_0(r)^2 Z_0(r)^2 dt^2 + \frac{dr^2}{B_0(r)} + r^2 d\Omega^2 \quad (21)$$

is also a perfect fluid sphere. That is, the mapping

$$\mathbf{T}_2(\sigma, \epsilon): \{\zeta_0, B_0\} \mapsto \{\zeta_0 Z_0(\zeta_0, B_0), B_0\} \quad (22)$$

takes perfect fluid spheres into perfect fluid spheres. Furthermore a second application of the transformation does not yield new information, \mathbf{T}_2 is idempotent in the sense that

$$\begin{aligned} \mathbf{T}_2(\sigma_n, \epsilon_n) \circ \cdots \circ \mathbf{T}_2(\sigma_3, \epsilon_3) \circ \mathbf{T}_2(\sigma_2, \epsilon_2) \circ \mathbf{T}_2(\sigma_1, \epsilon_1) \\ = \mathbf{T}_2(\sigma_n \cdots \sigma_3 \sigma_2 \sigma_1, \epsilon_{n \dots 321}), \end{aligned} \quad (23)$$

where

$$\begin{aligned} \epsilon_{n \dots 321} &= (\epsilon_1 \sigma_2 \sigma_3 \cdots \sigma_n) + (\sigma_1^{-1} \epsilon_2 \sigma_3 \cdots \sigma_n) \\ &+ (\sigma_1^{-1} \sigma_2^{-1} \epsilon_3 \cdots \sigma_n) + \cdots \\ &+ (\sigma_1^{-1} \sigma_2^{-1} \sigma_3^{-1} \cdots \epsilon_n). \end{aligned} \quad (24)$$

Furthermore, Theorem 2 is invertible (as long as $\sigma \neq 0$):

$$[\mathbf{T}_2(\sigma, \epsilon)]^{-1} = \mathbf{T}_2(1/\sigma, -\epsilon). \quad (25)$$

Proof for Theorem 2.— The proof of Theorem 2 is based on the technique of “reduction in order.” Assuming that $\{\zeta_0(r), B_0(r)\}$ solves Eq. (5), write

$$\zeta_1(r) = \zeta_0(r) Z_0(r). \quad (26)$$

and demand that $\{\zeta_1(r), B_0(r)\}$ also solves Eq. (5). We find

$$(r^2 \zeta_0 B_0' + 4r^2 \zeta_0' B_0 - 2r \zeta_0 B_0) Z_0' + (2r^2 \zeta_0 B_0) Z_0'' = 0, \quad (27)$$

which is an ordinary homogeneous second-order differential equation, depending only on Z_0' and Z_0'' . (So it can be viewed as a first-order homogeneous differential equation in Z_0' , which is solvable.) Separating the unknown variable to one side,

$$\frac{Z_0''}{Z_0'} = -\frac{1}{2} \frac{B_0'}{B_0} - 2 \frac{\zeta_0'}{\zeta_0} + \frac{1}{r}. \quad (28)$$

Rewrite $Z_0''/Z_0' = d \ln(Z_0')/dt$, and integrate twice over both sides of Eq. (28), to obtain

$$Z_0 = \sigma + \epsilon \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)}}, \quad (29)$$

depending on the old solution $\{\zeta_0(r), B_0(r)\}$, and two arbitrary integration constants σ and ϵ .

To see that the transformation \mathbf{T}_2 defined in Theorem 2 is idempotent we first show

$$\mathbf{T}_2 \circ \mathbf{T}_2 \triangleq \mathbf{T}_2, \quad (30)$$

and then iterate. The precise two-step composition rule is

$$\mathbf{T}_2(\sigma_2, \epsilon_2) \circ \mathbf{T}_2(\sigma_1, \epsilon_1) = \mathbf{T}_2\left(\sigma_2 \sigma_1, \epsilon_1 \sigma_2 + \frac{\epsilon_2}{\sigma_1}\right). \quad (31)$$

To see “idempotence,” note that for fixed $B_0(r)$ Eq. (5) has a solution space that is exactly two dimensional. Since the first application of \mathbf{T}_2 takes any specific solution and maps it into the full two-dimensional solution space, any subsequent application of \mathbf{T}_2 can do no more than move one around inside this two-dimensional solution space—physically this corresponds to a relabeling of parameters describing the perfect fluid metric you are dealing with, not the generation of new solutions. To be more explicit about this note that at step one

$$\zeta_0 \rightarrow \zeta_1 = \zeta_0 \left\{ \sigma_1 + \epsilon_1 \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)}} \right\}, \quad (32)$$

while at the second step

$$\zeta_1 \rightarrow \zeta_2 = \zeta_1 \left\{ \sigma_2 + \epsilon_2 \int \frac{rdr}{\zeta_1(r)^2 \sqrt{B_0(r)}} \right\}. \quad (33)$$

That is:

$$\begin{aligned} \zeta_2 = \zeta_0 & \left\{ \sigma_1 + \epsilon_1 \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)}} \right\} \left\{ \sigma_2 \right. \\ & \left. + \epsilon_2 \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)} [\sigma_1 + \epsilon_1 \int rdr / (\zeta_0(r)^2 \sqrt{B_0(r)})]^2} \right\}. \end{aligned} \quad (34)$$

But this can be rewritten as

$$\begin{aligned} \zeta_2 = \zeta_0 Z_0 & \left\{ \sigma_2 + \frac{\epsilon_2}{\epsilon_1} \int \frac{dZ_0}{Z_0^2} \right\} = \zeta_0 Z_0 \left\{ \sigma_2 - \frac{\epsilon_2}{\epsilon_1} \int d\left(\frac{1}{Z_0}\right) \right\} \\ = \zeta_0 Z_0 & \left\{ \sigma_2 - \frac{\epsilon_2}{\epsilon_1} \left[\frac{1}{Z_0} - \frac{1}{\sigma_1} \right] \right\}. \end{aligned} \quad (35)$$

Therefore

$$\zeta_2 = \zeta_0 \left\{ -\frac{\epsilon_2}{\epsilon_1} + \left[\sigma_2 + \frac{\epsilon_2}{\epsilon_1} \frac{1}{\sigma_1} \right] Z_0 \right\}. \quad (36)$$

That is

$$Z_1 = -\frac{\epsilon_2}{\epsilon_1} + \left[\sigma_2 + \frac{\epsilon_2}{\epsilon_1} \frac{1}{\sigma_1} \right] Z_0, \quad (37)$$

from which the composition law

$$\mathbf{T}_2(\sigma_2, \epsilon_2) \circ \mathbf{T}_2(\sigma_1, \epsilon_1) = \mathbf{T}_2\left(\sigma_2 \sigma_1, \epsilon_1 \sigma_2 + \frac{\epsilon_2}{\sigma_1}\right) \quad (38)$$

follows immediately. (Note that the composition law for \mathbf{T}_2 is actually a statement about applying reduction of order to second-order ODEs, it is not specifically a statement about perfect fluid spheres, though that is how we will use it in the current article.) The general composition law then follows by induction. To find the inverse transformation we choose $\sigma_2 = 1/\sigma_1$ and $\epsilon_1 = -\epsilon_2$, for which

$$\mathbf{T}_2(1/\sigma_1, -\epsilon_1) \circ \mathbf{T}_2(\sigma_1, \epsilon_1) = \mathbf{T}_2(1, 0) = \mathbf{I}. \quad (39)$$

□

Comment.— As other special cases of the composition law we also mention the results that

$$\prod_{i=1}^n \mathbf{T}_2(1, \epsilon_i) = \mathbf{T}_2\left(1, \sum_{i=1}^n \epsilon_i\right), \quad (40)$$

and

$$\begin{aligned} \mathbf{T}_2(\sigma, \epsilon)^n = \mathbf{T}_2(\sigma^n, \epsilon[\sigma^{n-1} + \sigma^{n-3} \dots + \sigma^{-(n-3)} \\ + \sigma^{-(n-1)}]). \end{aligned} \quad (41)$$

Now as long as $\sigma > 1$ then for sufficiently large n we see

$$\mathbf{T}_2(\sigma, \epsilon)^n \approx \mathbf{T}_2(\sigma^n, \sigma^{n-1} \epsilon) = \sigma^{n-1} \mathbf{T}_2(\sigma, \epsilon) \triangleq \mathbf{T}_2(\sigma, \epsilon), \quad (42)$$

where at the last step we have used the fact that the overall multiplicative factor σ^{n-1} can simply be reabsorbed into a redefinition of the time coordinate. Because of this result, we see that for fixed $\sigma > 1$ (and fixed but arbitrary ϵ) repeated numerical applications of $\mathbf{T}_2(\sigma, \epsilon)$ will have a well-defined limit. In Fig. 2 we have tested the composition law numerically.

A strictly limited version of Theorem 2, with little comment on its importance, can be found in [3]. Specifically, after several manipulations, changes of notation, and a change of coordinate system, the transformation exhibited in Eq. (16.12) of [3] can be seen to be equivalent to the subcase $\sigma = 0, \epsilon = 1$ of transformation \mathbf{T}_2 above.

For some purposes it is more useful to rephrase Theorem 2 as below:

Corollary 2.—Let $\{\zeta_a, B_0\}$ and $\{\zeta_b, B_0\}$ both represent perfect fluid spheres, then for all p and q

$$\{p\zeta_a + q\zeta_b, B_0\} \quad (43)$$

is also a perfect fluid sphere. Furthermore, for fixed B_0 all perfect fluid spheres can be written in this form.

Proof.—The result is automatic once you note that for fixed B_0 the ODE for ζ is second-order linear and homogeneous. □

We again defer extensive discussion of the application of this theorem and its corollary until Sec. III, at this stage mentioning only as a very simple example that \mathbf{T}_2 applied to either the Einstein static universe or the anti-de Sitter universe results in Schwarzschild's stellar solution




FIG. 2 (color online). The solid lines show $\zeta(r)$ for 10 reapplications of Theorem 2 onto the Minkowski metric. The dashed line corresponds to a single application with the specific choice for σ_{once} and ϵ_{once} determined by the composition law of Theorem 2. It can be seen that 10 applications of Theorem 2 can be reexpressed by a single application.

(position-independent density). Similarly, Corollary 2 applied to a combination of the Einstein static universe and anti-de Sitter space is another way of obtaining Schwarzschild's stellar solution.

Having now found the first and second generating theorems makes it useful to define two new theorems by composing them. Take a perfect fluid sphere solution $\{\zeta_0, B_0\}$. Applying Theorem 1 onto it gives us a new perfect fluid sphere $\{\zeta_0, B_1\}$. The new B_1 is given in Eq. (7). If we now continue by applying Theorem 2, again we get a new solution $\{\tilde{\zeta}, B_1\}$, where $\tilde{\zeta}$ now depends on the new B_1 . All together we can consider this as a single process, by introducing the following theorem:

Theorem 3.—If $\{\zeta_0, B_0\}$ denotes a perfect fluid sphere, then for all σ , ϵ , and λ , the three-parameter geometry defined by

$$ds^2 = -\zeta_0(r)^2 \left\{ \sigma + \epsilon \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r) + \lambda \Delta_0(r)}} \right\}^2 dt^2 + \frac{dr^2}{B_0(r) + \lambda \Delta_0(r)} + r^2 d\Omega^2 \quad (44)$$

is also a perfect fluid sphere, where Δ_0 is

$$\Delta_0(r) = \left(\frac{\zeta_0(r)}{\zeta_0(r) + r\zeta_0'(r)} \right)^2 r^2 \times \exp \left[2 \int \frac{\zeta_0'(r)}{\zeta_0(r)} \frac{\zeta_0(r) - r\zeta_0'(r)}{\zeta_0(r) + r\zeta_0'(r)} dr \right]. \quad (45)$$

That is

$$\mathbf{T}_3 = \mathbf{T}_2 \circ \mathbf{T}_1 : \{\zeta_0, B_0\} \mapsto \{\zeta_0, B_0 + \lambda \Delta_0(\zeta_0)\} \mapsto \{\zeta_0 Z_0(\zeta_0, B_0 + \lambda \Delta_0(\zeta_0)), B_0 + \lambda \Delta_0(\zeta_0)\}. \quad (46)$$

Now, instead of starting with Theorem 1 we could first apply Theorem 2 on $\{\zeta_0, B_0\}$. This gives us a new perfect fluid sphere $\{\zeta_1, B_0\}$, where $\zeta_1 = \zeta_0 Z_0(\zeta_0, B_0)$ is given by Eq. (29). Continuing with Theorem 1 leads to $\{\zeta_1, \tilde{B}\}$, where \tilde{B} depends on the new ζ_1 . Again, we can consider this as a single process, by introducing the following theorem:

Theorem 4.—If $\{\zeta_0, B_0\}$ denotes a perfect fluid sphere, then for all σ , ϵ , and λ , the three-parameter geometry defined by

$$ds^2 = -\zeta_0(r)^2 \left\{ \sigma + \epsilon \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)}} \right\}^2 dt^2 + \frac{dr^2}{B_0(r) + \lambda \Delta_0(\zeta_1, r)} + r^2 d\Omega^2 \quad (47)$$

is also a perfect fluid sphere, where $\Delta_0(\zeta_1, r)$ is defined as

$$\Delta_0(\zeta_1, r) = \left(\frac{\zeta_1(r)}{\zeta_1(r) + r\zeta_1'(r)} \right)^2 r^2 \times \exp \left[2 \int \frac{\zeta_1'(r)}{\zeta_1(r)} \frac{\zeta_1(r) - r\zeta_1'(r)}{\zeta_1(r) + r\zeta_1'(r)} dr \right], \quad (48)$$

depending on $\zeta_1 = \zeta_0 Z_0$, and where as before

$$Z_0(r) = \sigma + \epsilon \int \frac{rdr}{\zeta_0(r)^2 \sqrt{B_0(r)}}. \quad (49)$$

That is

$$\mathbf{T}_3 = \mathbf{T}_1 \circ \mathbf{T}_2 : \{\zeta_0, B_0\} \mapsto \{\zeta_0 Z_0(\zeta_0, B_0), B_0\} \mapsto \{\zeta_0 Z_0(\zeta_0, B_0), B_0 + \lambda \Delta_0(\zeta_0 Z_0(\zeta_0, B_0))\}. \quad (50)$$

Theorem 3 and Theorem 4 are in general distinct, which can be traced back to the fact that Theorem 1 and Theorem 2 do not in general commute. Furthermore Theorem 3 and Theorem 4 are in general not idempotent

$$\mathbf{T}_3 \not\equiv \mathbf{T}_4; \quad \mathbf{T}_3 \circ \mathbf{T}_3 \not\equiv \mathbf{T}_3; \quad \mathbf{T}_4 \circ \mathbf{T}_4 \not\equiv \mathbf{T}_4. \quad (51)$$

The best way to verify this is to try a few specific examples. There may be some specific and isolated special metrics for which Theorem 3 and Theorem 4 happen to be degenerate, or idempotent, and finding such metrics is important for our classification program.

B. Formal properties of the generating theorems

The solution-generating theorems we have developed interact in a number of interesting ways and exhibit numerous formal properties that will be useful when classifying generic perfect fluid spheres. To start with, Theorem 3 (\mathbf{T}_3) and Theorem 4 (\mathbf{T}_4) can be expressed in terms of Theorem 1 (\mathbf{T}_1) and Theorem 2 (\mathbf{T}_2):

$$\mathbf{T}_3(g) := (\mathbf{T}_2 \circ \mathbf{T}_1)g = \mathbf{T}_2(\mathbf{T}_1(g)); \quad (52)$$

$$\mathbf{T}_4(g) := (\mathbf{T}_1 \circ \mathbf{T}_2)g = \mathbf{T}_1(\mathbf{T}_2(g)).$$

where g is a metric representing a perfect fluid sphere. Having this in mind, some of the new solutions generated by starting from some specific solution can be identified. For example:

$$\mathbf{T}_4 \circ \mathbf{T}_1 \equiv \mathbf{T}_1 \circ \mathbf{T}_2 \circ \mathbf{T}_1 \equiv \mathbf{T}_1 \circ \mathbf{T}_3, \quad (53)$$

or

$$\mathbf{T}_3 \circ \mathbf{T}_3 \equiv \mathbf{T}_2 \circ \mathbf{T}_1 \circ \mathbf{T}_2 \circ \mathbf{T}_1 \equiv \mathbf{T}_2 \circ \mathbf{T}_4 \circ \mathbf{T}_1. \quad (54)$$

The idempotence of \mathbf{T}_1 and \mathbf{T}_2 in this formalism is:

$$(\mathbf{T}_1 \circ \mathbf{T}_1)g = \mathbf{T}_1(\mathbf{T}_1(g)) \triangleq \mathbf{T}_1(g); \quad (55)$$

$$(\mathbf{T}_2 \circ \mathbf{T}_2)g = \mathbf{T}_2(\mathbf{T}_2(g)) \triangleq \mathbf{T}_2(g).$$

Taken together, it is possible to simplify all formulae wherever \mathbf{T}_1 and \mathbf{T}_2 appear next to each other more than once. The following examples should demonstrate how this

TABLE I. Some well-known perfect fluid spheres and their corresponding metrics. Note that we have often reparametrized these metrics to make them easier to deal with, and so their appearance (but not the substance) may differ from other sources [1–3]. Metric S1 is a special case of K-O III, Tolman V, and Tolman VI notable for its extreme simplicity.

Name	Metric
Minkowski	$-dt^2 + dr^2 + r^2d\Omega^2$
Einstein static	$-dt^2 + (1 - r^2/R^2)^{-1}dr^2 + r^2d\Omega^2$
de Sitter	$-(1 - r^2/R^2)dt^2 + (1 - r^2/R^2)^{-1}dr^2 + r^2d\Omega^2$
Schwarzschild Interior	$-(A - B\sqrt{1 - r^2/R^2})^2dt^2 + (1 - r^2/R^2)^{-1}dr^2 + r^2d\Omega^2$
Schwarzschild Exterior	$-(1 - 2m/r)^2dt^2 + (1 - 2m/r)^{-1}dr^2 + r^2d\Omega^2$
S1	$-r^4dt^2 + dr^2 + r^2d\Omega^2$
K-O III	$-(A + Br^2)^2dt^2 + dr^2 + r^2d\Omega^2$
Kuch1 Ib	$-(Ar + Br \ln r)^2dt^2 + 2dr^2 + r^2d\Omega^2$
B-L	$-A(r^2/a^2)dt^2 + 2(1 + r^2/a^2)^{-1}dr^2 + r^2d\Omega^2$
Tolman IV	$-B^2(1 + r^2/a^2)dr^2 + \frac{1+2r^2/a^2}{(1-r^2/b^2)(1+r^2/a^2)}dr^2 + r^2d\Omega^2$
Tolman IV ($b \rightarrow \infty$)	$-B^2(1 + r^2/a^2)dt^2 + \frac{1+2r^2/a^2}{1+r^2/a^2}dr^2 + r^2d\Omega^2$
Tolman V	$-B^2r^{2(1+n)}dt^2 + (2 - n^2)[1 - Ar^{2(2-n^2)/(2+n)}]^{-1}dr^2 + r^2d\Omega^2$
Tolman V ($A \rightarrow 0$)	$-B^2r^{2(1+n)}dt^2 + (2 - n^2)dr^2 + r^2d\Omega^2$
Tolman VI	$-(Ar^{1-n} + Br^{1+n})^2dt^2 + (2 - n^2)dr^2 + r^2d\Omega^2$
Tolman VII	$-B^2 \cos\{\ln[\sqrt{1 - 2r^2/a^2 + r^4/b^4} + r^2/a^2 - b^2/a^2]^{1/2} + \theta\}^2 + (1 - 2r^2/a^2 + r^4/b^4)^{-1}dr^2 + r^2d\Omega^2$
Tolman VIII	$-A^2r^{2(n-1)(n-4)/n}(\frac{-n^2}{(n^2-4n+2)(n^2-8n+8)} + Br^{-(n^2-8n+8)/n} + Cr^{-2(n^2-4n+2)/n})$ $+ (\frac{-n^2}{(n^2-4n+2)(n^2-8n+8)} + Br^{-(n^2-8n+8)/n} + Cr^{-2(n^2-4n+2)/n})^{-1}dr^2 + r^2d\Omega^2$
Kuch 68 II	$-(1 - 2m/r)dt^2 + [(1 - 2m/r)(1 + C(2r - 2m)^2)]^{-1}dr^2 + r^2d\Omega^2$
Kuch 68 I	$-(A\sqrt{1 - 2m/r} + B[r^2/m^2 + 5r/m - 30 + 15\sqrt{1 - 2m/r} \ln\{1 - r/m - \sqrt{r(r - 2m)/m}\}])^2dt^2$ $+ (1 - 2m/r)^{-1}dr^2 + r^2d\Omega^2$
M-W III	$-Ar(r - a) + \frac{7/4}{1 - r^2/a^2}dr^2 + r^2d\Omega^2$
Kuch I b	$-r^2[A + \text{Barctanh}(a/\sqrt{a^2 + r^2})]^2dt^2 + 2(1 + r^2/a^2)^{-1}dr^2 + r^2d\Omega^2$
Heint IIa ($C=0$)	$-(1 + ar^2)^3dt^2 + [(1 + ar^2)/(1 - ar^2/2)]dr^2 + r^2d\Omega^2$
Heint IIa	$-(1 + ar^2)^3dt^2 + [1 - \frac{3ar^2}{2(1+ar^2)} + \frac{Cr^2}{(1+ar^2)\sqrt{1+4ar^2}}]^{-1}dr^2 + r^2d\Omega^2$

works:

$$\begin{aligned}
 (\mathbf{T}_2 \circ \mathbf{T}_3)g &\equiv (\mathbf{T}_2 \circ \mathbf{T}_2 \circ \mathbf{T}_1)g \triangleq (\mathbf{T}_2 \circ \mathbf{T}_1)g \equiv \mathbf{T}_3(g); \\
 (\mathbf{T}_1 \circ \mathbf{T}_4)g &\equiv (\mathbf{T}_1 \circ \mathbf{T}_1 \circ \mathbf{T}_2)g \triangleq (\mathbf{T}_1 \circ \mathbf{T}_2)g \equiv \mathbf{T}_4(g),
 \end{aligned}
 \tag{56}$$

and in the same way

$$\begin{aligned}
 \mathbf{T}_4 \circ \mathbf{T}_3 &\equiv \mathbf{T}_1 \circ \mathbf{T}_2 \circ \mathbf{T}_2 \circ \mathbf{T}_1 \triangleq \mathbf{T}_1 \circ \mathbf{T}_2 \circ \mathbf{T}_1 \equiv \mathbf{T}_1 \circ \mathbf{T}_3 \\
 &\equiv \mathbf{T}_4 \circ \mathbf{T}_1; \\
 \mathbf{T}_3 \circ \mathbf{T}_4 &\equiv \mathbf{T}_2 \circ \mathbf{T}_1 \circ \mathbf{T}_1 \circ \mathbf{T}_2 \triangleq \mathbf{T}_2 \circ \mathbf{T}_1 \circ \mathbf{T}_2 \\
 &\equiv \mathbf{T}_2 \circ \mathbf{T}_4 \equiv \mathbf{T}_3 \circ \mathbf{T}_2.
 \end{aligned}
 \tag{57}$$

These relationships can be used to structure the solution set generated starting from any particular perfect fluid sphere, and moreover to classify which metrics can be produced by our theorems, and ones which cannot. For example, the idempotence property of Theorem 1 and Theorem 2 en-

ables us to divide the class of perfect fluid spheres into seed metrics and nonseed metrics. Seed metrics can never be generated by using one of the two theorems \mathbf{T}_1 or \mathbf{T}_2 , while nonseed metrics are connected to other simpler metrics via one of these theorems. We formalize this in the following subsection.

C. Seed and nonseed metrics

Definition (Seed metric).—Take a metric g (or a parametrized class of metrics) and apply Theorem 1 or Theorem 2 on it. Three different cases are possible:

- (i) Each of the applications supplies us with a new solution. [$\mathbf{T}_1(g) \not\equiv g \not\equiv \mathbf{T}_2(g)$.] We define a metric with this behavior as a seed metric. (We shall soon see several examples of this behavior.)
- (ii) Only one of the applications supplies us with a new solution, while the other one gives us the same metric we started with. [$\mathbf{T}_1(g) \triangleq g$ or $\mathbf{T}_2(g) \triangleq g$.] These metrics are nonseed metrics. (We shall soon see several examples of this behavior.)

TABLE II. Some apparently new perfect fluid spheres and their corresponding metrics. Sometimes the relevant integrals cannot be done in elementary form. When they can be done they are explicitly shown.

Name	Metric
Martin 1	$-(Ar + Br \ln r)^2 dt^2 + 2 \frac{2A+2B \ln r+B}{2A+2B \ln r+B-Cr^2} dr^2 + r^2 d\Omega^2$
Martin 2	$-Ar(r-a)dt^2 + \frac{7}{4}(1 - \frac{r^2}{a^2} - B \frac{(r-a)r^{7/3}}{(4r-3a)^{4/3}})^{-1} dr^2 + r^2 d\Omega^2$
Martin 3	$-(1 + ar^2)^2 dt^2 + [1 - br^2/(1 + 3ar^2)^{2/3}]^{-1} dr^2 + r^2 d\Omega^2$
P1	$-(1 + ar^2)^2 [A + B \int \frac{(1+ar^2)^{-2}}{\sqrt{1-br^2/(1+3ar^2)^{2/3}}} r dr]^2 dt^2 + [1 - br^2/(1 + 3ar^2)^{2/3}]^{-1} dr^2 + r^2 d\Omega^2$
P2	$-(Ar + Br \ln r)^2 [\sigma + \epsilon \int (Ar + Br \ln r)^{-2} (2 \frac{2A+2B \ln r+B}{2A+2B \ln r+B-Cr^2})^{-1/2} r dr]^2 dt^2 + 2 \frac{2A+2B \ln r+B}{2A+2B \ln r+B-Cr^2} dr^2 + r^2 d\Omega^2$
P3	$-(1 + ar^2)^3 [A + B \frac{(5+2ar^2)\sqrt{1-ar^2/2}}{(1+ar^2)^{3/2}}]^2 dt^2 + [(1 + ar^2)/(1 - ar^2/2)] dr^2 + r^2 d\Omega^2$
P4	$-r^4 (A + B \int \frac{dr}{r^3 \sqrt{1+\lambda r^{2/3}}})^2 dt^2 + (1 + \lambda r^{2/3})^{-1} dr^2 + r^2 d\Omega^2$
P4	$-r^4 (A + B [\frac{15}{16} \lambda^3 \tanh^{-1}(1/\sqrt{1 + \lambda r^{2/3}}) - \frac{1}{16} \sqrt{1 + \lambda r^{2/3}} (8r^{-2} - 10\lambda r^{-4/3} + 15\lambda^2 r^{-2/3})])^2 dt^2 + (1 + \lambda r^{2/3})^{-1} dr^2 + r^2 d\Omega^2$
P5	$-r(A\sqrt{r-a} + B\sqrt{r+a})^2 + \frac{7/4}{1-r^2/a^2} dr^2 + r^2 d\Omega^2$
P6	$-r(r-a)(A + B \int \frac{dr}{(r-a)\sqrt{1-\frac{r^2}{a^2}-B\frac{(r-a)r^{7/3}}{(4r-3a)^{4/3}}}})^2 + \frac{7}{4}(1 - \frac{r^2}{a^2} - B \frac{(r-a)r^{7/3}}{(4r-3a)^{4/3}})^{-1} dr^2 + r^2 d\Omega^2$
P7	$-B^2(1 + r^2/a^2)[A + B \int \frac{\sqrt{a^2-2r^2}}{\sqrt{b^2-r^2(a^2-r^2)^{3/2}}} r dr]^2 dt^2 + \frac{1+2r^2/a^2}{(1-r^2/b^2)(1+r^2/a^2)} dr^2 + r^2 d\Omega^2$
P8	$-(1 + ar^2)^3 [A + B \int \frac{r dr}{(1+ar^2)^2 \sqrt{1-\frac{3ar^2}{2(1+ar^2)} + \frac{Cr^2}{(1+ar^2)\sqrt{1+4ar^2}}}}]^2 dt^2 + [1 - \frac{3ar^2}{2(1+ar^2)} + \frac{Cr^2}{(1+ar^2)\sqrt{1+4ar^2}}]^{-1} dr^2 + r^2 d\Omega^2$

(iii) Both applications give us the same metric we started with. [$\mathbf{T}_1(g) \triangleq g \triangleq \mathbf{T}_2(g)$.] Metrics of this type are fixed points of the transformation process and we then also have $\mathbf{T}_3(g) \triangleq g \triangleq \mathbf{T}_4(g)$.

While we have encountered numerical examples that seem to exhibit this fixed-point behavior, we have no analytic proof for the existence of nonobvious fixed-point metrics. There is one obvious but not particularly useful example of a fixed-point class of metrics. If we take the ODE in Eq. (4), and write down its most general solution as a functional of the arbitrary parameters $\zeta(r)$, then any of our solution-generating theorems applied to this most general solution will at most move us around in the parameter space characterizing the most general solution—the most general solution of Eq. (4), or equivalently Eq. (5), is thus an infinite-parameter fixed point of the generating theorems. But apart from this obvious example, it is unclear whether other fixed-point classes of metric exist.

We can nevertheless develop several formal lemmata regarding fixed-point metrics. For instance

Lemma 1.—Suppose we have a metric such that $\forall \sigma, \epsilon, \lambda$

$$\mathbf{T}_3(\sigma, \epsilon, \lambda)g \triangleq g, \tag{58}$$

where we recall that \triangleq denotes equality up to redefinition of parameters. Then in particular

$$\mathbf{T}_1(\lambda)g \triangleq g \triangleq \mathbf{T}_2(\sigma, \epsilon)g, \tag{59}$$

and so

$$\mathbf{T}_4(\sigma, \epsilon, \lambda)g \triangleq g. \tag{60}$$

Conversely, suppose we have a metric such that

$$\mathbf{T}_4(\sigma, \epsilon, \lambda)g \triangleq g, \tag{61}$$

then

$$\mathbf{T}_1(\lambda)g \triangleq g \triangleq \mathbf{T}_2(\sigma, \epsilon)g, \tag{62}$$

and so

$$\mathbf{T}_3(\sigma, \epsilon, \lambda)g \triangleq g. \tag{63}$$

Proof.—Trivial, note that $\mathbf{T}_3(\sigma, \epsilon, \lambda = 0) = \mathbf{T}_2(\sigma, \epsilon)$ and $\mathbf{T}_3(\sigma = 0, \epsilon = 0, \lambda) = \mathbf{T}_1(\lambda)$. Then recall $\mathbf{T}_4 = \mathbf{T}_1 \circ \mathbf{T}_2$. Similarly for the converse. \square

Lemma 2.—Suppose we have a metric g such that

$$\mathbf{T}_4 g \triangleq \mathbf{T}_3 g \tag{64}$$

and then define g' by

$$\mathbf{T}_4 g \triangleq g' \triangleq \mathbf{T}_3 g. \tag{65}$$

Then we have

$$\mathbf{T}_4 g' \triangleq g' \triangleq \mathbf{T}_3 g' \tag{66}$$

so that g' is a “fixed point” of both \mathbf{T}_3 and \mathbf{T}_4 .

TABLE III. Seed solutions and their generalizations derived via Theorems 1–4. The notation “[integral]” denotes a metric so complicated that explicitly writing out the relevant integral is so tedious that it does not seem worthwhile.

Seed	Theorem 1	Theorem 2	Theorem 3	Theorem 4
Minkowski	Einstein static	K-O III	interior Schwarzschild	Martin 3
exterior Schwarzschild	Kuch68 II	Kuch 86 I	[integral]	[integral]
de Sitter	Tolman IV	interior Schwarzschild	P7	[integral]
Tolman V ($A = 0$)	Tolman V	Tolman VI	Wyman III	Wyman IIa
S1	Tolman V ($n = +1$)	K-O III	P4	Martin 3
M-W III	Martin 2	P5	P6	[integral]
Heint IIa ($C=0$)	Heint IIa	P3	P8	[integral]

Note

$$\mathbf{T}_3(\sigma, \epsilon, \lambda) \sim \mathbf{T}_4(\sigma, \epsilon, \lambda), \quad (71)$$

$$\begin{aligned} \mathbf{T}_{4g'} &\triangleq \mathbf{T}_4 \circ \mathbf{T}_{3g} \triangleq \mathbf{T}_1 \circ \mathbf{T}_2 \circ \mathbf{T}_{1g} \triangleq \mathbf{T}_1 \circ \mathbf{T}_{3g} \\ &\triangleq \mathbf{T}_1 \circ \mathbf{T}_{4g} \triangleq \mathbf{T}_{4g} = g' \end{aligned} \quad (67)$$

and similarly for \mathbf{T}_3 :

$$\begin{aligned} \mathbf{T}_{3g'} &\triangleq \mathbf{T}_3 \circ \mathbf{T}_{4g} \triangleq \mathbf{T}_2 \circ \mathbf{T}_1 \circ \mathbf{T}_{2g} \triangleq \mathbf{T}_2 \circ \mathbf{T}_{4g} \\ &\triangleq \mathbf{T}_2 \circ \mathbf{T}_{3g} \triangleq \mathbf{T}_{3g} = g' \end{aligned} \quad (68)$$

□

Several other formal theorems along these lines can be constructed, but these seem the most important results.

Finally, among the formal properties enjoyed by the generating theorems, we mention the fact that Theorems 3 and 4 are “conjugate” to each other in the following sense

$$\begin{aligned} \mathbf{T}_4 &\equiv \mathbf{T}_1 \circ \mathbf{T}_2 = \mathbf{T}_1 \circ \mathbf{T}_2 \circ \mathbf{T}_1 \circ [\mathbf{T}_1]^{-1} \\ &= \mathbf{T}_1 \circ \mathbf{T}_3 \circ [\mathbf{T}_1]^{-1}, \end{aligned} \quad (69)$$

and similarly (when the appropriate inverse $[\mathbf{T}_2]^{-1}$ exists)

$$\begin{aligned} \mathbf{T}_3 &\equiv \mathbf{T}_2 \circ \mathbf{T}_1 = \mathbf{T}_2 \circ \mathbf{T}_1 \circ \mathbf{T}_2 \circ [\mathbf{T}_2]^{-1} \\ &= \mathbf{T}_2 \circ \mathbf{T}_4 \circ [\mathbf{T}_2]^{-1}. \end{aligned} \quad (70)$$

We can write this as

$$[r(r\zeta_0 Z_0)']\Delta' + [2r^2(\zeta_0 Z_0)'' - 2(r\zeta_0 Z_0)']\Delta + [r^2\zeta_0 B_0' + 4r^2\zeta_0' B_0 - 2r\zeta_0 B_0]Z_0' + 2r^2\zeta_0 B_0 Z_0'' = 0. \quad (72)$$

Note that if $\Delta = 0$ this becomes Eq. (13), while if $Z_0 = 1$ this becomes Eq. (27). In general, this ODE of first-order in Δ , and—as long Z is not a constant—inhomogeneous. In terms of Δ this ODE can be solved explicitly and the result stated as a new theorem:

Theorem 5.—Suppose $\{\zeta_0, B_0\}$ describes a perfect fluid sphere, and let $Z_0(r)$ be arbitrary. Define

$$\begin{aligned} \Delta(\lambda, r) &= \Delta_0(r) \left\{ \lambda - \int \frac{[(4r^2\zeta_0' B_0 + r^2\zeta_0 B_0' - 2r\zeta_0 B_0)Z_0' + 2r^2\zeta_0 B_0 Z_0'']\{\zeta_0 + r\zeta_0'\}^2}{r^3(r\zeta_0 Z_0)'\zeta_0^2 Z_0^2} \right. \\ &\quad \left. \times \exp\left[-2 \int \frac{(\zeta_0 Z_0)'}{\zeta_0 Z_0} \frac{\zeta_0 Z_0 - r(\zeta_0 Z_0)'}{\zeta_0 Z_0 + r(\zeta_0 Z_0)'} dr\right] dr \right\}, \end{aligned} \quad (73)$$

where

$$\Delta_0(r) = \left(\frac{\zeta_0 Z_0}{\zeta_0 Z_0 + r(\zeta_0 Z_0)'} \right)^2 r^2 \times \exp \left\{ 2 \int \frac{(\zeta_0 Z_0)' \zeta_0 Z_0 - r(\zeta_0 Z_0)'}{\zeta_0 Z_0 \zeta_0 Z_0 + r(\zeta_0 Z_0)'} dr \right\}. \quad (74)$$

Then for all λ , the geometry defined by an arbitrary chosen $Z_0(r)$ and setting

$$ds^2 = -\zeta_0(r)^2 Z_0(r)^2 dt^2 + \frac{dr^2}{B_0(r) + \Delta(\lambda, r)} + r^2 d\Omega^2 \quad (75)$$

corresponds to a perfect fluid sphere. That is, the mapping

$$\mathbf{T}_5(\lambda): \{\zeta_0, B_0\} \mapsto \{\zeta_0 Z_0, B_0 + \Delta(\lambda, \zeta_0)\} \quad (76)$$

takes perfect fluid spheres into perfect fluid spheres.

Note that if $Z_0(r) = 1$ this simply reduces to Theorem 1. Rearranging Eq. (72) in terms of Z_0 leads to a second-order inhomogenous differential equation, which cannot in general be solved for a prescribed Δ , *unless one imposes further constraints*. So further exploration in that direction is moot. There is however a related theorem (which may be easier to understand) in terms of parametric derivatives:

Theorem 6.—Let $\{\zeta(\mu), B(\mu)\}$ denote a one-parameter class of perfect fluid spheres, so that the differential Eq. (4) is satisfied for all μ . Then

$$\begin{aligned} [r(r\zeta)'] \left(\frac{dB}{d\mu} \right)' + [2r^2 \zeta'' - 2(r\zeta)'] \left(\frac{dB}{d\mu} \right) + 2r^2 B \left(\frac{d\zeta}{d\mu} \right)'' \\ + (r^2 B' - 2rB) \left(\frac{d\zeta}{d\mu} \right)' + (rB' - 2B + 2) \left(\frac{d\zeta}{d\mu} \right) = 0. \end{aligned} \quad (77)$$

In particular if $d\zeta/d\mu = 0$ this reduces to the ODE (13), while if $dB/d\mu = 0$ this reduces to the ODE (5). This is simply an alternative viewpoint on the previous theorem, emphasizing the differential equation to be solved.

We again defer extensive discussion to the next section, but at this stage point out that if we invoke Theorem 5 and apply it to Minkowski space, then making the choice $Z_0 = 1 + r^2/a^2$ leads to the general Tolman IV metric—that is:

$$\mathbf{T}_5(\text{Minkowski}; Z_0 = 1 + r^2/a^2) = (\text{Tolman IV}). \quad (78)$$

TABLE IV. Nonseed solutions and their generalizations.

Base	Theorem 1	Theorem 2	Theorem 3	Theorem 4
Tolman IV	Tolman IV	P7	P7	[integral]
B-L	B-L	Kuchb I b	Kuchb I b	[integral]
Heint IIa	Heint IIa	P8	P8	[integral]
Tolman VI	Wyman IIa	Tolman VI	[integral]	[integral]
Kuch1 Ib	Martin 1	Kuch1 Ib	P2	Martin 1
K-O III	Martin 3	K-O III	P1	Martin 3

Even before we systematically start our classification efforts, it is clear that the solution-generating theorems we have established will interrelate *many* of the standard perfect fluid spheres.

E. Formal properties of the linking theorems

Before turning to issues of systematic classification of perfect fluid metrics, we wish to establish a few formal properties of the linking theorems. To simplify the notation, let us define the differential expression

$$\begin{aligned} D(\zeta, B) &\equiv [r(r\zeta)']B' + [2r^2 \zeta'' - 2(r\zeta)']B + 2\zeta \\ &= 2r^2 B \zeta'' + (r^2 B' - 2rB)\zeta' + (rB' - 2B + 2)\zeta. \end{aligned} \quad (79)$$

Then the condition for a perfect fluid sphere is simply

$$D(\zeta, B) = 0. \quad (80)$$

Now define

$$D(\zeta_0, B_0 + \Delta_0) = D(\zeta_0, B_0) + D_1(\zeta_0; \Delta_0). \quad (81)$$

Then it is easy to check that

$$D_1(\zeta_0; \Delta_0) = [r(r\zeta_0)']\Delta_0' + [2r^2 \zeta_0'' - 2(r\zeta_0)']\Delta_0. \quad (82)$$

The ODE for Theorem 1, where we assume $\{\zeta_0, B_0\}$ is a perfect fluid sphere, is then

$$D_1(\zeta_0; \Delta_0) = 0. \quad (83)$$

Now define

$$D(\zeta_0 Z_0, B_0) = Z_0 D(\zeta_0, B_0) + D_2(\zeta_0, B_0; Z_0), \quad (84)$$

then it is easy to check that

$$\begin{aligned} D_2(\zeta_0, B_0; Z_0) &= (r^2 \zeta_0 B_0' + 4r^2 \zeta_0' B_0 - 2r \zeta_0 B_0) Z_0' \\ &+ (2r^2 \zeta_0 B_0) Z_0''. \end{aligned} \quad (85)$$

The ODE for Theorem 2, where we assume $\{\zeta_0, B_0\}$ is a perfect fluid sphere, is then

$$D_2(\zeta_0, B_0; Z_0) = 0. \quad (86)$$

Now let us consider any simultaneous shift in ζ and B , as considered in Theorem 5. We have

$$\begin{aligned} D(\zeta_0 Z_0, B_0 + \Delta_0) &= D(\zeta_0 Z_0, B_0) + D_1(\zeta_0 Z_0; \Delta_0) \\ &= Z_0 D(\zeta_0, B_0) + D_2(\zeta_0, B_0; Z_0) \\ &+ D_1(\zeta_0 Z_0; \Delta_0). \end{aligned} \quad (87)$$

But now let us write

$$D_1(\zeta_0 Z_0; \Delta_0) = Z_0 D_1(\zeta_0; \Delta_0) + D_{12}(\zeta_0; Z_0, \Delta_0), \quad (88)$$

where a brief computation yields

$$\begin{aligned} D_{12}(\zeta_0; Z_0, \Delta_0) &= r^2 \zeta_0 Z_0' \Delta_0' + [2r^2 \zeta_0 Z_0'' + 4r^2 \zeta_0' Z_0' \\ &- 2r \zeta_0 Z_0'] \Delta_0. \end{aligned} \quad (89)$$

Then all in all

$$D(\zeta_0 Z_0, B_0 + \Delta_0) = Z_0 D(\zeta_0, B_0) + Z_0 D_1(\zeta_0; \Delta_0) + D_{12}(\zeta_0; Z_0, \Delta_0) + D_2(\zeta_0, B_0; Z_0). \quad (90)$$

So if ζ_0, B_0 and $\{\zeta_0 Z_0, B_0 + \Delta_0\}$ are both perfect fluid spheres we must have

$$Z_0 D_1(\zeta_0; \Delta_0) + D_{12}(\zeta_0; Z_0, \Delta_0) + D_2(\zeta_0, B_0; Z_0) = 0. \quad (91)$$

This is

$$\begin{aligned} & Z_0 \times (\text{ODE for Theorem 1}) + (\text{cross term}) \\ & + (\text{ODE for Theorem 2}) \\ & = 0. \end{aligned} \quad (92)$$

The cross term vanishes if either $Z_0 = \text{constant}$ or $\Delta_0 = 0$ in which case we recover the usual Theorem 1 and Theorem 2. If we do things in the opposite order, then

$$D(\zeta_0 Z_0, B_0 + \Delta_0) = Z_0 D(\zeta_0, B_0 + \Delta_0) + D_2(\zeta_0, B_0 + \Delta_0; Z_0) \quad (93)$$

$$= Z_0 D(\zeta_0, B_0) + Z_0 D_1(\zeta_0, \Delta_0) + D_2(\zeta_0, B_0 + \Delta_0; Z_0). \quad (94)$$

We now have to compute

$$D_2(\zeta_0, B_0 + \Delta_0; Z_0) = D_2(\zeta_0, B_0; Z_0) + D_{21}(\zeta_0, B_0; Z_0, \Delta_0), \quad (95)$$

and after a brief calculation

$$\begin{aligned} D_{21}(\zeta_0, B_0; Z_0, \Delta_0) &= 2r^2 \zeta_0 \Delta_0 Z_0'' \\ &+ [r^2 \zeta_0 \Delta_0' + 4r^2 \zeta_0' \Delta_0 - 2r \zeta_0 \Delta_0] Z_0' \\ &= D_{12}(\zeta_0, B_0; Z_0, \Delta_0). \end{aligned} \quad (96)$$

Thus the cross term is the same, no matter how you calculate it, and we still have the identity

$$D(\zeta_0 Z_0, B_0 + \Delta_0) = Z_0 D(\zeta_0, B_0) + Z_0 D_1(\zeta_0; \Delta_0) + D_{12}(\zeta_0; Z_0, \Delta_0) + D_2(\zeta_0, B_0; Z_0). \quad (97)$$

Provided ζ_0, B_0 and $\{\zeta_0 Z_0, B_0 + \Delta_0\}$ are both perfect fluid spheres we again deduce

$$Z_0 D_1(\zeta_0; \Delta_0) + D_{12}(\zeta_0; Z_0, \Delta_0) + D_2(\zeta_0, B_0; Z_0) = 0. \quad (98)$$

Now this gives us another way of looking at Theorem 3 and Theorem 4. For Theorem 3 we first apply Theorem 1 so we have the two equations

$$D_1(\zeta_0; \Delta_0) = 0; \quad (99)$$

and

$$D_2(\zeta_0, B_0 + \Delta_0; Z_0) \equiv D_{12}(\zeta_0; Z_0, \Delta_0) + D_2(\zeta_0, B_0; Z_0) = 0. \quad (100)$$

Conversely, for Theorem 4 where we first apply Theorem 2 we see that we need to solve

$$D_2(\zeta_0, B_0; Z_0) = 0; \quad (101)$$

and

$$D_1(\zeta_0 Z_0; \Delta_0) \equiv Z_0 D_1(\zeta_0; \Delta_0) + D_{12}(\zeta_0; Z_0, \Delta_0) = 0. \quad (102)$$

For Theorem 5 we pick Z_0 arbitrarily, and solve the single ODE

$$D_1(\zeta_0 Z_0; \Delta_0) + D_2(\zeta_0, B_0; Z_0) = 0. \quad (103)$$

This is a single first-order linear inhomogeneous ODE for Δ_0 , and hence solvable. (In particular this makes it clear that Theorem 5 is an *inhomogeneous* version of Theorem 1 with a carefully arranged ‘‘source term’’ $D_2(\zeta_0, B_0; Z_0)$. While Theorem 5 is not idempotent it does satisfy the important formal property that:

Lemma 3.—

$$\mathbf{T}_1 \circ \mathbf{T}_5 \triangleq \mathbf{T}_5, \quad (104)$$

which, in particular, tells us that the output from Theorem 5 is never a seed metric.

Applying Theorem 5 we need to solve

$$D_1(\zeta_0 Z_0; \Delta_0) + D_2(\zeta_0, B_0; Z_0) = 0, \quad (105)$$

in order to map

$$\{\zeta_0, B_0\} \mapsto \{\zeta_0 Z_0, B_0 + \Delta_0\}. \quad (106)$$

Now apply Theorem 1 to $\{\zeta_0 Z_0, B_0 + \Delta_0\}$, this means we have to solve the homogeneous ODE

$$D_1(\zeta_0 Z_0; \Delta_1) = 0. \quad (107)$$

But then, using properties of first-order ODEs

$$\begin{aligned} \{D_1(\zeta_0 Z_0; \Delta_0) + D_2(\zeta_0, B_0; Z_0) = 0\} \oplus \{D_1(\zeta_0 Z_0; \Delta_1) = 0\} \\ \Rightarrow \{D_1(\zeta_0 Z_0; \Delta_0 + \Delta_1) + D_2(\zeta_0, B_0; Z_0) = 0\} \end{aligned} \quad (108)$$

which is the ODE from Theorem 5 back again. \square

(The net result of this observation, as we shall see in the next section, is that Theorem 5 can be used to connect one seed metric with the ‘‘descendants’’ of another seed metric.)

III. CLASSIFYING PERFECT FLUID SPHERES

We will now see the power of these transformation theorems (solution-generating theorems) by using them in a number of different ways: to generate several new perfect fluid spheres, to relate various perfect previously known fluid spheres to each other, and to classify the




FIG. 4. Numerical implementations of Theorem 3 and Theorem 4 can be used to investigate their behavior for n applications, as the number of applications becomes large. For specific choices of parameters λ_i , σ_i and ϵ_i successive applications of Theorem 3 and Theorem 4 appear to be approaching a limit, in the sense that both metric components seem to converge toward fixed points.

the Minkowski seed metric results in the Schwarzschild Interior geometry, whereas application of Theorem 1 after Theorem 2 gives us the Martin 3 solution, and the three-parameter Schwarzschild interior solution is quite distinct from the three-parameter Martin 3 solution.

Indeed, it seems as if it is possible to create endless new solutions for a perfect fluid sphere out of the Minkowski metric (or any other of the seed metrics). After several iterations the calculations become more complex, and can no longer be carried out analytically. We then resort to numeric computation to find out whether Theorem 3 and Theorem 4 have some sort of numerical limit, a numerical fixed point, or not. Depending on the choices made for the parameters $(\lambda, \sigma, \epsilon)$ both theorems converge very quickly. Specifically, we used MATLAB to numerically analyze the evolution of $\zeta(r)$ and $B(r)$ after applying Theorem 3 several times to the Minkowski metric. Numerical results are summarized in Fig. 4. This figure indicates that Theorem 3 and Theorem 4 both appear to have well-defined numerical limits, though we have no direct analytic solution for the fixed-point metric.

IV. DISCUSSION

Using Schwarzschild coordinates we have developed several transformation theorems that map perfect fluid spheres into perfect fluid spheres, and have used these transformations as a basis for classifying different types of perfect fluid sphere solutions. If we apply these theo-

rems on a known perfect fluid sphere, different solutions are often obtained. While some of the solutions are already known in the literature, most of them are novel. Therefore, we have developed a tool to generate new solutions for a perfect fluid sphere, which does not require us to directly solve the Einstein equations. (Of course the whole procedure was set up in such a way that we are implicitly and indirectly solving the Einstein equations, but the utility of the transformation theorems is that one does not have to go back to first principles for each new calculation.)

In addition, we have also established several relationships among the generating theorems. Previously, all metrics seemed to have nothing more in common than representing a perfect fluid sphere. We have developed the concept of a seed metric, which is one that cannot be generated by our theorems. Starting with a seed metric and applying our theorems, it is possible to structure, therefore to visualize, the relationship between all metrics generated from a given seed metric in a graph. Based on this example it is possible to create endless new solutions out of the Minkowski metric. We also used a numeric program to investigate whether Theorem 3 and Theorem 4 have fixed-point limits. Both seem to converge very quickly.

In summary, the class of static perfect fluid sphere solutions to the Einstein equations exhibits a deep and somewhat unexpected structure—the transformation theorems of this article provide a new and distinct way of looking at the problem, and yield a new way of viewing the interrelationships between different static fluid spheres.

- [1] M.S.R. Delgaty and K. Lake, *Comput. Phys. Commun.* **115**, 395 (1998).
- [2] M.R. Finch and J.E.F. Skea, "A Review of the Relativistic Static Fluid Sphere" (unpublished); <http://www.dft.if.uerj.br/usuarios/JimSkea/papers/pfrev.ps>
- [3] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations*, (Cambridge University Press, Cambridge, England, 2003).
- [4] D. Martin and M. Visser, *Classical Quantum Gravity* **20**, 3699 (2003).
- [5] H. A. Buchdahl, *Phys. Rev.* **116**, 1027 (1959); *Astrophys. J.* **146**, 275 (1966).
- [6] H. Bondi, *Mon. Not. R. Astron. Soc.* **107**, 410 (1947); **282**, 303 (1964).
- [7] M. Wyman, *Phys. Rev.* **75**, 1930 (1949).
- [8] S. Berger, R. Hojman, and J. Santamarina, *J. Math. Phys. (N.Y.)* **28**, 2949 (1987).
- [9] S. Rahman and M. Visser, *Classical Quantum Gravity* **19**, 935 (2002).
- [10] K. Lake, *Phys. Rev. D* **67**, 104015 (2003).
- [11] D. Martin and M. Visser, *Phys. Rev. D* **69**, 104028 (2004).