

Studying W^+W^- production at the Fermilab Tevatron with SHERPA

Tanju Gleisberg, Frank Krauss, Andreas Schälicke, Steffen Schumann, and Jan-Christopher Winter

Institute for Theoretical Physics, D-01062 Dresden, Germany

(Received 16 April 2005; published 25 August 2005)

The merging procedure of tree-level matrix elements with the subsequent parton shower as implemented in SHERPA will be studied for the example of W boson pair production at the Fermilab Tevatron. Comparisons with fixed order calculations at leading and next-to-leading order in the strong coupling constant and with other Monte Carlo simulations validate once more the impact and the quality of the merging algorithm and its implementation.

DOI: [10.1103/PhysRevD.72.034028](https://doi.org/10.1103/PhysRevD.72.034028)

PACS numbers: 13.85.-t, 13.85.Qk, 13.87.-a

I. INTRODUCTION

Studying the production of W pairs at collider experiments offers a great possibility for tests of the gauge sector of the standard model, that has been extensively investigated by the LEP2 collaborations [1–5]. Tests in this channel are quite sensitive, because there is a destructive interference of two contributions: a t -channel contribution, where both W bosons couple to incoming fermions, and an s -channel contribution, where the W bosons emerge through a triple gauge coupling, either γW^+W^- or ZW^+W^- . New physics beyond the standard model could easily manifest itself, either through new particles propagating in the s channel, like, for instance, a Z' particle in L-R symmetric models [6–9], or through anomalous triple gauge couplings, which could be loop-induced, mediated by heavy virtual particles running in the loop. In [10–12] the most general form of an effective Lagrangian for such interactions has been developed and discussed. Such tests of anomalous triple gauge couplings have been performed both at LEP2 [13–16] and at Tevatron, Run I [17–20] and at Run II [21]. Both scenarios could clearly modify the total cross section or, at least, lead to different distributions of the final state particles. In addition, W pairs, possibly in association with jets, represent a background to a number of relevant other processes, such as the production of top quarks, the production of a Higgs boson with a mass above roughly 135 GeV, or the production of supersymmetric particles, such as charginos or neutralinos [22,23]. Accordingly, there are a number of calculations and programs dealing with this process. At next-to-leading order (NLO) in the strong coupling constant, W pair production has been calculated by [24–26]. In addition, a number of programs have been made available, allowing the user to implement phase space cuts and to generate single events. First of all, there are fixed order calculations. At leading

order (LO), i.e. at tree-level, they are usually performed through automated tools, called matrix element or parton-level generators. Examples for such programs include COMPHEP [27], GRACE/GR@PPA [28,29], MADGRAPH/MADEVENT [30,31], ALPGEN [32], and AMEGIC++ [33]. At NLO, the program MCFM [34] provides cross sections and distributions for this process. Apart from such fixed order calculations, multipurpose event generators such as PYTHIA [35,36] or HERWIG [37,38] play a major role in the experimental analyses of collider experiments. They proved to be extremely successful in describing global features of such processes, like, for instance, the transverse momenta or rapidity distributions of the bosons. They are usually based on exact tree-level matrix elements for the production and decay of the boson pair, supplemented with a parton shower. The latter takes proper care of multiple parton emission and resums the corresponding leading and some of the subleading Sudakov logarithms. In view of the need for increasing precision, recently two approaches have been developed that incorporate higher order corrections into the framework of multipurpose event generators. The first one, called MC@NLO, provides a method to consistently match NLO calculations for specific processes with the parton shower [39,40]. The idea of this approach is to organize the counterterms necessary to cancel real and virtual infrared divergencies in such a way that the first emission of the parton shower is recovered. Of course, this method depends to some extent on the details of the parton shower, and it has some residual dependence on the process in question. So far, MC@NLO has been implemented in conjunction with HERWIG [41] for the following processes: production of W and Z bosons, or pairs of these bosons [39], production of the Higgs boson, production of heavy quarks [40]. An alternative approach is to consistently combine tree-level matrix elements for different multiplicities of additional jets and to merge them with the parton shower. This approach has been presented for the first time for the case of e^+e^- annihilations into jets [42]; later it has been extended to hadronic collisions [43] and it has been reformulated to a merging procedure with a dipole shower in [44]. The idea underlying this method is to separate the kinematical range of parton emission by a k_{\perp} -algorithm

*Electronic address: tanju@theory.phy.tu-dresden.de†Electronic address: krauss@theory.phy.tu-dresden.de<http://www.physik.tu-dresden.de/~krauss/hep/>‡Electronic address: dreas@theory.phy.tu-dresden.de§Electronic address: steffen@theory.phy.tu-dresden.de||Electronic address: winter@physik.tu-dresden.de

[45–47] into a regime of jet production, covered by the appropriate matrix elements, and a regime of jet evolution, covered by the respective shower. Then, the matrix elements are reweighted through Sudakov form factors and hard emissions in the parton shower leading to a jet are vetoed such that there is only a residual dependence on the jet resolution cut. This method is one of the cornerstones of the new event generator SHERPA [48]; it has been validated for the cases of e^+e^- annihilations into jets [49,50] and for the production of single vector bosons at the Fermilab Tevatron [51] and the CERN LHC [52]. In this publication this series of studies will be continued with an investigation of W pair production at the Fermilab Tevatron, Run II, where both W bosons decay leptonically, i.e. $p\bar{p} \rightarrow W^+W^- + X \rightarrow e^+\mu^-\nu_e\bar{\nu}_\mu + X$.¹ Input parameters used throughout this publication and the specifics, how the SHERPA runs have been obtained, are listed in the appendix, see Apps. A and C. After some consistency—including scale variation—checks of the merging algorithm in Sec. II, results obtained with SHERPA will be confronted with those from an NLO calculation provided by MCFM, cf. Sec. III. Then, in Sec. IV some exemplary results of SHERPA are compared with those obtained from other event generators, in particular, with those from PYTHIA and MC@NLO. A summary closes this publication.

II. CONSISTENCY CHECKS

In this section some sanity checks of the merging algorithm for the case of W pair production are presented. For this, first, the dependence of different observables on the key parameters of the merging procedure, namely, the internal matrix-element parton-shower separation scale Q_{cut} and the highest multiplicity n_{max} of included tree-level matrix elements, is examined. Secondly, the sensitivity of the results with respect to changes in the renormalization scale μ_R and the factorization scale μ_F will be discussed. All distributions shown in this section are inclusive results at the hadron level, where restrictive jet and lepton cuts have been applied, for details on the cuts cf. App. C. In all cases, the distributions are normalized to one using the respective total cross section as delivered by the merging algorithm.

A. Impact of the phase space separation cut

First of all, the impact of varying the jet resolution cut Q_{cut} is studied. SHERPA results have been obtained with an inclusive 2jet production sample, i.e. tree-level matrix elements up to two additional QCD emissions have been combined and merged with the parton shower. In all figures presented here the solid line shows the total inclusive result as obtained by SHERPA for the respective resolution cut

Q_{cut} . The reference curve drawn as a dashed line has been obtained as the mean of five different runs, where the resolution cut has been gradually increased, $Q_{\text{cut}} = 10, 15, 30, 50$ and 80 GeV. The shaded (colored) curves represent the contributions stemming from the different matrix-element final-state multiplicities. Results are shown for three different resolution cuts, namely $Q_{\text{cut}} = 15, 30$ and 80 GeV. It should be noted that the change of the rate predicted by the merging procedure under Q_{cut} variation has been found to be very small, although it is a leading order prediction only. Nevertheless, by varying the separation cut between 10 GeV and 80 GeV, the deviation of the total rate amounts to 2.4% only.

As a first result, consider the p_T distribution of the W^+ boson, presented in Fig. 1. The distributions become slightly softer for increasing cuts. However, this observable is very stable under variation of Q_{cut} with maximal deviations on the $\pm 5\%$ level only. The shape of the W^+ boson's p_T is already described at LO (using a parton shower only). As it can be seen from the figure, this LO dominance is nicely kept by the SHERPA approach under Q_{cut} variation. There the 1jet and 2jet contributions are reasonably—for the 80 GeV run, even strongly—suppressed with respect to the leading contribution. In Fig. 2 the transverse momentum spectrum of the W^+W^- system is depicted. Here, deviations show up, but they do not amount to more than $\pm 20\%$. Thus, the QCD radiation pattern depends only mildly on Q_{cut} (indicated by a vertical dashed-dotted line), which at the same time has been varied by nearly 1 order of magnitude. For $Q_{\text{cut}} = 15$ GeV the matrix-element domain is enhanced with respect to the reference resulting in a harder p_T tail. In contrast by using $Q_{\text{cut}} = 80$ GeV the hard tail of the diboson transverse momentum is underestimated with respect to the reference, since the parton shower attached only to the lowest order matrix element starts to fail in the description of high- p_T QCD radiation at $p_T \approx 30$ GeV. At $Q_{\text{cut}} = 80$ GeV a smooth transition is required. The higher order matrix elements then stop the decrease in the p_T prediction. In previous publications it turned out that differential jet rates most accurately probe the merging algorithm, since they most suitably reflect the interplay of the matrix elements and the parton shower in describing QCD radiation below and above the jet resolution cut. Results obtained with the Run II k_{\perp} -algorithm using $R = 1$ are shown for the $1 \rightarrow 0, 2 \rightarrow 1$ and $3 \rightarrow 2$ transition in the left, middle and right panels of Fig. 3, respectively. The value for the internal cut increases from $Q_{\text{cut}} = 15$ GeV (top) to $Q_{\text{cut}} = 80$ GeV (bottom). Compared with the p_T^{WW} spectra, similar characteristics of deviations from the reference curve appear. However, here, they are moderately larger reaching up to $\pm 30\%$. The dashed-dotted vertical line again marks the position of Q_{cut} , which also pictures the separation of the n jet from the $n + 1$ jet contribution. Small holes visible around the respective separation cuts are due to a mismatch

¹Singly resonant diagrams contributing to the parton level processes of $p\bar{p} \rightarrow e^+\mu^-\nu_e\bar{\nu}_\mu + X$ have been included.




FIG. 1 (color online). The p_T distribution of the W^+ boson and its dependence on Q_{cut} , chosen to be 15, 30 and 80 GeV (from top to bottom). The solid line shows the SHERPA prediction obtained with $n_{\text{max}} = 2$, the dashed line is the reference obtained as the mean of different Q_{cut} runs and the shaded (colored) lines underneath indicate the different multiplicity contributions. The lower part of the plots exhibits the normalized difference of the prediction with respect to the reference. Cuts and input parameters are specified in the appendices.




FIG. 2 (color online). The p_T distribution of the W^+W^- system under merging scale variation. The cut indicated through a vertical dashed-dotted line has been chosen as $Q_{\text{cut}} = 15, 30$ and 80 GeV (from top to bottom). The solid line shows the SHERPA prediction obtained with $n_{\text{max}} = 2$, the dashed line is the reference obtained as the mean of different Q_{cut} runs and the shaded (colored) lines underneath indicate the different multiplicity contributions. The lower part of the plots exhibits the normalized difference of the prediction with respect to the reference. Cuts and input parameters are specified in the appendices.




FIG. 3 (color online). Differential $1 \rightarrow 0$ jet rate Q_1 , $2 \rightarrow 1$ jet rate Q_2 and $3 \rightarrow 2$ jet rate Q_3 (left to right) for the SHERPA $n_{\max} = 2$ configuration. The cut has been chosen to be 15, 30 and 80 GeV (from top to bottom). The solid line shows the total result, the dashed line is the reference obtained as the mean of different Q_{cut} runs and the shaded (colored) lines underneath indicate the different multiplicity contributions. The vertical dashed-dotted line indicates the separation cut position. The lower part in all plots pictures the normalized difference of the corresponding prediction with respect to the reference. For input parameters and cuts, see Apps. A and C.

of matrix-element and parton-shower kinematics. For $Q_{\text{cut}} = 80$ GeV these holes are much more pronounced, reflecting the failure of the parton shower in filling the hard p_T emission phase space appropriately. Taken together, the

deviations found are very moderate; however, in certain phase space regions they may reach up to 30%. This is satisfactory, since the merging algorithm guarantees Q_{cut} independence on the leading logarithmic accuracy only.




FIG. 4 (color online). The p_T distribution of the W^+ boson in dependence on the variation of the maximal jet number. The comparison is to a (dashed) reference curve obtained with $n_{\max}^{\text{ref}} = n_{\max} - 1$. The cut has been chosen to be 15 GeV. In both plots the solid line shows the total result obtained with SHERPA. The shaded (colored) lines underneath indicate the different multiplicity contributions. The lower part in both plots visualizes the normalized difference of the corresponding prediction with respect to the reference. For input parameters and cuts, see Apps. A and C.

The residual dependence of the results on Q_{cut} may be exploited to tune the perturbative part of the Monte Carlo event generator.

B. Impact of the maximal number of included matrix elements

The approach of varying the maximal jet number n_{\max} can be exploited to further scrutinize the merging procedure. In all cases considered here, Q_{cut} has been fixed to $Q_{\text{cut}} = 15$ GeV. This maximizes the impact of higher order matrix elements. In spite of this, for very inclusive observables, the rates differ very mildly, the change is less than 2%. In Fig. 4, once more the transverse momentum distribution of the W^+ gauge boson is presented, illustrating that the treatment of the highest multiplicity matrix elements (for more details cf. [50,51]) completely compensates for the missing 2jet matrix element in the $n_{\max} = 1$ case. The behavior is almost unaltered when changing from the $n_{\max} = 1$ to the $n_{\max} = 2$ prediction (cf. the right panel). In contrast, $n_{\max} = 0$ yields a considerably softer distribution (cf. the left panel). Lepton p_T spectra show similar characteristics like the W^+ distribution. However, there are a number of observables, which turned out to be rather stable under the variation of n_{\max} , such as the pseudorapidity spectra of the W^+ boson, the positron and muon or correlations between the leptons, e.g. the $\Delta\phi$ or ΔR distribution. In these cases, deviations turn out to be smaller than $\pm 5\%$ in total, i.e. when considering the change between the pure shower and the inclusive 3jet production performance of SHERPA. Even the pseudorapid-




FIG. 5 (color online). The H_T distribution and its dependence on the variation of the maximal jet number. The separation cut has been chosen to be 15 GeV. The solid line shows the SHERPA prediction obtained with $n_{\max} = 1$, the dashed and the dotted lines stand for the $n_{\max} = 2$ and $n_{\max} = 3$ prediction, respectively; the dashed-dotted curve pictures the pure shower performance of SHERPA starting off with the lowest order matrix element. The lower part of the plot shows the normalized differences with respect to the $n_{\max} = 1$ case. For input parameters and cuts, see Apps. A and C.

ity spectra of the resolved jets are rather unaffected. In contrast, three more observables are presented showing a sizeable ($< \pm 30\%$) or even strong ($\approx \pm 100\%$) dependence on the variation of the maximal jet number. The H_T distribution is depicted in Fig. 5, where throughout this

paper H_T is defined as the scalar sum of all lepton and jet transverse momenta. The inclusive p_T spectra of associated jets are exhibited in Fig. 6, where the upper and lower panel of Fig. 6 show the spectra of the hardest and the second hardest jet, respectively. Owing to the nature of these three observables to be sensitive on extra jet emissions, predictions—as expected—become harder with the increase of n_{\max} . However, a stabilization of the predictions is clearly found with the inclusion of more higher order matrix elements, which describe real QCD emissions.

C. Effects of renormalization and factorization scale variations

In the following the impact of renormalization and factorization scale variations is discussed. For the SHERPA merging approach, this variation (also cf. [52]) is performed by multiplying all scales with a constant factor in all coupling constants and PDFs, which are relevant for the matrix-element evaluation, the Sudakov weights and for the parton-shower evolution. For this study, the SHERPA samples are produced with $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV. In all figures the solid line represents SHERPA's default scale




FIG. 6 (color online). SHERPA predictions of the inclusive p_T of the associated jets considered in dependence on the variation of the maximal jet number. The spectra of the hardest and the second hardest jet are depicted in the upper and the lower panel, respectively. The jet resolution cut has been taken to be 15 GeV. The solid line shows the result of the $n_{\max} = 1$ sample, the dashed and the dotted lines stand for the $n_{\max} = 2$ and $n_{\max} = 3$ sample, respectively; the dashed-dotted curve depicts the pure shower performance. The lower part of the plot shows the normalized differences with respect to the $n_{\max} = 1$ case. For the jet definition, the Run II k_{\perp} -algorithm with $R = 0.7$ and $p_T^{\text{jet}} > 15$ GeV has been used. For more details, see Apps. A and C.




FIG. 7 (color online). The p_T distribution of the W^+ boson under scale variations. All predictions stem from SHERPA with $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV. The solid line shows the prediction under default scale choices for the merging procedure. For the dashed and the dotted curve, all scales for the coupling constants and PDFs have been multiplied by 0.5 and 2.0, respectively. The lower part of the plot presents the normalized differences with respect to the default choice. Input parameters and cuts are given in Apps. A and C.

choices, whereas the dashed and the dotted curve show the outcome for scale multiplications by 0.5 and 2.0, respectively. The total rate as provided by the merging algorithm is again remarkably stable, varying with respect to the default only by $\pm 4.2\%$, thereby increasing for smaller scales. The transverse momentum distribution of the W^+ boson is investigated in Fig. 7. Scale variations slightly distort the shape, shifting it towards harder p_T for smaller scales and vice versa. The effect is more pronounced in the H_T distribution, shown in Fig. 8, and in the transverse momentum distribution of the diboson system, depicted in Fig. 9. However, the deviations maximally found reach up to $\pm 30\%$. In contrast to the findings stated so far, jet transverse momentum spectra do not feature shape distortions under scale variations. The pattern found from these investigations can be explained as follows. The single matrix-element contributions—here the 0jet and 1jet contributions—have their own rate and shape dependencies under scale variations. In their interplay these differences transfer to changing the admixture of the single contributions. Hence, shape modifications can appear as soon as different phase space regions are dominated by a single contribution. This also explains the behavior found for jet




FIG. 8 (color online). The H_T distribution and its dependence on the variation of μ_R and μ_F in the merging prescription. Fixing $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV, the solid line shows the prediction under default scale choices. For the dashed and the dotted curve, all scales for the coupling constants and PDFs have been multiplied by 0.5 and 2.0, respectively. The lower part of the plot presents the normalized differences with respect to the default choice. Input parameters and analysis cuts are given in Apps. A and C.




FIG. 9 (color online). The p_T distribution of the W pair under variation of μ_R and μ_F . Fixing $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV, the solid line shows the prediction under default scale choices. The dashed and the dotted curve is generated when all scales used for the coupling constants and PDFs have been multiplied by 0.5 and 2.0, respectively. The lower part of the plot presents the normalized differences with respect to the default choice. Input parameters (including a primordial k_{\perp} smearing) and cuts are given in Apps. A and C.

p_T s. In the case studied here, they are solely described by the 1jet matrix element with the parton shower attached, thus, their different rates cancel out due to normalization and their shapes are not affected. Taken together, the dependencies found here, together with the ones on Q_{cut} and n_{\max} , yield an estimate for the uncertainty related to the SHERPA predictions.

III. SHERPA COMPARISON WITH MCFM

In this section, the focus shifts from internal sanity checks to comparisons with a full NLO calculation. For this, the MCFM program [34] has been used. In both, MCFM and SHERPA the CKM matrix has been taken diagonal, and no b quarks are considered in the partonic initial state of the hard process. If not stated otherwise, in MCFM the renormalization and factorization scale have been chosen as $\mu_R = \mu_F = M_W$, according to the choice made in [34]. For more details on the input parameters and setups, see Apps. A and B. In the following the results of MCFM are confronted with those of SHERPA (using $Q_{\text{cut}} = 15$ GeV) obtained at the parton-shower level. Furthermore, for this analysis, realistic experimental cuts (cf. App. C) have been




FIG. 10 (color online). Normalized H_T distribution. SHERPA results are shown for $n_{\max} = 1$ (bright solid line) and $n_{\max} = 2$ (bright dashed line) and compared to the QCD NLO result of MCFM (solid line). The LO result with the same scale choice is depicted as a thin dashed line. A difference plot with the MCFM NLO prediction as reference is given within the figure.

applied and all distributions have been normalized to one. First the H_T distribution, depicted in Fig. 10, is considered. Clearly, higher order corrections affect the H_T shape. This is due to two reasons. First of all, the additional QCD radiation may manifest itself as jets, which thus contribute to H_T . Otherwise the additional partons still form a system against which the W pair may recoil. Quanti-

tatively, the inclusion of NLO results in a shift of the H_T distribution at harder values by up to 20%; in SHERPA this trend is amplified by roughly the same amount. The differences between MCFM and SHERPA, however, are due to the different scale choices in both codes. In MCFM all scales have been fixed to $\mu = M_W$, whereas, forced by the merging procedure, in SHERPA the scales are set dynamically. In view of the scale variation results discussed in the previous section for H_T (cf. Figure 8) deviations of this magnitude owing to different scale choices are possible. The impact of scale variations on the shape of the same observable is quantified in Fig. 11. This time, however, the SHERPA result with $n_{\max} = 1$ is compared to NLO results obtained from MCFM with scale choices in the range $\mu_R = \mu_F = M_W \dots 4M_W$ and with a LO result taken at $\mu_R = \mu_F = 2M_W$. Obviously, the smaller choice of scale results in the MCFM outcome to be closer to the one of SHERPA. As expected, in comparison to the scale variation results found for SHERPA, the shape uncertainties of the full NLO prediction due to varying the scales are smaller. In Fig. 12, H_T is depicted again, this time for the case of exclusive $p\bar{p} \rightarrow e^+ \mu^- \nu_e \bar{\nu}_\mu$ production. There, the real part of the NLO correction in MCFM is constrained such that it does not produce an extra jet (for jet definition, see App. C). In




FIG. 11 (color online). Normalized H_T distribution. Here both, the renormalization and factorization scale of the NLO calculation have been varied in the range $\mu_R = \mu_F = M_W \dots 4M_W$, indicated by the shaded area. These MCFM results are compared with the leading order result at $\mu_R = \mu_F = 2M_W$ (thin dashed line) and with the result of SHERPA where $n_{\max} = 1$ (bright solid line). The lower part of the plot shows the normalized differences with respect to the SHERPA result.

SHERPA the 0jet matrix element with the parton shower attached is considered exclusively, i.e. the parton shower is now forced not to produce any jet at all. In this case, the higher order corrections lead to a softer H_T distribution




FIG. 12 (color online). Normalized H_T distribution in exclusive $p\bar{p} \rightarrow e^+ \mu^- \nu_e \bar{\nu}_\mu$ production. The SHERPA result (bright solid line) is obtained with $n_{\max} = 0$ and a parton shower constrained not to produce any extra jets. This result is compared with the MCFM result at NLO in α_s (solid line) and with the LO result (thin dashed line). The latter two are taken for the default scale choices. The lower part of the plot shows the normalized differences with respect to the NLO result obtained from MCFM.




FIG. 13 (color online). Normalized p_T distribution of the W pair. The MCFM result at $\mu_R = \mu_F = M_W$ (thick line) is contrasted with the predictions made by SHERPA both at the matrix-element level (dark solid line) and at the parton-shower level with $n_{\max} = 0$ (bright dotted line) and $n_{\max} = 1$ (bright solid line). A primordial k_\perp smearing has been used to obtain the SHERPA shower results.

compared to the leading order prediction, and the results of MCFM and SHERPA show the same deviations as before (cf. Figure 10).

The effect of QCD radiation is best observed in the p_T distribution of the W pair, depicted in Fig. 13. Clearly, without any radiation, the p_T of the W pair is exactly zero, and only the emission of partons leads to a recoil of the diboson system. In the NLO calculation of MCFM, however, the spectrum is therefore described at lowest order, in this particular case taken at $\mu_R = \mu_F = M_W$. In contrast, in the SHERPA matrix-element result, subjected to the explicit jet cut, Sudakov form factors and α_s reweighting are applied with a variable scale choice, explaining the differences between the two matrix-element type results in this figure. Contrasting this with the parton-shower approach, it is clear that parton emission through the shower alone is not sufficient to generate sizeable p_T of the W pair in the hard region. For this, the corresponding matrix element has to be employed, leading to a very good agreement with the MCFM outcome in the high- p_T tail of the distribution. In the soft regime the result of the bare MCFM matrix element is unphysical. Because of the cascade emission of soft and collinear partons, SHERPA accounts for resummation effects, which clearly yield the depopulation of the softest- p_T region.

Another way to look at the effects of QCD radiation is to consider the relative angle between the two W bosons,² see

²The angle is measured in the frame, where the W^+W^- system rests at the beam axis, i.e. the diboson system is corrected on its initial \hat{z} boost.




FIG. 14 (color online). Normalized $\Delta\Phi_{WW}^*$ distribution of the W boson system. The MCFM result (thick line) is contrasted with results from SHERPA at the parton-shower level with $n_{\max} = 0$ (bright dotted line), $n_{\max} = 1$ (bright solid line) and $n_{\max} = 2$ (bright dashed line). Again a primordial k_\perp smearing has been used. Additionally, the dashed curve represents a prediction obtained with MC@NLO. The lower part of the plot shows the normalized differences with respect to the result of MCFM.

Fig. 14. Of course, when they decay into leptons plus neutrinos this is not an experimental observable, on the generator level, however, it is very nice to visualize the effect of QCD radiation in this way. Without any QCD radiation, the two W s would be oriented back-to-back, at $\Delta\Phi_{WW}^* = \pi$. Including QCD radiation, this washes out, as depicted in the figure. Again, resummation effects alter the result of the matrix element alone by decreasing the amount of softest radiation, this time corresponding to the back-to-back region around $\Delta\Phi^* \approx \pi$. The effect of high- p_T radiation can be clearly seen for small $\Delta\Phi^*$ by comparing the different n_{\max} predictions of SHERPA. The larger n_{\max} is chosen, the harder the prediction for small $\Delta\Phi^*$. On the other hand to better value the influence of the parton shower a prediction made by MC@NLO (see App. B) has been included. For a wide region of $\Delta\Phi^*$, it well agrees with the SHERPA result for $n_{\max} = 1$.

Figures 15 and 16 exhibit the transverse momentum distributions of the W^+ and of the e^+ produced in its decay, respectively. Only mild deviations less than 10% between MCFM and SHERPA are found, which again can be traced back to different scale choices in both approaches. These differences recur as and, therefore, explain part of the deviations found in the H_T spectrum, cf. Fig. 10. As expected, the inclusion of the 2jet contribution in SHERPA gives no further alterations of the $n_{\max} = 1$ result. Of course, the different radiation patterns also have some minor effects on the η distribution of the W^+ depicted in Fig. 17. In the $\Delta R_{e\mu}$ distribution presented in Fig. 18, the NLO result of MCFM and the parton-shower level results of




FIG. 15 (color online). Normalized transverse momentum distribution of the W^+ boson. The results of SHERPA for $n_{\max} = 1$ (bright solid line) and for $n_{\max} = 2$ (bright dashed line) are compared with the QCD NLO result obtained by MCFM (solid line) and with the LO result (thin dashed line) for the default scale choices, i.e. $\mu_R = \mu_F = M_W$. Within the plot the normalized differences with respect to the NLO result of MCFM are given.

SHERPA are in nearly perfect agreement with each other. Higher order effects tend to change the shape of the LO prediction with respect to the NLO one by roughly 10%. The interesting observation here is that this change is




FIG. 16 (color online). Normalized transverse momentum distribution of the e^+ produced in the decay of the W^+ . The results of SHERPA for $n_{\max} = 1$ (bright solid line) and for $n_{\max} = 2$ (bright dashed line) are confronted with the QCD NLO result obtained by MCFM (solid line) and with the LO result (thin dashed line). For the latter two, the scales are again fixed according to the default choices, i.e. $\mu_R = \mu_F = M_W$. Within the plot the normalized differences with respect to the NLO result of MCFM are shown.




FIG. 17 (color online). Normalized η distribution of the W^+ boson. The SHERPA results for $n_{\max} = 1$ (bright solid line) and $n_{\max} = 2$ (bright dashed line) are confronted with those of MCFM (solid line) and with the LO result (thin dashed line). Again, in the latter two the scales are chosen as $\mu_R = \mu_F = M_W$. The normalized differences with respect to the NLO result of MCFM are also shown.

seemingly not related to the transverse hardness of a jet system against which the W pair recoils. This gives rise to the assumption that the change with respect to the LO result is due to some altered spin structure in the $2 \rightarrow 5$ matrix element.




FIG. 18 (color online). Normalized ΔR distribution between the two charged leptons, the positron and the muon, emerging from the W decays. SHERPA results for $n_{\max} = 1$ (bright solid line) and $n_{\max} = 2$ (bright dashed line) are compared to those predicted by MCFM (solid line). The LO result with the same scale choice, is shown as a dashed line. The lower part of the plot shows the normalized differences with respect to the NLO result of MCFM.




FIG. 19 (color online). Normalized H_T distribution obtained from PYTHIA (dotted line), MC@NLO (dashed line) and SHERPA (solid line). For the generation of the SHERPA sample, $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV have been chosen. The lower part of the plot exhibits the normalized differences with respect to the SHERPA prediction. Input parameters and the employed cuts are specified in the Apps. A and C .

IV. COMPARISON WITH OTHER EVENT GENERATORS

In this section a comparison of SHERPA with other hadron-level event generators, in particular PYTHIA and MC@NLO will be discussed. Details on how their respective samples have been produced can be found in the Apps. A and B. The SHERPA samples have been generated with $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV. The comparison is again on inclusive distributions—normalized to one—under the influence of realistic experimental cuts, for details see App. C.

A. Comparison of the QCD activity

As before, the starting point is the discussion of the radiation activity predicted by the various codes. In Fig. 19, results for the H_T observable obtained from PYTHIA, MC@NLO and SHERPA are displayed. The predictions of the former two codes nicely agree with each other. Similar to the SHERPA MCFM comparison, SHERPA again predicts a slightly harder spectrum, with relative deviations of up to 20%. Closer inspection of the reason for the differences in the H_T spectrum reveals that the agreement of PYTHIA and MC@NLO is presumably a little bit accidental. A first hint into that direction can be read off Fig. 20, where the p_T spectrum of the W pair is displayed. In the region of low p_T (up to 100 GeV), the results of MC@NLO




FIG. 20 (color online). Normalized p_T distribution of the W^+W^- system. Results from PYTHIA (dotted line), MC@NLO (dashed line) and SHERPA (solid line) are compared. For the generation of the latter, $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV have been chosen. The lower part of the plot presents the normalized differences with respect to the SHERPA prediction. Input parameters (including a primordial k_{\perp} smearing) and the employed cuts are specified in the Apps. A and C .




FIG. 21 (color online). Difference of the scalar transverse momenta of the two W bosons, $|p_T^{W^+} - p_T^{W^-}|$. The predictions compared are: PYTHIA given as a dotted curve, MC@NLO depicted by the dashed line and SHERPA in inclusive 1jet production at $Q_{\text{cut}} = 15$ GeV drawn as a bright solid line as well as SHERPA in pure shower performance shown as a solid line. The lower part of the plot shows the normalized differences with respect to the SHERPA prediction with $n_{\max} = 1$. Input parameters and the employed cuts are summarized in the Apps. A and C .




FIG. 22 (color online). Transverse momentum distributions of the associated jets, in the upper panel, the inclusive p_T of the hardest jet is depicted, whereas in the lower panel that one of the second hardest jet is displayed. Again, results from PYTHIA are given by the dotted lines, MC@NLO results are represented as dashed lines and SHERPA results are the solid lines. For the generation of the latter, $n_{\max} = 1$ and $Q_{\text{cut}} = 15$ GeV have been used. The top solid line in the lower panel corresponds to the SHERPA result obtained with $n_{\max} = 2$. The lower part of both plots shows the normalized differences with respect to the SHERPA $n_{\max} = 1$ performance. The input parameters and the employed cuts are summarized in the Apps. A and C.

and SHERPA are in fairly good agreement,³ and sizeable differences larger than 10% appear only for $p_T > 100$ GeV. In contrast, the PYTHIA result for this observable shows a significant enhancement of the low- p_T region and stays well below the other predictions for $p_T > 10$ GeV. This comparison of the three differential cross sections

³Apart from the very soft region, where the difference is due to parton-shower cutoff effects in HERWIG.

clearly underlines that the three codes differ in their description of the QCD emissions.

Figure 21 depicts the norm of the scalar difference of the transverse momenta of the W^+ and W^- gauge boson, $|p_T^{W^+} - p_T^{W^-}|$. This observable is sensitive to higher order effects, since at LO it merely has a delta peak at $p_T = 0$ GeV. Again, the hardest prediction is delivered by SHERPA with $n_{\max} = 1$, results from MC@NLO, PYTHIA, and the pure shower performance of SHERPA are increasingly softer. For $|\Delta p_T| > 60$ GeV, this observable seems to depend more and more on the quality of modelling the hardest emission, which is intrinsically better described by MC@NLO and by SHERPA with $n_{\max} = 1$. The fact that the PYTHIA shower performs better than the pure SHERPA shower for high p_T differences can be traced back to the choice of starting scale for the shower evolution, which is either $s_{p\bar{p}}$ (PYTHIA) or s_{WW} (SHERPA).

In fact, differences appear in the p_T distributions of the hardest two jets, see Fig. 22. The upper part of this figure depicts the transverse momentum spectrum of the hardest jet. Surprisingly, although MC@NLO contains a matrix element for the emission of an extra jet, its p_T distribution is considerably softer (by up to 40%) than the result of SHERPA generated with $n_{\max} = 1$. This trend is greatly amplified when going to the spectrum of the second hardest jet. There, clear shape differences of the order of a factor 2 between the SHERPA 1jet sample and MC@NLO show up for $p_T \approx 180$ GeV. The surprise according to this figure is that PYTHIA and SHERPA using $n_{\max} = 1$ almost agree on the p_T distribution of the second jet, although they were different for the hardest jet. At that point it should be noted that the second jet in both cases, PYTHIA and SHERPA with $n_{\max} = 1$, is produced by the parton shower only. Given the drastically larger shower start scale of PYTHIA, it seems plausible to achieve to some extent a compensation for the intrinsic parton-shower deficiencies in filling the hard emission phase space.⁴ However, in the very moment, SHERPA events are generated with appropriate matrix elements, i.e. with $n_{\max} = 2$, this distribution is dramatically different for the three codes with deviations larger than a factor 2 for $p_T \approx 120$ GeV.

Taken together, these findings hint that the three codes differ in their modelling of the QCD activity, especially in those of the hardest QCD emission. For MC@NLO and SHERPA the latter can be traced back to the different ansatz in including the matrix element for this emission, where again different scale choices may trigger effects on the 20% level.

B. Comparison of lepton observables

Finally, the leptons in the final state as described by the three event generators PYTHIA, MC@NLO and SHERPA will

⁴PYTHIA's ability to account for harder second jets with respect to MC@NLO is a hint for the similarity of their H_T predictions.




FIG. 23 (color online). Normalized p_T spectrum of the positron. Results of PYTHIA (dotted line) and SHERPA (solid line) including spin correlations are confronted with those obtained from MC@NLO (dashed line) and with results from SHERPA, where spin correlations have been switched off (bright dashed line). All predictions are generated without the use of cuts. The vertical dashed-dotted line is added to indicate the position of the usually employed lepton p_T cut. For input parameters, see App. A. The lower part of the plot shows the normalized differences with respect to the SHERPA prediction including spin correlations.

be investigated. There, some significant differences appear between SHERPA and PYTHIA on the one hand, and MC@NLO on the other hand. These differences are due to the fact that at the moment spin correlations of the W decay products are not implemented in MC@NLO.⁵ To validate that effects are indeed due to the lack of spin correlations, SHERPA samples have been prepared, where these correlations are artificially switched off. Furthermore, in order to quantify these effects without any bias, results have been obtained without the application of any lepton and jet cuts.

The impact of the lack of spin correlations already becomes visible in one-particle observables, such as the p_T or the η spectrum of the positron produced in the W^+ decay. These are shown in Figs. 23 and 24, respectively. Confronting the two methods with each other, which correctly respect spin correlations, for the transverse momentum distribution of the e^+ , the following pattern is revealed. Because of the consistent inclusion of higher order tree-level matrix elements, the SHERPA $n_{\max} = 1$ setup produces a considerably harder spectrum than PYTHIA. In contrast, the distributions with no spin correlations both result in an even harder high- p_T tail. They agree

⁵This situation is currently being cured by the authors of MC@NLO who prepare a new version of their code including spin correlations [53].




FIG. 24 (color online). Normalized η spectrum of the positron. Results of PYTHIA (dotted line) and SHERPA (solid line) including spin correlations are compared with those obtained from MC@NLO (dashed line) and with results from SHERPA, where spin correlations have been switched off (bright dashed line). All predictions are generated without any restriction. The vertical dashed-dotted lines are added to indicate the position of the usually employed lepton η cuts. For input parameters, see App. A. The lower part of the plot shows the normalized differences with respect to the SHERPA prediction including spin correlations.




FIG. 25 (color online). Normalized $\Delta\phi_{e\mu}$ distribution. Results of PYTHIA (dotted line) and SHERPA (solid line) including spin correlations are compared with those obtained from MC@NLO (dashed line) and with results from SHERPA, where spin correlations have been switched off (bright dashed line). All predictions are obtained without the use of cuts. For input parameters, see App. A. The lower part of the plot shows the normalized differences with respect to the SHERPA prediction including spin correlations.




FIG. 26 (color online). Normalized $\Delta R_{e\mu}$ distribution. Results of PYTHIA (dotted line) and SHERPA (solid line) including spin correlations are compared with those obtained from MC@NLO (dashed line) and with results from SHERPA, where spin correlations have been eliminated (bright dashed line). All predictions are obtained without the use of cuts. For input parameters, see App. A. The lower part of the plot shows the normalized differences with respect to the SHERPA prediction including spin correlations.

quite well up to $p_T = 60$ GeV, hence, this coincidence may be assigned to the lack of spin correlations in the gauge boson decays. Above that region, the MC@NLO spectrum again becomes softer with respect to the SHERPA prediction where the spin correlations have been

eliminated. The fact that all four predicted distributions alter in their shape is not solely triggered by the different spin correlation treatments, again, the different descriptions of QCD radiation clearly contribute to the deviations found. In contrast, a simpler pattern is found for the aforementioned η distribution of the e^+ . The results of PYTHIA and SHERPA with spin correlations on the one hand and of MC@NLO and SHERPA without spin correlations on the other hand show perfect agreement. Differences between the two spin correlation treatments may, thereby, easily reach up to 40%.

The influence of spin correlations can also be seen in observables based on two particle correlations. As two illustrative examples take the $\Delta\phi$ and the ΔR distribution of the e^+ and the μ^- produced in the decay of the two W bosons. Again, the corresponding spectra, which have been exhibited in Figs. 25 and 26, differ significantly in shape depending on whether spin correlations are taken into account or not.

The discussion of the impact of spin correlations is completed by exploring the influence of the application of experimental cuts (cf. App. C) on the shape of certain spectra. It is clear that superimposing specific jet and lepton cuts strongly affects the event sample. Here, the cuts are mainly on the η and the p_T of the leptons. In turn their distributions alter. The characteristics found for the cutfree case are not substantially changed by the applied cuts and by the renormalization of the spectra according to these given cuts indicated by the vertical lines in the Figs. 23 and 24. More interestingly, however, these distributions drive alterations to secondary observables. In the two-particle correlations mentioned before, the effects already present without applying cuts are enforced. The slopes of the $\Delta\phi$ distributions increase, amplifying the




FIG. 27 (color online). In the left and middle panel the p_T spectrum of the W^+ before and after the application of cuts is depicted, respectively. The right panel exhibits the η distribution of the W^+ under the influence of these cuts. The predictions compared are: PYTHIA (dotted line), SHERPA (solid line), MC@NLO (dashed line) and SHERPA without correlations in the boson decays (bright dashed line). For input parameters, see App. A. The lower part of the plots shows the normalized differences with respect to the SHERPA prediction including spin correlations.

difference between both sets of predictions, the ones with and without spin correlations. The main change in the ΔR spectrum is an additional deviation between 0.2 (the cut) and 2.0, such that now the no-spin-correlation results are roughly 20% above the other ones. The case is different for the pseudorapidity distribution of the W^+ boson. Without the application of cuts one starts off distributions that agree on the 10% level. This is severely changed by the introduction of the cuts, see the rightmost panel of Fig. 27. In contrast to the aforementioned two-particle correlations, here the predictions without spin correlations are well separated from the other ones only after the application of the cuts. As a last example, consider the transverse momentum distribution of the W^+ boson. Both types of predictions stemming from uncutted (left panel) and from samples analyzed with cuts (middle panel) are pictured in Fig. 27. The inclusion of cuts apparently brings MC@NLO and SHERPA including the full correlations into good agreement, but this clearly happened accidentally.

To summarize, the examples shown here, clearly hint that the superposition of spin correlations (or their absence) together with cuts triggers sizeable effects in both types of observables, such that have already shown deviations in the absence of cuts and, more crucially, such that have not. In specific cases, such as the p_T spectrum of the W^+ , this may possibly lead to misinterpretations of the results.

V. CONCLUSION

In this work, the merging procedure for multiparticle tree-level matrix elements and the parton shower implemented in SHERPA has been further validated; this time, the case of W pair production at the Fermilab Tevatron has been considered. First, it has been shown that the results obtained with SHERPA are widely independent of specific merging procedure details such as the choice of the merging scale and, for sufficiently inclusive observables, the number of extra jets covered by the tree-level matrix elements. In addition, it has been shown that the specific form of the spectra produced by SHERPA is nearly independent—with deviations less than 20%—of the choice of the factorization scale and the renormalization scale.

Having established the self-consistency of the SHERPA results, they have been compared to those from an NLO calculation provided through MCFM. There, good agreement of the two codes has been found, again on the 20% level. Thus it is fair to state that the SHERPA results for the shapes are within theoretical errors consistent with an NLO calculation. The inclusion of the parton shower connected with specific scale choices in SHERPA, however, produces a surplus of QCD radiation with respect to the single parton emission in the real part of the NLO correction in MCFM. Finally, the results of SHERPA have been compared with those of other hadron-level event generators, namely, with PYTHIA and MC@NLO. In this comparison it turned out that

SHERPA predicts a significant increase of QCD radiation with respect to the other two codes. For the p_T spectra of jets accompanying the two W bosons, the differences are dramatic in the high- p_T tails. In addition, the impact of spin correlations has been quantified. In the observables considered here, it reaches 20% . . . 50%. This may be even larger than the impact of higher order corrections.

ACKNOWLEDGMENTS

The authors are very grateful to Gerhard Soff. Without his constant encouragement and valuable support, they could not have gone so far in doing their research work. The authors would like to thank Stefan Höche for valuable collaboration on the development of SHERPA. Furthermore, they would like to thank Marc Hohlfeld (D0 Collaboration) for pleasant conversation on the experimental aspects of this work. The authors are also indebted to Torbjörn Sjöstrand, John Campbell, Tim Stelzer, and Stefano Frixione for helpful advice. Financial support by BMBF, DESY, and GSI is gratefully acknowledged.

APPENDIX A: INPUT PARAMETERS OF SHERPA

All SHERPA studies have been carried out with the cteq6l PDF set [54]. The value of α_s has been chosen according to the corresponding value of the selected PDF, namely $\alpha_s = 0.118$. The running of the strong coupling constant is determined by the corresponding two-loop equation, except for the SHERPA MCFM comparison. There an one-loop running has been employed for α_s . Jets or initial partons are defined by gluons and all quarks but the top quark; this one is allowed to appear within the matrix elements only through the coupling of the W boson with the b quark. In the SHERPA MCFM comparison SHERPA runs, however are restricted to the light-flavour sector, i.e. the g, d, u, s, c sector. In the matrix-element calculation the quarks are taken massless, only the shower will attach current masses to them. The shower cutoffs applied are 2 GeV and 1 GeV for the initial and the final-state emissions, respectively. If explicitly stated a primordial k_\perp Gaussian smearing has been employed with both, mean and standard deviation being equal to 0.8 GeV. The standard model input parameters are:

$$\begin{aligned} m_W &= 80.419 \text{ GeV}, & \Gamma_W &= 2.06 \text{ GeV}, \\ m_Z &= 91.188 \text{ GeV}, & \Gamma_Z &= 2.49 \text{ GeV}, \\ G_\mu &= 1.16639 \times 10^{-5} \text{ GeV}^{-2}, \\ \sin^2\theta_W &= 1 - m_W^2/m_Z^2, & \alpha_s &= 0.118. \end{aligned} \tag{A1}$$

The electromagnetic coupling is derived from the Fermi constant G_μ according to

$$\alpha_{\text{em}} = \frac{\sqrt{2}G_\mu M_W^2 \sin^2\theta_W}{\pi}. \tag{A2}$$

The constant widths of the electroweak gauge bosons are introduced through the fixed-width scheme. The CKM matrix has been always taken diagonal.

APPENDIX B: SETUPS FOR MCFM, MC@NLO AND PYTHIA

MCFM—The program version employed is MCFM V4.0. The process chosen is NPROC=61. The investigations have been restricted to the d, u, s, c quark sector. The PDF set used is cteq6l. The default scheme for defining the electroweak couplings has been used and their input values have been adjusted with the corresponding parameter settings given for SHERPA. The renormalization scale and the factorization scale are fixed and set to $\mu_R = \mu_F = M_W$.

MC@NLO—The program version used is MC@NLO 2.31. The process number is taken as IPROC=-12850, so that the underlying event has not been taken into consideration. The two W boson decays into leptons are steered by the two MODBOS variables being set to 2 and 3 for the first and the second choice, respectively. The lepton pairs have been generated in a mass window of

$$M_W - 40\Gamma_W < m_{l\nu} < M_W + 40\Gamma_W. \quad (\text{B1})$$

Again, the cteq6l PDF set as provided by MC@NLO's own PDF library is used. The weak gauge boson masses and widths are aligned to the settings used for the previous codes. All other parameters have been left unchanged with respect to their defaults.

PYTHIA—The PYTHIA version used is 6.214. The process $p\bar{p} \rightarrow W^+W^- + X$ is selected through MSUB(25)=1. The specific decay modes of the two W 's are picked by putting MDME(206,1)=2 and MDME(207,1)=3, where all other available modes are set to zero. The possibility of parton-shower emissions rights up to the limit, which has been proven to be

more convenient for jet production [55], is achieved with MSTP(68)=2. This increases the IS shower start scale in PYTHIA to $\sqrt{s} = 1960$ GeV and accounts for a reasonably higher amount of hard QCD radiation. For all comparisons here, the underlying event is switched off, other parameters are left to their default.

APPENDIX C: PHASE SPACE CUTS

Two different analyses are used for the comparisons of the results obtained throughout this publication. A simple analysis has been taken to verify the pure behavior of the considered programs. For this case, only jets are analyzed utilizing the Run II k_{\perp} clustering algorithm defined in [56] with a pseudocone size of $R = 1$. The jet transverse momentum has to be greater than 15 GeV. For more realistic experimental scenarios, an analysis applying jet and lepton cuts has been availed. Then, the pseudocone size of the jet algorithm has been set to $R = 0.7$, and the jets have to fulfil the following constraints on the pseudorapidity and the transverse momentum,

$$|\eta^{\text{jet}}| < 2.0, \quad p_T^{\text{jet}} > 15 \text{ GeV}. \quad (\text{C1})$$

For the charged leptons the cuts on these observables are given by

$$|\eta^{\text{lep}}| < 1.0, \quad p_T^{\text{lep}} > 20 \text{ GeV}, \quad (\text{C2})$$

however, a cut on the missing transverse energy has not been introduced. There is a final selection criteria corresponding to the separation of the leptons from each other and from the jets,

$$\Delta R_{ll} > 0.2, \quad \Delta R_{lj} > 0.4. \quad (\text{C3})$$

-
- [1] LEP Collaboration, hep-ex/0312023.
 - [2] R. Barate *et al.* (ALEPH Collaboration), Phys. Lett. B **484**, 205 (2000).
 - [3] G. Abbiendi *et al.* (OPAL Collaboration), Phys. Lett. B **493**, 249 (2000).
 - [4] J. Abdallah *et al.* (DELPHI Collaboration), Eur. Phys. J. C **34**, 127 (2004).
 - [5] P. Achard *et al.* (L3 Collaboration), Phys. Lett. B **600**, 22 (2004).
 - [6] J.C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974).
 - [7] R.N. Mohapatra and J.C. Pati, Phys. Rev. D **11**, 566 (1975).
 - [8] R.N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165 (1981).
 - [9] E. Eichten, I. Hinchliffe, K. D. Lane, and C. Quigg, Rev. Mod. Phys. **56**, 579 (1984); **58**, 1065 (1986).
 - [10] K.J.F. Gaemers and G.J. Gounaris, Z. Phys. C **1**, 259 (1979).
 - [11] K. Hagiwara, R. D. Peccei, D. Zeppenfeld, and K. Hikasa, Nucl. Phys. **B282**, 253 (1987).
 - [12] M.S. Bilenky, J.L. Kneur, F.M. Renard, and D. Schildknecht, Nucl. Phys. **B409**, 22 (1993).
 - [13] P. Abreu *et al.* (DELPHI Collaboration), Phys. Lett. B **459**, 382 (1999).
 - [14] A. Heister *et al.* (ALEPH Collaboration), Eur. Phys. J. C **21**, 423 (2001).
 - [15] G. Abbiendi *et al.* (OPAL Collaboration), Eur. Phys. J. C **33**, 463 (2004).
 - [16] P. Achard *et al.* (L3 Collaboration), Phys. Lett. B **586**, 151 (2004).
 - [17] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **75**, 1017 (1995).
 - [18] S. Abachi *et al.* (D0 Collaboration), Phys. Rev. D **56**, 6742

- (1997).
- [19] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. D **58**, 031102 (1998).
- [20] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. D **58**, 051101 (1998).
- [21] D. Acosta *et al.* (CDF Collaboration), hep-ex/0501050.
- [22] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. Lett. **80**, 442 (1998).
- [23] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **80**, 5275 (1998).
- [24] J. Ohnemus, Phys. Rev. D **44**, 1403 (1991).
- [25] S. Frixione, Nucl. Phys. **B410**, 280 (1993).
- [26] J. Ohnemus, Phys. Rev. D **50**, 1931 (1994).
- [27] A. Pukhov *et al.*, hep-ph/9908288.
- [28] T. Ishikawa, T. Kaneko, K. Kato, S. Kawabata, Y. Shimizu, and H. Tanaka (MINAMI-TATEYA Group Collaboration), KEK Report No. KEK-92-19.
- [29] K. Sato *et al.*, in *Proceedings of the Seventh International Workshop on Advanced Computing and Analysis Techniques in Physics Research*, edited by P.C. Bhat and M. Kasemann (ACAT, Batavia, 2000); P.C. Bhat and M. Kasemann, AIP Conference Proceedings No. 583 (AIP, New York, 2001), p. 214.
- [30] T. Stelzer and W.F. Long, Comput. Phys. Commun. **81**, 357 (1994).
- [31] F. Maltoni and T. Stelzer, J. High Energy Phys. 02 (2003) 027.
- [32] M.L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A.D. Polosa, J. High Energy Phys. 07 (2003) 001.
- [33] F. Krauss, R. Kuhn, and G. Soff, J. High Energy Phys. 02 (2002) 044.
- [34] J.M. Campbell and R.K. Ellis, Phys. Rev. D **60**, 113006 (1999).
- [35] T. Sjöstrand, P. Eden, C. Friberg, L. Lönnblad, G. Miu, S. Mrenna, and E. Norrbin, Comput. Phys. Commun. **135**, 238 (2001).
- [36] T. Sjöstrand, L. Lönnblad, and S. Mrenna, hep-ph/0108264.
- [37] G. Corcella *et al.*, J. High Energy Phys. 01 (2001) 010.
- [38] G. Corcella *et al.*, hep-ph/0210213.
- [39] S. Frixione and B.R. Webber, J. High Energy Phys. 06 (2002) 029.
- [40] S. Frixione, P. Nason, and B.R. Webber, J. High Energy Phys. 08 (2003) 007.
- [41] S. Frixione and B.R. Webber, hep-ph/0402116.
- [42] S. Catani, F. Krauss, R. Kuhn, and B.R. Webber, J. High Energy Phys. 11 (2001) 063.
- [43] F. Krauss, J. High Energy Phys. 08 (2002) 015.
- [44] L. Lönnblad, J. High Energy Phys. 05 (2002) 046.
- [45] S. Catani, Y.L. Dokshitzer, M. Olsson, G. Turnock, and B.R. Webber, Phys. Lett. B **269**, 432 (1991).
- [46] S. Catani, Y.L. Dokshitzer, and B.R. Webber, Phys. Lett. B **285**, 291 (1992).
- [47] S. Catani, Y.L. Dokshitzer, M.H. Seymour, and B.R. Webber, Nucl. Phys. **B406**, 187 (1993).
- [48] T. Gleisberg, S. Höche, F. Krauss, A. Schälicke, S. Schumann, and J. Winter, J. High Energy Phys. 02 (2004) 056.
- [49] F. Krauss, A. Schälicke, and G. Soff, hep-ph/0503087.
- [50] A. Schälicke and F. Krauss, hep-ph/0503281.
- [51] F. Krauss, A. Schälicke, S. Schumann, and G. Soff, Phys. Rev. D **70**, 114009 (2004).
- [52] F. Krauss, A. Schälicke, S. Schumann, and G. Soff, hep-ph/0503280.
- [53] S. Frixione (private communication).
- [54] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P. Nadolsky, and W.K. Tung, J. High Energy Phys. 07 (2002) 012.
- [55] G. Miu and T. Sjöstrand, Phys. Lett. B **449**, 313 (1999).
- [56] G.C. Blazey *et al.*, hep-ex/0005012.