Solar neutrinos and 1-3 leptonic mixing

Srubabati Goswami¹ and Alexei Yu. Smirnov^{2,3}

¹Harish-Chandra Research Institute, Chhatnag Road, Jhusi, Allahabad 211 019, India

²The Abdus Salam International Centre for Theoretical Physics, I-34100, Trieste, Italy

³Institute for Nuclear Research, Russian Academy of Scienses, Moscow, Russia

(Received 16 February 2005; revised manuscript received 8 August 2005; published 22 September 2005)

Effects of the 1-3 leptonic mixing on the solar neutrino observables are studied and the signatures of nonzero θ_{13} are identified. For this we have rederived the formula for 3ν -survival probability including all relevant corrections and constructed the isocontours of observables in the $\sin^2\theta_{12}-\sin^2\theta_{13}$ plane. Analysis of the solar neutrino data gives $\sin^2\theta_{13} = 0.007^{+0.080}_{-0.007}$ (90% C.L.) for $\Delta m^2 = 8 \times 10^{-5} \text{ eV}^2$. The combination of the ratio CC/NC at Sudbury Neutrino Observatory (SNO) and gallium production rate selects $\sin^2\theta_{13} = 0.017 \pm 0.026$ (1 σ). The global fit of all oscillation data leads to zero best value of $\sin^2\theta_{13}$. The sensitivity (1 σ error) of future solar neutrino studies to $\sin^2\theta_{13}$ can be improved down to 0.01–0.02 by precise measurements of the *pp*-neutrino flux and the CC/NC ratio as well as spectrum distortion at high (E > 4 MeV) energies. Combination of experimental results sensitive to the low and high energy parts of the solar neutrino spectrum resolves the degeneracy of angles θ_{13} and θ_{12} . Comparison of $\sin^2\theta_{13}$ as well as $\sin^2\theta_{12}$ measured in the solar neutrinos and in the reactor/accelerator experiments may reveal new effects which can not be seen otherwise.

DOI: 10.1103/PhysRevD.72.053011

I. INTRODUCTION

Determination of the 1-3 mixing parametrized by the angle θ_{13} is of great importance for phenomenology, for future experimental programs, and eventually, for understanding the underlying physics.

At present, for $\sin^2 \theta_{13}$ we only have the upper bounds. The CHOOZ [1] reactor experiment (see also PaloVerde [2]) gives

$$\sin^2 \theta_{13} \le 0.036, \quad 90\% \text{ C.L.}$$
 (1)

for $\Delta m_{13}^2 = 2.5 \times 10^{-3} \text{ eV}^2$. The recent global fit of the oscillation results which includes also the solar salt phase II SNO, KamLAND, and atmospheric neutrino data leads to [3]

$$\sin^2 \theta_{13} \le 0.026 \ (0.047), \quad 90\% \ (3\sigma) \ C.L.$$
 (2)

According to [4] the 90% C.L. bound is $\sin^2 \theta_{13} \le 0.024$. The limit (2) is slightly better than the earlier (before SNO salt phase II) limits [5–7] $\sin^2 \theta_{13} \le 0.031$ (0.055). In [8] even stronger 3σ upper bound is quoted: $\sin^2 \theta_{13} \le 0.041$.

The combined fit of the CHOOZ, K2K, and atmospheric neutrino results (without solar ones) gives [3]

$$\sin^2 \theta_{13} \le 0.032 \ (0.060), \quad 90\% \ (3\sigma) \text{ C.L.}$$
 (3)

(See also [9].) Comparison of the bounds (2) and (3) shows that effect of inclusion of the solar neutrino data in the global fit is weak but not negligible: the bounds improve by $\sim 25\%$.

Future long baseline experiments are geared towards precision measurement of θ_{13} . MINOS [10] and CERN to Gran-Sasso [11] alone will be able to only slightly improve the limit (2) down to

$$\sin^2 \theta_{13} \le 0.015, \quad 90\% \text{ C.L.}$$
 (4)

PACS numbers: 14.60.Pq

J-PARC accelerator experiment [12] will have substantially higher sensitivity:

$$\sin^2 \theta_{13} \le 0.0015, \quad 90\% \text{ C.L.},$$
 (5)

but when all correlations and degeneracies of parameters are taken into account the estimated sensitivity reduces to $\sin^2 \theta_{13} \approx 0.004$ (90% C.L.) [13].

The Double-CHOOZ reactor experiment [14] will be able to improve the bound down to

$$\sin^2 \theta_{13} = 0.008, \quad 90\% \text{ C.L.}$$
 (6)

for $\Delta m^2 = 2.4 \times 10^{-3} \text{ eV}^2$. It provides a clean channel of measurement of θ_{13} complementary to accelerator measurements [13,15].

Detection of neutrinos from the Galactic supernova can, in principle, probe $\sin^2\theta_{13}$ down to $10^{-5} \cdot 10^{-6}$. The problem here is large uncertainties in the original neutrino fluxes. Essentially one can find whether $\sin^2\theta_{13}$ is larger or smaller than 10^{-3} . Both the upper and the lower bounds on $\sin^2\theta_{13}$ may be established if $\sin^2\theta_{13}$ is in the range $10^{-5} \cdot 10^{-3}$ [16,17].

Ultimate sensitivity, $\sin^2 \theta_{13} = 10^{-4} \cdot 10^{-5}$, can be achieved at the neutrino factories [18].

Influence of the 1-3 mixing on the detected solar neutrino fluxes has been considered in a number of publications before [7,19–34]. The present solar neutrino data are not precise enough to really probe $\sin^2\theta_{13}$ in the allowed range (2) and (3), but on the other hand the bounds obtained using the solar data only are not much weaker than those from the reactor and atmospheric neutrino results. Indeed, recent analysis of the solar neutrino data alone [3] gives

$$\sin^2 \theta_{13} \le 0.038 \ (0.062), \quad 90\% \ (2\sigma) \text{ C.L.}, \quad (7)$$

and very weak 3σ bound. Inclusion of the latest KamLAND data improves the latter bound substantially [3]

$$\sin^2 \theta_{13} \le 0.035 \ (0.075), \quad 90\% \ (3\sigma) \text{ C.L.}$$
 (8)

(This can be compared with (3).)

The 1-3 mixing taken at the present upper bound modifies the allowed region of the parameters $\sin^2\theta_{12}$ - Δm_{21}^2 obtained from the global fit. It shifts the region toward larger Δm_{21}^2 and smaller $\sin^2\theta_{12}$, however the shift of the best-fit point is within 1σ [7].

It was noticed that the Earth matter regeneration effect has strong dependence on θ_{13} , as $P \propto \cos^6 \theta_{13}$ [35]. This can be used to determine θ_{13} in future solar neutrino experiments [35,36]. Measurements of the *pp*-neutrino fluxes will also improve the sensitivity to $\sin^2 \theta_{13}$ [21].

In this paper we study in details possible effects of 1-3 mixing on the solar neutrino observables. We analyze the present determination of θ_{13} and θ_{12} and evaluate the sensitivity of future solar neutrino experiments to θ_{13} .

The paper is organized as follows. We first (Sec. II) rederive the ν_e -survival probability including all relevant corrections and estimating accuracy of the approximations. We consider dependence of the probability on $\sin^2\theta_{13}$. In Sec. III we find simple analytic expressions for the probability in the high and low energy limits. In Sec. IV we introduce the $\sin^2\theta_{12} - \sin^2\theta_{13}$ plots and derive the equations for contours of constant values of observables in this plot. In Sec. V we find the allowed region of parameters $\sin^2\theta_{12}$ and $\sin^2\theta_{13}$ and consider the bounds on these parameters from individual measurements. In Sec. VI the sensitivity of future, in particular, the low energy (LowNu) solar neutrino experiments to 1-3 mixing, is estimated. We discuss possible implications of future measurements of θ_{13} in Sec. VII. Our results are summarized in Sec. VIII.

II. THE SURVIVAL PROBABILITY AND θ_{13}

The analytic formula which describes conversion of the solar neutrinos in the case of three neutrino mixing and mass split hierarchy, $\Delta m_{31}^2 \gg \Delta m_{21}^2$, has been derived a long time ago [31,32]. Here Δm_{31}^2 and Δm_{21}^2 are the mass splits associated to 1-3 mixing and to the main channel of the solar neutrino conversion correspondingly. The formula was analyzed and further corrected recently [33–35].

In view of future precision measurements one needs to have an accurate formula for the probability with all relevant corrections included and errors due to approximation made estimated. For this reason we rederive the formula in a simple and adequate way which employs features of the large mixing angle (LMA) Mikheyev-Smirnov-Wolfenstein (MSW) solution. We will use a very high level of adiabaticity [37–40] in whole 3ν system. Because of loss of coherence between the mass eigenstates, ν_i (i = 1, 2, 3), the ν_e survival probability can be written as

$$P_{ee} = \sum_{i=1,2,3} P_{ei}^{\text{sun}} P_{ie}^{\text{Earth}},\tag{9}$$

where P_{ei}^{sun} is the probability of $\nu_e \rightarrow \nu_i$ conversion in the Sun and P_{ie}^{Earth} is the $\nu_i \rightarrow \nu_e$ oscillation probability inside the Earth. Strong adiabaticity of the neutrino conversion in the Sun leads immediately to

$$P_{ei}^{\rm sun} = |U_{ei}^{m0}|^2, \tag{10}$$

where U_{ei}^{m0} is the *ei*-element of neutrino mixing matrix in matter in the production point. Indeed, $U_{ei}^{m0} \equiv \langle \nu_e | \nu_{mi} \rangle$ determines the admixture of the eigenstate, ν_{mi} , in the neutrino state in the initial point. Then due to adiabaticity ν_{mi} transforms to ν_i when neutrino propagates to the surface of the Sun. Corrections due to the adiabaticity violation are negligible [40]: $\Delta P_{ee}/P_{ee} \sim$ $\gamma^2 \cos 2\theta_{12}/4\sin^2\theta_{12}$, where γ is the adiabaticity parameter. For E = 10 MeV we find $\Delta P_{ee}/P_{ee} \sim 10^{-8}$.

Combining (9) and (10) we obtain

$$P_{ee} = \sum_{i=1,2,3} |U_{ei}^{m0}|^2 P_{ie}^{\text{Earth}}.$$
 (11)

In the absence of the Earth matter effect (the day signal) we have

$$P_{ie}^{\text{Earth}} = |U_{ei}|^2, \tag{12}$$

where U_{ei} are the elements of the vacuum mixing matrix.

In what follows we will neglect the Earth matter effect on the 1-3 mixing which is determined by $2EV_E/\Delta m_{13}^2 < 0.005$, (here V_E is the typical potential inside the Earth). Therefore

$$P_{3e}^{\text{Earth}} \approx |U_{e3}|^2. \tag{13}$$

The Earth matter effect on two other probabilities can be described by the term F_{reg} defined as the deviation of probability P_{2e}^{Earth} from the no-oscillation one:

$$P_{2e}^{\text{Earth}} = |U_{e2}|^2 + F_{\text{reg}}.$$
 (14)

Then the unitarity condition, $\sum_{i=1,2,3} P_{ie}^{\text{Earth}} = 1$ and (13) lead to

$$P_{1e}^{\text{Earth}} = |U_{e1}|^2 - F_{\text{reg}}.$$
 (15)

Plugging (12), (13), and (15) into (11) we obtain

$$P_{ee} = \sum_{i=1,2,3} |U_{ei}^{m0}|^2 |U_{ei}|^2 + F_{\text{reg}}(|U_{e2}^{m0}|^2 - |U_{e1}^{m0}|^2).$$
(16)

We will use the standard parametrization of the mixing matrix in terms of mixing angles:

SOLAR NEUTRINOS AND 1-3 LEPTONIC MIXING

$$|U_{e1}| \equiv \cos\theta_{13}\cos\theta_{12}, \qquad |U_{e2}| \equiv \cos\theta_{13}\sin\theta_{12}, |U_{e3}| \equiv \sin\theta_{13},$$
(17)

and similar definition holds for the matrix elements in matter with substitution $\theta_{13} \rightarrow \theta_{13}^{m0}$ and $\theta_{12} \rightarrow \theta_{12}^{m0}$. The angle θ_{12} is identified with the "solar mixing angle."

In terms of mixing angles the probability (16) can be rewritten as

$$P_{ee} = \cos^2 \theta_{13}^{m0} \cos^2 \theta_{13} P_{ad} + \sin^2 \theta_{13}^{m0} \sin^2 \theta_{13} - F_{reg} \cos^2 \theta_{13}^{m0} \cos^2 \theta_{12}^{m0},$$
(18)

where P_{ad} is the usual two neutrino adiabatic probability:

$$P_{ad} \equiv \sin^2 \theta_{12} + \cos^2 \theta_{12}^{m0} \cos^2 \theta_{12}.$$
 (19)

For $F_{\text{reg}} = 0$ the probability (18) coincides with the one obtained in [32].

Let us find the mixing angle θ_{13}^{m0} in matter. The Hamiltonian of 3ν system can be written in the flavor basis as

$$H = U_{23}U_{13}U_{12}H^{d}U_{12}^{\dagger}U_{13}^{\dagger}U_{23}^{\dagger} + V_{3}.$$
 (20)

Here $H^d = \text{diag}(0, \Delta m_{21}^2/2E, \Delta m_{13}^2/2E), \quad V_3 \equiv \text{diag}(V, 0, 0), \text{ and } V \equiv \sqrt{2}G_F \rho Y_e, \text{ where } \rho \text{ is the matter density and } Y_e \text{ is the electron number density fraction.}$

Performing rotations $U_{23}U_{13}$ we arrive at the basis (ν'_e , ν'_2 , ν'_3) in which the Hamiltonian becomes

$$H = U_{12}H^d U_{12}^{\dagger} + U_{13}^{\dagger} V U_{13}.$$
 (21)

If the matter effect is small we can neglect mixing of the third state (off-diagonal terms of the matrix $U_{13}^{\dagger}VU_{13}$). This state then decouples and for the rest of the system the Hamiltonian becomes

$$H_2 = U_{12} H^d U_{12}^{\dagger} + \cos^2 \theta_{13} V_2, \qquad (22)$$

where $V_2 \equiv \text{diag}(V, 0)$.

In the lowest approximation the matter effect on the 1-3 mixing can be found making an additional 1-3 rotation which eliminates the 1-3 elements in (21). The angle of rotation, θ'_{13} , is given by

$$\tan 2\theta_{13}' = \frac{V \sin 2\theta_{13}}{\Delta m_{13}^2 / (2E) - V \cos 2\theta_{13} + \Delta m_{21}^2 / (2E) \sin^2 \theta_{12}}$$

\$\approx \sin 2\theta_{13}\epsilon\$,

(23)

where

$$\epsilon_{13} \equiv \frac{2EV}{\Delta m_{13}^2} = 0.062 \left(\frac{E}{10 \text{ MeV}}\right) \left(\frac{\rho Y_e}{100 \text{ g/cc}}\right) \left(\frac{2.5 \times 10^{-3} \text{ eV}^2}{\Delta m_{13}^2}\right).$$
(24)

So, approximately, $\theta'_{13} = \theta_{13} \epsilon_{13}$.

2 EV

The additional 1-3 rotation generates negligible 2-3 $elements^1$

$$H_{23} = \frac{V}{4}\sin 2\theta_{13}\sin 2\theta_{12}\frac{\Delta m_{21}^2}{\Delta m_{13}^2}.$$
 (25)

It also leads to corrections to the 1-2 block of the Hamiltonian:

$$H_2 = U_{12}H^d U_{12}^{\dagger} + \cos^2\theta_{13}(1 - \sin^2\theta_{13}\epsilon_{13})V_2.$$
 (26)

In the first order in $2VE/\Delta m_{13}^2$ the rotation θ'_{13} decouples the third state. So, the 1-3 mixing angle in matter is given by

$$\theta_{13}^{m} = \theta_{13} + \theta_{13}' + O(\theta_{13}' \epsilon_{13}), \qquad (27)$$

and using (23) we obtain

$$\sin^2\theta_{13}^m \approx \sin^2\theta_{13}(1+2\epsilon_{13}) + O(\sin^2\theta_{13}\epsilon_{13}^2, \sin^4\theta_{13}\epsilon_{13}).$$
(28)

(See also [33].) For typical energy E = 10 MeV the relative matter correction is (10-15)%. The correction is negligible at low energies.

The 1-2 mixing in matter is determined from diagonalization of (22) or (26). Neglecting very small correction ($\sim \epsilon_{13} \sin^2 \theta_{13}$) in (26) one obtains for θ_{12}^m the standard 2ν formula with substitution $V \rightarrow V \cos^2 \theta_{13}$.

Let us find dependence of F_{reg} on the 1-3 mixing explicitly. The probability of $\nu_2 \rightarrow \nu_e$ conversion can be written in terms of the amplitudes of transitions between the mass states $\nu_2 \rightarrow \nu_i$, A_{i2} , as

$$P_{2e} = |A_{12}U_{e1} + A_{22}U_{e2}|^2$$

= $\cos^2\theta_{13}|A_{12}\cos\theta_{12} + A_{22}\sin\theta_{12}|^2.$ (29)

Since in the absence of matter effect $A_{12} = 0$ and $A_{22} = 1$, in the lowest approximation in $A_{12} \propto V$, we obtain

$$F_{\rm reg} = \cos^2 \theta_{13} \sin 2\theta_{12} \text{Re}A_{12}.$$
 (30)

The transition amplitude A_{12} should be found by solving the evolution equation with the Hamiltonian (26). It is proportional to the potential: $A_{12} \propto V \cos^2 \theta_{13}$, and consequently, F_{reg} (30) can be rewritten as

$$F_{\rm reg} = \cos^4 \theta_{13} f_{\rm reg} + O(f_{\rm reg}^2), \tag{31}$$

¹Elimination of this term requires an additional 2-3 rotation by the angle $\theta'_{23} = 2EH_{23}/\Delta m^2_{31} < 10^{-4}$.

where

$$f_{\rm reg} \equiv \sin 2\theta_{12} \text{Re}A_{12}^{\prime} \tag{32}$$

is the ν_e -regeneration factor given by the 2ν amplitude $A'_{12} \equiv A_{12}/\cos^2\theta_{13}$, and it does not depend on θ_{13} .

The regeneration factor integrated with the exposure function over the zenith angle can be estimated as

$$\bar{f}_{\rm reg} \sim \frac{EV_0 \sin^2 2\theta_{12}}{4\Delta m_{21}^2},$$
 (33)

where V_0 is the potential at the surface of the Earth. Indeed, the integrated effect is mainly due to the oscillations in the mantle of the Earth along the trajectories which are not too close to the horizontal ones. (For the core crossing trajectories the effect of the core is attenuated [41].) In turn, for the mantle trajectories the adiabaticity is nearly fulfilled [40] and therefore characteristics of oscillations (averaged probability and depth of oscillations) are determined by the potentials in the initial and final points of the trajectory, that is, by the potential near the surface of the Earth (for numerical consideration see [42]).

Inserting the expression for F_{reg} into (18) we obtain

$$P_{ee} = \cos^2 \theta_{13}^{m0} \cos^2 \theta_{13} (P_{ad} - \cos^2 \theta_{13} \cos^2 \theta_{12}^{m0} f_{reg}) + \sin^2 \theta_{13}^{m0} \sin^2 \theta_{13}.$$
(34)

Taking the mixing angle θ_{13}^m in the first order in ϵ_{13} we find

$$P_{ee} \approx \cos^4 \theta_{13} (1 - 2\tan^2 \theta_{13} \epsilon_{13})$$
$$\times (P_{ad} - \cos^2 \theta_{13} \cos 2\theta_{12}^{m0} f_{reg}) + \sin^4 \theta_{13}. \quad (35)$$

Notice that the prefactor $(1 - 2\tan^2 \theta_{13} \epsilon_{13}(E))$ which describes the matter effect on the 1-3 mixing differs in the high and low energy limits.

The expression (34) can be rewritten as

$$P_{ee} \cong \cos^2 \theta_{13}^m \cos^2 \theta_{13} P_{ee}^{(2)} (\Delta m^2, \theta_{12}, \theta_{13}) + \sin^2 \theta_{13}^m \sin^2 \theta_{13},$$
(36)

where

$$P_{ee}^{(2)} = P_{ad} - \cos^2 \theta_{13} \cos^2 \theta_{12}^{m0} f_{\text{reg}}$$

is the total ν_e survival probability for 2ν mixing which includes both the conversion in the Sun and the oscillations in the Earth. P_{ad} depends on θ_{13} via the effective matter potential $V\cos^2\theta_{13}$.

The probability should be averaged over the distribution of the neutrino sources in the Sun. For the *K* component of the solar neutrino spectrum ($K \equiv pp$, Be, pep, B, N, O) the averaged probability can be written as [40]

$$P_{ee,K}^{(2)} = \sin^2\theta + \cos^2\theta_m(V_K)\cos^2\theta - 0.5\delta_K\cos^2\theta_m(V_K)$$
$$\times \cos^2\theta - \cos^2\theta_m(V_K)\cos^2\theta_{13}f_{\text{reg}}, \qquad (37)$$

where V_K is the effective value of the matter potential in

the region of production of the *K* neutrino component. δ_K is an additional correction due to the integration over the production region [40], and in what follows we will neglect it.

Notice there are three features of dynamics of propagation which allow one to immediately derive the 3ν survival probability in terms of the 2ν probabilities:

1) Adiabaticity of propagation of the third neutrino;

2) Negligible Earth matter effect on 1-3 mixing;

3) Averaging of oscillations and loss of coherence of the third mass eigenstate.

Essentially this means that the third neutrino eigenstate decouples from dynamics and propagates independently. Its contribution to the probability adds incoherently. Therefore the problem is reduced to a two neutrino problem and additional factors in (34) are just projection of the flavor basis onto the basis of states which contains the decoupling state ν_{3m} . In (36) factors $\sin^2 \theta_{13}^{m0}$ and $\cos^2 \theta_{13}^{m0}$ are from the projection in the initial state, whereas the factors $\sin^2 \theta_{13}$ and $\cos^2 \theta_{13}$ are from projection in the final state.

Up to negligible corrections the expression for the probability (36) can be rewritten in form

$$P_{ee} \cong (1 - \sin^2 \tilde{\theta}_{13}^2)^2 P_{ee}^{(2)}(\Delta m^2, \theta_{12}, \tilde{\theta}_{13}) + \sin^4 \tilde{\theta}_{13}, \quad (38)$$

where

$$\sin^2 \tilde{\theta}_{13} \equiv \sin^2 \theta_{13} (1 + \epsilon_{13}), \tag{39}$$

which coincides with the form without matter corrections to θ_{13} .² This means that the matter effect on the 1-3 mixing in the probability can be accounted by renormalization of $\sin^2\theta_{13}$. Inversely, using usual formula for the probability without matter corrections in the analysis of the data we determine $\tilde{\theta}_{13}$ averaged over the relevant energy interval. Then the true vacuum mixing angle will be smaller:

$$\sin^2\theta_{13} = \sin^2\tilde{\theta}_{13}(1 - \bar{\boldsymbol{\epsilon}}_{13}). \tag{40}$$

Here $\bar{\epsilon}_{13} = \langle \epsilon_{13} \rangle_E$ is the value averaged over the relevant energy range. Renormalization depends on the neutrino energy: $\bar{\epsilon}_{13} \sim 0.05$ at high energies and $\bar{\epsilon}_{13} \sim 0.002$ in the *pp*-neutrino range, and the latter can be neglected.

III. TWO LIMITS

As we will see, the expected sensitivity of solar neutrino studies to the 1-3 mixing will not be better than $\sigma(\sin^2\theta_{13}) \sim 0.01$. According to (36) that would correspond to the change of the survival probability

$$\delta P \approx -2P\sigma(\sin^2\theta_{13}) \sim 1\%. \tag{41}$$

²Strictly, this is correct in the high energy or low energy limits, where $P_{ee}^{(2)} \sim P_{ad}$ depends very weakly on the potential. At low energies the matter corrections are negligible anyway (see Eq. (24).

Therefore in the present discussions we can neglect various corrections to the probability which are much smaller that 0.01. In particular,

- (a) the term $\sin^4 \theta_{13} < 0.002$ can be neglected in Eq. (36);
- (b) the matter effect on the 1-3 mixing is small and can be taken into account as a small renormalization of sin²θ₁₃ (see Sec. II);
- (c) we will neglect $f_{\rm reg}^2 < 10^{-3}$ corrections in $F_{\rm reg}$.

In qualitative consideration we will use simplified expressions for the survival probability obtained in the limits of high and low energies. In fact, all relevant fluxes are either in the low (pp, Be, N, O, pep) or in the high, E > 5 MeV (B, hep), energy limits. The expansion parameter equals

$$\eta \equiv \frac{\Delta m_{21}^2}{2EV} \tag{42}$$

at high energies and $1/\eta$ at low energies.³

1) In the high energy limit, performing expansion of (37) in η we find the survival probability

$$P_{ee}^{(h)} \approx \cos^4 \theta_{13} \sin^2 \theta_{12} + \frac{1}{4} \cos^2 \theta_{12} \sin^2 2\theta_{12} \eta^2 + \cos^6 \theta_{13} f_{\text{reg}}.$$
 (43)

Here the first term is due to the nonoscillatory adiabatic conversion. In the second term, which describes the effect of averaged oscillations, dependence on θ_{13} is absent. In contrast, the third (regeneration) term has the strongest dependence on θ_{13} [35]. It originates from the common factor $\cos^4\theta_{13}$ and the effective potential in the 3ν case which contains another $\cos^2\theta_{13}$ factor.

In the lowest order in $\sin^2\theta_{13}$ we can rewrite (43) as

$$P_{ee}^{(h)} \approx \sin^2 \theta_{12} + \frac{1}{4} \cos 2\theta_{12} \sin^2 2\theta_{12} \eta^2 + f_{\text{reg}} - \sin^2 \theta_{13} (2\sin^2 \theta_{12} + 3f_{\text{reg}}).$$
(44)

Neglecting the oscillation corrections (second term) and the double suppressed term $\sim \sin^2 \theta_{13} f_{reg}$ we obtain

$$P_{ee}^{(h)} \approx (1 - 2\sin^2\theta_{13})\sin^2\theta_{12} + f_{\text{reg.}}$$
 (45)

2) For the low energy neutrinos (E < 2 MeV), the matter effect is small and in the lowest order in η^{-1} we have

$$P_{ee}^{(l)} \approx \cos^4 \theta_{13} (1 - 0.5 \sin^2 2\theta_{12}) - 0.5 \cos^6 \theta_{13} \cos 2\theta_{12} \sin^2 2\theta_{12} \eta^{-1}.$$
(46)

Here the first term is the effect of averaged vacuum oscillations. The second one gives the matter effect correction which contains strong dependence on θ_{13} . An additional second power of $\cos \theta_{13}$ originates from the matter potential. The regeneration effect is neglected. Numerical calculations confirm that the linear in $\eta^{-1} \propto E$ approximation (46) works well up to 2 MeV.

In the lowest order in $\sin^2 \theta_{13}$ we find from (46)

$$P_{ee}^{(l)} \approx 1 - 0.5 \sin^2 2\theta_{12} - 0.5 \cos 2\theta_{12} \sin^2 2\theta_{12} \eta^{-1} -$$
(47)

$$-\sin^2\theta_{13}[2 - \sin^2 2\theta_{12} - 1.5\cos 2\theta_{12}\sin^2 2\theta_{12}\eta^{-1}].$$
(48)

For very low energies (pp-neutrino range) the matter corrections can be neglected, so that

$$P_{ee}^{(l)} \cong (1 - 2\sin^2\theta_{13})(1 - 0.5\sin^22\theta_{12}).$$
(49)

From Eqs. (43)–(48) we conclude the following.

- (a) The dependences of the probability on θ_{12} for the high energies and low energies are opposite: $P_{ee}^{(l)}$ increases whereas $P_{ee}^{(h)}$ decreases with decrease of θ_{12} .
- (b) In contrast, $P_{ee}^{(l)}$ and $P_{ee}^{(h)}$ both decrease with increase of θ_{13} . So, θ_{12} and θ_{13} correlate at low energies and anticorrelate at high energies. For the ⁸B neutrinos an increase of θ_{13} can be compensated by increase of θ_{12} , whereas at low energies (for the *pp* neutrinos), θ_{12} should decrease to compensate an increase of θ_{13} .
- (c) Both for the high and low neutrino energies the dependence of probability on Δm_{21}^2 appears via small corrections only, and therefore, is weak.

Apparently strong dependence on θ_{13} cancels in the ratio of probabilities (or the ratio of the corresponding observables):

$$\frac{P_{ee}^{(h)}}{P_{ee}^{(l)}} = \frac{\sin^2\theta_{12} + 0.25\cos^{-4}\theta_{13}\cos2\theta_{12}\sin^22\theta_{12}\eta^2 + \cos^2\theta_{13}f_{\text{reg}}}{1 - 0.5\sin^22\theta_{12}}.$$
(50)

Notice that dependence of the ratio on θ_{13} appears via small corrections and it can be neglected in the first approximation. So, measurements of the ratio gives $\sin^2 \theta_{12}$. Then one of the probabilities (in low or high limits) can be used to find $\sin^2 \theta_{13}$.

³Notice that in [40] we use different definition of η : $\eta \rightarrow 1/\eta'$.

IV. $\sin^2 \theta_{12}$ - $\sin^2 \theta_{13}$ PLOTS AND ISOCONTOURS OF OBSERVABLES

There are two important features of the problem:

- (a) Weak dependence of the probability and observables on Δm_{21}^2 within the allowed region;
- (b) $\sin^2\theta_{12} \sin^2\theta_{13}$ degeneracy.

Let us consider these features in some details.

1) The KamLAND spectral data has shown high precision in determination of Δm_{21}^2 . The relative error

$$\delta_{\Delta} \equiv \frac{\delta(\Delta m_{21}^2)}{\Delta m_{21}^2} \tag{51}$$

is about $\delta_{\Delta} = 12\%$ (3 σ) at the present, and with 3 kTy statistics from KamLAND this can further reduce down to 7% [43].

As follows from (43) and (48), variation of the probability in the high energy limit with Δm_{21}^2 equals

$$\delta P_{ee}^{(h)} = \left[\frac{1}{2}\eta^2 \cos 2\theta_{12} \sin^2 2\theta_{12} - \cos^6 \theta_{13} f_{\text{reg}}\right] \delta_{\Delta}.$$
 (52)

For E = 10 MeV and $\delta_{\Delta} \sim 0.1$, we find $\delta P \sim 0.005$.

In the low energy part, variations of the probability with Δm_{21}^2 are even weaker:

$$\delta P_{ee}^{(l)} \approx \frac{1}{2} \eta^{-1} \cos^6 \theta_{13} \cos 2\theta_{12} \sin^2 2\theta_{12} \delta_{\Delta}.$$
 (53)

For E = 0.4 MeV we find $\delta P_{ee}^{(3)} = 0.011 \delta_{\Delta} \approx 0.001$.

2) According to the discussion in Sec. II, in the first approximation, all experiments which probe the high en-

ergy part of the spectrum are sensitive to the same combination of mixing angles θ_{12} and θ_{13} . A similar statement is true for the low energy experiments but the combination of mixing angles is different. Therefore to determine $\sin^2 \theta_{12}$ and $\sin^2 \theta_{13}$ separately one needs to employ both types of experiments.

In view of these two features, we will present results of our study in $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane for fixed values of Δm_{21}^2 .

Let us construct the contours of constant values of various observables in the $\sin^2\theta_{12} - \sin^2\theta_{13}$ plane. Essentially, the isocontours for a given observable *X* coincide with the contours of constant survival probability averaged over the relevant energy range with the corresponding cross section:

$$\langle R_{ee} \rangle_X = C_X. \tag{54}$$

The survival probabilities found in the high and low energy limits determine two different types of the isocontours. For the high energy observables (E > 5 MeV), using Eq. (44), we find an analytic expression for the contours of constant *X*:

$$\sin^2 \theta_{13} \approx \frac{\sin^2 \theta_{12} + 0.25 \cos 2\theta_{12} \sin^2 2\theta_{12} \eta_X^2 + \langle f_{\text{reg}} \rangle_X - C_X}{2 \sin^2 \theta_{12} + 3 \langle f_{\text{reg}} \rangle_X},$$
(55)

where

$$\eta_X^2 \equiv \left\langle \left(\frac{\Delta m^2}{2EV_{\rm B}}\right)^2 \right\rangle_X \tag{56}$$

is averaged over the neutrino energy value of η^2 folded with the cross section, energy resolution of detector, etc. In the allowed range of $\sin^2 \theta_{12}$ (with the average value of mixing denoted by $\bar{\theta}_{12}$) the strongest dependence comes from the denominator of (55):

$$\sin^2 \theta_{13} \sim A(\sin^2 \theta_{12} - C_X), \qquad A = (2\sin^2 \bar{\theta}_{12} + 3f_{\text{reg}})^{-1}.$$
 (57)

So, in the first approximation the contours are straight lines. This reproduces well the results of numerical calculations presented in Secs. V and VI. Notice that $\sin^2 \theta_{13}$ increases with $\sin^2 \theta_{12}$.

For the low energy observables Y using (48), we obtain expressions for the isocontours as

$$\sin^2\theta_{13} \approx \frac{1 - 0.5\sin^22\theta_{12} - 0.5\cos2\theta_{12}\sin^22\theta_{12}\eta_Y^{-1} - C_Y}{2 - \sin^22\theta_{12} - 1.5\cos2\theta_{12}\sin^22\theta_{12}\eta_Y^{-1}},$$
(58)

where

$$\eta_Y^{-1} \equiv \left\langle \frac{2EV}{\Delta m_{12}^2} \right\rangle_Y. \tag{59}$$

In the rough approximation

$$\sin^2 \theta_{13} \propto 1 - C_Y - 2\sin^2 \theta_{12} (1 - \sin^2 \theta_{12}) \tag{60}$$

and it has minimum at $\sin^2 \theta_{12} = 0.5$. Other terms in (58) shift this minimum to smaller values of $\sin^2 \theta_{12}$. So, in the region of interest, $\sin^2 \theta_{12} \sim 0.3$, the lines of constant

values of observables have negative slope: $\sin^2 \theta_{13}$ decreases with $\sin^2 \theta_{12}$.

V. CONSTRAINTS ON THE MIXING PARAMETERS FROM THE PRESENT EXPERIMENTS

In this section we describe the results of our numerical computations.

We first perform the global analysis of different data samples. Some details of the analysis follow. The CHOOZ

SOLAR NEUTRINOS AND 1-3 LEPTONIC MIXING

 χ^2 depends on Δm_{31}^2 and in our fit we vary this parameter freely in the range allowed by the zenith angle Super-Kamiokande atmospheric neutrino data [44]. The details of procedure for the solar and CHOOZ data can be found in [28] while the KamLAND analysis follows the method outlined in [6,29]. We use the KamLAND data from the latest version of paper [45] where a new source of the background has been taken into account.

For the solar neutrinos we include the data on total rates from the radiochemical experiments Cl [46] and Ga (SAGE, GALLEX, and GNO combined) [47], the Super-Kamiokande day-night spectrum data [48], the SNO charged current (CC), neutrino-electron scattering (ES), and neutral current (NC) rates from the salt phase I, the combined CC + ES + NC energy spectrum from the pure D_2O phase [49] as well as the day-night spectrum data of CC, ES, and NC events from the salt phase II [50].

The ⁸B neutrino flux normalization, $f_{\rm B} \equiv F_{\rm B}/F_{\rm B}^{\rm SSM}$, where $F_{\rm B}$ is the true flux and $F_{\rm B}^{\rm SSM}$ is the flux predicted in the standard solar model (SSM) [51], is left as a free parameter in the fit. Essentially $f_{\rm B}$ is fixed by the NC data from the SNO experiment. For the other fluxes and their uncertainties we use the predictions from the SSM [51].

In Fig. 1 we show the allowed regions of the parameters $\sin^2\theta_{12} \cdot \sin^2\theta_{13}$ (for two different values of Δm_{12}^2) which follow from the global fit of all available solar neutrino data as well as data from reactor experiments KamLAND and CHOOZ. The contours are marginalized over $f_{\rm B}$. The global minimum has been obtained by letting all the pa-



FIG. 1 (color online). The 90%, 95%, 99%, and 99.73% C.L. allowed regions of mixing angles from the global χ^2 analysis of the solar neutrino, KamLAND and CHOOZ neutrino data. The left panel is for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$ and the right panel is for $\Delta m_{21}^2 = 9 \times 10^{-5} \text{ eV}^2$. The global best-fit point obtained by letting all the parameters (including Δm_{21}^2) vary freely is marked by the crossed circle.

rameters (including Δm_{21}^2) vary freely in the fit. With this, the values of parameters in the global minimum come at

$$\Delta m_{21}^2 = 8.0 \times 10^{-5} \text{ eV}^2, \qquad \sin^2 \theta_{12} = 0.31, \\ \sin^2 \theta_{13} = 0.00, \qquad f_{\text{B}} = 0.84.$$
(61)

The two panels in Fig. 1 correspond to two fixed illustrative values of Δm_{21}^2 from the current allowed range. The contours are plotted with respect to the global minimum (61) using the definition of $\Delta \chi^2$ which corresponds to 3 parameters. The upper bound, $\sin^2 \theta_{13} < 0.055$ (99% C.L.) is tighter than the one obtained without including the phase II SNO salt data (0.061). According to Fig. 1 with increase of Δm_{21}^2 the allowed region shifts to smaller $\sin^2 \theta_{12}$; with increase of $\sin^2 \theta_{13}$ the value of $\sin^2 \theta_{12}$ in minimum of χ^2 practically does not change.

We have performed the global fit of the solar neutrino data only. The best-fit values of parameters are found to be

$$\Delta m_{21}^2 = 6.4 \times 10^{-5} \text{ eV}^2, \qquad \sin^2 \theta_{12} = 0.33, \\ \sin^2 \theta_{13} = 0.01, \qquad f_{\text{B}} = 0.84.$$
(62)

Now the 1-3 mixing is nonzero but statistically insignificant. The solar χ^2 is somewhat flat in the region $\sin^2 \theta_{13} =$ 0-0.03. For $\sin^2 \theta_{13} = 0.01$ and 0.02 we obtain the $\chi^2_{min} =$ 113.84 and 114.04, correspondingly.

Notice that we find small boron neutrino flux: $f_{\rm B} = 0.84$ in the units of flux predicted in [51]. Recent calculations of fluxes with new radiative opacities and the heavy element abundances give an even smaller value $f_{\rm B} = 0.78$ [52].

In Fig. 2 we show the constant χ^2 contours obtained from global analysis of the solar neutrino data only and when Δm_{21}^2 is fixed. The contours have been calculated with respect to the global minimum (62) and using the $\Delta \chi^2$ values corresponding to 3 parameters. We also show the local minima found for fixed values of Δm_{21}^2 . In particular, for $\Delta m_{21}^2 = 8 \times 10^{-5}$ eV2 we obtain

$$\sin^2 \theta_{13} = 0.007^{+0.080}_{-0.007}$$
 (90% C.L.). (63)

The best-fit value of θ_{13} changes with Δm_{21}^2 very weakly, e.g., for $\Delta m_{21}^2 = 9 \times 10^{-5} \text{ eV}^2$ we find $\sin^2 \theta_{13} = 0.01$. The Fig. 2 shows also that at present the solar data alone has a weak sensitivity to $\sin^2 \theta_{13}$: the 90% C.L. bound equals $\sin^2 \theta_{13} \leq 0.087$. Inversely, the values of $\sin^2 \theta_{13}$ allowed by the global fit (2) have weak impact on the global analysis of the solar neutrino data. From Fig. 2 we find that with increase of $\sin^2 \theta_{13}$ from 0 to 0.05 the best-fit value and 90% allowed region of $\sin^2 \theta_{12}$ shift to larger values by $\Delta \sin^2 \theta_{12} = 0.02$ only which is smaller than 1σ error and in agreement with earlier estimations in [22].

Let us now consider bounds on mixing angles θ_{12} and θ_{13} from individual observables.

1) At present only the gallium experiments can be considered as the low energy experiments. For the Ge production rate, Q_{Ge} , we have



FIG. 2 (color online). The 90%, 95%, 99%, and 99.73% C.L. allowed regions of mixing angles from the global χ^2 analysis of the solar neutrino data only. The left panel is for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$ and the right panel is for $\Delta m_{21}^2 = 9 \times 10^{-5} \text{ eV}^2$. The global best-fit point obtained by letting all the parameters vary freely is marked by \oplus . Also shown by crosses are the local best-fit points obtained at fixed Δm_{21}^2 .

$$Q_{\text{Ge}} = \left[P_{ee}^{pp} Q_{\text{Ge}}^{pp} + P_{ee}^{pep} Q_{\text{Ge}}^{pep} + P_{ee}^{\text{Be}} Q_{\text{Ge}}^{\text{Be}} + P_{ee}^{\text{CNO}} Q_{\text{Ge}}^{\text{CNO}} + P_{ee}^{\text{B}} f_{\text{B}} Q_{\text{Ge}}^{\text{B}}\right], \tag{64}$$

where Q_{Ge}^{i} is the contribution to the rate from the *i*th component of the solar neutrino flux according to the SSM, P_{ee}^{i} is the corresponding effective (averaged over the energy range) survival probability. The rate has the contributions both from the low and high energy parts of the spectrum, though the latter (from boron neutrinos) is much smaller. We can subtract the boron neutrino effect, $Q_{Ge}^{B,exp}$, using experimental results from SNO: $Q_{Ge}^{l} \equiv Q_{Ge} - Q_{Ge}^{B,exp}$. Taking the ⁸B flux from SNO as an input, we find that the ⁸B neutrino contribution equals $Q_{Ge}^{B,exp} = 4.2_{-0.7}^{+1.4}$ SNU and then $Q_{Ge}^{l} = 63.9_{-5.2}^{+5.1}$ SNU for the rest of the neutrinos [47].

The conversion effect on Q_{Ge}^{l} is described by the probability in low energy limit:

$$Q_{\text{Ge}}^{l} = (Q_{\text{Ge}}^{\text{SSM}} - Q_{\text{Ge}}^{\text{B}}) [\cos^{4}\theta_{13}(1 - 0.5\sin^{2}2\theta_{12}) - \cos^{6}\theta_{13}\cos^{2}2\theta_{12}\sin^{2}2\theta_{12}\eta_{\text{Ge}}^{-1}], \qquad (65)$$

where

$$\eta_{\rm Ge}^{-1} \equiv \frac{1}{Q_{\rm Ge}^{\rm SSM} - Q_{\rm Ge}^{\rm B}} (\eta_{pp}^{-1} Q_{\rm Ge}^{pp} + \eta_{pep}^{-1} Q_{\rm Ge}^{pep} + \eta_{\rm Be}^{-1} Q_{\rm Ge}^{\rm Be} + \eta_{\rm CNO}^{-1} Q_{\rm Ge}^{\rm CNO}).$$
(66)

At the same time we find that dependence of the total rate Q_{Ge} on f_{B} and the high energy conversion effect is very weak: 10% variations of f_{B} would produce change of Q_{Ge} about 0.4 SNU. Therefore results of numerical calculations are given in Figs. 3–5, for Q_{Ge} and $f_{\text{B}} = 1$. We show the contours of constant total rate Q_{Ge} which correspond to the present central value as well as to 1σ and 2σ deviations from it. The contours are well described by the analytical expression given in (58) with $\eta_Y^{-1} = \eta_{\text{Ge}}^{-1}$ and $C_Y \equiv C_{\text{Ge}}$.

For zero 1-3 mixing the central value $Q_{\text{Ge}} = 68.1$ SNU corresponds to $\sin^2 \theta_{12} = 0.34$, and the best-fit value $\sin^2 \theta_{12} = 0.33$ is accepted at 0.5σ level. For a given Q_{Ge} , $\sin^2 \theta_{13}$ increases with decrease of $\sin^2 \theta_{12}$. Taking the best-fit value $\sin^2 \theta_{12} = 0.33$ (see Eq. (62)) and $Q_{\text{Ge}} = 68.1$ SNU we find $\sin^2 \theta_{13} = 0.004$.

2) The CC/NC ratio of events at SNO is given by the integrated (with cross section) survival probability at high energies: $CC/NC = \langle P_{ee}^{(h)} \rangle$. The isocontours are described approximately by Eq. (55) with $C_X \equiv C_{\text{SNO}} = CC/NC$. The results of exact numerical calculations are shown in Fig. 3. The isocontours are constructed for the present central value of CC/NC as well as for its 1σ and 2σ deviations. At $\sin^2\theta_{12} = 0.33$ the central value of CC/NC gives $\sin^2\theta_{13} = 0.046$.

As seen from Fig. 3, the combination of SNO CC/NC and Ga results prefers the nonzero but statistically insig-



FIG. 3 (color online). The isocontours of the CC/NC ratio at SNO and Ge production rate in the $\sin^2\theta_{12} - \sin^2\theta_{13}$ plane for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$. Shown are the isocontours for the central values of observables and the $\pm 1\sigma$ and $\pm 2\sigma$ deviations. We take $f_{\rm B} = 1$. The solid contour bounds the 99% C.L. allowed area from the global fit of all data.

SOLAR NEUTRINOS AND 1-3 LEPTONIC MIXING



FIG. 4 (color online). The isocontours of $R_{\rm SK}$ and Ge production rate in the $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane for $\Delta m_{21}^2 = 8 \times 10^{-5}$ eV². Shown are the contours for the central values and $\pm 1\sigma$, $\pm 2\sigma$ deviations from these central values. We take $f_{\rm B} = 1$. The solid contour bounds the 99% C.L. allowed area from the global fit of all data.

nificant 1-3 mixing. The isocontours of the central observed values of CC/NC and Q_{Ge} cross at

$$\sin^2\theta_{13} = 0.017 \pm 0.026 \tag{67}$$

which is about 0.7σ deviation from zero. Here 1σ is evaluated from the Fig. 3 using CC/NC and $Q_{\rm Ge}$ measurements.

The CC/NC ratio is free of the ⁸B neutrino normalization factor $f_{\rm B}$. Isocontours for the Ge production rate are constructed for $f_{\rm B} = 1$, i.e., for the SSM value of the ⁸B flux. The dependence of $Q_{\rm Ar}$ on $f_{\rm B}$ essentially disappears if one uses for the boron neutrino flux the value measured by NC in SNO or by Super-Kamiokande (SK). In any case the influence of uncertainty related to $f_{\rm B}$ on the intersection of isocontours is weak. Larger uncertainty (~ 1 SNU) follows from the Be neutrino flux contribution. That leads to $\Delta \sin^2 \theta_{13} \sim$ 0.01. Future Borexino measurements can reduce this uncertainty.

3) The Super-Kamiokande rate of the νe -events. The ratio of the observed to SSM predicted event rates is given by

$$R_{\rm SK} = f_{\rm B}[\langle P_{ee} \rangle (1-r) + r], \qquad (68)$$

where $r \equiv \sigma_{\nu_{\mu}}/\sigma_{\nu_{e}}$ is the ratio of the $\nu_{e} - e$ and $\nu_{\mu} - e$ cross sections. The lines of constant $R_{\rm SK}$ coincide with the lines $\langle P_{ee} \rangle = C_{\rm SK} = \text{const}$, where according to (68)



FIG. 5 (color online). The isocontours of the Ar and Ge production rates in the $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$. Shown are the contours for the central values as well as $\pm 1\sigma$ and $\pm 2\sigma$ deviations. The contours have been calculated for $f_{\rm B} = 1$.

$$C_{\rm SK} = \frac{R_{\rm SK}/f_B - r}{1 - r}.$$
 (69)

The lines are given by the Eq. (55) with $C_X = C_{SK}$ and the averaging (integration) $\langle ... \rangle_X$ should be done according to the SK experiment characteristics.

The isocontours of the SK rate in the $\sin^2\theta_{12} - \sin^2\theta_{13}$ plane are presented in Fig. 4 for $f_{\rm B} = 1$. They correspond to the SK central value $R_{\rm SK} = 0.41$ as well as the 1σ and 2σ deviations. According to Fig. 4, for $\sin^2\theta_{13} = 0$ the ratio $R_{\rm SK} = 0.41$ is achieved at $\sin^2\theta_{12} \approx 0.21$ which is about 2σ below the best global value. In turn, the best global value $\sin^2\theta_{12} = 0.33$ and $R_{\rm SK} = 0.41$ lead to $\sin^2\theta_{13} = 0.17$.

With decrease of $f_{\rm B}$ the corresponding isocontours shift to larger θ_{12} and smaller θ_{13} . For arbitrary f_B one can still use the grid of contours calculated for $f_{\rm B} = 1$, but with changed values $R_{\rm SK}^{(1)}$. Indeed, according to (69), for given values $R_{\rm SK}$ and $f_{\rm B}$ one should take the contour $R_{\rm SK}^{(1)} = R_{\rm SK}/f_{\rm B}$.

The combination of SK and Ga data (crossing point of the contours which correspond to the central experimental values) selects $\sin^2 \theta_{13} = 0.053$ as is seen from Fig. 4. For $f_{\rm B} = 0.84$ the best-fit SK value, $R_{\rm SK} = 0.41$, corresponds to the isocontour with $R_{\rm SK}^{(1)} = 0.49$. The intersection of this contour with the isocontour $Q_{\rm Ge} = 68.1$ SNU occurs at much smaller 1-3 mixing: $\sin^2 \theta_{13} = -0.004$.

4) The argon production rate in the Homestake experiment is determined by SRUBABATI GOSWAMI AND ALEXEI YU. SMIRNOV

$$Q_{\rm Ar} = [P_{ee}^{\rm Be} Q_{\rm Ar}^{\rm Be} + P_{ee}^{pep} Q_{\rm Ar}^{pep} + P_{ee}^{\rm CNO} Q_{\rm Ar}^{\rm CNO} + P_{ee}^{\rm B} f_{\rm B} Q_{\rm Ar}^{\rm B}].$$
(70)

According to the SSM $Q_{\rm Ar}$ is dominated by the high energy (boron) neutrinos. Therefore the contours of constant rate (Fig. 5) have typical slope of the high energy observables. We show these contours for $f_{\rm B} = 1$.

For $\sin^2 \theta_{13} = 0$ and the best global value of $\sin^2 \theta_{12}$ the rate $Q_{\rm Ar}$ is about 3.15 SNU which is 2.5 σ above the experimental result. The combination of Cl and Ga results prefers large 1-3 mixing: $\sin^2 \theta_{13} = 0.1$ which deviates from zero by $\sim 2.5\sigma$.

For the arbitrary $f_{\rm B}$ again one needs to rescale $Q_{\rm Ar}^{\rm B} \rightarrow Q_{\rm Ar}^{\rm B}/f_{\rm B}$. Thus, for $f_{\rm B} = 0.84$ and $Q_{\rm Ar} = 2.55$ SNU the isocontour with $Q_{\rm Ar} \approx 3.04$ SNU should be used. It intersects with isocontour $Q_{\rm Ge} = 68.1$ SNU at $\sin^2\theta_{13} = 0.032$.

Summarizing, for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$, the combination of SNO and Ga data selects $\sin^2 \theta_{13} = 0.017$, which is practically independent of f_{B} . For $f_{\text{B}} = 0.84$ (which follows from the global fit) the combination of SK and Ga

results leads to $\sin^2 \theta_{13} = -0.004$ and the combination of Cl and Ga data selects $\sin^2 \theta_{13} \sim 0.032$, however statistical significance of the latter is low. The average from these three combinations is about $\sin^2 \theta_{13} \sim 0.01$. It agrees well with results of fit of all the solar data for the same Δm_{21}^2 . For larger $f_{\rm B}$ the larger values of $\sin^2 \theta_{13}$ would be preferable.

Notice that available spectral information does not give strong restriction on $\sin^2 \theta_{13}$. In fact, flatness of the measured spectrum of events at SNO and SK may even favor nonzero $\sin^2 \theta_{13}$. Indeed, at high energies an increase of $\sin^2 \theta_{13}$ implies increase of $\sin^2 \theta_{12}$ and therefore flattening of the spectrum.

The results on 1-3 mixing described above are in agreement with the best-fit value from the global analysis of all solar neutrino data, $\sin^2 \theta_{13} = 0.01$, which is realized, however, for smaller Δm_{21}^2 (62). Inclusion of the CHOOZ result diminishes $\sin^2 \theta_{13}$ to practically zero.

Let us consider dependence of the isocontours on Δm_{21}^2 . Using (55) we find for the high energy part:

$$\delta(\sin^2\theta_{13}) \approx \frac{1}{2} \bigg[\sin^2\theta_{13} + \frac{C_X - \sin^2\theta_{12} - 3\langle f_{\rm reg} \rangle_X (0.5 - \sin^2\theta_{13})}{\sin^2\theta_{12} + 1.5\langle f_{\rm reg} \rangle_X} \bigg] \delta_\Delta, \tag{71}$$

where we have expressed η in terms of $\sin^2 \theta_{13}$.

In the low energy region according to (58)

$$\delta(\sin^2 \theta_{13}) \approx \frac{\eta_Y^{-1}}{4} \frac{\cos 2\theta_{12} \sin^2 2\theta_{12}}{1 - 0.5 \sin^2 2\theta_{12}} \delta_{\Delta}, \qquad (72)$$

and the prefactor is small: $\delta(\sin^2\theta_{13}) \approx 0.011\delta_{\Delta}$.

In Fig. 6 we show results of numerical computations of the isocontours for Ga rate and CC/NC ratio for three different values of Δm_{21}^2 . In agreement with our qualitative consideration a shift of the low energy contours is much weaker. Variations of Δm_{21}^2 from 7 to $9 \times 10^{-5} \text{ eV}^2$ (which is about 25% change) gives $\delta(\sin^2\theta_{13}) = 0.008$ and $\delta(\sin^2\theta_{12}) = 0.012$. Reduction of the uncertainty in Δm_{21}^2 down to 14% will result in $\delta(\sin^2\theta_{13}) = 0.004$ and $\delta(\sin^2\theta_{12}) = 0.007$. This is much smaller than the expected sensitivity of future solar neutrino studies. The contours have been obtained for $f_{\rm B} = 1$. We have found that 10% variations of $f_{\rm B}$ produce shift of the Ge lines by $\delta(\sin^2\theta_{12}) = 0.004$, and the SNO contours do not change at all.

Let us consider the effect of matter corrections to the 1-3 mixing. As we have established in Sec. II, the effect is reduced to renormalization of $\sin^2\theta_{13}$ according to Eq. (40). True value of $\sin^2\theta_{13}$ is smaller than the one extracted from the data without taking into account matter correction. The renormalization is negligible for the low energy observables and it is of the order 5% at high energies. So, the isocontours of high energy observables should be shifted down by a factor ~0.95. This, in turn

leads to the decrease of $\sin^2 \theta_{13}$ in the intersection points: e.g., instead of $\sin^2 \theta_{13} = 0.017$ one finds 0.016.

The figures presented in this section allow us to estimate relevance of the nonzero 1-3 mixing for the solar neutrino.



FIG. 6 (color online). The isocontours of CC/NC = 0.34 at SNO and $Q_{\text{Ge}} = 68.1$ SNU in Ga experiments in the $\sin^2\theta_{12} \cdot \sin^2\theta_{13}$ plane for different values of Δm_{21}^2 : $\Delta m_{21}^2 = 9 \times 10^{-5} \text{ eV}^2$ —the dotted lines; $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$ —the solid lines; $\Delta m_{21}^2 = 7 \times 10^{-5} \text{ eV}^2$ —the dashed lines.

For a given CC/NC ratio an increase of $\sin^2 \theta_{13}$ from 0 to 0.05 (~ 3σ upper bound) corresponds to decrease the Ge production rate by 2σ or $\Delta Q_{\text{Ge}} \sim -8$ SNU. Inversely, for a given Q_{Ge} , the same increase of $\sin^2 \theta_{13}$ leads to 1.8σ decrease of the ratio CC/NC: Δ CC/NC ~ -0.05. The same change of $\sin^2 \theta_{13}$ diminishes the SK rate by 4σ : $\Delta R_{\text{SK}} = -0.04$. So, the 1σ variations of θ_{13} produce about 1σ or smaller variations of observables. As we have discussed in the beginning of previous section, the impact of θ_{13} on the global fit of the solar neutrino data is much weaker: 3σ increase of $\sin^2 \theta_{13}$ leads to only 0.7σ increase of $\sin^2 \theta_{12}$.

VI. SENSITIVITY OF THE FUTURE SOLAR NEUTRINO EXPERIMENTS TO 1-3 MIXING

As we have shown, some sensitivity of the solar neutrino data to the 1-3 mixing appears essentially due to the combination of the SNO and SK results from the onehand side and the gallium results from the other side. Future high precision measurements will have better sensitivity. To evaluate this sensitivity and to study possible implications of measurements we have constructed fine grids of the isocontours of various observables. We will use the expected 1σ errors, σ_{13} , in determination of $\sin^2\theta_{13}$ as a measure of the sensitivity. 2σ and 3σ errors can also be estimated from the grids of isocontours we present. In the first approximation $n\sigma$ error is given by $n\sigma_{13}$.

1) In Fig. 7 we show the fine grid of isocontours for the CC/NC ratio and the Ge production rate. The SNO



FIG. 7 (color online). The isocontours of the CC/NC ratio at SNO (dotted lines) and Ge production rate Q_{Ge} (solid line) with reduced steps 0.01 and 2.6 SNU, respectively, in the $\sin^2\theta_{12} - \sin^2\theta_{13}$ plane. Δm_{21}^2 is held fixed at $8 \times 10^{-5} \text{ eV}^2$.

(phase III) will measure the NC event rate with 6% accuracy [53]. If we assume a reduced error of 5% for the CC data then this makes the CC/NC error as 8% which can be compared to the present error of 11%. (This corresponds to reduction of the absolute error in the CC/NC ratio down to 0.024 from the current value of 0.035.)

According to Fig. 7, the current Ge production rate accuracy is $\Delta Q_{\text{Ge}} \approx 4$ SNU. Using this accuracy and the expected SNO-III error, we find from Fig. 7 the sensitivity $(1\sigma \text{ error})$ to $\sin^2\theta_{13}$ in the region $\sin^2\theta_{13} \sim 0 - 0.03$:

$$\sigma_{13}(\text{CC/NC} + \text{Ga}) = 0.023. \tag{73}$$

It is only slightly better than the present one (67).

If we take 5% accuracy for future possible measurements of CC/NC ratio and the Ge production rate error 2.6 SNU (the later is slightly larger than the present systematic error [47], and it could correspond to future hypothetical large mass gallium experiment for which statistical error is substantially smaller than the systematic one; this can be considered as the ultimate sensitivity of the gallium experiments), then according to Fig. 7

$$\sigma_{13}(\text{CC/NC} + \text{Ga}) = 0.016.$$
 (74)

In this case the value $\sin^2 \theta_{13} = 0.017$ would have only 1σ significance, and in the case of very small θ_{13} the upper bound will be $\sin^2 \theta_{13} < 0.024$ (0.048) at 90% (3 σ).

2) The Borexino and KamLAND plan to measure signal from the ⁷Be neutrinos. The isocontours can be well described by the low energy formula (58). Taking into account contribution from the nonelectron (converted) neutrinos we find the isocontours, R_{Borexino} , which coincide with isocontours $\langle P_{ee} \rangle = C_{\text{Be}}$, where $C_{\text{Be}} = (R_{\text{Borexino}} - r)(1 - r)^{-1}$.

The results of calculations are shown in Fig. 8. The Borexino contours have similar form to those for Ga rate.

Borexino is expected to accumulate 44 000 events in 4 years of time (~ 30 events/day) for LMA MSW solution. This corresponds to a statistical error of only 0.5% [54]. But the accuracy of measurement will be dominated by the background and systematics. Furthermore, the SSM prediction for the ⁷Be flux has 10% uncertainty which will add to the total error unless one keeps the normalization of ⁷Be flux as a free parameter to be determined from the Borexino experiment itself with a higher accuracy. Using Fig. 8 we estimate that a 5% accuracy of measurements of the rate at Borexino will contribute to improvement of sensitivity of the solar studies to $\sin^2 \theta_{13}$.

3) New low energy solar neutrino experiments will be able to measure the pp-neutrino flux with high accuracy [55]. Two kinds of experiments are being discussed [56]: one—using the charged current reactions (LENS, MOON, SIREN) and the other—using the neutrino-electron scattering process (XMASS, CLEAN, HERON, MUNU, GENIUS).

SRUBABATI GOSWAMI AND ALEXEI YU. SMIRNOV



FIG. 8 (color online). The isocontours of Borexino rate suppression with respect to the SSM prediction in the $\sin^2\theta_{12} - \sin^2\theta_{13}$ plane for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$.

For the *pp*-neutrinos we can use the isocontour equation for the low energies as in (58) with substitution $C_Y = C_{pp} = \langle P_{ee} \rangle_{pp}$ and $\eta_{pp}^{-1} = \langle EV/\Delta m_{21}^2 \rangle_{pp}$. Practically, η_{pp}^{-1} is negligible.



FIG. 9 (color online). The isocontours of the CC/NC ratio (dashed lines) and νe scattering rate of pp-neutrinos in the $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane for $\Delta m_{21}^2 = 8 \times 10^{-5} \text{ eV}^2$.

In Fig. 9 we show the contours of constant rate of the $\nu - e$ scattering events due to the *pp*-neutrinos. The neutral current contribution is also included. According to Fig. 9, future measurements of the CC/NC ratio with 5% accuracy and the *pp*-rate with 2% accuracy will have a sensitivity (1 σ error)

$$\sigma_{13}(\text{CC/NC} + pp) = 0.015.$$
 (75)

Consequently, the value $\sin^2 \theta_{13} = 0.034$ will be established with 2.3 σ significance, and in the case of very small θ_{13} the upper bounds $\sin^2 \theta_{13} < 0.022 \ (0.045)$ at 90% (3 σ) can be achieved.

4) Future Megaton scale water Cherenkov detectors HyperKamiokande [57] and UNO [58] will be able to measure the ratio $R_{\nu e}$ with accuracy $\Delta R_{\nu e} \sim 0.003$. We construct the fine grid of the $R_{\nu e}$ isocontours for $f_{\rm B} = 1$ (Fig. 10). According to Fig. 10 these measurements of $R_{\nu e}$ and the *pp*-rate (2% accuracy) would have a sensitivity (1 σ error)

$$\sigma_{13}(\text{CC/NC} + pp) = 0.011.$$
 (76)

However, the problem here is the poor knowledge of $f_{\rm B}$. It is difficult to expect that accuracy of the theoretical predictions will be better than 10%. Comparable accuracy will be achieved in the direct measurements of the neutral currents. The global fit of the data which include the

FIG. 10 (color online). The isocontours of the neutrinoelectron scattering rate for the *pp*-neutrinos (solid line) and high energy ⁸B neutrinos (dotted line) in the $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane. The isorates for the high energy neutrinos are drawn with an step 0.003 corresponding to the expected statistical error of a Megaton water detector like HyperKamiokande. Δm_{21}^2 is held fixed at 8×10^{-5} eV² and $f_{\rm B} = 1$.

spectral information and the regeneration effect with free $f_{\rm B}$ may give better accuracy.

So, we conclude that the future solar neutrino studies may reach a sensitivity $\sin^2 \theta_{13} \sim 0.015 - 0.020 \ (1\sigma)$, as compared with the present sensitivity 0.05-0.06. It seems that even combined fit of several future precision measurements will not go down to 0.01 which may be considered as an ultimate sensitivity of solar neutrino measurements of $\sin^2 \theta_{13}$.

On the basis of studies performed in Secs. V and VI we can identify signatures of the nonzero $\sin^2 \theta_{13}$. In general, they show up as a mismatch of the low and high energy measurements. The signatures include

(a) Small value of the Ge production rate Q_{Ge} . For a given ratio CC/NC one can find the critical value of $Q_{\text{Ge}}^c(\text{CC/NC})$, so that inequality

$$Q_{\rm Ge} < Q_{\rm Ge}^c (\rm CC/NC) \tag{77}$$

will testify for the nonzero value of $\sin^2 \theta_{13}$. From the Fig. 7 we find $Q_{Ge}^c(0.30) = 76$ SNU, $Q_{Ge}^c(0.32) = 74$ SNU, $Q_{Ge}^c(0.34) = 72$ SNU, $Q_{Ge}^c(0.36) = 70$ SNU, etc. The larger the difference in (77), the larger $\sin^2 \theta_{13}$. In fact, the present data satisfy the inequality (77), however the statistical significance is low. Apparently if CC/NC and Q_{Ge} are measured with high accuracy, the presence of nonzero $\sin^2 \theta_{13}$ can be established.

The inequality (77) can be inverted, so that small values of CC/NC for a given Q_{Ge} ,

$$\operatorname{CC}/\operatorname{NC} < \operatorname{CC}/\operatorname{NC}^{c}(Q_{\operatorname{Ge}}), \tag{78}$$

testify for nonzero θ_{13} .

(b) Small Borexino rate:

$$R_{\rm Bor} < R^c_{\rm Bor}({\rm CC/NC}). \tag{79}$$

The critical Borexino rates for different values of CC/NC equal: $R_{Bor}^{c}(0.280) = 0.71$, $R_{Bor}^{c}(0.305) = 0.69$, $R_{Bor}^{c}(0.325) = 0.67$, $R_{Bor}^{c}(0.340) = 0.66$, $R_{Bor}^{c}(0.36) = 0.645$.

(c) Small νe event rate produced by the *pp*-neutrinos:

$$R_{pp} < R_{pp}^c (\text{CC/NC}). \tag{80}$$

According to Fig. 10 the critical values $R_{pp}^{c}(CC/NC)$ equal 0.75 (0.294), 0.73(0.318), 0.71(0.343), 0.69(0.373).

(d) Weak spectrum distortion of the boron neutrinos. The spectrum distortion can be characterized, e.g., by the ratio of suppressions at low and high energies $r_R \equiv R_{\nu_e} (E < 7 \text{ MeV})/R_{\nu_e} (E > 7 \text{ MeV})$. There is certain critical value of the distortion, r_R^c , which depends on CC/NC, so that weaker distortion will testify for nonzero 1-3 mixing. Indeed, for fixed CC/NC with increase of $\sin^2 \theta_{13}$, the 1-2 mixing, $\sin^2 \theta_{12}$, should increase (45) and therefore, the up turn of spectrum at low energies should be weaker.

VII. IMPLICATIONS OF THE SOLAR NEUTRINO MEASUREMENTS OF θ_{13}

Let us first compare sensitivities of the future solar neutrino experiments and future reactor and accelerator experiments given in Eqs. (4)–(6). With the expected accuracy of SNO-III and 2% accuracy of the pp-neutrino flux measurements the solar neutrino experiments can reach sensitivity comparable to the one of the forthcoming MINOS and CNGS experiments. This sensitivity is at the level of the present bound from the global fit of all oscillation data and it is certainly lower than the sensitivity of future measurements.

The solar neutrino sensitivity would be comparable to that of J-PARC and Double-CHOOZ if the pp-neutrino flux is measured with 1% accuracy and CC/NC ratio—with 3% accuracy which looks rather unrealistic now.

Notice that the survival probability P_{ee} does not depend on the *CP* phase δ , and therefore the problem of degeneracy with δ [59] does not arise. So, the solar neutrinos studies can provide a clean channel for determination of θ_{13} like the reactor neutrinos. Therefore, in spite of the lower sensitivity the solar neutrino data will contribute somehow to resolution of degeneracy of these parameters in the future global fits.

It seems that the reactor and accelerator experiments will obtain results much before future high statistics solar neutrino experiments will start to operate. In this case possible implications of the solar neutrino measurements can include

1) Precise determination of $\sin^2 \theta_{12}$: resolution of the $\theta_{12} - \theta_{13}$ degeneracy.

2) Comparison of results on θ_{13} (as well as on θ_{12}) from the solar neutrinos and reactor/accelerator experiments. The outcome could be further confirmation of the reactor/accelerator results by solar neutrinos, or establishing certain discrepancy which may imply new physics beyond the LMA solution of the solar neutrino problem.

Let us consider these two points in some details.

A. Resolving degeneracy of angles

Because of the strong $\sin^2\theta_{12} - \sin^2\theta_{13}$ degeneracy unknown value of θ_{13} leads to uncertainty in the determination of $\sin^2\theta_{12}$ and vice versa. Thus, in the CC/NC measurements, inclusion of the 1-3 mixing produces an increase of $\sin^2\theta_{12}$ by 0.04 - 0.05 (that is, by 20%) if $\sin^2\theta_{13}$ increases from 0 to 0.06 (see Fig. 7). In the *pp*-neutrino flux measurements $\sin^2\theta_{12}$ decreases by 0.06 - 0.08, when $\sin^2\theta_{13}$ increases from 0 to 0.06 (see Fig. 9). So, precise determination of $\sin^2\theta_{12}$ ($\sin^2\theta_{13}$) from the solar neutrino data is not possible unless $\sin^2\theta_{13}$ ($\sin^2\theta_{12}$) is measured or strongly restricted. As follows from our analysis a combination of high statistics measurement at high and low energies can remove the degeneracy.

B. Confronting the solar and reactor/accelerator results

If nonzero $\sin^2 \theta_{13}$ is measured, our analysis will allow to understand relevance of the 1-3 mixing to the precision solar neutrino studies. For fixed Δm_{21}^2 , the 1-3 mixing will change determination of θ_{12} .

Notice that dependence of sensitivity of the solar neutrino studies to θ_{13} on the mass Δm_{31}^2 is very weak (via matter corrections to 1-3 mixing) in contrast to laboratory measurements. So, for small Δm_{13}^2 the former will have some advantage.

If θ_{13} effect is not found in future laboratory experiments and the upper bound at the level $\sin^2 \theta_{13} < 0.01$ is established, the effect of 1-3 mixing will be irrelevant even for measurements with the 1% accuracy. The uncertainty of future measurements of $\sin^2 \theta_{12}$ due to unknown θ_{13} will be reduced down to 0.01. Furthermore, the precise measurements of $\sin^2 \theta_{12}$ by CC/NC will have the smallest uncertainty due to larger slope of the isolines.

The analysis performed in this paper will allow us to understand new physics if inconsistencies in results of measurements will show up. Implications will depend on the character of results, and in the positive case, on particular value of $\sin^2 \theta_{13}$. Comparing the laboratory bound with the results of solar neutrino studies one can realize new physics effects which may show up in the solar neutrinos only and cannot be found otherwise. Let us consider some examples.

Suppose that in the standard 3ν analysis precision solar neutrino measurements will give a nonzero value of $\sin^2 \theta_{13}$ which is larger than the upper bound from the reactor/ accelerator experiments. Such a situation may indicate some additional suppression of the solar neutrino flux at low energies. Indeed, this would shift the low isocontours to the left down. It would be equivalent of taking the low isocontours in our plots (e.g., in Fig. 3) with larger survival probability. The intersection with high energy isocontours will shift down and therefore the inferred value of $\sin^2 \theta_{13}$ will be reduced. This situation can be reproduced in the presence of sterile neutrino with $\Delta m^2 < 10^{-5}$ eV² [60]. In this case the extracted value of $\sin^2 \theta_{12}$ reduces too.

Agreement between the solar and laboratory measurements can be also achieved by an additional suppression of the high energy (boron) flux due to flavor conversion, so that CC/NC lines move to the right down. In this case also the intersection of the high and low isocontours moves down and the inferred value of $\sin^2\theta_{12}$ increases. Such an effect can be produced by certain nonstandard neutrino interactions.

The value of $\sin^2 \theta_{13}$ extracted from the solar neutrino studies can be diminished if both the low and high energy fluxes are reduced by some additional mechanism. In fact, this imitates the effect of the 1-3 mixing and can be achieved by additional nonstandard interactions.

Let us consider the opposite situation: $\sin^2 \theta_{13}$ is discovered in the reactor/accelerator experiments, but the solar

neutrino data prefer zero or smaller value of $\sin^2 \theta_{13}$. This can be explained if the survival probability is larger than the one predicted for given values of θ_{12} and θ_{13} . Again some scenarios with modified dynamics of conversion due to new interactions of neutrinos can reproduce such a situation.

Similarly discrepancy can be realized comparing values of the 1-2 mixing extracted from the solar neutrino data and reactor/accelerator experiments. Indeed, the $\sin^2\theta_{12}$ can be measured with 3% accuracy at 1σ (1 d.o.f.) in the reactor experiments with baseline 50-70 km [43,61,62] provided that the 1-3 mixing is determined (restricted) accurately enough from independent measurements. Since a nonzero θ_{13} drives the 1-2 mixing to smaller values in such an experiment, combination with high energy solar neutrino experiments for which nonzero θ_{13} drives $\sin^2 \theta_{12}$ to higher values can resolve the θ_{13} - θ_{12} degeneracy. This is similar to combination of low energy and high energy solar neutrino experiments as discussed in the present article. In this case one can compare the $(\sin^2\theta_{12} - \sin^2\theta_{13})$ allowed regions from the solar neutrino studies and from reactor experiments.

VIII. CONCLUSION

1) We have derived formula for the survival probability in the three neutrino context (35) which includes all corrections relevant for the future precision of measurements. We estimated the accuracy of approximation made.

2) In the three generation context in the first approximation (neglecting the matter effect on θ_{13}) the solar neutrino probabilities are functions of the mass split Δm_{21}^2 and the mixing angles: θ_{12} and θ_{13} . The split Δm_{21}^2 is already determined with a high precision by current data. Furthermore, the survival probability has a weak dependence on Δm_{21}^2 in the allowed region. In view of this we have constructed various plots in $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane which clearly demonstrate the degeneracy of these mixing angles and also show how the uncertainty in one is going to affect the precision determination of the other. Synergy between the high and low energy solar neutrino flux measurement provides a better constraint on θ_{13} and allows us to resolve the degeneracy of these mixing angles.

3) Among the existing solar neutrino experiments only the Ga experiment is sensitive to the low energy part of the spectrum and, in particular, to the *pp*-neutrinos. Therefore we construct the isorates for Ga experiments in the $\sin^2\theta_{12}$ - $\sin^2\theta_{13}$ plane and investigate how the combinations of these with the isorates of Cl, SK, and CC/NC ratio in SNO constrain θ_{13} .

At present the sensitivity of the solar neutrino experiments to θ_{13} is rather weak and no indications of nonzero value has been obtained. Our analysis of all solar neutrino data leads to statistically insignificant nonzero best-fit value: $\sin^2 \theta_{13} = 0.01$. Similar value, $\sin^2 \theta_{13} \sim 0.007$, follows from the fit of the solar neutrino data for $\Delta m^2 = 8 \times 10^{-5} \text{ eV}^2$ selected by the KamLAND experiment. The combination of the CC/NC ratio at SNO and Q_{Ge} rate, which are essentially independent of the solar neutrino fluxes, gives $\sin^2 \theta_{13} = 0.017 \pm 0.026 (1\sigma)$.

The global fit of all existing data (including reactor, accelerator, and atmospheric neutrino results) leads to $\sin^2 \theta_{13} = 0.00$ as the best-fit point and to the upper bounds $\sin^2 \theta_{13} = 0.041 (0.061)$ at 90% (3 σ) from the two parameter plots in $\sin^2 \theta_{12} - \sin^2 \theta_{13}$ plane.

4) The precision of θ_{13} determination will improve with future accurate measurements of the *pp*-neutrino flux, the CC/NC ratio, and the energy spectrum of events at E > 4 MeV. We find that measurements with 5% error in the CC/NC ratio and 2% error in the *pp*-neutrino flux in *ve* scattering will have a sensitivity (1 σ error) $\sin^2 \theta_{13} \sim 0.015$ (1 σ). The value $\sin^2 \theta_{13} \sim 0.01$ (1 σ) looks like the ultimate sensitivity which could be reached in the solar neutrino experiments of the next generation.

Since the probability involved is P_{ee} the problem of degeneracy with *CP* phase δ does not appear and solar neutrinos can provide a clean channel for θ_{13} measurements like the reactor neutrinos.

5) It is likely that new accelerator and reactor results will be obtained first. Comparison of the results from solar neutrinos and accelerator/reactor experiments may confirm each other and further improve determination of $\sin^2\theta_{13}$ or reveal some discrepancy which will indicate a new physics beyond usual interpretation of the solar neutrino anomaly.

Determination of $\sin^2 \theta_{13}$ will eliminate or reduce the θ_{12} - θ_{13} degeneracy thus improving measurements of 1-2 mixing. Inversely, the better determination of $\sin^2 \theta_{12}$ can help in improving the precision of the $\sin^2 \theta_{13}$ measurements in the solar neutrino studies.

Comparison of the results of $\sin^2 \theta_{13}$ (as well as $\sin^2 \theta_{12}$) measurements in solar neutrinos and in reactor and accelerator experiments may uncover new interesting physics beyond LMA MSW picture which cannot be found otherwise.

ACKNOWLEDGMENTS

S. G. would like to thank The Abdus Salam International Centre for Theoretical Physics for hospitality and A. Bandyopadhyay and S. Choubey for using some of the codes developed with them. The authors are grateful to E. Lisi for valuable comments.

- M. Apollonio *et al.*, Phys. Lett. B 466, 415 (1999); Eur. Phys. J. C 27, 331 (2003).
- [2] F. Boehm et al., Phys. Rev. D 62, 072002 (2000).
- [3] G. L. Fogli, E. Lisi, A. Marrone, and A. Palazzo, hep-ph/ 0506083.
- [4] A. Strumia and F. Vissani, hep-ph/0503246.
- [5] J. N. Bahcall, M. C. Gonzalez-Garcia, and C. Pena-Garay, J. High Energy Phys. 08 (2004) 016.
- [6] A. Bandyopadhyay et al., Phys. Lett. B 608, 115 (2005).
- [7] M. Maltoni, T. Schwetz, M. A. Tortola, and J. W. F. Valle, New J. Phys. 6, 122 (2004).
- [8] M.C. Gonzalez-Garcia, hep-ph/0410030.
- [9] S. Goswami, Nucl. Phys. (Proc. Suppl.) 143, 121 (2005).
- [10] R. Saakian, Nucl. Phys. (Proc. Suppl.) 111, 169 (2002).
- [11] http://proj-cngs.web.cern.ch/proj-cngs/.
- [12] http://jkj.tokai.jaeri.go.jp/.
- [13] P. Huber, M. Lindner, T. Schwetz, and W. Winter, Nucl. Phys. B665, 487 (2003).
- [14] F. Ardellier et al., hep-ex/0405032.
- [15] H. Minakata, H. Sugiyama, O. Yasuda, K. Inoue, and F. Suekane, Phys. Rev. D 68, 033017 (2003).
- [16] C. Lunardini and A. Y. Smirnov, J. Cosmol. Astropart. Phys. 06 (2003) 009.
- [17] G. L. Fogli, E. Lisi, A. Mirizzi, and D. Montanino, Nucl. Phys. (Proc. Suppl.) 145, 343 (2005).
- [18] M. Lindner, Int. J. Mod. Phys. A **18**, 3921 (2003) and references therein.
- [19] G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, and A. Palazzo, hep-ph/0104221.

- [20] G. L. Fogli, G. Lettera, E. Lisi, A. Marrone, A. Palazzo, and A. Rotunno, Phys. Rev. D 66, 093008 (2002); G. L.
 Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, and A. M. Rotunno, Phys. Rev. D 69, 017301 (2004).
- [21] J. N. Bahcall and C. Pena-Garay, J. High Energy Phys. 11 (2003) 004.
- [22] M. C. Gonzalez-Garcia and C. Pena-Garay, Phys. Lett. B 527, 199 (2002).
- [23] M. C. Gonzalez-Garcia and C. Pena-Garay, Phys. Rev. D 68, 093003 (2003).
- [24] A.B. Balantekin and H. Yuksel, J. Phys. G 29, 665 (2003).
- [25] M. Maltoni, T. Schwetz, M. A. Tortola, and J. W. F. Valle, Phys. Rev. D 68, 113010 (2003); M. C. Gonzalez-Garcia, M. Maltoni, C. Pena-Garay, and J. W. F. Valle, Phys. Rev. D 63, 033005 (2001).
- [26] P. C. de Holanda and A. Yu. Smirnov, Astropart. Phys. 21, 287 (2004).
- [27] P.C. de Holanda and A.Yu. Smirnov, J. Cosmol. Astropart. Phys. 02 (2003) 001,; P.C. de Holanda and A.Yu. Smirnov, Phys. Rev. D 66, 113005 (2002).
- [28] A. Bandyopadhyay, S. Choubey, S. Goswami, and K. Kar, Phys. Rev. D 65, 073031 (2002).
- [29] A. Bandyopadhyay, S. Choubey, R. Gandhi, S. Goswami, and D. P. Roy, J. Phys. G 29, 2465 (2003); A. Bandyopadhyay, S. Choubey, R. Gandhi, S. Goswami, and D. P. Roy, Phys. Lett. B 559, 121 (2003).
- [30] A. Bandyopadhyay, S. Choubey, S. Goswami, S. T. Petcov, and D. P. Roy, Phys. Lett. B 583, 134 (2004).

- [31] S.C. Lim, in Proceedings of BNL Neutrino Workshop, Upton, NY, 1987, edited by M. Murtagh (unpublished), BNL-52079, C87/02/05.
- [32] X. Shi and D. N. Schramm, Phys. Lett. B 283, 305 (1992);
 X. Shi, D. N. Schramm, and J. N. Bahcall, Phys. Rev. Lett. 69, 717 (1992).
- [33] G. L. Fogli, E. Lisi, and A. Palazzo, Phys. Rev. D 65, 073019 (2002).
- [34] C. S. Lim, K. Ogure, and H. Tsujimoto, Phys. Rev. D 67, 033007 (2003).
- [35] M. Blennow, T. Ohlsson, and H. Snellman, Phys. Rev. D 69, 073006 (2004).
- [36] E. Kh. Akhmedov, M. A. Tortola, and J. W. F. Valle, J. High Energy Phys. 05 (2004) 057.
- [37] S. P. Mikheyev and A. Yu. Smirnov, Yad. Fiz. 42, 1441 (1985) [Sov. J. Nucl. Phys. 42, 913 (1985)]; Nuovo Cimento Soc. Ital. Fis. C 9, 17 (1986); S. P. Mikheyev and A. Yu. Smirnov, Zh. Eksp. Teor. Fiz. 91, 7 (1986) [Sov. Phys. JETP 64, 4 (1986)].
- [38] A. Messiah, in Proceedings of the 6th Moriond Workshop On Massive Neutrino in Particle Physics and Astrophysics, 1986, edited by O. Fackler and J. Tran Thanh Van (unpublished), p. 373.
- [39] W. Haxton, Phys. Rev. Lett. 57, 1271 (1986); S. J. Parke, Phys. Rev. Lett. 57, 1275 (1986).
- [40] P. C. de Holanda, W. Liao, and A. Y. Smirnov, Nucl. Phys. B702, 307 (2004).
- [41] A. N. Ioannisian and A. Y. Smirnov, Phys. Rev. Lett. 93, 241801 (2004).
- [42] M. C. Gonzalez-Garcia, C. Pena-Garay, and A. Y. Smirnov, Phys. Rev. D 63, 113004 (2001).
- [43] A. Bandyopadhyay, S. Choubey, S. Goswami, and S.T. Petcov, Phys. Rev. D 72, 033013 (2005).
- [44] E. Kearns, talk at Neutrino 2004, Paris, http:// neutrino2004.in2p3.fr.
- [45] T. Araki *et al.* (KamLAND Collaboration), Phys. Rev. Lett. **94**, 081801 (2005).
- [46] B. T. Cleveland et al., Astrophys. J. 496, 505 (1998).
- [47] J. N. Abdurashitov *et al.* (SAGE Collaboration), Zh. Eksp. Teor. Fiz. **122**, 211 (2002) [J. Exp. Theor. Phys. **95**, 181 (2002)]; W. Hampel *et al.* (GALLEX Collaboration), Phys. Lett. B **447**, 127 (1999); M. Altmann *et al.* (GNO

Collaboration), Phys. Lett. B 616, 174 (2005).

- [48] S. Fukuda *et al.* (Super-Kamiokande Collaboration), Phys. Lett. B **539**, 179 (2002).
- [49] Q.R. Ahmad *et al.* (SNO Collaboration), Phys. Rev. Lett. **89**, 011302 (2002); S.N. Ahmed *et al.* (SNO Collaboration), Phys. Rev. Lett. **92**, 181301 (2004).
- [50] B. Aharmim et al. (SNO Collaboration), nucl-ex/0502021.
- [51] J. N. Bahcall and M. H. Pinsonneault, Phys. Rev. Lett. 92, 121301 (2004).
- [52] J.N. Bahcall, S. Basu, and A.M. Serenelli, astro-ph/ 0502563.
- [53] K. Graham, talk at NOON 2004, February, 2004, Tokyo, Japan, http://www-sk.icrr.u-tokyo.ac.jp/noon2004/; H. Robertson for the SNO Collaboration, Talk given at TAUP 2003, University of Washington, Seattle, Washington, http://mocha.phys.washington.edu/taup2003.
- [54] C. Galbiati, talk at Neutrino 2004, Paris, http:// neutrino2004.in2p3.fr.
- [55] M. Nakahata, talk given at the Int. Workshop on Neutrino Oscillations and their Origin (NOON2004), 2004, Tokyo, Japan (unpublished); S. Schönert, talk at Neutrino 2002, Munich, Germany (unpublished).
- [56] See e.g., Y. Suzuki, Nucl. Phys. (Proc. Suppl.) 143, 27 (2005).
- [57] T. Nakaya, Nucl. Phys. (Proc. Suppl.) 118, 210 (2003); K. Nakamura, talk at the workshop "Neutrinos and Implications for Physics Beyond the Standard Model" 2002, http://insti.physics.sunysb.edu/itp/conf/neutrino/ talks/nakamura.pdf.
- [58] UNO Proto-collaboration, UNO Whitepaper: Physics Potential and Feasibility of UNO, Report No. SBHEP-01-03, 2000, http://nngroup.physics.sunysb.edu/uno/; see also C. K. Jung, hep-ex/0005046.
- [59] V. Barger, D. Marfatia, and K. Whisnant, Phys. Rev. D 65, 073023 (2002).
- [60] P.C. de Holanda and A.Y. Smirnov, Phys. Rev. D 69, 113002 (2004).
- [61] A. Bandyopadhyay, S. Choubey, and S. Goswami, Phys. Rev. D 67, 113011 (2003).
- [62] H. Minakata, H. Nunokawa, W.J.C. Teves, and R. Zukanovich Funchal, Phys. Rev. D 71, 013005 (2005).