

Cherenkov radiation from e^+e^- pairs and its effect on ν_e induced showers

Sourav K. Mandal, Spencer R. Klein, and J. David Jackson

Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

(Received 8 June 2005; published 11 November 2005)

We calculate the Cherenkov radiation from an e^+e^- pair at small separations, as occurs shortly after a pair conversion. The radiation is reduced (compared to that from two independent particles) when the pair separation is smaller than the wavelength of the emitted light. We estimate the reduction in light in large electromagnetic showers, and discuss the implications for detectors that observe Cherenkov radiation from showers in the Earth's atmosphere, as well as in oceans and Antarctic ice.

DOI: [10.1103/PhysRevD.72.093003](https://doi.org/10.1103/PhysRevD.72.093003)

PACS numbers: 41.60.Bq, 95.55.Vj

I. INTRODUCTION

Cherenkov radiation from relativistic particles has been known for over 70 years [1]. However, to date, almost all studies have concentrated on the radiation from individual particles. Frank [2], Eidman [3] and Balazs [4] considered the Cherenkov radiation from electric and magnetic dipoles, but only in the limit of vanishing separations d . Their work was nicely reviewed by Jelley [5].

Several more recent calculations have considered Cherenkov radiation from entire electromagnetic showers, in the coherent or almost coherent limit [6]. The fields from the e^+ and e^- largely cancel, and the bulk of the coherent radiation is due to the net excess of e^- over e^+ (the Askaryan effect) [7]. Hadronic showers produce radiation through the same mechanism [8]. Coherent radiation occurs when the wavelength of the radiation is large compared to the radial extent of the shower; for real materials, this only occurs for radio waves.

Here, we consider another case, the reduction of radiation from slightly-separated oppositely-charged comoving pairs. This includes e^+e^- pairs produced by photon conversion. When high-energy photons convert to e^+e^- pairs, the pair opening angle is small and the e^+ and e^- separate slowly.

Near the pair, the electric and magnetic fields from the e^+ and e^- must be considered separately. However, for an observer far away from the pair (compared to the pair separation d), the electric and magnetic fields from the e^+ and e^- largely cancel. Cherenkov radiation is produced at a distance of the order of the photon wavelength Λ from the charged particle trajectory. So, for $d < \Lambda$, cancellation reduces the Cherenkov radiation from a pair to below that for two independent particles. For a typical pair opening angle m/k , where k is the photon energy and m the electron mass, without multiple scattering, $\Lambda > d$ for a distance $k\Lambda/m$. For blue light ($\Lambda = 400$ nm) from a 1 TeV pair, the radiation is reduced until the pair travels a distance of 40 cm (neglecting multiple scattering).

A similar cancellation effect was observed for energetic (~ 100 GeV) e^+e^- pairs in nuclear emulsions [9]. Ionization from newly created e^+e^- pairs is reduced

when the pair separation is less than the screening distance for ionization in the target.

In this paper, we calculate the Cherenkov radiation from e^+e^- pairs, simulate optical radiation from pairs follow realistic trajectories, and consider the radiation from electromagnetic showers. We consider two classes of experiments: underwater/in-ice neutrino observatories and air Cherenkov telescopes.

II. CHERENKOV RADIATION FROM PAIRS

Cherenkov radiation from closely spaced e^+e^- pairs can be derived by extending the derivation for point charges, by replacing a point charge with an oppositely charged, separated pair. We sketch the derivation for radiation from point charges, review previous work on radiation from infinitesimal dipoles, and derive the expression for Cherenkov radiation from a closely-spaced comoving pair.

We follow the notation and derivation from Ref. [10]. In Fourier space, the charge density ρ and current density \vec{J} from a point-charge ze propagating with speed v in the x_1 direction can be written as

$$\rho(\vec{k}, \omega) = \frac{ze}{2\pi} \delta(\omega - k_1 v); \quad \vec{J}(\vec{k}, \omega) = \vec{v} \rho(\vec{k}, \omega), \quad (1)$$

where \vec{k} is the wave vector and ω the photon energy. This current deposits energy into the medium through electromagnetic interactions. We use Maxwell's equations beyond a radius a around the particle track, where a is comparable to the average atomic separation. Then, by conservation of energy, the Cherenkov radiation power is equal to the energy flow through a cylinder of this radius, giving

$$\left(\frac{dE}{dx}\right) = -caRe \int_0^\infty B_3^*(\omega) E_1(\omega) d\omega. \quad (2)$$

E_1 is the component of \vec{E} parallel to the particle track, and B_3 is the component of \vec{B} in the x_3 direction, evaluated at an impact parameter b at a point with $x_2 = b$, $x_3 = 0$. We omit the time-phase factors for brevity.

Using the wave equations in a dielectric medium and the definition of fields, then integrating over momenta, which

eliminates the space-phase factors, one finds

$$E_1(\omega) = -\frac{ize\omega}{v^2} \left(\frac{2}{\pi}\right)^{1/2} \left[\frac{1}{\epsilon(\omega)} - \beta^2 \right] K_0(\lambda b) \quad (3)$$

where

$$\lambda^2 = \frac{\omega^2}{v^2} [1 - \beta^2 \epsilon(\omega)].$$

Similarly,

$$\begin{aligned} E_2(\omega) &= \frac{ze}{v} \left(\frac{2}{\pi}\right) \frac{\lambda}{\epsilon(\omega)} K_1(\lambda b); \\ B_3(\omega) &= \epsilon(\omega) \beta E_2(\omega). \end{aligned} \quad (4)$$

K_0 and K_1 are the zeroth and first order modified Bessel functions of the second kind.

The far-field radiation depends on the asymptotic form of the energy deposition at $|\lambda a| \gg 1$. For $\beta > 1/\sqrt{\epsilon(\omega)}$ for real $\epsilon(\omega)$, λ is completely imaginary. The asymptotic contribution of the Bessel functions in the integrand of dE/dx is finite, giving the well-known expression for the Cherenkov radiation

$$\left(\frac{dE}{dx}\right) = \frac{(ze)^2}{c^2} \int_{\epsilon(\omega) > 1/\beta^2} \omega \left(1 - \frac{1}{\beta^2 \epsilon(\omega)}\right) d\omega. \quad (5)$$

Note how a has dropped out ([10], Ch. 13). The derivation of this Cherenkov radiation may be expanded to give the field from a pair.

The radiation from an e^+e^- pairs depends on two parameters: the separation d and the angle between the direction of motion and the orientation of the pair. For relativistic pairs created by photon conversion, the transverse (to the direction of motion) separation is important; the longitudinal separation of a highly relativistic pair can be neglected, due to Lorentz length contraction.

Balazs [4] provided an expression for Cherenkov radiation from an infinitesimal dipole D oriented transverse to its momentum. The fields are approximated by a linear Taylor expansion of the corresponding point-charge fields:

$$E_1^{(D)}(\omega) = -d \frac{\partial E_1(\omega)}{\partial x_2}; \quad B_3^{(D)}(\omega) = -d \frac{\partial B_3(\omega)}{\partial x_2}$$

where d is the effective pair separation, so $D = ze d$. Then, following the same steps as in the point-charge case, Balazs finds

$$\left(\frac{dE}{dx}\right) = \frac{1}{2} \frac{D^2}{c^4} \int_{\epsilon(\omega) > 1/\beta^2} \epsilon(\omega) \omega^3 \left(1 - \frac{1}{\beta^2 \epsilon(\omega)}\right)^2 d\omega. \quad (6)$$

For a point dipole oriented parallel to its direction of motion, the radiation is negligible for $\beta \lesssim 1$ [5].

To compute the Cherenkov radiation for finite separations d , let us consider a pair moving in the $+x$ direction. The pair lies entirely in the transverse plane y - z , with the line between them making an angle α with respect to the y axis. Then, generalizing Eq. (1), the charge density from

the pair is

$$\rho(\vec{k}, \omega) = \frac{ze}{2\pi} \delta(\omega - k_1 v) [e^{-i(k_2 y_+ - k_3 z_+)} - e^{-i(k_2 y_- - k_3 z_-)}].$$

The two charges have positions relative to the center of mass

$$\begin{aligned} y_+ &= \frac{d}{2} \cos \alpha, & z_+ &= -\frac{d}{2} \sin \alpha, \\ y_- &= -\frac{d}{2} \cos \alpha, & z_- &= \frac{d}{2} \sin \alpha. \end{aligned}$$

The angle α is the relative azimuth between the line connecting the two charges and the azimuth of observation.

The generalization of $E_1(\omega)$ of Eq. (3) is

$$\begin{aligned} E_1(\omega) &= \frac{-ize\omega}{v^2} \sqrt{\frac{\pi}{2}} \left(\frac{1}{\epsilon(\omega)} - \beta^2 \right) \\ &\quad \times [K_0(\lambda b_-) - K_0(\lambda b_+)], \end{aligned} \quad (7)$$

where

$$b_{\pm} = \sqrt{\frac{d^2}{4} \sin^2 \alpha + \left(b \pm \frac{d}{2} \cos \alpha\right)^2}.$$

As before, we take $|\lambda a| \gg 1$ and $a < b$, so we need only consider $d \ll b$; there is little interference for $d \gtrsim b$. Therefore, we can simplify using

$$b_{\pm} \simeq b \pm \frac{d}{2} \cos \alpha.$$

Then, as before, considering completely imaginary λ and $|\lambda a| \gg 1$,

$$\begin{aligned} E_1(\omega) &= \frac{2ze\omega}{c^2} \left(1 - \frac{1}{\beta^2 \epsilon(\omega)}\right) \\ &\quad \times \sqrt{\frac{i}{|\lambda|}} \frac{e^{i\lambda b}}{b} \sin \left[\frac{d}{2} |\lambda| \cos \alpha \right] \end{aligned} \quad (8)$$

and a similar expression for $B_3(\omega)$. Here we have taken $b_- \simeq b_+ \simeq b$ in the denominator.

At $\alpha = \pm \pi/2$, $E_1(\omega) = 0$. The Cherenkov radiation is no longer symmetric about the direction of motion, and vanishes at right angles to the direction of the dipole. As the charge separation increases (or the wavelength decreases), the angular distribution evolves from two wide lobes into a many-lobed structure, as shown in Fig. 1. After integration over even a narrow range of ω or d , the angular distribution becomes an almost-complete disk, with two narrow zeroes remaining at a direction perpendicular to the dipole vector.

After assembling the pieces, and averaging over α , we find the generalization of Eq. (5),

$$\begin{aligned} \left(\frac{dE}{dx}\right) &= \frac{(ze)^2}{c^2} \int_{\epsilon(\omega) > 1/\beta^2} d\omega \omega \left(1 - \frac{1}{\beta \epsilon(\omega)}\right) \\ &\quad \times 2[1 - J_0(\lambda d)]. \end{aligned}$$




FIG. 1. The azimuthal angular distribution (transverse to the direction of motion) of Cherenkov radiation for 500 nm photons from a pair of charges oriented as shown in the figure. Distributions are shown for pair separations 100 nm (solid line), 1 μm (dashed line) and 5 μm (dotted line), with $\sqrt{\epsilon(\omega)} = n = 1.3$ and $\beta = 1$.

Here J_0 is the zeroth order Bessel function of the first kind. For $\lambda d \ll 1$, this reproduces Eq. (6). For $\lambda d \gg 1$, the dE/dx is twice that expected for an independent particle [Eq. (5)]. The transition is shown in Fig. 2. As the emission wavelength Λ approaches d , the pair spectrum converges to the point-charge spectrum in an oscillatory fashion, characteristic of the Bessel function. For certain values of λd , the radiation exceeds that of two independent charged particles.




FIG. 2. The spectrum of Cherenkov radiation at $\beta = 1$, $\sqrt{\epsilon(\omega)} = n = 1.3$. Solid line is for e^+e^- with the particles considered independently, and the dashed lines are for pairs treated coherently, with separations 100 nm, 1 μm and 5 μm .

For the remainder of the paper, we assume that media satisfy $\sqrt{\epsilon(\omega)} = n$, where n is independent of frequency. In realistic detection media, any variation of n with frequency is small, and would have little effect on Cherenkov radiation from relativistic particles.

With real e^+e^- pairs, two effects should be considered. Electromagnetic radiation is not emitted instantaneously, but occurs while the radiating particles travel a distance known as the formation length, l_f . For Cherenkov radiation, $l_f = \Lambda/\sin^2(\theta_C) = \Lambda\epsilon\beta^2/(\epsilon\beta^2 - 1)$ [11], depends only on the Cherenkov emission angle, θ_C and the photon wavelength; l_f depends only slightly (through β) on the electron energy.

While the pair is covering the distance l_f , the pair separation will change by an amount $\Delta d = l_f \sin(\theta)$, where θ is the angle between the e^+ and e^- velocity vectors. Since θ is of order $1/\gamma$, $\Delta d/d \ll 1$, so the change in separation is not significant.

Second, the Cherenkov radiation produced at a point (x -coordinate) depends on the fields emitted by the charged particles at earlier times, when d may be different than at the point of radiation. For full rigor, these retarded separations should be used in the calculation. Again, this has a negligible effect on the results.

III. RADIATION FROM e^+e^- PAIRS IN SHOWERS

Many experiments study Cherenkov radiation from large electromagnetic showers. The radiation from a shower may be less than would be expected if every particle were treated as independent. We use a simple simulation to consider 300 to 800 nm radiation from electromagnetic showers. This frequency range is typical for photomultiplier based Cherenkov detectors; at longer wavelength, there is little radiation, while shorter wavelength light is absorbed by the glass in the phototube.

We simulated 1000 γ conversions to e^+e^- pairs with total energies from 10^8 to 10^{20} eV. Pairs were produced with the energy partitioned between the e^+ and e^- following the Bethe-Heitler differential cross section $d\sigma \approx E_{\pm}(1 - E_{\pm})$, where E_{\pm} is the electron (or positron energy) [12]. At high energies in dense media (above 10^{16} eV in water or ice), the Landau-Pomeranchuk-Migdal (LPM) effect becomes important, and more asymmetric pairs predominate [13]. The pairs are generated with initial opening angle of m/k ; the fixed angle is a simplification, but the pair separation is dominated by multiple scattering, so it has little effect on our results.

The e^- and e^+ are tracked through a water medium (with $n = \sqrt{\epsilon} = 1.3$) in steps of $0.02X_0$, where X_0 is the radiation length, 36.1 cm in water. At each step, the particles multiple scatter, following a Gaussian approximation ([14], Ch. 27). The particles radiate bremsstrahlung photons, using a simplified model where photon emission follows a Poisson distribution, with mean free path X_0 .




FIG. 3. Average Cherenkov photon emission rate for pairs with energies from 10^{10} (leftmost, dot-dashed curve) to 10^{15} eV (rightmost, dotted curve) vs the distance traveled by the pair in water, relative to emission from two independent particles.

Although this model has almost no soft bremsstrahlung, soft emission has little effect on Cherenkov radiation, since the electron or positron velocity is only slightly affected.

At each step, we compute the Cherenkov radiation for each pair. They are treated coherently when $d < 2\lambda$; at larger separations the particles radiate independently.

As shown in Fig. 3, the particles in lower energy pairs ($< 10^{10}$ eV) radiate almost independently. In contrast, the radiation from very high-energy pairs ($> 10^{15}$ eV) is largely suppressed. The broad excursions slightly above unity occur when $J_0(\lambda d) > 1$ for many of the scattered pairs.

IV. IMPLICATIONS FOR EXPERIMENTS

At least two types of astrophysical observatories depend on Cherenkov radiation. Water and ice based neutrino observatories observe Cherenkov radiation from the charged particles produced in neutrino interactions, and air Cherenkov telescopes look for γ -ray induced electromagnetic showers in the Earth's atmosphere.

Current neutrino observatories can search for electron neutrinos with energies above 50 TeV (for ν_μ , the threshold is much lower) [15]. They use large arrays of photomultiplier tubes to observe the Cherenkov radiation from ν_e induced showers. For water, $n \approx 1.3$, Fig. 3 shows that $\lambda d < 1$ while the pair travels significant distances. Ice is similar to water, with a slightly lower density; n of ice depends on its structure, and is typically ≈ 1.29 [16].

To quantify the effect of Cherenkov radiation from ν_e interactions, we use a toy model of an electromagnetic shower. The shower evolves through generations, with each generation having twice as many particles as the preceding generation, with half the energy. Each generation evolves over a distance of X_0 ; other simulations have

evolved generations over a shorter distance $(\ln 2)X_0$, leading to a more compact shower [17]. In these showers, most of the particles are produced in the last radiation lengths.

Figure 4 shows the Cherenkov radiation expected from a model 10^{20} eV shower with coherent Cherenkov radiation (solid line) and in a model where all particles radiate independently (dotted line). This model does not include the LPM effect, so it should be considered only illustrative. The LPM effect lengthens the high-energy (above a few 10^{15} eV) portion of the shower. By spreading the shower longitudinally, the LPM effect will give the electrons and positrons more time to separate, and so will somewhat lessen the difference between the two results. However, it is clear from Fig. 4 that coherence has a significant effect for the first ≈ 22 generations. Since the front of the shower contains relatively few particles, it will not affect the measured energy; the change in number of radiated photons (and hence on the energy measurement) should be less than 1%.

However, the suppression will affect the apparent length of the shower. For the first ≈ 8 generations, the shower will emit less light than a single charged particle. Because of the LPM effect, each of these generations (with mean particle energy E_g above a few 10^{15} eV) develop over a distance $X = X_0 \sqrt{E_g / 5E_{LPM}}$, where $E_{LPM} = 278$ TeV for water is the effective LPM energy [17], greatly elongating the shower. So, the first 8 generations include most of the length of the shower. So, the suppression of Cherenkov radiation hides the initial shower development, making the shower appear considerably more compact. The reduction in early-stage radiation should help in separating electron cascades from muon-related backgrounds, especially muons that undergo hard interactions, and lose a large fraction of their energy.




FIG. 4. Cherenkov radiation from a 10^{20} eV shower in water, using the Heitler toy model, versus shower depth (smoothed). The two curves compare the radiation for e^+e^- calculated as independent particles and as coherent pairs.

Atmospheric Cherenkov telescopes like the Whipple observatory study astrophysical γ rays with energies from 100 GeV to 10 TeV. These telescopes observe Cherenkov radiation from pairs in the upper atmosphere; for a 1 TeV shower, the maximum particle density occurs at an altitude of 8 km above sea level (asl) [18], where the density is about 1/3 that at sea level. Since $n - 1$ depends linearly on the density, at 8 km asl $n - 1 \approx 1 \times 10^{-4}$, so for 500 nm photons radiated from ultrarelativistic particles, $\lambda d < 1$ only for $d < 6 \mu\text{m}$. In this low-density medium, the effect of the pair opening angle is significant and multiple scattering is less important. Pairs with $k < 1$ TeV will separate by 30 μm in a distance less than 30 meters; at 8 km asl, this is 3% of a radiation length. This distance is too short to affect the radiation pattern from the shower.

Cherenkov radiation is also used in lead-glass block calorimetry, and in Cherenkov counters for particle identification; their response to photon conversions may be affected by this coherence.

Although the reactions are slightly different, a similar analysis applies to the reduction of ionization by e^+e^- pairs. Perkins observed that the ionization from pairs with mean energy 180 GeV in emulsion was suppressed for the first $\approx 250 \mu\text{m}$ after the pairs were created [9]. With $X_0 = 3$ cm (typical for emulsion), the e^+ and e^- trajectories will be about 4 nm apart after traveling 250 μm . For relativistic

particles, the screening distance (effective range for dE/dx) is determined by the plasma frequency of the medium, ω_p . For silver bromide, the dominant component of emulsion, $\hbar\omega_p = 48$ eV [10] (in a complete emulsion, $\hbar\omega_p$ will be slightly lower). This yields a screening distance $c/\omega_p = 4$ nm, which is very close to the calculated separation.

V. CONCLUSION

We have calculated the Cherenkov radiation from e^+e^- pairs as a function of the pair separation d . When $d^2 < v^2/(\omega^2[1 - \beta^2\epsilon(\omega)])$, the radiation is suppressed compared to that from two independent particles.

This suppression affects the radiation from electromagnetic showers in dense media. Although the total radiation from a shower is not affected, emission from the front part of the shower is greatly reduced; this will affect studies of the shower development, and may affect measurements of the position of the shower.

ACKNOWLEDGMENTS

This work was funded by the U.S. National Science Foundation under Grant No. OPP-0236449 and the U.S. Department of Energy under Contract No. DE-AC-76SF00098.

-
- [1] P. A. Cherenkov, Dokl. Akad. Nauk SSSR **2**, 451 (1934); S. I. Vavilov, Dokl. Akad. Nauk SSSR **2**, 457 (1934); I. M. Frank and Ig. Tamm, Dokl. Akad. Nauk SSSR **14**, 109 (1937).
 - [2] I. M. Frank, Zh. Fiz. SSR **7**, 49 (1943).
 - [3] V. Eidman, in Transactions of the Gorky Research Physiotechical Institute and Radiophysical Faculty of the Gorky State University, Scientific Notes 30, 1956.
 - [4] N. L. Balazs, Phys. Rev. **104**, 1220 (1956).
 - [5] J. V. Jelley, *Cherenkov Radiation and Its Applications* (Pergamon Press, New York, 1958).
 - [6] E. Zas, F. Halzen, and T. Stanev, Phys. Rev. D **45**, 362 (1992); J. Alvarez-Muniz *et al.*, Phys. Rev. D **68**, 043001 (2003); S. Razzaque *et al.*, Phys. Rev. D **69**, 047101 (2004); A. R. Beresnyak, astro-ph/0310295.
 - [7] G. A. Askaryan and B. A. Dolgoshein, JETP Lett. **25**, 213 (1977).
 - [8] J. Alvarez-Muniz and E. Zas, Phys. Lett. B **434**, 396 (1998).
 - [9] D. H. Perkins, Philos. Mag. **46**, 1146 (1955).
 - [10] J. D. Jackson, *Classical Electrodynamics* (John Wiley & Sons, New York, 1998), 3rd ed.
 - [11] M. S. Zolotarev and K. T. McDonald, physics/0003096.
 - [12] H. A. Bethe and W. Heitler, Proc. R. Soc. A **146**, 83 (1934).
 - [13] S. Klein, Rev. Mod. Phys. **71**, 1501 (1999); S. Klein, hep-ex/0402028.
 - [14] S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
 - [15] M. Ackermann, Astropart. Phys. **22**, 127 (2004); Icecube Collaboration, IceCube Preliminary Design Document, 2001. Available at www.icecube.wisc.edu
 - [16] P. B. Price and K. Woschnagg, Astropart. Phys. **15**, 97 (2001).
 - [17] S. Klein, astro-ph/0412546.
 - [18] C. M. Hoffman, C. Sinnis, P. Fleury, and M. Punch, Rev. Mod. Phys. **71**, 897 (1999).