

Unified framework for symmetry breaking in SO(10)K. S. Babu,¹ Iliia Gogoladze,^{2,*} Pran Nath,^{3,4} and Raza M. Syed³¹*Department of Physics, Oklahoma State University, Stillwater, Oklahoma 74078, USA*²*Department of Physics, University of Notre Dame, Notre Dame, Indiana 46556, USA*³*Department of Physics, Northeastern University, Boston, Massachusetts 02115-5000, USA*⁴*Max Planck Institute for Physics, Fohringer Ring 6, D-80805, Munich, Germany*

(Received 30 June 2005; published 16 November 2005)

A new SO(10) unified model is proposed based on a one-step breaking of SO(10) to the standard model gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ using a single 144 of Higgs. The symmetry breaking occurs when the SU(5) 24-plet component of 144 develops a vacuum expectation value. Further, it is possible to obtain from the same 144 a light Higgs doublet necessary for electroweak symmetry breaking using recent ideas of string vacua landscapes and fine-tuning. Thus the breaking of SO(10) down to $SU(3)_C \times U(1)_{em}$ can be accomplished with a single Higgs. We analyze this symmetry breaking pattern in the nonsupersymmetric as well as in the supersymmetric SO(10) model. In this scenario masses of the quarks and leptons arise via quartic couplings. We show that the resulting mass pattern is consistent with experimental data, including neutrino oscillations. The model represents an alternative to the currently popular grand unified scenarios.

DOI: [10.1103/PhysRevD.72.095011](https://doi.org/10.1103/PhysRevD.72.095011)

PACS numbers: 12.10.Dm, 12.10.Kt

I. INTRODUCTION

In any grand unified theory (GUT) understanding the Higgs sector is not an easy task. Usually these models require the existence of more than one Higgs multiplet. In the minimal SU(5) GUT one employs one adjoint 24-plet (Σ) and a fundamental 5-plet to break the GUT symmetry down to $SU(3)_C \times U(1)_{em}$. The Yukawa couplings of the 5-plet Higgs also generate quark and lepton masses. The Higgs sector becomes somewhat more complicated in larger GUT structures such as SO(10) [1] since there is a larger symmetry that needs to be broken. Conventional SO(10) models employ at least two different Higgs representations to break the symmetry down to $SU(3)_C \times SU(2)_L \times U(1)_Y$ (a 16 or a 126 to change rank, and one of 45, 54, or a 210 to break the symmetry down further [2]). To achieve electroweak symmetry breaking and to generate quark and lepton masses an additional 10-dimensional representation is also needed. A minimal SO(10) model studied recently, for example, utilizes one 10, one 126, and one 210 Higgs representations to achieve symmetry breaking and to generate masses for the quarks, leptons, and the neutrinos [3].

In this paper we discuss the following question: Is it possible to achieve SO(10) symmetry breaking all the way down to the $SU(3)_C \times U(1)_{em}$ with a single Higgs representation? We find that this is indeed the case if one employs a 144-plet of Higgs of SO(10). The 144-plet is contained in the product 10×16 , and thus carries one vector and one spinor index. An interesting property of the 144-plet which makes such a symmetry breaking chain possible is that it contains in it an SU(5) adjoint with a U(1) charge, as well as standard model Higgs doublet fields.

This can be seen from the following decomposition of 144 under $SU(5) \times U(1)$ subgroup of SO(10)

$$144 = \bar{5}(3) + 5(7) + 10(-1) + 15(-1) + 24(-5) + 40(-1) + \overline{45}(3). \quad (1)$$

It is significant that the SU(5) adjoint 24(-5) above also carries a U(1) charge. Once the standard model singlet in it acquires a vacuum expectation value (VEV), it would change the rank of the group, leading to a one-step breaking of SO(10) down to $SU(3)_C \times SU(2)_L \times U(1)_Y$. The submultiplets $\bar{5}(3)$, $5(7)$, and $\overline{45}(3)$ all contain fields with identical quantum numbers as the standard model Higgs doublet. If one combination of doublets from these submultiplets is made light by fine-tuning, it can be used for electroweak symmetry breaking. Such fine-tuning is justified in the context of the multiple vacua of the string landscapes, which has been widely discussed recently. Although consistency of the first stage of symmetry breaking requires the mass-squared of all the physical Higgs particles to be positive (including that of the light Higgs doublet), we show that radiative corrections involving the Higgs self-couplings can turn the mass-squared of the light Higgs doublet negative, facilitating the second stage of symmetry breaking.

Realistic fermion masses can be obtained within this minimal scenario. Recall that the fermions of each family belong to the 16-dimensional spinor representation of SO(10). Under $SU(5) \times U(1)$ subgroup the 16 decomposes as follows

$$16 = 10(-1) + \bar{5}(3) + 1(-5). \quad (2)$$

Fermion masses will arise from quartic couplings of the $16_i 16_j (144_{144})$ and $16_i 16_j (144^* 144^*)$. These couplings

*On leave of absence from Andronikashvili Institute of Physics, GAS, 380077 Tbilisi, GA.

would lead to Dirac masses for all the fermions as well as large Majorana masses for the right-handed neutrinos. Since the light Higgs doublet is a linear combination of doublets from 5 and 45 of SU(5), the resulting mass pattern is not that of minimal SU(5) and is consistent with experimental data, including neutrino oscillations.

The outline of the rest of the paper is as follows: In Sec. II we discuss symmetry and mass growth in the SU(5) × U(1) language. In Sec. III we discuss the techniques of calculation for the analysis of 144 and $\overline{144}$ -plet couplings using the method developed in Ref. [4].¹ Here also we discuss the set of couplings $(144 \times \overline{144})_1$, $(144 \times \overline{144})_1(144 \times \overline{144})_1$, $(144 \times \overline{144})_{45}(144 \times \overline{144})_{45}$, and $(144 \times \overline{144})_{210}(144 \times \overline{144})_{210}$, where $(144 \times \overline{144})_1$ means a singlet in the $(144 \times \overline{144})$ coupling and $(144 \times \overline{144})_1 \times (144 \times \overline{144})_1$ are couplings generated by integrating out the singlet field, and similarly for couplings with subscripts 45 and 210. These couplings are needed in the computation of symmetry breaking which is then analyzed. In Sec. IV Higgs phenomenon and mass growth are analyzed for the breaking of SO(10). Here it is shown that within the landscape scenario with fine-tuning [7,8] one gets a pair of light Higgs doublets exactly as in the minimal supersymmetric standard model (MSSM) while the Higgs triplets and other modes are either absorbed or become super heavy. In Sec. V couplings of quarks and leptons are discussed and it is shown that such couplings are quartic in nature. As an illustration the couplings involving $(16 \times 16)_{10}(144 \times 144)_{10}$ and $(16 \times 16)_{10}(\overline{144} \times \overline{144})_{10}$ are explicitly discussed. It is shown that the resulting masses and mixings are consistent with experimental data. Conclusions are given in Sec. VI.

II. SYMMETRY BREAKING AND MASS GROWTH WITH 144 IN THE SU(5) × U(1) LANGUAGE

Analysis of the symmetry breaking and of fermion mass generation with a 144 of Higgs turns out to be rather complicated. Before we delve into the full detail in the SO(10) language, which is presented in the next section, we analyze here these issues in the simpler SU(5) × U(1) subgroup language. We will present our analysis in a non-supersymmetric model. Generalization to supersymmetry (SUSY) require the addition of a $\overline{144}$ chiral multiplet, so that the flatness of the D-term potential can be maintained at the GUT scale, leaving supersymmetry intact at that scale. The analysis in this section would also hold for SUSY models with some redefinitions of parameters, provided that the 144* field is identified with the $\overline{144}$ of the SUSY SO(10) model.

¹For further application of the technique of Ref. [4], see Ref. [5]. For later works using other techniques for the computation of SO(10) couplings, see Ref. [6].

A. One-step GUT symmetry breaking

Since in the SU(5) × U(1) decomposition of SO(10) the 144 contains an SU(5) adjoint carrying a nonzero U(1) charge (see Eq. (1)), it is instructive to analyze symmetry breaking of SU(5) × U(1) with a complex adjoint Σ . One can construct such a representation from two adjoint representations of SU(5): $\Sigma = \Sigma_1 + i\Sigma_2$. Then Σ is not self-adjoint, and we denote the conjugate of Σ as Σ^\dagger .

The most general SU(5) × U(1) invariant renormalizable potential involving the Σ and Σ^\dagger fields is

$$V = -M^2 \text{tr}(\Sigma \Sigma^\dagger) + \frac{\kappa_1}{2} \text{tr}(\Sigma^2 \Sigma^{\dagger 2}) + \frac{\kappa_2}{2} (\text{tr}(\Sigma \Sigma^\dagger))^2 + \frac{\kappa_3}{2} \text{tr}(\Sigma^2) \text{tr}(\Sigma^{\dagger 2}) + \frac{\kappa_4}{2} \text{tr}(\Sigma \Sigma^\dagger \Sigma \Sigma^\dagger). \quad (3)$$

Among the possible local minima is the one which preserves the standard model gauge symmetry given by the vacuum structure

$$\langle \Sigma \rangle = \langle \Sigma^\dagger \rangle = v \text{diag}(2, 2, 2, -3, -3). \quad (4)$$

For some range of the parameters of the potential, this minimum will be the global minimum. Minimization of the potential gives

$$v^2 = \frac{M^2}{7(\kappa_1 + \kappa_4) + 30(\kappa_2 + \kappa_3)}. \quad (5)$$

Clearly this VEV structure breaks SU(5) down to SU(3)_C × SU(2)_L × U(1)_Y. Further, since Σ is charged under the U(1), its VEV breaks this U(1). Thus we see that the SU(5) × U(1) symmetry is broken down to the SM gauge symmetry in one step with one complex adjoint Higgs field. This can also be verified directly by computing the gauge boson masses. The physical Higgs boson masses can all be made positive for some range of parameters of the potential. It is then clear that if an SO(10) representation contains submultiplets which transform under SU(5) × U(1) symmetry as an adjoint carrying U(1) charge, then there is the possibility that this Higgs field can break SO(10) all the way down to the SM gauge symmetry in one step. We observe that the 144-plet of SO(10) is the simplest representation which has this property.² Technical details of this assertion in the SO(10) language is postponed to the next section.

Identical conclusions can be arrived at for the case of supersymmetric SU(5) × U(1) model. The potential of Eq. (3) will become the superpotential (with Σ^* replaced by a chiral superfield $\overline{\Sigma}$), the couplings κ_i will be non-renormalizable operators with inverse dimensions of mass, and the mass term M^2 will be replaced by M . Thus we conclude that in SUSY SO(10) a 144 + $\overline{144}$ pair of chiral

²The next simplest possibility is to have a 560 of SO(10), which contains a 24(-5), 1(-5), as well as $\overline{5}(3)$, 45(7), and $\overline{45}(3)$ under SU(5) × U(1).

superfields can break SO(10) in one step down to the supersymmetric standard model gauge group.

B. Electroweak symmetry breaking

Having recognized that a single 144-plet can break non-supersymmetric SO(10) down to the SM in one step, we turn our attention to the subsequent electroweak symmetry breaking. As noted earlier, the 144-plet also contains fields which have the same quantum numbers as the SM Higgs doublet. We explain how these doublet fragments from the 144-plet can be used for the purpose of electroweak symmetry breaking, thus making the model very economical.

An immediate question that can be raised is how to obtain negative mass-squared for the light Higgs doublet of the SM arising from the 144-plet. Consistency of the GUT symmetry breaking would require positivity of the mass-squared of all the physical Higgs bosons, including that of the light SM doublet. If the surviving symmetry and spectrum below the GUT scale are corresponding to those of the SM, there would be no interactions that turn the Higgs mass-squared negative needed for electroweak symmetry breaking. In analogy to the stop squark quartic couplings turning the Higgs mass-squared negative in the supersymmetric SM, we observe that the quartic couplings between the light Higgs doublet and any fragment of the 144-plet with mass an order of magnitude or so below the GUT scale can turn the Higgs doublet mass-squared negative.³ We illustrate this mechanism with a simple SU(5) toy model with an adjoint Higgs field below.

Consider a toy model with global SU(5) symmetry broken spontaneously by an adjoint Higgs field Σ . The most general renormalizable potential for this field is given by

$$V = m^2 \text{Tr}(\Sigma^2) + \frac{\kappa_1}{2} \text{tr}(\Sigma^4) + \frac{\kappa_2}{2} (\text{tr}(\Sigma^2))^2 + \mu \text{tr}(\Sigma^3). \quad (6)$$

One possible VEV structure is as shown in Eq. (4). Minimization of the potential of Eq. (6) gives

$$m^2 = -(7\kappa_1 + 30\kappa_2)v^2 + \frac{3}{2}\mu v. \quad (7)$$

The (3,2, -5/6) and (3*,2,5/6) components of Σ (under the surviving SU(3) \times SU(2) \times U(1) symmetry) are Goldstone bosons, while the (8,1,0), (1,3,0), and the (1,1,0) components have masses given, respectively, by

$$m_8^2 = \frac{15}{2}\mu v + 5\kappa_1 v^2, \quad m_3^2 = -\frac{15}{2}\mu v + 20\kappa_1 v^2, \\ m_1^2 = -\frac{3}{2}\mu v + (14\kappa_1 + 60\kappa_2)v^2. \quad (8)$$

For a range of parameters, all three squared-masses can be chosen positive, establishing the consistency of symmetry breaking.

It is possible by fine-tuning to make the SU(2)_L triplet field to be much lighter than the SU(5) breaking scale, while keeping the other two components heavy. We wish to ask if such a finely tuned triplet can subsequently break SU(2)_L further down to U(1)_L. This would, however, require that m_3^2 turn negative at lower scales even though it starts off as being positive at the high scale. Consider the case when κ_1 and μ/v are somewhat smaller than 1 (say of order 0.01), while κ_2 is of order one. Then m_3 and m_8 are generically an order of magnitude below the GUT scale v , while m_1 is of order v . In the momentum range below v and above m_8 , the mass parameters m_3^2 and m_8^2 will evolve, while the singlet decouples. The renormalization group equations (RGE) for the running of these mass parameters are found to be

$$\frac{dm_8^2}{dt} = \frac{1}{8\pi^2} \left[\frac{15}{8}(\mu + 4\kappa_1 v)^2 + \frac{5}{2}(\kappa_1 + 2\kappa_2)m_8^2 + \frac{3}{2}\kappa_2 m_3^2 \right], \quad (9)$$

$$\frac{dm_3^2}{dt} = \frac{1}{8\pi^2} \left[4\kappa_2 m_8^2 + \frac{5}{2} \left(\frac{1}{2}\kappa_1 + \kappa_2 \right) m_3^2 \right],$$

where $t = \log(Q)$, where Q is the renormalization group scale. With κ_2 being of order one and $\kappa_1, \mu/v \ll 1$, we see from these equations that if m_3^2 at the scale v starts off being smaller than m_8^2 , it can turn negative in going down from v to the mass scale m_8 . The mass parameter m_8^2 will remain positive in this case. This example shows that fine-tuning of the weak triplet can be done at the scale m_8 in such a way that its squared-mass turns negative at lower energy scales.

In analogy with this example, if any fragment of the 144-plet of SO(10) remains somewhat lighter than the GUT scale, then the quartic couplings involving that fragment and the Higgs doublet would turn the doublet mass-squared negative.⁴ It is interesting to note that if such fragments from the 144 that survive below the GUT scale are the color octet(s) and the weak triplet(s), unification of the three SM gauge couplings will occur nicely at a scale of ($10^{16} - 10^{17}$) GeV [10]. The above analysis and conclusions are in the context of a non-SUSY SO(10) scenario. In the SUSY SO(10) case the top-stop Yukawa couplings will

³Yukawa couplings between the Higgs doublet and fermions cannot turn the Higgs mass-squared negative. Cubic self-couplings (which are not allowed in the SM Higgs doublet) and/or quartic/cubic scalar couplings involving other fields are necessary for this to happen.

⁴This statement does not contradict Michel-Radicati theorem [9] as we are using radiative corrections to turn the doublet mass-squared negative.

turn the Higgs doublet mass-square negative in the usual way.

C. Doublet-triplet splitting

We denote the components of the 144 multiplet by Q 's. As can be seen from Eq. (1) the Higgs fields reside in the multiplets $Q_i(\bar{5}) + Q^i(5) + Q_j^i(24) + Q_{ij}^k(\bar{45})$. Similarly we denote the components of the 144* multiplet by P 's

$$\begin{aligned}
 V_{DT} = & m^2 \text{tr}(Q_i P^i) + m^2 \text{tr}(Q^i P_i) + m^2 \text{tr}(Q_{ij}^k P_k^{ij}) + \lambda_1 \text{tr}(Q_i Q^i) \text{tr}(Q_j^k Q_k^j) + \lambda_2 \text{tr}(P_i P^i) \text{tr}(P_j^k P_k^j) + \lambda_3 \text{tr}(Q_i P^i) \text{tr}(Q_j^k P_k^j) \\
 & + \lambda_3' \text{tr}(Q^i P_i) \text{tr}(Q_j^k P_k^j) + \lambda_4 \text{tr}(Q_{ij}^k P_k^{ij}) \text{tr}(Q_n^m P_m^n) + \eta_1 \text{tr}(Q_i Q_j^i Q_k^j Q^k) + \eta_2 \text{tr}(P_i P_j^i P_k^j P^k) + \eta_3 \text{tr}(Q_i P_j^i Q_k^j P^k) \\
 & + \eta_3' \text{tr}(P_i Q_j^i P_k^j Q^k) + \eta_4 \text{tr}(Q_{ij}^k Q_k^i P_m^j P^m) + \eta_4' \text{tr}(P_k^{ij} P_i^k P_j^m P_m) + \eta_5 \text{tr}(Q_{ij}^k Q_k^i Q_m^j Q^m) + \eta_5' \text{tr}(P_k^{ij} Q_k^i P_m^j Q_m) \\
 & + \eta_6 \text{tr}(P_k^{ij} Q_{lm}^k Q_l^i P_j^m),
 \end{aligned} \tag{10}$$

where λ_i and η_i are dimensionless couplings which represent different types of contractions of indices. It is easily checked that each term in the potential has an overall zero U(1) quantum number. There are several Higgs doublets and Higgs triplets and antitriplets in this model. Thus one set of Higgs doublets and triplets and antitriplets arise from Q_i, Q^i, P_i, P^i . Specifically the doublets are $Q_\alpha, Q^\alpha, P_\alpha, P^\alpha$, while the triplets (antitriplets) are $Q^a, P^a (Q_a, P_a)$, where $a = 1, 2, 3$ is the color index, and $\alpha = 1, 2$ is the SU(2) index. Additional Higgs doublets, triplets, and antitriplets arise from the 45 of SU(5) (from the $\bar{144}$ -plet) and from the $\bar{45}$ (from the 144-plet). We discuss the decomposition of these below. The 45 of SU(5) embedded in $\bar{144}$ has the following SU(2) \times SU(3) \times U(1)_Y decomposition

$$45 = (2, 1)(3) + (1, 3)(-2) + (3, 3)(-2) + (1, \bar{3})(8) + (2, \bar{3})(-7) + (1, \bar{6})(-2) + (2, 8)(3). \tag{11}$$

One notices that while there is only one SU(2) Higgs doublet (\tilde{P}^α), there is one SU(3)_C Higgs triplet \tilde{P}^a and one antitriplet \tilde{P}_a . Similarly, for the $\bar{45}$ embedded in 144 one has one SU(2) Higgs doublet \tilde{Q}_α , one SU(3)_C Higgs triplet \tilde{Q}^a , and one antitriplet \tilde{Q}_a . The above analysis shows that the Higgs doublet mass matrix will be 3×3 while the Higgs triplet mass matrix will be 4×4 . We focus first on the Higgs doublet mass matrix after Q_j^i and P_j^i develop VEVs. We display the mass matrix for the Higgs doublets in the basis where the rows are $(P_\alpha, Q_\alpha, h_\alpha)$ and the columns by $(P^\alpha, Q^\alpha, h^\alpha)$

$$\begin{bmatrix} 30\lambda_2 v^2 + 9\eta_2 v^2 & m^2 + 30\lambda_3' v^2 + 9\eta_3' v^2 & v^2 \eta_4' c \\ m^2 + 30\lambda_3 v^2 + 9\eta_3 v^2 & 30\lambda_1 v^2 + 9\eta_1 v^2 & v^2 \eta_5' c \\ v^2 \eta_4 c & v^2 \eta_5 c & m^2 + 30\lambda_4 v^2 + \frac{21}{4} \eta_6 v^2 \end{bmatrix}, \tag{12}$$

where $c = -15\sqrt{3}/2\sqrt{2}$. Next, we display the mass matrix for the Higgs triplets in the basis where the rows are labeled $(P_a, Q_a, \tilde{Q}_a, \tilde{P}_a)$ and the columns by $(P^a, Q^a, \tilde{P}^a, \tilde{Q}^a)$. In this basis the Higgs triplet mass matrix is

$$\begin{bmatrix} 30\lambda_2 v^2 + 4\eta_2 v^2 & m^2 + 30\lambda_3' v^2 + 4\eta_3' v^2 & \tilde{c} \eta_4' v^2 & 0 \\ m^2 + 30\lambda_3 v^2 + 4\eta_3 v^2 & 30\lambda_1 v^2 + 4\eta_1 v^2 & \tilde{c} \eta_5' v^2 & 0 \\ \tilde{c} \eta_4 v^2 & \tilde{c} \eta_5 v^2 & m^2 + 30\lambda_4 v^2 + 2\eta_6 v^2 & 0 \\ 0 & 0 & 0 & m^2 + 30\lambda_4 v^2 + 9\eta_6 v^2 \end{bmatrix}, \tag{13}$$

where $\tilde{c} = 5\sqrt{2}$. Equation (12) has several arbitrary parameters $\lambda_1, \lambda_2, \eta_1, \eta_2$, etc., and one can utilize these to make the determinant of Eq. (12) vanish, while the determinant of Eq. (13) is nonvanishing. Because of this, and the fact that the Clebsch coefficients are different in Eq. (12) compared to Eq. (13), one can arrange for a pair of light Higgs doublets while keeping the Higgs triplets heavy. As will be shown in Eqs. (50) and (51) from the full SO(10) analysis, we can identify one of the superposition of the doublet fragments as the SM Higgs doublet while keeping

and in this case the relevant Higgs multiplets will be $P_i(\bar{5}) + P^i(5) + P_j^i(24) + P_k^{ij}(45)$. The $\bar{5}, 5$, and 45 representations contain SU(2)_L doublets and SU(3)_C triplets. Before studying the doublet-triplet splitting in the full SO(10) theory, here we analyze this possibility within the simpler SU(5) \times U(1) theory, but with couplings motivated by the full SO(10) theory. The relevant potential that we consider is given by

the Higgs color triplets fragments all super heavy. It is true that one must fine-tune in order to have the Higgs doublet light, but we find it very interesting that with a single 144-plet complete breaking of the SO(10) symmetry down to the residual SU(3)_C \times U(1)_{em} can be achieved.

D. Fermion mass growth

For the fermion masses we have the following quartic coupling allowed by gauge invariance, in terms of

SU(5) \times U(1) decomposition

$$W = \frac{h_1}{M} \epsilon_{ijkln} 10^{ij} 10^{kl} \Sigma_m^n Q^m + \frac{h_2}{M} \epsilon_{ijknm} 10^{ij} 10^{kl} \Sigma_l^n Q^m + \frac{h_3}{M} 10^{ij} \bar{5}_i \Sigma_j^m \bar{P}_m + \frac{h_4}{M} 10^{ij} \bar{5}_i \Sigma_l^j \bar{P}_j, \quad (14)$$

where 10^{ij} and $\bar{5}_i$ are the 10-plet and $\bar{5}$ -plet of quark-lepton fields and M is the super heavy mass introduced in Eq. (3). From Eq. (14) we see that it is possible to realize Georgi-Jarlskog type relations with appropriate choice of the h_i couplings even without the 45 of SU(5) acquiring electro-weak VEV. In general, there are additional terms involving the 45 of SU(5), which would provide additional freedom since the Higgs doublet will now be a linear combination of 5 and $\bar{45}$.

III. CALCULATIONAL TECHNIQUES FOR THE FULL SO(10) ANALYSIS

In this section we discuss the breaking of SO(10) down to SU(3)_C \times U(1)_{em} in a single step by a single pair of 144 + $\bar{144}$. Our analysis will be valid for the supersymmetric SO(10) model as well as for the nonsupersymmetric model. In the latter case one simply identifies $\bar{144}$ with the 144* field. However, for formal reasons we consider a single pair of 160 + $\bar{160}$ where the additional 16 + $\bar{16}$ that reside in 160 + $\bar{160}$ are needed for consistency as we will see below. The analysis involving the 144 + $\bar{144}$ is rather intricate and we use the techniques developed recently in Refs. [4] based on the oscillator method of Refs. [11,12] to compute the desired couplings. There are no cubic couplings of quarks and leptons with the 144 + $\bar{144}$ of Higgs and one needs quartic couplings to grow quark and lepton masses in this scheme. We discuss now the basic ingredients of the model. We begin by displaying the particle content of 144 + $\bar{144}$ in multiplets of SU(5). For $\bar{144}$ -plet one has

$$\bar{144} = \bar{5}(\mathcal{P}_i) + 5(\mathcal{P}^i) + \bar{10}(\mathcal{P}_{ij}) + \bar{15}(\mathcal{P}_{ij}^{(S)}) + 24(\mathcal{P}_j^i) + \bar{40}(\mathcal{P}_{ijk}^l) + 45(\mathcal{P}_k^{ij}), \quad (15)$$

where the subscripts and superscripts i, j, k, \dots are SU(5) indices which take on the values 1, \dots , 5. Similarly, for 144 we find

$$144 = 5(\mathcal{Q}^i) + \bar{5}(\mathcal{Q}_i) + 10(\mathcal{Q}^{ij}) + 15(\mathcal{Q}_{(S)}^{ij}) + 24(\mathcal{Q}_j^i) + 40(\mathcal{Q}_i^{jk}) + \bar{45}(\mathcal{Q}_{jk}^i). \quad (16)$$

To make progress we need a field theoretic description of the 144 and $\bar{144}$ -plet of fields. Possible candidates are the vector-spinors $|\Psi_{(\pm)\mu}\rangle$. However, an unconstrained vector-spinor has $16 \times 10 = 160$ independent components and is reducible. Thus irreducible vector-spinors with dimension-

ality 144 must be gotten by removing 16 of the 160 components of the unconstrained vector-spinor. We define the 144-dimensional constrained vector-spinors $|\Psi_{(\pm)\mu}\rangle$ by imposition of the constraint

$$\Gamma_\mu |\Psi_{(\pm)\mu}\rangle = 0, \quad (17)$$

where Γ_μ satisfy a rank-10 Clifford algebra

$$\{\Gamma_\mu, \Gamma_\nu\} = 2\delta_{\mu\nu} \quad (18)$$

and where μ, ν are the SO(10) indices and take on the values 1, \dots , 10.

We discuss now further the relationship of $|\Psi_{(\pm)\mu}\rangle$ and $|\Psi_{(\pm)\mu}\rangle$. We begin by writing the 160 and $\bar{160}$ component spinor:

$$|\Psi_{(+)\dot{a}\mu}\rangle = |0\rangle \mathbf{P}_{\dot{a}\mu} + \frac{1}{2} b_i^\dagger b_j^\dagger |0\rangle \mathbf{P}_{\dot{a}\mu}^{ij} + \frac{1}{24} \epsilon^{ijklm} b_j^\dagger b_k^\dagger b_l^\dagger b_m^\dagger |0\rangle \mathbf{P}_{\dot{a}\mu}^{ijklm}, \quad (19)$$

$$|\Psi_{(-)\dot{b}\mu}\rangle = b_1^\dagger b_2^\dagger b_3^\dagger b_4^\dagger b_5^\dagger |0\rangle \mathbf{Q}_{\dot{b}\mu} + \frac{1}{12} \epsilon^{ijklm} b_k^\dagger b_l^\dagger b_m^\dagger |0\rangle \mathbf{Q}_{\dot{b}\mu}^{ijklm} + b_i^\dagger |0\rangle \mathbf{Q}_{\dot{b}\mu}^i, \quad (20)$$

where the Latin letters i, j, k, l, m, \dots are SU(5) indices and the Greek letters μ, ν, ρ, \dots represent SO(10) indices. The Latin subscripts $\dot{a}, \dot{b}, \dot{c}, \dot{d} (= 1, 2, 3)$ are reserved for generation indices. The reducible fields appearing in Eqs. (19) and (20) can be identified in SU(5) notation as follows [4]:

$$10 = 5 + \bar{5}: \quad \mathbf{P}_\mu = (\mathbf{P}_{c_k}, \mathbf{P}_{\bar{c}_k}) \equiv (\mathbf{P}^k, \mathbf{P}_k), \quad (21)$$

$$\bar{10} = 5 + \bar{5}: \quad \mathbf{Q}_\mu = (\mathbf{Q}_{c_k}, \mathbf{Q}_{\bar{c}_k}) \equiv (\mathbf{Q}^k, \mathbf{Q}_k),$$

$$100 = \bar{50} + 50: \quad \mathbf{P}_\mu^{ij} = (\mathbf{P}_{c_k}^{ij}, \mathbf{P}_{\bar{c}_k}^{ij}) \equiv (\mathbf{R}^{[ij]k}, \mathbf{R}_k^{[ij]}),$$

$$\bar{100} = \bar{50} + 50: \quad \mathbf{Q}_{\mu ij} = (\mathbf{Q}_{ijc_k}, \mathbf{Q}_{ij\bar{c}_k}) \equiv (\mathbf{S}_{[ij]}^k, \mathbf{S}_{[ij]k}),$$

$$50 = 45 + 5: \quad \mathbf{R}_k^{[ij]} = \mathbf{P}_k^{ij} + \frac{1}{4} (\delta_k^i \hat{\mathbf{P}}^j - \delta_k^j \hat{\mathbf{P}}^i),$$

$$\bar{50} = \bar{40} + \bar{10}: \quad \mathbf{R}^{[ij]k} = \epsilon^{ijlmn} \mathbf{P}_{lmn}^k + \epsilon^{ijklm} \hat{\mathbf{P}}_{lm},$$

$$50 = 40 + 10: \quad \mathbf{S}_{[ij]k} = \epsilon_{ijlmn} \mathbf{Q}_k^{lmn} + \epsilon_{ijklm} \hat{\mathbf{Q}}^{lm},$$

$$\bar{50} = \bar{45} + \bar{5}: \quad \mathbf{S}_{[jk]}^i = \mathbf{Q}_{jk}^i + \frac{1}{4} (\delta_k^i \hat{\mathbf{Q}}_j - \delta_j^i \hat{\mathbf{Q}}_k), \quad (22)$$

where $[ij]$ stands for anticommutation in ij , i.e., $[ij] = -[ji]$,

$$\begin{aligned}
 \overline{50} &= 25 + \overline{25} : \quad \mathbf{P}_{i\mu} = (\mathbf{P}_{ic_k}, \mathbf{P}_{i\overline{c}_k}) \equiv (\mathbf{R}_i^k, \mathbf{R}_{ik}), \\
 50 &= 25 + 25 : \quad \mathbf{Q}_{i\mu}^i = (\mathbf{Q}_{ic_k}^i, \mathbf{Q}_{c_k}^i) \equiv (\mathbf{S}_i^k, \mathbf{S}^{ik}), \\
 25 &= 24 + 1 : \quad \mathbf{R}_j^i = \mathbf{P}_j^i + \frac{1}{5} \delta_j^i \widehat{\mathbf{P}}, \\
 \overline{25} &= \overline{10} + \overline{15} : \quad \mathbf{R}_{ij} = \frac{1}{2} (\mathbf{P}_{ij} + \mathbf{P}_{ij}^{(S)}), \\
 25 &= 24 + 1 : \quad \mathbf{S}_j^i = \mathbf{Q}_j^i + \frac{1}{5} \delta_j^i \widehat{\mathbf{Q}}, \\
 25 &= 10 + 15 : \quad \mathbf{S}^{ij} = \frac{1}{2} (\mathbf{Q}^{ij} + \mathbf{Q}_{(S)}^{ij}).
 \end{aligned} \tag{23}$$

Further, b_i^\dagger and b_i^\dagger ($i = 1, 2, \dots, 5$) are the fermionic creation and annihilation operators and obey the anticommutation rules [11]

$$\{b_i, b_j^\dagger\} = \delta_i^j; \quad \{b_i, b_j\} = 0; \quad \{b_i^\dagger, b_j^\dagger\} = 0 \tag{24}$$

and the SU(5) singlet state $|0\rangle$ satisfies $b_i|0\rangle = 0$. SO(10) invariance requires the following constraints in general

$$\Gamma_\mu |\Psi_{(+)\mu}\rangle = |\Psi'_{(-)}\rangle, \tag{25}$$

$$\Gamma_\mu |\Psi_{(-)\mu}\rangle = |\Psi'_{(+)}\rangle, \tag{26}$$

where

$$\begin{aligned}
 |\Psi'_{(-)}\rangle &= b_1^\dagger b_2^\dagger b_3^\dagger b_4^\dagger b_5^\dagger |0\rangle \widehat{\mathbf{P}} + \frac{1}{12} \epsilon^{ijklm} b_k^\dagger b_l^\dagger b_m^\dagger |0\rangle \\
 &\quad \times (\mathbf{P}_{ij} + 6\widehat{\mathbf{P}}_{ij}) + b_i^\dagger |0\rangle (\mathbf{P}^i + \widehat{\mathbf{P}}^i),
 \end{aligned} \tag{27}$$

$$\begin{aligned}
 |\Psi'_{(+)}\rangle &= |0\rangle \widehat{\mathbf{P}} + \frac{1}{2} b_i^\dagger b_j^\dagger |0\rangle (\mathbf{Q}^{ij} + 6\widehat{\mathbf{Q}}^{ij}) \\
 &\quad + \frac{1}{24} \epsilon^{ijklm} b_j^\dagger b_k^\dagger b_l^\dagger b_m^\dagger |0\rangle (\mathbf{Q}_i + \widehat{\mathbf{Q}}_i).
 \end{aligned} \tag{28}$$

We note in passing that to get the 144 and $\overline{144}$ spinors, $|Y_{(\pm)\mu}\rangle$, we need to impose Eq. (17). This constrain will require setting $|\Psi'_{(\pm)}\rangle = 0$ and thus require the following constraints

$$\begin{aligned}
 \widehat{\mathbf{P}} &= 0, \quad \widehat{\mathbf{P}}^i = -\mathbf{P}^i, \quad \widehat{\mathbf{P}}_{ij} = -\frac{1}{6} \mathbf{P}_{ij}, \\
 \widehat{\mathbf{Q}} &= 0, \quad \widehat{\mathbf{Q}}_i = -\mathbf{Q}_i, \quad \widehat{\mathbf{Q}}^{ij} = -\frac{1}{6} \mathbf{Q}^{ij}.
 \end{aligned} \tag{29}$$

Thus we have

$$|Y_{(\pm)\mu}\rangle = (|\Psi_{(\pm)\mu}\rangle)_{\text{constraint of Eq.(29)}}. \tag{30}$$

However, as we stated already we will be dealing with the full $160 + \overline{160}$ multiplets.

To normalize the SU(5) fields contained in the tensor, $|\Psi_{(\pm)\mu}\rangle$, we carry out a field redefinition

$$\begin{aligned}
 \{1\} : \quad \widehat{\mathbf{P}} &= \sqrt{5} \widehat{\mathcal{P}}, \quad \{\overline{5}\} : \quad \mathbf{P}_i = \mathcal{P}_i, \\
 \{5\} : \quad \mathbf{P}^i &= \mathcal{P}^i, \quad \widehat{\mathbf{P}}^i = 2\widehat{\mathcal{P}}^i, \\
 \{\overline{10}\} : \quad \mathbf{P}_{ij} &= \sqrt{2} \mathcal{P}_{ij}, \quad \widehat{\mathbf{P}}_{ij} = \frac{1}{2\sqrt{3}} \widehat{\mathcal{P}}_{ij}, \\
 \{\overline{15}\} : \quad \mathbf{P}_{ij}^{(S)} &= \sqrt{2} \mathcal{P}_{ij}^{(S)}, \quad \{24\} : \quad \mathbf{P}_j^i = \mathcal{P}_j^i, \\
 \{\overline{40}\} : \quad \mathbf{P}_{ijk}^l &= \frac{1}{6} \mathcal{P}_{ijk}^l, \quad \{45\} : \quad \mathbf{P}_k^{ij} = \mathcal{P}_k^{ij}.
 \end{aligned} \tag{31}$$

$$\begin{aligned}
 \{1\} : \quad \widehat{\mathbf{Q}} &= \sqrt{5} \widehat{\mathcal{Q}}, \quad \{5\} : \quad \mathbf{Q}^i = \mathcal{Q}^i, \\
 \{\overline{5}\} : \quad \mathbf{Q}_i &= \mathcal{Q}_i, \quad \widehat{\mathbf{Q}}_i = 2\widehat{\mathcal{Q}}_i, \\
 \{10\} : \quad \mathbf{Q}^{ij} &= \sqrt{2} \mathcal{Q}_{ij}, \quad \widehat{\mathbf{Q}}^{ij} = \frac{1}{2\sqrt{3}} \widehat{\mathcal{Q}}^{ij}, \\
 \{15\} : \quad \mathbf{Q}_{(S)}^{ij} &= \sqrt{2} \mathcal{Q}_{(S)}^{ij}, \quad \{24\} : \quad \mathbf{Q}_j^i = \mathcal{Q}_j^i, \\
 \{40\} : \quad \mathbf{Q}_i^{ijk} &= \frac{1}{6} \mathcal{Q}_i^{ijk}, \quad \{\overline{45}\} : \quad \mathbf{Q}_j^k = \mathcal{Q}_j^k.
 \end{aligned} \tag{32}$$

In terms of the normalized fields, the kinetic energy of the 160 and $\overline{160}$, i.e., $-\langle \partial_A \Psi_{(\pm)\mu} | \partial^A \Psi_{(\pm)\mu} \rangle$, where A is the Lorentz index, takes the form

$$\begin{aligned}
 \mathcal{L}_{\text{kin}}^{\overline{160}} &= -\partial_A \widehat{\mathcal{P}}^\dagger \partial_A \widehat{\mathcal{P}} - \partial_A \mathcal{P}_i^\dagger \partial^A \mathcal{P}_i - \partial_A \mathcal{P}^{i\dagger} \partial^A \mathcal{P}^i \\
 &\quad - \partial_A \widehat{\mathcal{P}}^{i\dagger} \partial^A \widehat{\mathcal{P}}^i - \frac{1}{2!} \partial_A \mathcal{P}_{ij}^\dagger \partial^A \mathcal{P}_{ij} \\
 &\quad - \frac{1}{2!} \partial_A \widehat{\mathcal{P}}_{ij}^\dagger \partial^A \widehat{\mathcal{P}}_{ij} - \frac{1}{2!} \partial_A \mathcal{P}_{ij}^{(S)\dagger} \partial^A \mathcal{P}_{ij}^{(S)} \\
 &\quad - \partial_A \mathcal{P}_j^{i\dagger} \partial^A \mathcal{P}_j^i - \frac{1}{3!} \partial_A \mathcal{P}_{ijk}^\dagger \partial^A \mathcal{P}_{ijk}^l \\
 &\quad - \frac{1}{2!} \partial_A \mathcal{P}_k^{ij\dagger} \partial^A \mathcal{P}_k^{ij},
 \end{aligned} \tag{33}$$

$$\begin{aligned}
 \mathcal{L}_{\text{kin}}^{160} &= -\partial_A \widehat{\mathcal{Q}}^\dagger \partial_A \widehat{\mathcal{Q}} - \partial_A \mathcal{Q}^{i\dagger} \partial^A \mathcal{Q}^i - \partial_A \mathcal{Q}_i^\dagger \partial^A \mathcal{Q}_i \\
 &\quad - \partial_A \widehat{\mathcal{Q}}_i^\dagger \partial^A \widehat{\mathcal{Q}}_i - \frac{1}{2!} \partial_A \mathcal{Q}^{ij\dagger} \partial^A \mathcal{Q}^{ij} \\
 &\quad - \frac{1}{2!} \partial_A \widehat{\mathcal{Q}}^{ij\dagger} \partial^A \widehat{\mathcal{Q}}^{ij} - \frac{1}{2!} \partial_A \mathcal{Q}_{(S)}^{ij\dagger} \partial^A \mathcal{Q}_{(S)}^{ij} \\
 &\quad - \partial_A \mathcal{Q}_j^{i\dagger} \partial^A \mathcal{Q}_j^i - \frac{1}{3!} \partial_A \mathcal{Q}_i^{ijk\dagger} \partial^A \mathcal{Q}_i^{ijk} \\
 &\quad - \frac{1}{2!} \partial_A \mathcal{Q}_{ij}^{k\dagger} \partial^A \mathcal{Q}_{ij}^k.
 \end{aligned} \tag{34}$$

IV. SYMMETRY BREAKING

In this section we discuss how SO(10) breaks to $SU(3)_C \times SU(2)_L \times U(1)_Y$. For this purpose we consider a superpotential of the form

$$\begin{aligned}
W = & M(\overline{160}_H \times 160_H) + \frac{\lambda_1}{M'}(\overline{160}_H \times 160_H)_1 \\
& \times (\overline{160}_H \times 160_H)_1 + \frac{\lambda_{45}}{M'}(\overline{160}_H \times 160_H)_{45} \\
& \times (\overline{160}_H \times 160_H)_{45} + \frac{\lambda_{210}}{M'}(\overline{160}_H \times 160_H)_{210} \\
& \times (\overline{160}_H \times 160_H)_{210}. \tag{35}
\end{aligned}$$

There are of course many more terms that one can add to Eq. (35) but we consider only the terms displayed in Eq. (35) for simplicity. The relevant terms in the superpotential that accomplish symmetry breaking are

$$W_{SB} = M\mathbf{Q}_\mu^i \mathbf{P}_{i\mu} + \alpha_1 \mathbf{Q}_\mu^i \mathbf{P}_{i\mu} \mathbf{Q}_\nu^j \mathbf{P}_{j\nu} + \alpha_2 \mathbf{Q}_\mu^i \mathbf{P}_{j\mu} \mathbf{Q}_\nu^j \mathbf{P}_{j\nu}, \tag{36}$$

where

$$\begin{aligned}
\alpha_1 &= \frac{1}{M'} \left(-2\lambda_1 - \lambda_{45} + \frac{1}{6}\lambda_{210} \right), \\
\alpha_2 &= -\frac{1}{M'} (4\lambda_{45} + \lambda_{210}). \tag{37}
\end{aligned}$$

Expanding into the irreducible components we find

$$\begin{aligned}
W = & M\mathcal{Q}_j^i \mathcal{P}_i^j + \alpha_1 \mathcal{Q}_j^i \mathcal{P}_i^j \mathcal{Q}_k^l \mathcal{P}_k^l + \alpha_2 \mathcal{Q}_k^i \mathcal{P}_j^k \mathcal{Q}_i^j \mathcal{P}_i^l \\
& + M\widehat{\mathcal{Q}} \widehat{\mathcal{P}} + 2\left(\alpha_1 + \frac{2}{5}\alpha_2\right) \mathcal{Q}_i^k \mathcal{P}_k^l \widehat{\mathcal{Q}} \widehat{\mathcal{P}} \\
& + \frac{2}{\sqrt{5}} \alpha_2 \mathcal{Q}_k^i \mathcal{P}_j^k \mathcal{Q}_i^j \widehat{\mathcal{P}} + \frac{2}{\sqrt{5}} \alpha_1 \mathcal{Q}_k^i \mathcal{P}_j^k \mathcal{P}_i^j \widehat{\mathcal{Q}} \\
& + \frac{1}{5} \alpha_2 \mathcal{Q}_i^k \mathcal{Q}_k^l \widehat{\mathcal{P}} \widehat{\mathcal{P}} + \frac{1}{5} \alpha_1 \mathcal{P}_i^k \mathcal{P}_k^l \widehat{\mathcal{Q}} \widehat{\mathcal{Q}} \\
& + \left(\alpha_1 + \frac{1}{5}\alpha_2\right) \widehat{\mathcal{Q}} \widehat{\mathcal{P}} \widehat{\mathcal{Q}} \widehat{\mathcal{P}}. \tag{38}
\end{aligned}$$

In the minimization we look for solutions of the type

$$\begin{aligned}
\langle \mathcal{Q}_j^i \rangle &= q \text{diag}(2, 2, 2, -3, -3), \\
\langle \mathcal{P}_j^i \rangle &= p \text{diag}(2, 2, 2, -3, -3). \tag{39}
\end{aligned}$$

One finds the following results from the minimization of the potential

$$\begin{aligned}
Mp + 2p^2q(30\alpha_1 + 7\alpha_2) + 2\left(\alpha_1 + \frac{2}{5}\alpha_2\right)pQ_0P_0 \\
+ \frac{1}{75}\alpha_2qP_0^2 + \frac{26}{5\sqrt{5}}\alpha_2(pQ_0 + 2qP_0)p = 0, \tag{40}
\end{aligned}$$

$$\begin{aligned}
Mq + 2q^2p(30\alpha_1 + 7\alpha_2) + 2\left(\alpha_1 + \frac{2}{5}\alpha_2\right)qQ_0P_0 \\
+ \frac{1}{75}\alpha_2pQ_0^2 + \frac{26}{5\sqrt{5}}\alpha_2(qP_0 + 2pQ_0)q = 0, \tag{41}
\end{aligned}$$

$$\begin{aligned}
\left[60\left(\alpha_1 + \frac{2}{5}\alpha_2\right)qp + M\right]P_0 + \frac{2}{5}\alpha_2p^2Q_0 + \frac{156}{\sqrt{5}}\alpha_2p^2q \\
+ 2\left(\alpha_1 + \frac{1}{5}\alpha_2\right)Q_0P_0^2 = 0, \tag{42}
\end{aligned}$$

and

$$\begin{aligned}
\left[60\left(\alpha_1 + \frac{2}{5}\alpha_2\right)qp + M\right]Q_0 + \frac{2}{5}\alpha_2q^2P_0 + \frac{156}{\sqrt{5}}\alpha_2pq^2 \\
+ 2\left(\alpha_1 + \frac{1}{5}\alpha_2\right)Q_0^2P_0 = 0, \tag{43}
\end{aligned}$$

where $Q_0 = \langle \widehat{\mathcal{Q}} \rangle$ and $P_0 = \langle \widehat{\mathcal{P}} \rangle$. The D-flatness condition $\langle 144 \rangle = \langle \overline{144} \rangle$ gives $q = p$. With the above vacuum expectation value, spontaneous breaking occurs so that $\text{SO}(10) \rightarrow \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y$. We note that the VEVs Q_0 and P_0 do not play a role in the above breakdown as this breakdown will occur even when $Q_0 = 0 = P_0$.

V. HIGGS PHENOMENON AND MASS GROWTH

We outline here the Higgs phenomenon and the mass growth associated with the spontaneous breaking given by Eqs. (40)–(43). The fields that participate in the Higgs phenomenon include the 45 vector super multiplet that belongs to the adjoint representation of SO(10) and the $144 + \overline{144}$ chiral superfields. For the analysis of the Higgs phenomenon it is useful to decompose the 45-plet of SO(10) in multiplets of SU(5) so that

$$45 = 1 + 10 + \overline{10} + 24. \tag{44}$$

After spontaneous symmetry breaking the 10_{45} massless vector super multiplet absorbs the 10_{144} chiral multiplet to become a 10_{45} massive vector super multiplet with spins $(1, \frac{1}{2}, 0)$. Similarly, the $\overline{10}_{45}$ vector super multiplet absorbs the $\overline{10}_{\overline{144}}$ chiral multiplet to become the $\overline{10}_{45}$ massive vector super multiplet. Now the 24-plet of SU(5) decomposes under $\text{SU}(3)_C \times \text{SU}(2)_L$ as follows

$$24 = (8, 1) + (\overline{3}, 2) + (3, 2) + (1, 3) + (1, 1). \tag{45}$$

After spontaneous breaking the super vector multiplets with the quantum numbers $(\overline{3}, 2) + (3, 2)$ absorb one linear combination of the chiral multiplets $((\overline{3}, 2) + (3, 2))_{144}$ and $((\overline{3}, 2) + (3, 2))_{\overline{144}}$ becoming a massive $(\overline{3}, 2) + (3, 2)$ vector super multiplets while the orthogonal linear combination of $((\overline{3}, 2) + (3, 2))_{144}$ and $((\overline{3}, 2) + (3, 2))_{\overline{144}}$ which is not absorbed becomes massive. The vector super multiplets corresponding to $(8, 1) + (1, 3) + (1, 1)$ remain massless. The chiral super multiplets corresponding to $(8, 1) + (1, 3)$ become massive. (The (1,1) components of 24_{144} and $24_{\overline{144}}$ require special treatment and we return to it below). Thus we have accounted for the mass growth of the $10 + \overline{10}$ vector super multiplet and the mass growth of the 12 components $(\overline{3}, 2) + (3, 2)$ of the 24-plet vector super multiplet. This leaves us to discuss mass growth of the singlet

vector super multiplet in Eq. (44). This mass growth comes about by absorption of the chiral superfield combination $(\frac{2}{5}\Sigma_a^i - \frac{3}{5}\Sigma_\alpha^i)$ where the repeated indices are summed (a is the color index which takes on values 1, 2, 3 and α is the SU(2) index and takes on values 4, 5), and Σ_j^i is a linear combination of \mathcal{Q}_j^i and \mathcal{P}_j^i . Since Σ is traceless the above equals $\Sigma_a^i = -\Sigma_\alpha^i$. Thus the singlet vector super multiplet absorbs a linear combination of \mathcal{P}_a^i and \mathcal{Q}_a^i becoming a singlet massive vector super multiplet while the orthogonal combination of the chiral superfields \mathcal{P}_a^i and \mathcal{Q}_a^i becomes massive. Thus after the symmetry breaking and Higgs phenomenon only the (1, 8) + (1, 3) + (1, 1) vector super-

multiplet remains massless and the remaining components of 45 of the vector super multiplet become massive. Similarly, all the unabsorbed components of the 144 + $\overline{144}$ become massive. At this stage the gauge group SO(10) has broken down to $SU(3)_C \times SU(2)_L \times U(1)_Y$. To accomplish the breaking of the electroweak symmetry we need a pair of Higgs doublets. Such a possibility arises for $\mathcal{Q}_\alpha, \mathcal{P}^\alpha$.

To exhibit this we need to compute masses for $\mathcal{Q}_i, \mathcal{P}^i$. It is also instructive to compute masses for $\mathcal{Q}^i, \mathcal{P}_i, \widehat{\mathcal{Q}}_i, \widehat{\mathcal{P}}^i, \mathcal{Q}_{jk}^i$, and \mathcal{P}_k^{ij} . The relevant terms in the superpotential are

$$\begin{aligned} W_{\text{mass}} = & \left[M + \frac{1}{M'} \left(-4\lambda_1 + 6\lambda_{45} - \frac{1}{3}\lambda_{210} \right) \langle \mathbf{S}_n^m \mathbf{R}_m^n \rangle \right] (\mathcal{Q}_i \mathcal{P}^i + \mathcal{Q}^i \mathcal{P}_i) - \left[\frac{8}{3} \frac{\lambda_{210}}{M'} \langle \mathbf{S}_m^i \mathbf{R}_j^m \rangle \right] \mathcal{Q}_i \mathcal{P}^j \\ & + \left[\frac{1}{M'} \left(8\lambda_{45} - \frac{2}{3}\lambda_{210} \right) \langle \mathbf{S}_m^i \mathbf{R}_j^m \rangle \right] \mathbf{S}_{[in]}^l \mathbf{R}_l^{[nj]} + \left[-\frac{1}{2}M + \frac{1}{M'} \left(2\lambda_1 + \lambda_{45} - \frac{1}{6}\lambda_{210} \right) \langle \mathbf{S}_n^m \mathbf{R}_m^n \rangle \right] \mathbf{S}_{[ij]}^k \mathbf{R}_k^{[ij]}. \end{aligned} \quad (46)$$

Equation (46) makes it apparent why for technical reasons we need to keep the 160 + $\overline{160}$ multiplet. To see this let us set all the couplings λ to zero in Eq. (46) so that the only terms surviving are proportional to M . Next suppose we impose on $\mathbf{S}_{[ij]}^k$ and $\mathbf{R}_k^{[ij]}$ the constraint of Eq. (29) so that we are strictly considering only the 144 + $\overline{144}$ multiplet. Then we see that the last line of Eq. (46) contributes an additional mass term $-\frac{M}{4}\mathcal{Q}_i \mathcal{P}^i$ while there is no such term for $\mathcal{Q}^i \mathcal{P}_i$. This additional term is clearly not desired and the reason for its appearance is that we are identifying the 5-plet in $\mathbf{R}_k^{[ij]}$ with \mathcal{P}^i and $\overline{5}$ -plet in $\mathbf{S}_{[ij]}^k$ with \mathcal{Q}_i because of Eq. (29) which feeds in the undesired additional term. Thus for book keeping we must not impose the constraint of Eq. (29) in the beginning. Returning to Eq. (46), the corresponding mass terms in the Lagrangian are given by

$$\mathcal{L} = \left| \frac{\partial W_{\text{mass}}}{\partial \mathcal{Q}_i} \right|^2 + \left| \frac{\partial W_{\text{mass}}}{\partial \mathcal{P}^i} \right|^2 + \left| \frac{\partial W_{\text{mass}}}{\partial \mathcal{Q}^i} \right|^2 + \left| \frac{\partial W_{\text{mass}}}{\partial \mathcal{P}_i} \right|^2 + \left| \frac{\partial W_{\text{mass}}}{\partial \mathbf{S}_{[ij]}^k} \right|^2 + \left| \frac{\partial W_{\text{mass}}}{\partial \mathbf{R}_k^{[ij]}} \right|^2. \quad (47)$$

Explicitly we have

$$\begin{aligned} \mathcal{L} = & |\sigma|^2 (\mathcal{P}_i \mathcal{P}_i^\dagger + \mathcal{Q}^i \mathcal{Q}^{i\dagger}) + |\sigma + \omega_1 \beta|^2 (\mathcal{P}^\alpha \mathcal{P}^{\alpha\dagger} + \mathcal{Q}_\alpha \mathcal{Q}_\alpha^\dagger) + |\sigma + \omega_2 \beta|^2 (\mathcal{P}^a \mathcal{P}^{a\dagger} + \mathcal{Q}_a \mathcal{Q}_a^\dagger) \\ & + \frac{1}{2} \left[|\rho - \omega_1 \gamma|^2 + 3 \left| \rho - \frac{1}{2}(\omega_1 + \omega_2) \gamma \right|^2 \right] (\widehat{\mathcal{P}}^\alpha \widehat{\mathcal{P}}^{\alpha\dagger} + \widehat{\mathcal{Q}}_\alpha \widehat{\mathcal{Q}}_\alpha^\dagger) + \left[|\rho - \omega_1 \gamma|^2 + \left| \rho - \frac{1}{2}(\omega_1 + \omega_2) \gamma \right|^2 \right] \\ & \times (\widehat{\mathcal{P}}^a \widehat{\mathcal{P}}^{a\dagger} + \widehat{\mathcal{Q}}_a \widehat{\mathcal{Q}}_a^\dagger) + |\rho - \omega_1 \gamma|^2 (\mathcal{P}_k^{\alpha\beta} \mathcal{P}_k^{\alpha\beta\dagger} + \mathcal{Q}_{\alpha\beta}^k \mathcal{Q}_{\alpha\beta}^{k\dagger}) + |\rho - \omega_2 \gamma|^2 (\mathcal{P}_k^{ab} \mathcal{P}_k^{ab\dagger} + \mathcal{Q}_{ab}^k \mathcal{Q}_{ab}^{k\dagger}) \\ & + 2 \left| \rho - \frac{1}{2}(\omega_1 + \omega_2) \gamma \right|^2 (\mathcal{P}_k^{a\alpha} \mathcal{P}_k^{a\alpha\dagger} + \mathcal{Q}_{\alpha\alpha}^k \mathcal{Q}_{\alpha\alpha}^{k\dagger}) + \left[\left| \rho - \frac{1}{2}(\omega_1 + \omega_2) \gamma \right|^2 - |\rho - \omega_1 \gamma|^2 \right] \\ & \times (\mathcal{P}_\alpha^{aa} \widehat{\mathcal{P}}^{a\dagger} + \mathcal{Q}_{\alpha\alpha}^a \widehat{\mathcal{Q}}_a^\dagger + \text{H.c.}) + \left[\left| \rho - \frac{1}{2}(\omega_1 + \omega_2) \gamma \right|^2 - |\rho - \omega_2 \gamma|^2 \right] (\mathcal{P}_a^{\alpha\alpha} \widehat{\mathcal{P}}^{\alpha\dagger} + \mathcal{Q}_{\alpha\alpha}^a \widehat{\mathcal{Q}}_a^\dagger + \text{H.c.}), \end{aligned} \quad (48)$$

where

$$\begin{aligned}
\sigma &= M + \frac{qP}{M'} \left(-4\lambda_1 + 6\lambda_{45} - \frac{1}{3}\lambda_{210} \right) \left(30 + \frac{P_0 Q_0}{pq} \right), \\
\rho &= -\frac{1}{2}M + \frac{qP}{M'} \left(2\lambda_1 + \lambda_{45} - \frac{1}{6}\lambda_{210} \right) \left(30 + \frac{P_0 Q_0}{pq} \right), \\
\omega_1 &= qp \left[9 - \frac{3}{\sqrt{5}} \left(\frac{P_0}{p} + \frac{Q_0}{q} \right) + \frac{1}{5} \frac{P_0 Q_0}{pq} \right], \\
\omega_2 &= qp \left[4 + \frac{2}{\sqrt{5}} \left(\frac{P_0}{p} + \frac{Q_0}{q} \right) + \frac{1}{5} \frac{P_0 Q_0}{pq} \right], \\
\gamma &= \frac{1}{M'} \left(8\lambda_{45} - \frac{2}{3}\lambda_{210} \right), \\
\beta &= -\frac{8}{3} \frac{\lambda_{210}}{M'}.
\end{aligned} \tag{49}$$

The masses for the fields P^i , Q_i , Q^i , P_i , Q_{ij}^k , and P_k^{ij} can be computed from Eqs. (40)–(43) and (48). It is easily checked that the masses vanish unless one has a nonvanishing λ_{45} or λ_{210} . A scrutiny of the mass growth above shows that the masses of the Higgs doublets Q_α , P^α are split from the Higgs triplets Q_a , P^a . Indeed this allows one to fine-tune the masses of the Higgs doublets to zero by the condition

$$\begin{aligned}
\frac{MM'}{qp} + (-120\lambda_1 + 180\lambda_{45}) \left(1 + \frac{1}{30} \frac{P_0 Q_0}{pq} \right) \\
+ \lambda_{210} \left[-34 + \frac{8}{\sqrt{5}} \left(\frac{P_0}{p} + \frac{Q_0}{q} \right) - \frac{13}{15} \frac{P_0 Q_0}{pq} \right] = 0. \tag{50}
\end{aligned}$$

With this constraint one finds that the Higgs doublets Q_α , P^α are massless while the Higgs triplets Q_a , P^a are massive. Thus the Higgs triplet mass M_{H_3} of the fields Q_a , P^a , under the constraint that the Higgs doublet Q_α , P^α be massless, is given by

$$M_{H_3} = \frac{40}{3} \frac{qP}{M'} \left[1 - \frac{1}{\sqrt{5}} \left(\frac{P_0}{p} + \frac{Q_0}{q} \right) \right] \lambda_{210}. \tag{51}$$

We note that it is not possible to achieve a doublet-triplet splitting for the multiplets Q^i and P_i . Thus the doublet-triplet splitting we are considering is unique. Further, we find that Q^i , P_i develop a mass

$$M_{Q^i, P_i} = 24 \frac{qP}{M'} \left[1 - \frac{1}{3\sqrt{5}} \left(\frac{P_0}{p} + \frac{Q_0}{q} \right) + \frac{1}{45} \frac{P_0 Q_0}{pq} \right] \lambda_{210}. \tag{52}$$

In the above Q_0 and P_0 are crucial in getting a pair of light Higgs doublets. Thus suppose we have $Q_0 = 0 = P_0$ in Eqs. (40)–(43). In this case one finds an additional constraint. Thus from Eqs. (42) and (43) one finds that $Q_0 = 0 = P_0$ imply that $\alpha_2 = 0$. Under this constraint Eq. (50) is not consistent with Eqs. (40) and (41). i.e., one cannot find a pair of light Higgs doublets. We note that this result is a consequence of considering only a limited number of couplings in Eq. (35). Inclusion of a larger set of couplings should allow one to get consistent solutions without inclusion of the singlet VEVs.

To summarize the results thus far we have here a complete breaking of SO(10) to $SU(3)_C \times SU(2)_L \times U(1)_Y$. To get a pair of light Higgs we need to invoke a string landscape scenario which has been extensively discussed recently [7,8]. Effectively in this framework we use a fine-tuning to keep one pair of Higgs doublets light while keeping the Higgs triplets heavy. With the usual radiative electroweak symmetry breaking mechanism, the light doublets of Higgs can develop VEVs breaking the $SU(2)_L \times U(1)_Y$ symmetry down to $U(1)_{em}$. Thus with the above mechanism one can break SO(10) down to $SU(3)_C \times U(1)_{em}$ with just one pair of $144 + \overline{144}$ of Higgs. A similar analysis shows that one gets masses for the remaining parts of the $144 + \overline{144}$ chiral multiplets not absorbed by the vector bosons which become heavy. The heavy spectrum of this model differs significantly from the standard SU(5) and the standard SO(10) models. It consists of several parts: (i) the super heavy lepto-quarks associated with the breaking of the $SO(10) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$, (ii) the super heavy Higgs triplet field, (iii) the components of 24-plet fields P_j^i , Q_j^i which are unabsorbed by the Higgs phenomenon and become super heavy, (iv) the remaining components of $144 + \overline{144}$, aside from a 24-plets of SU(5) each in 144 and $\overline{144}$ discussed in (iii) and excluding the Higgs doublets Q_α , P^α , which become super heavy, and (v) $16 + \overline{16}$ -plet of super heavy fields. Gauge coupling unification will require a careful analysis of contribution of each of the above. We do not address this question further here.

We discuss briefly the issue of split vs nonsplit supersymmetry breaking. Equation (50) is a tree-level relation and its imposition to achieve light Higgs doublets presumes that the loop corrections to the Higgs masses are small. Specifically it requires that the scale of supersymmetry breaking is not high, e.g., the masses of squarks are at the electroweak scale and not at the GUT scale. An alternative possibility is that one may impose Eq. (50) but with loop correction included. In this case we can allow for split supersymmetry scenario where the masses of the squarks are superheavy while gaugino masses may lie at the electroweak scale. Thus the model we are considering can accommodate a split or a nonsplit supersymmetry breaking scenario.

VI. COUPLINGS OF QUARKS AND LEPTONS WITH 144 AND $\overline{144}$ OF HIGGS

The 144 and $\overline{144}$ -plets of Higgs have no SO(10) invariant trilinear coupling with the 16 -plet of matter. However one can write quartic couplings of the type $(1/M_P)(16 \times 16)(144 \times 144)$ and $(1/M_P)(16 \times 16)(144 \times \overline{144})$, where M_P is a super heavy mass. These interactions will generate effective cubic couplings with the light Higgs doublets Q_α , P^α after spontaneous breaking of SO(10) discussed in Secs. III and IV. The size of these couplings is

$O(M_G/M_P)$, where M_G is the grand unification scale. The most general set of quartic couplings involving quarks and leptons are⁵

$$\begin{aligned} (16 \times 16)_{10}(144 \times 144)_{10}, \quad (16 \times 16)_{10}(\overline{144} \times \overline{144})_{10}, \\ (16 \times 16)_{\overline{126}}(144 \times 144)_{126}, \\ (16 \times 16)_{\overline{126}}(\overline{144} \times \overline{144})_{126}. \end{aligned} \quad (53)$$

The couplings $(16 \times 16)_{10}(144 \times 144)_{10}$ and $(16 \times 16)_{10}(\overline{144} \times \overline{144})_{10}$ arise from the following structures

$$\begin{aligned} \mathbf{W}^{(10)} = & \frac{1}{2} \Phi_{\nu\mathcal{U}} \mathcal{M}_{\mathcal{U}\mathcal{U}}^{(10)} \Phi_{\nu\mathcal{U}} \\ & + h_{\dot{a}\dot{b}}^{(10)} \langle \Psi_{(+)\dot{a}\mu}^* | B \Gamma_\nu | Y_{(+)\dot{b}\mu} \rangle k_{\dot{c}\dot{d}}^{(10)} \Phi_{\nu\mathcal{U}} \\ & + f_{\dot{a}\dot{b}}^{(10)} \langle \Psi_{(+)\dot{a}}^* | B \Gamma_\nu | \Psi_{(+)\dot{b}} \rangle l_{\dot{c}\dot{d}}^{(10)} \Phi_{\nu\mathcal{U}} \\ & + \bar{h}_{\dot{a}\dot{b}}^{(10)} \langle Y_{(-)\dot{a}\mu}^* | B \Gamma_\nu | Y_{(-)\dot{b}\mu} \rangle \bar{k}_{\dot{c}\dot{d}}^{(10)} \Phi_{\nu\mathcal{U}}, \end{aligned} \quad (54)$$

where the indices $\mathcal{U}, \mathcal{U}'$ run over several Higgs represen-

$$\begin{aligned} \mathbf{W}^{(16 \times 16)_{10}(\overline{144} \times \overline{144})_{10}} = & -2\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)} \langle \Psi_{(+)\dot{a}}^* | B \Gamma_\rho | \Psi_{(+)\dot{b}} \rangle \langle Y_{(+)\dot{c}\nu}^* | \Gamma_\rho | Y_{(+)\dot{d}\nu} \rangle \\ = & -4\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)} [\langle \Psi_{(+)\dot{a}}^* | B b_i | \Psi_{(+)\dot{b}} \rangle \langle Y_{(+)\dot{c}\nu}^* | B b_i^\dagger | Y_{(+)\dot{d}\nu} \rangle + \langle \Psi_{(+)\dot{a}}^* | B b_i^\dagger | \Psi_{(+)\dot{b}} \rangle \langle Y_{(+)\dot{c}\nu}^* | B b_i | Y_{(+)\dot{d}\nu} \rangle] \\ = & 2\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)(+)} [\epsilon_{ijklmn} \mathbf{M}_{\dot{a}i}^T \mathbf{M}_{\dot{b}j}^{ij} \mathbf{P}_{\dot{c}\mu}^{kT} \mathbf{P}_{\dot{d}\mu}^{mn} - 8\mathbf{M}_{\dot{a}i}^T \mathbf{M}_{\dot{b}j}^{ij} \mathbf{P}_{\dot{c}\mu}^T \mathbf{P}_{\dot{d}\mu} + \epsilon_{ijklmn} \mathbf{M}_{\dot{a}}^{klT} \mathbf{M}_{\dot{b}}^{mn} \mathbf{P}_{\dot{c}\mu}^T \mathbf{P}_{\dot{d}\mu}^{ij} \\ & - 8\mathbf{M}_{\dot{a}j}^T \mathbf{M}_{\dot{b}i} \mathbf{P}_{\dot{c}\mu}^T \mathbf{P}_{\dot{d}\mu}^{ij}] \end{aligned} \quad (57)$$

and

$$\begin{aligned} \mathbf{W}^{(16 \times 16)_{10}(144 \times 144)_{10}} = & -2\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)} \langle \Psi_{(+)\dot{a}}^* | B \Gamma_\rho | \Psi_{(+)\dot{b}} \rangle \langle Y_{(-)\dot{c}\nu}^* | B \Gamma_\rho | Y_{(-)\dot{d}\nu} \rangle \\ = & -4\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)} [\langle \Psi_{(+)\dot{a}}^* | B b_i | \Psi_{(+)\dot{b}} \rangle \langle Y_{(-)\dot{c}\nu}^* | B b_i^\dagger | Y_{(-)\dot{d}\nu} \rangle + \langle \Psi_{(+)\dot{a}}^* | B b_i^\dagger | \Psi_{(+)\dot{b}} \rangle \langle Y_{(-)\dot{c}\nu}^* | B b_i | Y_{(-)\dot{d}\nu} \rangle] \\ = & 2\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)(+)} [8\mathbf{M}_{\dot{a}i}^T \mathbf{M}_{\dot{b}j}^{ij} \mathbf{Q}_{\dot{c}\mu}^{kT} \mathbf{Q}_{\dot{d}\mu}^{lm} - 8\mathbf{M}_{\dot{a}i}^T \mathbf{M}_{\dot{b}j} \mathbf{Q}_{\dot{c}\mu}^{iT} \mathbf{Q}_{\dot{d}\mu} - \mathbf{M}_{\dot{a}}^{ijT} \mathbf{M}_{\dot{b}}^{kl} \mathbf{Q}_{\dot{c}\mu}^T \mathbf{Q}_{\dot{d}\mu}^{lm} + \mathbf{M}_{\dot{a}}^{ijT} \mathbf{M}_{\dot{b}}^{kl} \mathbf{Q}_{\dot{c}\mu}^T \mathbf{Q}_{\dot{d}\mu}^{lm} \\ & - \mathbf{M}_{\dot{a}}^{ijT} \mathbf{M}_{\dot{b}}^{kl} \mathbf{Q}_{\dot{c}\mu}^T \mathbf{Q}_{\dot{d}\mu}^{lm} + \epsilon_{ijklm} \mathbf{M}_{\dot{a}i}^T \mathbf{M}_{\dot{b}j} \mathbf{Q}_{\dot{c}\mu}^T \mathbf{Q}_{\dot{d}\mu}^{lm} + \epsilon_{ijklm} \mathbf{M}_{\dot{a}}^{ijT} \mathbf{M}_{\dot{b}}^{kl} \mathbf{Q}_{\dot{c}\mu}^T \mathbf{Q}_{\dot{d}\mu}^{lm}], \end{aligned} \quad (58)$$

and where

$$\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)} = f_{\dot{a}\dot{b}}^{(10)} h_{\dot{c}\dot{d}}^{(10)} l_{\dot{c}\dot{d}}^{(10)} \widetilde{\mathcal{M}}_{\dot{c}\dot{d}}^{(10)} k_{\dot{c}\dot{d}}^{(10)}, \quad \zeta_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)} = f_{\dot{a}\dot{b}}^{(10)} \bar{h}_{\dot{c}\dot{d}}^{(10)} l_{\dot{c}\dot{d}}^{(10)} \widetilde{\mathcal{M}}_{\dot{c}\dot{d}}^{(10)} \bar{k}_{\dot{c}\dot{d}}^{(10)}, \quad \widetilde{\mathcal{M}}^{(10)} = [\mathcal{M}^{(10)} + (\mathcal{M}^{(10)})^T]^{-1}. \quad (59)$$

We note that $\xi_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)(+)}$, and $\zeta_{\dot{a}\dot{b},\dot{c}\dot{d}}^{(10)(+)}$, are the same as defined by Eq. (59), provided we replace h 's, \bar{h} 's, f 's by $h^{(+)}$'s, $\bar{h}^{(+)}$'s, $f^{(+)}$'s, respectively, where $(+)$ indicates that the couplings are symmetric, i.e., $h_{\dot{a}\dot{b}}^{(10)(\pm)} = \frac{1}{2}(h_{\dot{a}\dot{b}}^{(10)} \pm h_{\dot{b}\dot{a}}^{(10)})$. The above couplings produce quark and lepton masses after GUT symmetry breaking followed by spontaneous breaking of the electroweak symmetry. A preliminary analysis shows that the relation on Yukawa couplings

⁵We have not included the $(16 \times 16)_{120}(144 \times 144)_{120}$ and $(16 \times 16)_{120}(\overline{144} \times \overline{144})_{120}$ couplings here since these couplings are antisymmetric in the generation indices and vanish with only a single $144 + \overline{144}$.

of the same kind, $\mathcal{M}^{(10)}$ represents the mass matrix and $f^{(10)}$, $k^{(10)}$, $\bar{k}^{(10)}$, and $l^{(10)}$ are constants. B is the usual SO(10) charge conjugation operator defined by $B = \prod_{\mu=\text{odd}} \Gamma_\mu$. The semispinors $\Psi_{(\pm)}$ transform as a $16(\overline{16})$ -dimensional irreducible representation of SO(10) and contain $1 + \bar{5} + 10(1 + 5 + \overline{10})$ in its SU(5) decomposition. $|\Psi_{(+)\dot{a}}\rangle$ is given by

$$\begin{aligned} |\Psi_{(+)\dot{a}}\rangle = & |0\rangle \mathbf{M}_{\dot{a}} + \frac{1}{2} b_i^\dagger b_j^\dagger |0\rangle \mathbf{M}_{\dot{a}}^{ij} \\ & + \frac{1}{24} \epsilon^{ijklm} b_j^\dagger b_k^\dagger b_l^\dagger b_m^\dagger |0\rangle \mathbf{M}_{\dot{a}i}. \end{aligned} \quad (55)$$

Elimination of the super heavy fields $\Phi_{\nu\mathcal{U}}$ using the F-flatness condition gives

$$\mathbf{W}_{\text{dim}-5}^{(10)} = \mathbf{W}^{(16 \times 16)_{10}(144 \times 144)_{10}} + \mathbf{W}^{(16 \times 16)_{10}(\overline{144} \times \overline{144})_{10}}, \quad (56)$$

where

such as $h_b = h_t$ does not hold. It would be interesting to study the quark-lepton textures [13–15] in this framework. Further, the preliminary analysis shows that baryon and lepton number violating dimension five operators in this theory are rather different than what one has in the usual SU(5) and SO(10) models [16–20]. Thus, for example, in SU(5) unified models the baryon and lepton number violating dimension five operators arise from the Higgs triplet exchange from pairs of $5 + \bar{5}$. In the present model there are several sources of baryon and lepton number violations, including Higgs triplets from 5 and $\bar{5}$ and from the exchange of $45 + \bar{45}$ present in $144 + \overline{144}$ of Higgs. A full analysis of this issue involves additional couplings where the mediation occurs via 120 , 126 , etc. and is beyond the

scope of this paper. An analysis of this will be given elsewhere [21]. We note here that if in Eq. (53) only couplings involving $\overline{144}$ are kept, the resulting fermion mass matrices will have the same structure as in [3] with a single 10 and one $\overline{126}$ coupling to fermions. Such matrices are fully consistent with experimental data.

VII. CONCLUSION

In conclusion we have investigated a new class of SO(10) models where the breaking of SO(10) down to $SU(3)_C \times SU(2)_L \times U(1)_Y$ can occur in a single step. Further, it is possible to achieve with fine-tuning, a pair of light Higgs doublets which is justifiable within a string based landscaped scenario (see also Ref. [22]). The light Higgs doublets allow one to break the electroweak symmetry $SU(2)_L \times U(1)_Y$ down to $U(1)_{em}$ and thus SO(10) can break to $SU(3)_C \times U(1)_{em}$. The cubic interactions of the light Higgs doublets with quarks and leptons arise from quartic interactions of the type (16.16)(144.144) and (16.16)($\overline{144}$. $\overline{144}$) after spontaneous breaking of SO(10). In this scenario the baryon and lepton number violating dimension five operators receive contributions not just from the conventional Higgs triplet fields but also from

the exchange of $45 + \overline{45}$ components of $144 + \overline{144}$. The above feature distinguishes the above scenario from the conventional models and would lead to different estimates on the proton lifetime and in conventional GUTs. A more detailed analysis of the quark-lepton masses as well as of the proton life time is outside the scope of this paper and will be dealt with elsewhere [21]. Finally we note that above the GUT scale one has a large number of degrees of freedom and the renormalization group evolution is very rapid and thus the theory becomes nonperturbative. We view this theory as descending directly from a theory of quantum gravity where the scale of quantum gravity (e.g., string scale) may lie close to the GUT scale.

ACKNOWLEDGMENTS

The work of K. B. is supported in part by DOE Grant No. DE-FG03-98ER-41076. The work of I. G. is supported in part by NSF Grant No. PHY-0098791. The work of P. N. and R. S. is supported in part by NSF Grant No. PHY-0139967. Part of this work was done while P. N. was visiting the Max Planck Institute, Munich. He acknowledges support from the Alexander von Humboldt Foundation and thanks the MPI for the hospitality extended him.

-
- [1] H. Georgi, in *Particles and Fields*, edited by C. E. Carlson (AIP, New York, 1975); H. Fritzsch and P. Minkowski, *Ann. Phys. (N.Y.)* **93**, 193 (1975).
- [2] R. Slansky, *Phys. Rep.* **79**, 1 (1981).
- [3] K. S. Babu and R. N. Mohapatra, *Phys. Rev. Lett.* **70**, 2845 (1993); B. Bajc, G. Senjanovic, and F. Vissani, *Phys. Rev. Lett.* **90**, 051802 (2003); H. S. Goh, R. N. Mohapatra, and S. Nasri, *Phys. Rev. D* **70**, 075022 (2004); S. Bertolini, M. Frigerio, and M. Malinsky, *Phys. Rev. D* **70**, 095002 (2004); K. S. Babu and C. Macesanu, hep-ph/0505200.
- [4] P. Nath and R. M. Syed, *Phys. Lett. B* **506**, 68 (2001); *Nucl. Phys.* **B618**, 138 (2001); *Nucl. Phys.* **B676**, 64 (2004).
- [5] S. Wiesenfeldt, *Phys. Rev. D* **71**, 075006 (2005).
- [6] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac, and N. Okada, *J. Math. Phys. (N.Y.)* **46**, 033505 (2005); C. S. Aulakh and A. Girdhar, *Nucl. Phys.* **B711**, 275 (2005).
- [7] S. Kachru, R. Kallosh, A. Linde, and S. P. Trivedi, *Phys. Rev. D* **68**, 046005 (2003); F. Denef and M. R. Douglas, *J. High Energy Phys.* 05 (2004) 072; N. Arkani-Hamed and S. Dimopoulos, *J. High Energy Phys.* 06 (2005) 073; M. Dine, E. Gorbatov, and S. Thomas, hep-th/0407043; I. Antoniadis and S. Dimopoulos, *Nucl. Phys.* **B715**, 120 (2005); C. Kokorelis, hep-th/0406258.
- [8] B. Kors and P. Nath, *Nucl. Phys.* **B711**, 112 (2005); K. S. Babu, T. Enkhbat, and B. Mukhopadhyaya, *Nucl. Phys.* **B720**, 47 (2005); K. R. Dienes, E. Dudas, and T. Gherghetta, *Phys. Rev. D* **72**, 026005 (2005).
- [9] L. Michel, *Contribution to Colloq. on Fundamental Interactions, in honor of Antoine Visconti, Marseille, France*, CERN-TH-2716 1979 (unpublished); L. Michel and L. Radicati, *Ann. Phys. (N.Y.)* **66**, 758 (1971).
- [10] C. Bachas, C. Fabre, and T. Yanagida, *Phys. Lett. B* **370**, 49 (1996).
- [11] R. N. Mohapatra and B. Sakita, *Phys. Rev. D* **21**, 1062 (1980).
- [12] F. Wilczek and A. Zee, *Phys. Rev. D* **25**, 553 (1982).
- [13] H. Georgi and C. Jarlskog, *Phys. Lett. B* **86**, 297 (1979).
- [14] P. Nath, *Phys. Rev. Lett.* **76**, 2218 (1996); *Phys. Lett. B* **381**, 147 (1996).
- [15] K. S. Babu and S. M. Barr, *Phys. Lett. B* **381**, 137 (1996).
- [16] S. Weinberg, *Phys. Rev. D* **26**, 287 (1982); N. Sakai and T. Yanagida, *Nucl. Phys.* **B197**, 533 (1982); S. Dimopoulos, S. Raby, and F. Wilczek, *Phys. Lett. B* **112**, 133 (1982); J. Ellis, D. V. Nanopoulos, and S. Rudaz, *Nucl. Phys.* **B202**, 43 (1982).
- [17] P. Nath, A. H. Chamseddine, and R. Arnowitt, *Phys. Rev. D* **32**, 2348 (1985); J. Hisano, H. Murayama, and T. Yanagida, *Nucl. Phys.* **B402**, 46 (1993); T. Goto, T. Nihei, and J. Arafune, *Phys. Rev. D* **52**, 505 (1995); K. S. Babu and S. M. Barr, *Phys. Lett. B* **381**, 137 (1996); P. Nath and R. Arnowitt, *Phys. At. Nucl.* **61**, 975 (1998); T. Goto and T. Nihei, *Phys. Rev. D* **59**, 115009 (1999); K. S. Babu and M. J. Strassler, hep-ph/9808447; B. Bajc, P. Fileviez Perez, and G. Senjanovic, *Phys. Rev. D* **66**, 075005 (2002).

- [18] V. Lucas and S. Raby, Phys. Rev. D **54**, 2261 (1996); **55**, 6986 (1997).
- [19] K. S. Babu, J. C. Pati, and F. Wilczek, Nucl. Phys. **B566**, 33 (2000); Phys. Lett. B **423**, 337 (1998).
- [20] K. S. Babu and S. M. Barr, Phys. Rev. D **48**, 5354 (1993); Z. Chacko and R. N. Mohapatra, Phys. Rev. D **59**, 011702 (1999); Phys. Rev. Lett. **82**, 2836 (1999); Z. Berezhiani, Z. Tavartkiladze, and M. Vysotsky, hep-ph/9809301; I. Gogoladze and A. Kobakhidze, Yad. Fiz. **60N1**, 136 (1997) [Phys. At. Nucl. **60**, 126 (1997)]. G. Altarelli, F. Feruglio, and I. Masina, J. High Energy Phys. **11** (2000) 040; T. Ibrahim and P. Nath, Phys. Rev. D **62**, 095001 (2000); Q. Shafi and Z. Tavartkiladze, Phys. Lett. B **487**, 145 (2000); N. Maekawa, Prog. Theor. Phys. **106**, 401 (2001); K. Turzynski, J. High Energy Phys. **10** (2002) 044.
- [21] K. S. Babu, I. Gogoladze, P. Nath, and R. M. Syed, in progress.
- [22] X. Calmet, Eur. Phys. J. C **41**, 245 (2005).