

Search for second-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

- A. Abulencia,²³ D. Acosta,¹⁷ J. Adelman,¹³ T. Affolder,¹⁰ T. Akimoto,⁵⁴ M. G. Albrow,¹⁶ D. Ambrose,¹⁶ S. Amerio,⁴² D. Amidei,³³ A. Anastassov,⁵¹ K. Anikeev,¹⁶ A. Annovi,⁴⁵ J. Antos,¹ M. Aoki,⁵⁴ G. Apollinari,¹⁶ J.-F. Arguin,³² T. Arisawa,⁵⁶ A. Artikov,¹⁴ W. Ashmanskas,¹⁶ A. Attal,⁸ F. Azfar,⁴¹ P. Azzi-Bacchetta,⁴² P. Azzurri,⁴⁵ N. Bacchetta,⁴² H. Bachacou,²⁸ W. Badgett,¹⁶ A. Barbaro-Galtieri,²⁸ V. E. Barnes,⁴⁷ B. A. Barnett,²⁴ S. Baroiant,⁷ V. Bartsch,³⁰ G. Bauer,³¹ F. Bedeschi,⁴⁵ S. Behari,²⁴ S. Belforte,⁵³ G. Bellettini,⁴⁵ J. Bellinger,⁵⁸ A. Belloni,³¹ E. Ben Haim,⁴³ D. Benjamin,¹⁵ A. Beretvas,¹⁶ J. Beringer,²⁸ T. Berry,²⁹ A. Bhatti,⁴⁹ M. Binkley,¹⁶ D. Bisello,⁴² M. Bishai,¹⁶ R. E. Blair,² C. Blocker,⁶ K. Bloom,³³ B. Blumenfeld,²⁴ A. Bocci,⁴⁹ A. Bodek,⁴⁸ V. Boisvert,⁴⁸ G. Bolla,⁴⁷ A. Bolshov,³¹ D. Bortoletto,⁴⁷ J. Boudreau,⁴⁶ S. Bourov,¹⁶ A. Boveia,¹⁰ B. Brau,¹⁰ C. Bromberg,³⁴ E. Brubaker,¹³ J. Budagov,¹⁴ H. S. Budd,⁴⁸ S. Budd,²³ K. Burkett,¹⁶ G. Busetto,⁴² P. Bussey,²⁰ K. L. Byrum,² S. Cabrera,¹⁵ M. Campanelli,¹⁹ M. Campbell,³³ F. Canelli,⁸ A. Canepa,⁴⁷ D. Carlsmith,⁵⁸ R. Carosi,⁴⁵ S. Carron,¹⁵ M. Casarsa,⁵³ A. Castro,⁵ P. Catastini,⁴⁵ D. Cauz,⁵³ M. Cavalli-Sforza,³ A. Cerri,²⁸ L. Cerrito,⁴¹ S. H. Chang,²⁷ J. Chapman,³³ Y. C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴⁵ G. Chlachidze,¹⁴ F. Chlebana,¹⁶ I. Cho,²⁷ K. Cho,²⁷ D. Chokheli,¹⁴ J. P. Chou,²¹ P. H. Chu,²³ S. H. Chuang,⁵⁸ K. Chung,¹² W. H. Chung,⁵⁸ Y. S. Chung,⁴⁸ M. Ciljak,⁴⁵ C. I. Ciobanu,²³ M. A. Ciocci,⁴⁵ A. Clark,¹⁹ D. Clark,⁶ M. Coca,¹⁵ A. Connolly,²⁸ M. E. Convery,⁴⁹ J. Conway,⁷ B. Cooper,³⁰ K. Copic,³³ M. Cordelli,¹⁸ G. Cortiana,⁴² A. Cruz,¹⁷ J. Cuevas,¹¹ R. Culbertson,¹⁶ D. Cyr,⁵⁸ S. DaRonco,⁴² S. D'Auria,²⁰ M. D'onofrio,¹⁹ D. Dagenhart,⁶ P. de Barbaro,⁴⁸ S. De Cecco,⁵⁰ A. Deisher,²⁸ G. De Lentdecker,⁴⁸ M. Dell'Orso,⁴⁵ S. Demers,⁴⁸ L. Demortier,⁴⁹ J. Deng,¹⁵ M. Deninno,⁵ D. De Pedis,⁵⁰ P. F. Derwent,¹⁶ C. Dionisi,⁵⁰ J. R. Dittmann,⁴ P. DiTuro,⁵¹ C. Dörr,²⁵ A. Dominguez,²⁸ S. Donati,⁴⁵ M. Donega,¹⁹ P. Dong,⁸ J. Donini,⁴² T. Dorigo,⁴² S. Dube,⁵¹ K. Ebina,⁵⁶ J. Efron,³⁸ J. Ehlers,¹⁹ R. Erbacher,⁷ D. Errede,²³ S. Errede,²³ R. Eusebi,⁴⁸ H. C. Fang,²⁸ S. Farrington,²⁹ I. Fedorko,⁴⁵ W. T. Fedorko,¹³ R. G. Feild,⁵⁹ M. Feindt,²⁵ J. P. Fernandez,⁴⁷ R. Field,¹⁷ G. Flanagan,³⁴ L. R. Flores-Castillo,⁴⁶ A. Foland,²¹ S. Forrester,⁷ G. W. Foster,¹⁶ M. Franklin,²¹ J. C. Freeman,²⁸ Y. Fujii,²⁶ I. Furic,¹³ A. Gajjar,²⁹ M. Gallinaro,⁴⁹ J. Galyardt,¹² J. E. Garcia,⁴⁵ M. Garcia Sciveres,²⁸ A. F. Garfinkel,⁴⁷ C. Gay,⁵⁹ H. Gerberich,²³ E. Gerchtein,¹² D. Gerdes,³³ S. Giagu,⁵⁰ G. P. di Giovanni,⁴³ P. Giannetti,⁴⁵ A. Gibson,²⁸ K. Gibson,¹² C. Ginsburg,¹⁶ N. Giokaris,¹⁴ K. Giolo,⁴⁷ M. Giordani,⁵³ M. Giunta,⁴⁵ G. Giurgiu,¹² V. Glagolev,¹⁴ D. Glenzinski,¹⁶ M. Gold,³⁶ N. Goldschmidt,³³ J. Goldstein,⁴¹ G. Gomez,¹¹ G. Gomez-Ceballos,¹¹ M. Goncharov,⁵² O. González,⁴⁷ I. Gorelov,³⁶ A. T. Goshaw,¹⁵ Y. Gotra,⁴⁶ K. Goulianatos,⁴⁹ A. Gresele,⁴² M. Griffiths,²⁹ S. Grinstein,²¹ C. Grossi-Pilcher,¹³ U. Grundler,²³ J. Guimaraes da Costa,²¹ C. Haber,²⁸ S. R. Hahn,¹⁶ K. Hahn,⁴⁴ E. Halkiadakis,⁴⁸ A. Hamilton,³² B.-Y. Han,⁴⁸ R. Handler,⁵⁸ F. Happacher,¹⁸ K. Hara,⁵⁴ M. Hare,⁵⁵ S. Harper,⁴¹ R. F. Harr,⁵⁷ R. M. Harris,¹⁶ K. Hatakeyama,⁴⁹ J. Hauser,⁸ C. Hays,¹⁵ H. Hayward,²⁹ A. Heijboer,⁴⁴ B. Heinemann,²⁹ J. Heinrich,⁴⁴ M. Hennecke,²⁵ M. Herndon,⁵⁸ J. Heuser,²⁵ D. Hidas,¹⁵ C. S. Hill,¹⁰ D. Hirschbuehl,²⁵ A. Hocker,¹⁶ A. Holloway,²¹ S. Hou,¹ M. Houlden,²⁹ S.-C. Hsu,⁹ B. T. Huffman,⁴¹ R. E. Hughes,³⁸ J. Huston,³⁴ K. Ikado,⁵⁶ J. Incandela,¹⁰ G. Introzzi,⁴⁵ M. Iori,⁵⁰ Y. Ishizawa,⁵⁴ A. Ivanov,⁷ B. Iyutin,³¹ E. James,¹⁶ D. Jang,⁵¹ B. Jayatilaka,³³ D. Jeans,⁵⁰ H. Jensen,¹⁶ E. J. Jeon,²⁷ M. Jones,⁴⁷ K. K. Joo,²⁷ S. Y. Jun,¹² T. R. Junk,²³ T. Kamon,⁵² J. Kang,³³ M. Karagoz-Unel,³⁷ P. E. Karchin,⁵⁷ Y. Kato,⁴⁰ Y. Kemp,²⁵ R. Kephart,¹⁶ U. Kerzel,²⁵ V. Khotilovich,⁵² B. Kilminster,³⁸ D. H. Kim,²⁷ H. S. Kim,²⁷ J. E. Kim,²⁷ M. J. Kim,¹² M. S. Kim,²⁷ S. B. Kim,²⁷ S. H. Kim,⁵⁴ Y. K. Kim,¹³ M. Kirby,¹⁵ L. Kirsch,⁶ S. Klimenko,¹⁷ M. Klute,³¹ B. Knuteson,³¹ B. R. Ko,¹⁵ H. Kobayashi,⁵⁴ K. Kondo,⁵⁶ D. J. Kong,²⁷ J. Konigsberg,¹⁷ K. Kordas,¹⁸ A. Korytov,¹⁷ A. V. Kotwal,¹⁵ A. Kovalev,⁴⁴ J. Kraus,²³ I. Kravchenko,³¹ M. Kreps,²⁵ A. Kreymer,¹⁶ J. Kroll,⁴⁴ N. Krumnack,⁴ M. Kruse,¹⁵ V. Krutelyov,⁵² S. E. Kuhlmann,² Y. Kusakabe,⁵⁶ S. Kwang,¹³ A. T. Laasanen,⁴⁷ S. Lai,³² S. Lami,⁴⁵ S. Lammel,¹⁶ M. Lancaster,³⁰ R. L. Lander,⁷ K. Lannon,³⁸ A. Lath,⁵¹ G. Latino,⁴⁵ I. Lazzizzera,⁴² C. Lecci,²⁵ T. LeCompte,² J. Lee,⁴⁸ J. Lee,²⁷ S. W. Lee,⁵² R. Lefèvre,³ N. Leonardo,³¹ S. Leone,⁴⁵ S. Levy,¹³ J. D. Lewis,¹⁶ K. Li,⁵⁹ C. Lin,⁵⁹ C. S. Lin,¹⁶ M. Lindgren,¹⁶ E. Lipeles,⁹ T. M. Liss,²³ A. Lister,¹⁹ D. O. Litvinsev,¹⁶ T. Liu,¹⁶ Y. Liu,¹⁹ N. S. Lockyer,⁴⁴ A. Loginov,³⁵ M. Loretz,⁴² P. Loverre,⁵⁰ R.-S. Lu,¹ D. Lucchesi,⁴² P. Lujan,²⁸ P. Lukens,¹⁶ G. Lungu,¹⁷ L. Lyons,⁴¹ J. Lys,²⁸ R. Lysak,¹ E. Lytken,⁴⁷ P. Mack,²⁵ D. MacQueen,³² R. Madrak,¹⁶ K. Maeshima,¹⁶ P. Maksimovic,²⁴ G. Manca,²⁹ F. Margaroli,⁵ R. Marginean,¹⁶ C. Marino,²³ A. Martin,⁵⁹ M. Martin,²⁴ V. Martin,³⁷ M. Martínez,³ T. Maruyama,⁵⁴ H. Matsunaga,⁵⁴ M. E. Mattson,⁵⁷ R. Mazini,³² P. Mazzanti,⁵ K. S. McFarland,⁴⁸ D. McGivern,³⁰ P. McIntyre,⁵² P. McNamara,⁵¹ R. McNulty,²⁹ A. Mehta,²⁹ S. Menzemer,³¹ A. Menzione,⁴⁵ P. Merkel,⁴⁷ C. Mesropian,⁴⁹ A. Messina,⁵⁰ M. von der Mey,⁸ T. Miao,¹⁶ N. Miladinovic,⁶ J. Miles,³¹ R. Miller,³⁴ J. S. Miller,³³ C. Mills,¹⁰ M. Milnik,²⁵ R. Miquel,²⁸ S. Miscetti,¹⁸ G. Mitselmakher,¹⁷ A. Miyamoto,²⁶ N. Moggi,⁵ B. Mohr,⁸ R. Moore,¹⁶ M. Morello,⁴⁵ P. Movilla Fernandez,²⁸ J. Mühlstädt,²⁸ A. Mukherjee,¹⁶ M. Mulhearn,³¹ Th. Muller,²⁵ R. Mumford,²⁴ P. Murat,¹⁶ J. Nachtman,¹⁶ S. Nahm,⁵⁹ I. Nakano,³⁹ A. Napier,⁵⁵

A. ABULENCIA *et al.*

PHYSICAL REVIEW D 73, 051102 (2006)

- D. Naumov,³⁶ V. Necula,¹⁷ C. Neu,⁴⁴ M. S. Neubauer,⁹ J. Nielsen,²⁸ T. Nigmanov,⁴⁶ L. Nodulman,² O. Norniella,³
 T. Ogawa,⁵⁶ S. H. Oh,¹⁵ Y. D. Oh,²⁷ T. Okusawa,⁴⁰ R. Oldeman,²⁹ R. Orava,²² K. Osterberg,²² C. Pagliarone,⁴⁵
 E. Palencia,¹¹ R. Paoletti,⁴⁵ V. Papadimitriou,¹⁶ A. Papikonomou,²⁵ A. A. Paramonov,¹³ B. Parks,³⁸ S. Pashapour,³²
 J. Patrick,¹⁶ G. Pauletta,⁵³ M. Paulini,¹² C. Paus,³¹ D. E. Pellett,⁷ A. Penzo,⁵³ T. J. Phillips,¹⁵ G. Piacentino,⁴⁵ J. Piedra,⁴³
 K. Pitts,²³ C. Plager,⁸ L. Pondrom,⁵⁸ G. Pope,⁴⁶ X. Portell,³ O. Poukhov,¹⁴ N. Pounder,⁴¹ F. Prakoshyn,¹⁴ A. Pronko,¹⁶
 J. Proudfoot,² F. Ptohos,¹⁸ G. Punzi,⁴⁵ J. Pursley,²⁴ J. Rademacker,⁴¹ A. Rahaman,⁴⁶ A. Rakitin,³¹ S. Rappoccio,²¹
 F. Ratnikov,⁵¹ B. Reisert,¹⁶ V. Rekovic,³⁶ N. van Remortel,²² P. Renton,⁴¹ M. Rescigno,⁵⁰ S. Richter,²⁵ F. Rimondi,⁵
 K. Rinnert,²⁵ L. Ristori,⁴⁵ W. J. Robertson,¹⁵ A. Robson,²⁰ T. Rodrigo,¹¹ E. Rogers,²³ S. Rolli,⁵⁵ R. Roser,¹⁶ M. Rossi,⁵³
 R. Rossin,¹⁷ C. Rott,⁴⁷ A. Ruiz,¹¹ J. Russ,¹² V. Rusu,¹³ D. Ryan,⁵⁵ H. Saarikko,²² S. Sabik,³² A. Safonov,⁷
 W. K. Sakamoto,⁴⁸ G. Salamanna,⁵⁰ O. Salto,³ D. Saltzberg,⁸ C. Sanchez,³ L. Santi,⁵³ S. Sarkar,⁵⁰ K. Sato,⁵⁴ P. Savard,³²
 A. Savoy-Navarro,⁴³ T. Scheidle,²⁵ P. Schlabach,¹⁶ E. E. Schmidt,¹⁶ M. P. Schmidt,⁵⁹ M. Schmitt,³⁷ T. Schwarz,³³
 L. Scodellaro,¹¹ A. L. Scott,¹⁰ A. Scribano,⁴⁵ F. Scuri,⁴⁵ A. Sedov,⁴⁷ S. Seidel,³⁶ Y. Seiya,⁴⁰ A. Semenov,¹⁴ F. Semeria,⁵
 L. Sexton-Kennedy,¹⁶ I. Sfiligoi,¹⁸ M. D. Shapiro,²⁸ T. Shears,²⁹ P. F. Shepard,⁴⁶ D. Sherman,²¹ M. Shimojima,⁵⁴
 M. Shochet,¹³ Y. Shon,⁵⁸ I. Shreyber,³⁵ A. Sidoti,⁴³ A. Sill,¹⁶ P. Sinervo,³² A. Sisakyan,¹⁴ J. Sjolin,⁴¹ A. Skiba,²⁵
 A. J. Slaughter,¹⁶ K. Sliwa,⁵⁵ D. Smirnov,³⁶ J. R. Smith,⁷ F. D. Snider,¹⁶ R. Snihur,³² M. Soderberg,³³ A. Soha,⁷
 S. Somalwar,⁵¹ V. Sorin,³⁴ J. Spalding,¹⁶ F. Spinella,⁴⁵ P. Squillacioti,⁴⁵ M. Stanitzki,⁵⁹ A. Staveris-Polykalas,⁴⁵
 R. St. Denis,²⁰ B. Stelzer,⁸ O. Stelzer-Chilton,³² D. Stentz,³⁷ J. Strologas,³⁶ D. Stuart,¹⁰ J. S. Suh,²⁷ A. Sukhanov,¹⁷
 K. Sumorok,³¹ H. Sun,⁵⁵ T. Suzuki,⁵⁴ A. Taffard,²³ R. Tafirout,³² R. Takashima,³⁹ Y. Takeuchi,⁵⁴ K. Takikawa,⁵⁴
 M. Tanaka,² R. Tanaka,³⁹ M. Tecchio,³³ P. K. Teng,¹ K. Terashi,⁴⁹ S. Tether,³¹ J. Thom,¹⁶ A. S. Thompson,²⁰
 E. Thomson,⁴⁴ P. Tipton,⁴⁸ V. Tiwari,¹² S. Tkaczyk,¹⁶ D. Toback,⁵² S. Tokar,¹⁴ K. Tollefson,³⁴ T. Tomura,⁵⁴ D. Tonelli,⁴⁵
 M. Tönnesmann,³⁴ S. Torre,⁴⁵ D. Torretta,¹⁶ S. Tourneur,⁴³ W. Trischuk,³² R. Tsuchiya,⁵⁶ S. Tsuno,³⁹ N. Turini,⁴⁵
 F. Ukegawa,⁵⁴ T. Unverhau,²⁰ S. Uozumi,⁵⁴ D. Usynin,⁴⁴ L. Vacavant,²⁸ A. Vaiciulis,⁴⁸ S. Vallecorsa,¹⁹ A. Varganov,³³
 E. Vataga,³⁶ G. Velev,¹⁶ G. Veramendi,²³ V. Veszpremi,⁴⁷ T. Vickey,²³ R. Vidal,¹⁶ I. Vila,¹¹ R. Vilar,¹¹ I. Vollrath,³²
 I. Volobouev,²⁸ F. Würthwein,⁹ P. Wagner,⁵² R. G. Wagner,² R. L. Wagner,¹⁶ W. Wagner,²⁵ R. Wallny,⁸ T. Walter,²⁵
 Z. Wan,⁵¹ M. J. Wang,¹ S. M. Wang,¹⁷ A. Warburton,³² B. Ward,²⁰ S. Waschke,²⁰ D. Waters,³⁰ T. Watts,⁵¹ M. Weber,²⁸
 W. C. Wester III,¹⁶ B. Whitehouse,⁵⁵ D. Whiteson,⁴⁴ A. B. Wicklund,² E. Wicklund,¹⁶ H. H. Williams,⁴⁴ P. Wilson,¹⁶
 B. L. Winer,³⁸ P. Wittich,⁴⁴ S. Wolbers,¹⁶ C. Wolfe,¹³ S. Worm,⁵¹ T. Wright,³³ X. Wu,¹⁹ S. M. Wynne,²⁹ A. Yagil,¹⁶
 K. Yamamoto,⁴⁰ J. Yamaoka,⁵¹ Y. Yamashita,³⁹ C. Yang,⁵⁹ U. K. Yang,¹³ W. M. Yao,²⁸ G. P. Yeh,¹⁶ J. Yoh,¹⁶ K. Yorita,¹³
 T. Yoshida,⁴⁰ I. Yu,²⁷ S. S. Yu,⁴⁴ J. C. Yun,¹⁶ L. Zanello,⁵⁰ A. Zanetti,⁵³ I. Zaw,²¹ F. Zetti,⁴⁵ X. Zhang,²³
 J. Zhou,⁵¹ and S. Zucchelli⁵

(CDF Collaboration)

¹Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China²Argonne National Laboratory, Argonne, Illinois 60439, USA³Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain⁴Baylor University, Waco, Texas 76798, USA⁵Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy⁶Brandeis University, Waltham, Massachusetts 02254, USA⁷University of California, Davis, Davis, California 95616, USA⁸University of California, Los Angeles, Los Angeles, California 90024, USA⁹University of California, San Diego, La Jolla, California 92093, USA¹⁰University of California, Santa Barbara, Santa Barbara, California 93106, USA¹¹Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain¹²Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA¹³Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA¹⁴Joint Institute for Nuclear Research, RU-141980 Dubna, Russia¹⁵Duke University, Durham, North Carolina 27708¹⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA¹⁷University of Florida, Gainesville, Florida 32611, USA¹⁸Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy¹⁹University of Geneva, CH-1211 Geneva 4, Switzerland²⁰Glasgow University, Glasgow G12 8QQ, United Kingdom²¹Harvard University, Cambridge, Massachusetts 02138, USA

SEARCH FOR SECOND-GENERATION SCALAR ...

PHYSICAL REVIEW D 73, 051102 (2006)

- ²²*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*
- ²³*University of Illinois, Urbana, Illinois 61801, USA*
- ²⁴*The Johns Hopkins University, Baltimore, Maryland 21218, USA*
- ²⁵*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- ²⁶*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan*
- ²⁷*Center for High Energy Physics, Kyungpook National University, Taegu 702-701; Seoul National University, Seoul 151-742; and SungKyunKwan University, Suwon 440-746, Korea*
- ²⁸*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- ²⁹*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁰*University College London, London WC1E 6BT, United Kingdom*
- ³¹*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
- ³²*Institute of Particle Physics, McGill University, Montréal, Canada H3A 2T8 and University of Toronto, Toronto, Canada M5S 1A7*
- ³³*University of Michigan, Ann Arbor, Michigan 48109, USA*
- ³⁴*Michigan State University, East Lansing, Michigan 48824, USA*
- ³⁵*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- ³⁶*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- ³⁷*Northwestern University, Evanston, Illinois 60208, USA*
- ³⁸*The Ohio State University, Columbus, Ohio 43210, USA*
- ³⁹*Okayama University, Okayama 700-8530, Japan*
- ⁴⁰*Osaka City University, Osaka 588, Japan*
- ⁴¹*University of Oxford, Oxford OX1 3RH, United Kingdom*
- ⁴²*University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- ⁴³*LPNHE-Université de Paris 6/IN2P3-CNRS*
- ⁴⁴*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁴⁵*Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy*
- ⁴⁶*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- ⁴⁷*Purdue University, West Lafayette, Indiana 47907, USA*
- ⁴⁸*University of Rochester, Rochester, New York 14627, USA*
- ⁴⁹*The Rockefeller University, New York, New York 10021, USA*
- ⁵⁰*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, University of Rome "La Sapienza," I-00185 Roma, Italy*
- ⁵¹*Rutgers University, Piscataway, New Jersey 08855, USA*
- ⁵²*Texas A&M University, College Station, Texas 77843, USA*
- ⁵³*Istituto Nazionale di Fisica Nucleare, University of Trieste, Udine, Italy*
- ⁵⁴*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁵⁵*Tufts University, Medford, Massachusetts 02155, USA*
- ⁵⁶*Waseda University, Tokyo 169, Japan*
- ⁵⁷*Wayne State University, Detroit, Michigan 48201, USA*
- ⁵⁸*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁵⁹*Yale University, New Haven, Connecticut 06520, USA*
- (Received 19 December 2005; published 1 March 2006)

Results on a search for pair production of second-generation scalar leptoquark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV are reported. The data analyzed were collected by the CDF detector during the 2002–2003 Tevatron Run II and correspond to an integrated luminosity of 198 pb^{-1} . Leptoquarks (LQ) are sought through their decay into (charged) leptons and quarks, with final state signatures represented by two muons and jets and one muon, large transverse missing energy and jets. We observe no evidence for LQ production and derive 95% C.L. upper limits on the LQ production cross sections as well as lower limits on their mass as a function of β , where β is the branching fraction for $\text{LQ} \rightarrow \mu q$.

DOI: 10.1103/PhysRevD.73.051102

PACS numbers: 14.80.-j, 13.85.Rm

The symmetry between leptons and quarks present in the standard model (SM) of particle physics, has lead to the theoretical speculation that a more fundamental force could operate at energy scales larger than the electroweak symmetry breaking scale, allowing for quark to lepton

transitions, mediated by new gauge bosons. Theories like grand unification or R-parity violating supersymmetric models introduce the idea of quark to lepton transitions [1]. Whenever quarks and leptons are allowed to couple directly to each other, new bosons carrying both lepton and

A. ABULENCIA *et al.*

baryon quantum numbers can also exist. They are called leptoquark (LQ) [2], they can have spin 0 (scalar LQ) or 1 (vector LQ) are color-triplet particles, and can either be produced singly or in pairs. Most of the other characteristics, such as weak isospin, electric charge and the coupling λ to lepton and quark are model dependent [3]. Their masses are not predicted. To accommodate experimental constraints on flavor changing neutral currents, LQ are assumed to couple to fermions of the same generation [3,4]. While first generation LQ have been extensively searched for at $e - p$, e^+e^- and $p\bar{p}$ colliders, via single and pair production, and strong limits on their production cross section and mass have been set [5,6], second and third generation LQ are detectable via pair production at hadronic colliders and can be singly produced at e^+e^- machines. Indeed upper limits on second-generation LQ production cross section and lower limits on their masses exist from LEP[7] and the Tevatron Run I[8,9], but they are usually weaker than the ones for first generation LQ. In this paper we present new results on a search for second-generation LQ pair production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. Assuming two different values for the LQ branching fraction β to muon and quark, we consider the following final state signatures: both LQ decaying into muons of opposite charge and quarks ($\beta = 1$), and one of the LQ decaying into a muon and a quark and the other into a neutrino and a quark ($\beta = 0.5$). Since we do not observe evidence for LQ production, we set an upper limit on the production cross section times branching fraction (Br) where $Br = \beta^2$ or $Br = 2\beta(1 - \beta)$. These results are then combined with the one from a search for scalar LQ pairs decaying into $\nu\nu qq$, resulting in jets and missing transverse energy topology[10]. In this way we can express our limits as a continuous function of the parameter β .

CDF is a general-purpose detector built to study the physics of $p\bar{p}$ collisions at the Tevatron accelerator at Fermilab and it is described in detail elsewhere [11]. We use a cylindrical coordinate system around the beampipe in which θ is the polar angle, ϕ is the azimuthal angle and $\eta = -\ln(\tan(\frac{\theta}{2}))$. The transverse energy is defined as $E_T = E \sin\theta$ and the transverse momentum is defined as $P_T = P \sin\theta$, where E is the energy measured by the calorimeter and P the momentum measured by the tracking system. The vector \vec{E}_T is equal to $-\sum E_T^i \vec{n}_i$ where \vec{n}_i is a unit vector that points from the interaction vertex to the i th calorimeter tower in the transverse plane. Its magnitude, \cancel{E}_T , is called transverse missing energy. \cancel{E}_T is corrected following the correction of jet energies, and if muons are identified in the event, \cancel{E}_T is corrected for the muon momenta.

The data used in the analysis were collected during the 2002–2003 Tevatron Run II. The integrated luminosity for this data sample is $198 \pm 12 \text{ pb}^{-1}$. Events are selected online by requiring track segments (“stubs”) reconstructed in the muon chambers and matched to individual tracks

PHYSICAL REVIEW D 73, 051102 (2006)

reconstructed in the central tracker (high P_T muon trigger). The efficiency of the trigger combinations used in the $\mu\mu jj$ and $\mu\nu jj$ analyses, measured using $Z \rightarrow \mu^+\mu^-$ data [12,13], is $\sim 90\%$, varying from about 87% to 95% depending on the type of muon chamber used to detect the candidate muon. Muons are selected as “tight” or “loose” (this second category being used in the $\mu\mu$ analysis only). A tight muon requires a reconstructed track segment in the muon chambers with positions well matched to the extrapolation of a single track, while a loose muon is the one selected by requiring only one isolated track. In both cases the energy deposition in the calorimeters must be consistent with that of a minimum-ionizing particle. We apply a cut on the χ^2 of the track fit to eliminate kaons and pions which have decayed in flight. The identification efficiency for muons has also been measured using data [12,13] and is approximately 90%, going from 89% to 95% for different types of detector and selection used to identify the candidate muon. The coordinate of the lepton (also assumed to be the event coordinate) along the beamline must fall within 60 cm of the center of the detector ($|z_{\text{vertex}}| < 60$ cm) to ensure a good energy measurement in the calorimeter. This cut has an efficiency of $(95 \pm 0.1(\text{stat}) \pm 0.5(\text{sys}))\%$, and is determined from studies of minimum bias events. The efficiencies of the identification cuts, the trigger selection and the vertex cut, measured using data are taken into account by using scale factors between data and Monte Carlo events. Jets are reconstructed using a cone of fixed radius $R = \sqrt{(\Delta\eta^2 + \Delta\phi^2)} = 0.7$ and for these analyses are required to be in the $|\eta| < 2.0$ range. Jets are calibrated as a function of η and E_T and their energy is corrected to the parton level [14]. Neutrinos produce missing transverse energy, \cancel{E}_T , which is measured by balancing the calorimeter energy in the transverse plane. The muon sample is heavily contaminated by events produced by cosmic rays interactions with the detector. Since these events do not originate from a common interaction vertex, the timing capability of the central outer tracker (COT) is used to reject events with two muon tracks, one of which travels toward the beam pipe. We also require that the muon track passes close to the beam line, within distances less than 0.02 cm (0.2 cm) for tracks with (without) silicon hits. In the analyses we are describing, the signal selection criteria are set according to the kinematic distribution (e.g. p_T of the muons and E_T of the jets) of decay products determined from Monte Carlo studies, optimized to eliminate background with a minimal loss of signal events [15].

In the dimuon + jets topology, from the inclusive muon triggers dataset we select events with two reconstructed isolated muons with $P_T > 25 \text{ GeV}/c$. The first muon is required to be tight, i.e. to have a stub associated to a track, while the second one can be without a stub (“stubbless”). Events are further selected if there are at least two jets with $E_T > 30$ and 15 GeV, respectively. In the search in the muon, neutrino and two jets topology, we select events

SEARCH FOR SECOND-GENERATION SCALAR ...

with one reconstructed tight muon with $P_T > 25 \text{ GeV}/c$. We veto events with a second loose or tight muon to be orthogonal to the previous selection. We then accept events where there is large missing transverse energy, $\cancel{E}_T > 60 \text{ GeV}$ and at least two jets with $E_T > 30 \text{ GeV}$.

The above datasets are composed predominantly of events coming from QCD production of Z/W bosons in association with jets and $t\bar{t}$ production (where one or both the W 's from top decay into muon and neutrino). To reduce these backgrounds we apply several cuts which depend on the final state topology.

(1) $\mu\mu$ analysis:

- (i) veto of events whose reconstructed dilepton mass falls in the window $76 < m_{\mu\mu} < 110 \text{ GeV}/c^2$ to remove the $Z +$ jets contribution and $m_{\mu\mu} < 15 \text{ GeV}/c^2$ to avoid contamination from J/Ψ and Υ production;
- (ii) $E_T(j_1) + E_T(j_2) > 85 \text{ GeV}$ and $E_T(e_1) + E_T(e_2) > 85 \text{ GeV}$;
- (iii) $\sqrt{(E_T(j_1) + E_T(j_2))^2 + (E_T(e_1) + E_T(e_2))^2} > 200 \text{ GeV}$. The effect of the last two cuts is shown in Fig. 1, where SM background is compared to a LQ signal for $M_{LQ} = 220 \text{ GeV}/c^2$.

(2) $\mu\nu$ analysis:

- (i) $\Delta\phi(\cancel{E}_T - \text{jet}) > 5^\circ$ to veto events where the transverse missing energy is mismeasured due to a mismeasure of the jet energy, and $\Delta\phi(\cancel{E}_T - \mu) < 175^\circ$ to ensure that the missing energy does not come from mismeasurement of the muon momentum;
- (ii) $E_T(j_1) + E_T(j_2) > 80 \text{ GeV}$;
- (iii) $M_T(\mu\nu) > 120 \text{ GeV}/c^2$ to reduce the $W + 2$ jets background;
- (iv) a mass-dependent cut consisting in selecting events falling in mass windows defined




FIG. 1. Graphical representation of the last two topological cuts applied in the $\mu\mu jj$ analysis as observed on MC events. The discrimination between SM background and LQ signal is evident.

around several LQ masses. We require that the reconstructed mass combinations of the jets, muon and \cancel{E}_T be consistent with those reconstructed from the LQ Monte Carlo. The ambiguity of the jet assignments allows for two different sets of reconstructed masses of the LQ pair in each event: $M(\mu - \text{jet}1)$, $M_T(\cancel{E}_T - \text{jet}2)$ and $M(\mu - \text{jet}2)$, $M_T(\cancel{E}_T - \text{jet}1)$ (when using \cancel{E}_T we obtain only a transverse mass, M_T). We build lineshapes of the mass distributions by matching the reconstructed objects to the generator level objects, to obtain a mean value of the reconstructed mass and its width. We then select events for which the following conditions apply: $|M(\mu, j_1) - M_{LQ}| < 2\sigma_1$ or $|M(\mu, j_2) - M_{LQ}| < 2\sigma_2$, where M_{LQ} is the mean of the reconstructed LQ distribution and $\sigma_{1,2}$ are the width parametrizations.

For the transverse mass distributions, we have chosen a mass-dependent lower cut denoted by $T_{1,2}^{\min}$: $T_1^{\min} = 20 + (M_{LQ} - 120) \text{ GeV}/c^2$, and $T_2^{\min} = 20 + (M_{LQ} - 120)/2 \text{ GeV}/c^2$ so that our cut is defined as: $M_T(\cancel{E}_T, j_1) > T_1^{\min}$ or $M_T(\cancel{E}_T, j_2) > T_2^{\min}$. In Fig. 2 we plot the mass distributions of the selected events (before the mass limit cut) compared to the signal distribution for $M_{LQ} = 180 \text{ GeV}/c^2$.




FIG. 2. Final mass distributions (see text) of the surviving events before the mass limit cut compared to the signal distribution for $M_{LQ} = 180 \text{ GeV}/c^2$.

TABLE I. Number of events surviving all cuts in the muon, missing energy and jets topology, compared with background expectations, as a function of the LQ mass (in GeV/c^2). Errors on the expectations are both statistical and systematic.

Mass	140	160	180	200	220
Wjj	0.9 ± 0.1	1.4 ± 0.1	1.4 ± 0.1	1.6 ± 0.1	1.6 ± 0.1
top	1.7 ± 0.2	1.8 ± 0.2	1.4 ± 0.2	1.0 ± 0.2	0.8 ± 0.2
Zjj	0.20 ± 0.01				
mujets	0.3 ± 0.3				
Total	3.1 ± 0.3	3.7 ± 0.4	3.2 ± 0.3	3.1 ± 0.3	2.9 ± 0.3
Data	3	4	2	0	0

We study the properties of the physics backgrounds by generating events corresponding to $Z/W + 2$ jets with ALPGEN [16] + HERWIG [17] (to perform parton showering) and $t\bar{t}$ with PYTHIA [18]. A complete simulation of the CDF II detector based on GEANT[19] and full event reconstruction is then performed. To normalize the number of simulated events to data we use the theoretical cross sections for $t\bar{t}$ from [20] and for $\gamma/Z \rightarrow \mu\mu + 2$ jets from [21]. The background arising from multijet events, where a jet is mismeasured as a muon or where the muon comes from pion decay (QCD/fake), is evaluated using data. In the $\mu\mu jj$ analysis we examine the data for same-sign events (events with two muons of the same charge) remaining after each kinematical cut. We estimate the background contribution to be twice the number of same-sign events, in the assumption that there is no evidence of LQ signal in these type of events (the LQ pair have opposite charge, giving rise to two opposite charge muons). In the $\mu\nu jj$ analysis the contribution from the QCD/fakes background is estimated by examining the phase space of the \cancel{E}_T vs. the muon fractional isolation for data events in which the muon isolation requirement is not enforced. Here the muon fractional isolation is defined as the ratio between the calorimetric energy not associated with the lepton in a cone of $\Delta R = 0.4$ around the lepton and the energy of the lepton. The following assumptions are made: since jets are produced in association with other particles, the isolation fraction of a jet will generally be larger than the one corresponding to a muon; there is no correlation between

the isolation of the muon and \cancel{E}_T , and in the region where the \cancel{E}_T is small and the isolation of the muon is large the LQ contribution is expected to be negligible (background-dominated region). With these assumptions, from the ratio of the number of events in the background-dominated regions we can extrapolate the contribution in the signal region. Other backgrounds from $b\bar{b}$, $Z \rightarrow \tau\bar{\tau}$, WW are negligible due to the muon isolation and large muon and jet transverse energy requirements. In the $\mu\mu$ channel the expected number of $Z + 2$ jets events is 1.7 ± 0.1 . The expected number of $t\bar{t}$ events is 0.22 ± 0.03 events. We estimate 1 ± 1 fake events. The overall background estimate is: 3 ± 1 events. In the $\mu\nu$ channel, the number of events in each mass region, compared with the background expectations is reported in Table I.

We check the prediction of our background sources with data in control regions where the background contribution is maximized. For the $\mu\mu$ analysis the region is defined by requiring two muons with $P_T > 25 \text{ GeV}/c$, $75 < m_{\mu\mu} < 105 \text{ GeV}/c^2$ and 2 jets with $E_T > 30, 15 \text{ GeV}$. We observe 110 events and expect 88 ± 10 . For the $\mu\nu$ analysis we ask for one muon with $P_T > 25 \text{ GeV}/c$, $\cancel{E}_T > 35 \text{ GeV}$ and 2 jets with $E_T > 30 \text{ GeV}$ and observe 203 events to be compared with a prediction of 221 ± 15 from SM sources.

The efficiency to detect our signal is obtained from MC simulated LQ (PYTHIA) events to account for kinematical and geometrical acceptance. The total efficiencies for a LQ signal are reported in Table II.

TABLE II. Efficiencies after all cuts, errors (statistical and systematic) and 95% C.L. upper limits on the production cross section \times branching fraction Br, as a function of M_{LQ} , for the two channels.

$M_{\text{LQ}}(\text{GeV}/c^2)$	$\mu\mu jj$		$\mu\nu jj$	
	ϵ	$\sigma \times \text{Br}(\text{pb})$	ϵ	$\sigma \times \text{Br}(\text{pb})$
100	0.020 ± 0.003	1.35	0.0050 ± 0.0005	-
120	0.05 ± 0.005	0.52	0.070 ± 0.005	0.86
160	0.13 ± 0.01	0.18	0.070 ± 0.005	0.73
200	0.19 ± 0.02	0.13	0.110 ± 0.005	0.41
220	0.21 ± 0.02	0.11	0.13 ± 0.01	0.24
240	0.24 ± 0.02	0.10	0.13 ± 0.01	0.24
260	0.26 ± 0.02	0.09	0.14 ± 0.01	0.21

SEARCH FOR SECOND-GENERATION SCALAR ...

PHYSICAL REVIEW D 73, 051102 (2006)

TABLE III. 95% C.L. lower limits on the second-generation scalar LQ mass (in GeV/c^2), as a function of β . The limit from CDF [9] ($\mu\mu jj$) Run I ($\sim 120 \text{ pb}^{-1}$) is also given.

β	$\mu\mu jj$	$\mu\nu jj$	$\nu\nu jj$	Combined	CDF Run I
0.01	114	125	...
0.05	110	133	...
0.1	...	137	...	143	...
0.2	...	155	...	157	...
0.3	100	162	...	176	...
0.4	152	168	...	200	...
0.5	171	170	...	208	...
0.6	184	168	...	213	...
0.7	196	162	...	217	...
0.8	206	155	...	221	...
0.9	215	137	...	224	...
1.0	224	226	202

The following systematic uncertainties are considered when calculating signal acceptance and background predictions: (i) luminosity: 6% (ii) choice of parton distribution functions: 2.1% (iii) statistical error of MC < 1.2% (iv) jet energy calibration scale < 1% (v) muon reconstruction : 0.8% (vi) z_{vertex} cut : 0.5%. (vii) initial and final state radiation 1.8%. After all selection cuts, 2 events remain in the $\mu\mu$ channel, while the number of events remaining in the $\mu\nu$ channel is reported in Table I.

In the analyses described above the number of events passing the selection cuts is consistent with the expected number of background events. The conclusion of the two searches is that there is no LQ signal: hence we derive an upper limit on the LQ production cross section at 95% confidence level. We use a Bayesian approach [22] with a flat probability distribution for the signal cross section and Gaussian distributions for acceptance and background uncertainties. The cross section limits are tabulated in Table II and the mass limits are tabulated in Table III. To compare our experimental results with the theoretical expectation, we use the next-to-leading order (NLO) cross section for scalar LQ pair production from [23] with CTEQ6 parton distribution functions [24].

The theoretical uncertainties correspond to the variations from $M_{\text{LQ}}/2$ to $2M_{\text{LQ}}$ of the renormalization scale μ used in the NLO QCD calculation. To set a limit on the LQ mass we compare our 95% CL upper experimental limit to the theoretical cross section for $\mu = 2M_{\text{LQ}}$, which is conservative as it corresponds to the lower value of the theoretical cross section. We find lower limits on $M(\text{LQ})$ at $224 \text{ GeV}/c^2$ ($\beta = 1$) and $170 \text{ GeV}/c^2$ ($\beta = 0.5$). They are reported in Fig. 3. To obtain the best limit however, we combine the results from the two decay channels just described with the result of a search for LQ in the case where the LQ pair decays to a neutrino and quark with branching ratio $\text{Br}(\text{LQ} \rightarrow \nu q) = 1.0$ [10]. The individual channel analyses are in fact optimized for fixed values of β (1,0.5,0) while in the combined analysis, due to the contributions of the different decay channels, the signal acceptance can be naturally expressed as a function of β . As for the treatment of uncertainties, the searches in the $\mu\mu jj$ and $\mu\nu jj$ channels use common criteria and sometime apply the same kind of requirements so the uncertainties in the acceptances are considered correlated. When calculating the limit combination including the $\nu\nu jj$ channel the uncertainties are considered uncorrelated. For each β value a 95% C.L. upper limit on the expected number of events is returned for each mass, and by comparing this to the theoretical expectation, lower limits on the LQ mass are set. The combined limit as a function of β is shown in Fig. 4, together with the individual channel limits. The combined mass limits are also tabulated in Table III.

The final result presented here is better than the results obtained with Tevatron Run I data[8]. This is mostly due to the small increase in the cross section as a function of the center of mass energy (from 1.8 to 1.96 TeV), and an increase in the muon acceptance. A comment is in order when comparing this result with the ones recently published [6] for first generation LQ and third generation LQ [25]. While the signatures of LQ production is very similar (high P_T leptons, large transverse missing energy and energetic jets) one has to consider the constraint on the LQ particle to only couple to same generation fermions. This implies different types of selection and exclude the




FIG. 3. 95% C.L. limit on the experimental cross section times branching ratio as a function of the LQ mass for the $\mu\nu jj$ and $\mu\mu jj$ channel. The NLO theoretical cross section is plotted for different values of the renormalization scale. Mass limits of $170 \text{ GeV}/c^2$ and $224 \text{ GeV}/c^2$ respectively are obtained.




FIG. 4 (color online). Leptoquark mass exclusion regions at 95% C.L. as function of $\text{Br}(\text{LQ} \rightarrow \mu\bar{q})$.

possibility of combining intergeneration results. Also, in the case of electrons and muons, as can be seen from the current results, the similarity in their acceptance results in similar cross section and mass limits, while the result is quite different in the case of third generation LQ, due to the much smaller acceptance for τ leptons. The results presented in this paper, as well as the ones recently published in [6], are obtained with a statistical sample corresponding to about twice the luminosity collected in Tevatron Run I. A future increase in the Run II luminosity by an order of magnitude is estimated to extend the LQ mass range to

$\sim 300 \text{ GeV}/c^2$ for the $\beta = 1$ case [26]. Substantially higher LQ masses will be explored at the future Large Hadron Collider (LHC)[27].

In conclusion, we have performed a search for pair production of second-generation scalar LQ in the dimuons + jets and muon, missing energy + jets topologies, using 198 pb^{-1} of proton-antiproton collision data recorded by the CDF experiment during Run II of the Tevatron. We combined these findings with the ones from a search in the $\cancel{E}_T + \text{jets}$ topology[10]. No evidence for LQ is observed. Assuming that the LQ decays to muon and quark with variable branching ratio β we exclude LQ with masses below $226 \text{ GeV}/c^2$ for $\beta = 1$, $208 \text{ GeV}/c^2$ for $\beta = 0.5$ and $143 \text{ GeV}/c^2$ for $\beta = 0.1$ at 95% C.L.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium fuer Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, UK; the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; and in part by the European Community's Human Potential Programme under contract HPRN-CT-20002, Probe for New Physics.

-
- [1] D. Acosta and S. K. Blessing, Annu. Rev. Nucl. Part. Sci. **49**, 389 (1999) and references therein.
 - [2] W. Buchmuller, R. Ruckl, and D. Wyler, Phys. Lett. B **191**, 442 (1987)**448**, 320(E) (1999).
 - [3] M. Leurer, Phys. Rev. D **49**, 333 (1994).
 - [4] S. Davidson, D. Bailey, and B. A. Campbell, Z. Phys. C **61**, 613 (1994).
 - [5] C. Adloff *et al.* (H1 Collaboration), Eur. Phys. J. C **11**, 447 (1999); **14**, 553(E) (2000); C. Adloff *et al.* (H1 Collaboration), Phys. Lett. B **523**, 234 (2001); J. Breitweg *et al.* (ZEUS Collaboration), Eur. Phys. J. C **16**, 253 (2000); J. Breitweg *et al.* (ZEUS Collaboration), Phys. Rev. D **63**, 052002 (2001); V. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **64**, 092004 (2001).
 - [6] V. Abazov *et al.*, (D0 Collaboration), Phys. Rev. D **71**, 071104 (2005); D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **72**, 051107 (2005).
 - [7] P. Abreu *et al.* (DELPHI Collaboration), Phys. Lett. B **446**, 62 (1999); M. Acciari *et al.* (L3 Collaboration), Phys. Lett. B **489**, 81 (2000); R. Barate *et al.* (ALEPH Collaboration), Eur. Phys. J. C **12**, 183 (2000); G. Abbiendi *et al.* (OPAL Collaboration), Phys. Lett. B **526**, 233 (2002).
 - [8] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **81**, 4806 (1998).
 - [9] S. Abachi *et al.* (D0 Collaboration), Phys. Rev. Lett. **84**, 2088 (2000); B. Abbot *et al.* (D0 Collaboration), Phys. Rev. Lett. **83**, 2896 (1999); S. Abachi *et al.* (D0 Collaboration), Phys. Rev. Lett. **75**, 3618 (1995).
 - [10] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 112001(E) (2005); **71**, 119901 (2005).
 - [11] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
 - [12] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **94**, 091803 (2005).
 - [13] M. Karagoz-Unel, AIP Conf. Proc. No. 753 (AIP, New York, 2005), p. 400.

SEARCH FOR SECOND-GENERATION SCALAR ...

- [14] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **74**, 2626 (1995).
- [15] D. Ryan, Ph.D. thesis, Tufts University, 2004.
- [16] M. L. Mangano *et al.*, J. High Energy Phys. 07 (2003) 001.
- [17] G. Corcella *et al.* J. High Energy Phys. 01 (2001) 10.
- [18] T. Sjostrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [19] R. Brun and F. Carminati, CERN Program Library Long Writeup W5013 , 1993.
- [20] N. Kidonakis and R. Vogt, Phys. Rev. **D68**, 114014 (2003); M. Cacciari *et al.*, J. High Energy Phys. 404 (2004) 68.
- [21] J. Campbell and R. K. Ellis, Phys. Rev. **D65**, 113007 (2002).
- [22] J. Conway, CERN 2000-005, 247, 2000 (unpublished).
- [23] M. Kramer *et al.*, Phys. Rev. Lett.**79**, 341 (1997).
- [24] J. Pumplin *et al.*, J. High Energy Phys. 07 (2002) 012.
- [25] A. Safonov (CDF Collaboration), Proceedings of the 8th International Workshop on Tau Lepton Physics (Tau04), Nara, Japan, 2004 (unpublished).
- [26] S. Rolli, Proceedings of the TeV4LHC Workshop, FNAL, 2004 (unpublished).
- [27] M. Kramer *et al.*, Phys. Rev. **D71**, 057503 (2005).

PHYSICAL REVIEW D **73**, 051102 (2006)