

Statistics of $SU(5)$ D -brane models on a type II orientifold

Florian Gmeiner^{1,2,*} and Maren Stein^{2,†}

¹Max-Planck-Institut für Physik, Föhringer Ring 6, 80805 München, Germany

²Arnold-Sommerfeld-Center for Theoretical Physics, Department für Physik, Ludwig-Maximilians-Universität München, Theresienstraße 37, 80333 München, Germany

(Received 8 March 2006; published 26 June 2006)

We perform a statistical analysis of models with $SU(5)$ and flipped $SU(5)$ gauge group in a type II orientifold setup. We investigate the distribution and correlation of properties of these models, including the number of generations and the hidden sector gauge group. Compared to the recent analysis [F. Gmeiner, R. Blumenhagen, G. Honecker, D. Lüst, and T. Weigand, *J. High Energy Phys.* 01 (2006) 004; F. Gmeiner, *Fortschr. Phys.* 54, 391 (2006).] of models with a standard model-like gauge group, we find very similar results.

DOI: [10.1103/PhysRevD.73.126008](https://doi.org/10.1103/PhysRevD.73.126008)

PACS numbers: 11.25.Mj, 11.25.Wx, 12.10.Dm

I. INTRODUCTION

Grand unified theories provide an interesting framework for unification of the strong and electroweak forces. The minimal simple Lie group that could be used to achieve this is $SU(5)$ [1] or, as a modification of this, flipped $SU(5) \times U(1)_X$ [2]. The latter is more interesting from a phenomenological point of view, because models based on this gauge group might survive the experimental limits on proton decay.

In this paper we continue with the analysis of [3,4], where the gauge sector statistics of a type II orientifold has been considered. Our approach is inspired by the statistical treatment of the string vacuum problem. For a recent review on the intersecting brane models we will be dealing with the reader might want to consult [5]. We report on a systematic computer analysis of intersecting brane models in a $T^6/\mathbb{Z}_2 \times \mathbb{Z}_2$ orientifold background [6] which extends the results published in [4] in the direction of grand unified theories. We focus on the frequency distributions of $SU(5)$ as well as flipped $SU(5)$ gauge groups. Explicit constructions of models of this type have already been performed in [7,8], nonsupersymmetric models have been constructed in [9].

II. SETUP AND METHODS

We work in the intersecting brane picture of type IIA, compactified on a toroidal orientifold of $T^6/\mathbb{Z}_2 \times \mathbb{Z}_2$. We use the setup and the notation of [4] and refer the reader to this paper for more details. In particular we are treating only factorizable branes, that can be expressed by their wrapping numbers on the three two-tori of T^6 .

The $D6$ -branes wrapping special Lagrangian three-cycles are parametrized by integer-valued coefficients X^I , Y^I , $I \in \{0, \dots, 3\}$. There are two different possibilities for the geometry of the three T^2 s, expressed in the three

variables $\beta_i \in \{1, 2\}$, $i \in \{1, 2, 3\}$, where a value of 2 stands for a tilted torus. Furthermore, we define a rescaling factor $c := \prod_{i=1}^3 \beta_i$.

There are three basic constraints to get consistent string vacua in our setup:

- (1) The supersymmetry conditions, written in our variables as

$$\sum_{I=0}^3 \frac{Y^I}{U_I} = 0, \quad \sum_{I=0}^3 X^I U_I > 0. \quad (1)$$

They assure that the D -branes wrap special Lagrangian cycles and exclude the appearance of antibranes. The U_I parametrize a rescaled version of the complex structure moduli, defined as $U_I = (U_0, U_i)$ with $U_0 := R_1^{(1)} R_1^{(2)} R_1^{(3)}$ and $U_i := R_1^{(i)} R_2^{(j)} R_2^{(k)}$, where $i, j, k \in \{1, 2, 3\}$ cyclic and $R_{1/2}^{(i)}$ are the radii of the two-torus i .

- (2) The tadpole cancellation condition for k stacks of N_a branes, given by

$$\sum_{a=1}^k N_a \vec{X}_a = \vec{L}, \quad (2)$$

where the L^I parametrize the orientifold charge. Concretely we have $\vec{L} = (8c, \{8\beta_i\})^T$.

- (3) An additional constraint from K -theory [10]:

$$\sum_{a=1}^k N_a Y_a^0 \in 2\mathbb{Z}, \quad \frac{\beta_i}{c} \sum_{a=1}^k N_a Y_a^i \in 2\mathbb{Z}. \quad (3)$$

Chiral matter in a bifundamental representation arises at the intersection of two stacks of branes with a multiplicity given by the intersection number

$$I_{ab} = \sum_{I=0}^3 (X_a^I Y_b^I - X_b^I Y_a^I). \quad (4)$$

Furthermore, we get symmetric and antisymmetric representations

$$\#\text{Sym}_a = \frac{1}{2}(I_{aa'} - I_{a06}), \quad \#\text{Anti}_a = \frac{1}{2}(I_{aa'} + I_{a06}). \quad (5)$$

*Electronic address: flo@mppmu.mpg.de

†Electronic address: mstein@theorie.physik.uni-muenchen.de

In the original $SU(5)$ construction, the standard model particles are embedded in a $\bar{\mathbf{5}}$ and a $\mathbf{10}$ representation of the unified gauge group as follows

$$\begin{aligned} SU(5) &\rightarrow SU(3) \times SU(2) \times U(1)_Y, \\ \bar{\mathbf{5}} &\rightarrow (\bar{\mathbf{3}}, \mathbf{1})_{2/3} + (\mathbf{1}, \mathbf{2})_{-1}, \\ \mathbf{10} &\rightarrow (\bar{\mathbf{3}}, \mathbf{1})_{-4/3} + (\mathbf{3}, \mathbf{2})_{1/3} + (\mathbf{1}, \mathbf{1})_2, \end{aligned} \quad (6)$$

where the hypercharge is generated by the $SU(3) \times SU(2)$ -invariant generator

$$Z = \text{diag}(-1/3, -1/3, -1/3, 1/2, 1/2). \quad (7)$$

In the flipped $SU(5)$ construction, the embedding is given by

$$\begin{aligned} SU(5) \times U(1)_X &\rightarrow SU(3) \times SU(2) \times U(1)_Y, \\ \bar{\mathbf{5}}_{-3} &\rightarrow (\bar{\mathbf{3}}, \mathbf{1})_{-4/3} + (\mathbf{1}, \mathbf{2})_{-1}, \\ \mathbf{10}_1 &\rightarrow (\bar{\mathbf{3}}, \mathbf{1})_{2/3} + (\mathbf{3}, \mathbf{2})_{1/3} + (\mathbf{1}, \mathbf{1})_0, \\ \mathbf{1}_5 &\rightarrow (\mathbf{1}, \mathbf{1})_2, \end{aligned} \quad (8)$$

including a right-handed neutrino. The hypercharge is in this case given by the combination $Y = -\frac{2}{5}Z + \frac{2}{5}X$.

We would like to realize models of both type within our orientifold setup. The $SU(5)$ case is simpler, since in principle it requires only two branes, a $U(5)$ brane a and a $U(1)$ brane b , which intersect such that we get the $\bar{\mathbf{5}}$ representation at the intersection. The $\mathbf{10}$ will be realized as the antisymmetric representation of the $U(5)$ brane. To get reasonable models, we have to require that the number of antisymmetric representations is equal to the number of $\bar{\mathbf{5}}$ representations,

$$I_{ab} = -\#\text{Anti}_a. \quad (9)$$

In a pure $SU(5)$ model one should also restrict to configurations with $\#\text{Sym}_a = 0$ to exclude $\mathbf{15}$ representations from the beginning. It has been proven in [7] that in this case no three generation models can be constructed. Besides, symmetric representations might also be interesting from a phenomenological point of view, thus we will include them in our discussion.

The flipped $SU(5)$ case is a bit more involved since in addition to the constraints of the $SU(5)$ case one has to make sure that the $U(1)_X$ stays massless and the $\bar{\mathbf{5}}$ and $\mathbf{10}$ will have the right charges. To achieve this, at least one additional brane c is needed. Generically, the $U(1)_X$ can be constructed as a combination of all $U(1)$ s present in the model

$$U(1)_X = \sum_{i=1}^k x_i U(1)_i. \quad (10)$$

The condition that the hypercharge should be massless can be formulated as

$$\sum_{a=1}^k x_a N_a \vec{Y}_a = 0, \quad (11)$$

with some unknown coefficients x_a .

This condition boils down to a system of linear equations which can be solved by a standard algorithm. In the case of models without symmetric representations of $SU(5)$, one can be almost sure to find a solution, given four or more hidden-sector brane stacks. In contrary for models including symmetric representations the probability for a massless $U(1)$ lies slightly below 50% almost regardless of the number of brane stacks in the hidden sector.

III. RESULTS

Having specified the additional constraints, we use the techniques developed in [4] to generate as many solutions to the tadpole, supersymmetry and K -theory conditions as possible. The requirement of a specific set of branes to generate the $SU(5)$ or flipped $SU(5)$ simplifies the computation and gives us the possibility to explore a larger part of the moduli space as compared to the analysis in [4].

Before conducting an analysis of the gauge sector properties of the models under consideration, we would like to check if the number of solutions decreases exponentially for large values¹ of the U_I . This has been observed in [4] for the general solutions. In Fig. 1 the number of solutions with and without symmetric representations are shown. The scaling holds in our present case as well, although the result is a bit obscured by the much smaller statistics. In total we found 6198 solutions without restrictions on the number of generations and the presence of symmetric representations. Excluding these representations reduces the number of solutions to 914. Looking at the flipped $SU(5)$ models, we found 3816 without the restriction to have a massless $U(1)_X$ and 1970 including this constraint. Imposing the condition to get no symmetric representations reduces the number of solutions further to 394.

The correct number of generations turns out to be the strongest constraint on the statistics in our previous work on standard model constructions. The $SU(5)$ case is not different in this aspect. In Fig. 2 we show the number of solutions for different numbers of generations. We did not find any solutions with three $\bar{\mathbf{5}}$ and $\mathbf{10}$ representations. This situation is very similar to the one we encountered in our previous analysis of models with a standard model gauge group [4]. An analysis of the models which have been explicitly constructed showed that they exist only for very large values of the complex structure parameters. The same is true in the present case. Because the number of models decreases rapidly for higher values of the parameters, we can draw the conclusion that these models are statistically heavily suppressed.

Comparing the standard and the flipped $SU(5)$ construction the result for models with one generation might be

¹“Large values” in our rescaled version of the complex structure parameters means a large difference between the radii of at least one of the three two-tori.




FIG. 1 (color online). Logarithmic plot of the number of solutions with an $SU(5)$ factor depending on the absolute value of the parameters U . The blue bars (left) show the result including models with symmetric representations of $SU(5)$. The red bars (right) represent only solutions without these representations.





FIG. 2 (color online). Plots of the number of solutions for different numbers of generations for $SU(5)$ (upper plot): models with (red, left bars) and without (red, right bars) symmetric representations of $SU(5)$; and flipped $SU(5)$ (lower plot): all models (red bars, left), models permitting a massless $U(1)$ (blue bars, middle) and massless solutions without symmetric representations (green bars, right).

surprising, since there are more one generation models in the flipped than in the standard case. This is due to the fact that there are generically different possibilities to realize the additional $U(1)_X$ factor for one geometrical setup, which we counted as distinct models.

As in the unflipped case the massless models without symmetric representations have a clear maximum at eight generations whereas for massless solutions including symmetric representations one or two generations prevail. The aforementioned different probability for finding a massless $U(1)$ in the case of models with and without symmetric representations can also be seen from Fig. 2.

Regarding the hidden sector, we found in total only four $SU(5)$ models which did not have a hidden sector at all—one with 4, two with 8 and one with 16 generations. In the flipped $SU(5)$ case such models do not exist at all.

The frequency distribution of properties of the hidden sector gauge group, the probability of finding a gauge group of specific rank M and the distribution of the total rank, are shown in Figs. 3 and 4. The distribution for individual gauge factors is qualitatively very similar to





FIG. 3 (color online). Logarithmic plots of the number of solutions with a specific rank M gauge factor in the hidden sector. The upper plot shows $SU(5)$, blue (left) and red (right) bars represent solutions with and without symmetric reps. of $SU(5)$; the lower plot shows flipped $SU(5)$ models, red bars (left) stand for the total number of solutions, blue (middle) and green (right) bars represent all massless solutions and those without symmetric reps. of $SU(5)$ respectively.




FIG. 4 (color online). Plots of the number of solutions for given values of the total rank of the hidden sector gauge group. The upper plot shows $SU(5)$, blue (left) and red (right) bars represent solutions with and without symmetric reps. of $SU(5)$; the lower one flipped $SU(5)$ models, as before all models (red bars, left), those satisfying the massless condition (blue bars, middle) and the massless ones without symmetric representations (green bars, right).

the one obtained for all possible solutions in previous work (see Figs. 7, 4 resp. of [4]). This is expected to be the case, since we found in an earlier analysis that the hidden sector statistics should be generic and, from a qualitative point of view, independent of the constraints on the visible sector. One remarkable difference between standard and flipped $SU(5)$ models is the lower probability for higher rank gauge groups. The massless models show no exceptional behavior as far as the gauge factors are concerned.

The total rank distribution for both, the standard and the flipped version, differs in one aspect from the one obtained in [4], namely, in the large fraction of hidden sector groups with a total rank of 10 or 9, respectively. This can be explained by just one specific construction which is possible for various values of the complex structure param-

eters. In this setup all but one of the hidden sector branes are on top of the orientifold planes on all three tori. If we exclude this specific feature of the $SU(5)$ construction, the remaining distribution shows the behavior estimated from the prior results.

This holds true for the massless models as well, where mainly solutions without symmetric representations contribute to the peak at a total rank of nine. Yet it is striking that no massless models with a total rank of three are found and that the massless models without symmetric representations exclusively appear with a rank of two, five or nine.

Note that while comparing the distributions one has to take into account that the total rank of the hidden sector gauge group in the $SU(5)$ case is lowered by the contribution from the visible sector branes to the tadpole cancellation conditions. In the flipped case, the additional $U(1)$ -brane contributes as well.

IV. CONCLUSIONS

In this paper we presented an analysis of a large number of $SU(5)$ and flipped $SU(5)$ models on a $T^6/\mathbb{Z}_2 \times \mathbb{Z}_2$ orientifold. Our analysis showed that three generation models with a minimal grand unified gauge group are heavily suppressed in this setup. This result was expected, since we know that the explicit construction of three generation $SU(5)$ models on this specific orientifold has turned out to be difficult. For models without symmetric representations it has been proven in [7] that there exist no models at all.

The analysis of the hidden sector showed that the frequency distributions of the total rank of the gauge group and of single gauge group factors are quite similar to the results obtained in [4]. Differences in the qualitative picture result from specific effects in the $SU(5)$ construction.

Comparing the results for the standard and flipped $SU(5)$ models with and without a massless $U(1)_X$, we find no significant differences. If we allow for symmetric representations, there is basically no additional suppression factor. If we restrict ourselves to models without these representations, flipped constructions are 3 times less likely than the standard ones.

ACKNOWLEDGMENTS

It is a pleasure to thank Carlo Angelantonj, Ralph Blumenhagen, Mirjam Cvetič, Gabriele Honecker, Tianjun Li, Dieter Lüst, and Timo Weigand for valuable discussions.

- [1] H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
- [2] S.M. Barr, Phys. Lett. **112B**, 219 (1982); J.P. Derendinger, J.E. Kim, and D.V. Nanopoulos, Phys. Lett. **139B**, 170 (1984).
- [3] R. Blumenhagen, F. Gmeiner, G. Honecker, D. Lüst, and T. Weigand, Nucl. Phys. **B713**, 83 (2005).
- [4] F. Gmeiner, R. Blumenhagen, G. Honecker, D. Lüst, and T. Weigand, J. High Energy Phys. 01 (2006) 004; F. Gmeiner, Fortschr. Phys. **54**, 391 (2006).
- [5] R. Blumenhagen, M. Cvetič, P. Langacker, and G. Shiu, J. High Energy Phys. 03 (2005) 050.
- [6] S. Förste, G. Honecker, and R. Schreyer, Nucl. Phys. **B593**, 127 (2001); M. Cvetič, G. Shiu, and A.M. Uranga, Nucl. Phys. **B615**, 3 (2001).
- [7] M. Cvetič, I. Papadimitriou, and G. Shiu, Nucl. Phys. **B659**, 193 (2003); **B696**, 298(E) (2004).
- [8] M. Axenides, E. Floratos, and C. Kokorelis, J. High Energy Phys. 10 (2003) 006; C.M. Chen, G.V. Kaniotis, V.E. Mayes, D.V. Nanopoulos, and J.W. Walker, Phys. Lett. B **611**, 156 (2005); **625**, 96 (2005); C.M. Chen, V.E. Mayes, and D.V. Nanopoulos, Phys. Lett. B **633**, 618 (2006).
- [9] R. Blumenhagen, B. Körs, D. Lüst, and T. Ott, Nucl. Phys. **B616**, 3 (2001); J.R. Ellis, P. Kanti, and D.V. Nanopoulos, Nucl. Phys. **B647**, 235 (2002).
- [10] A.M. Uranga, Nucl. Phys. **B598**, 225 (2001).