

Measurements of CP -violating asymmetries and branching fractions in B decays to ωK and $\omega\pi$

- B. Aubert,¹ R. Barate,¹ M. Bona,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulsom,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu. Todyshev,¹² D. S. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ D. Kovalskyi,¹⁷ J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroeseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ A. Dvoretskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² F. Winklmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ H. Jasper,²³ B. Spaan,²³ T. Brandt,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ W. F. Mader,²⁴ R. Nogowski,²⁴ A. Petzold,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵ G. R. Bonneau,²⁵ P. Grenier,^{25,*} E. Latour,²⁵ Ch. Thiebaux,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ A. I. Robertson,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ A. Petrella,²⁷ L. Piemontese,²⁷ E. Prencipe,²⁷ F. Anulli,²⁸ R. Baldini-Ferroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,†} M. Piccolo,²⁸ M. Rama,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. R. Gaillard,³² J. A. Nash,³² M. B. Nikolich,³² W. Panduro Vazquez,³² X. Chai,³³ M. J. Charles,³³ U. Mallik,³³ N. T. Meyer,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ A. V. Gritsan,³⁵ M. Fritsch,³⁶ G. Schott,³⁶ N. Arnaud,³⁷ M. Davier,³⁷ G. Grosdidier,³⁷ A. Höcker,³⁷ F. Le Diberder,³⁷ V. Lepeltier,³⁷ A. M. Lutz,³⁷ A. Oyanguren,³⁷ S. Pruvot,³⁷ S. Rodier,³⁷ P. Roudeau,³⁷ M. H. Schune,³⁷ A. Stocchi,³⁷ W. F. Wang,³⁷ G. Wormser,³⁷ C. H. Cheng,³⁸ D. J. Lange,³⁸ D. M. Wright,³⁸ C. A. Chavez,³⁹ I. J. Forster,³⁹ J. R. Fry,³⁹ E. Gabathuler,³⁹ R. Gamet,³⁹ K. A. George,³⁹ D. E. Hutchcroft,³⁹ D. J. Payne,³⁹ K. C. Schofield,³⁹ C. Touramanis,³⁹ A. J. Bevan,⁴⁰ F. Di Lodovico,⁴⁰ W. Menges,⁴⁰ R. Sacco,⁴⁰ C. L. Brown,⁴¹ G. Cowan,⁴¹ H. U. Flaecher,⁴¹ D. A. Hopkins,⁴¹ P. S. Jackson,⁴¹ T. R. McMahon,⁴¹ S. Ricciardi,⁴¹ F. Salvatore,⁴¹ D. N. Brown,⁴² C. L. Davis,⁴² J. Allison,⁴³ N. R. Barlow,⁴³ R. J. Barlow,⁴³ Y. M. Chia,⁴³ C. L. Edgar,⁴³ M. P. Kelly,⁴³ G. D. Lafferty,⁴³ M. T. Naisbit,⁴³ J. C. Williams,⁴³ J. I. Yi,⁴³ C. Chen,⁴⁴ W. D. Hulsbergen,⁴⁴ A. Jawahery,⁴⁴ C. K. Lae,⁴⁴ D. A. Roberts,⁴⁴ G. Simi,⁴⁴ G. Blaylock,⁴⁵ C. Dallapiccola,⁴⁵ S. S. Hertzbach,⁴⁵ X. Li,⁴⁵ T. B. Moore,⁴⁵ S. Saremi,⁴⁵ H. Staengle,⁴⁵ S. Y. Willocq,⁴⁵ R. Cowan,⁴⁶ K. Koeneke,⁴⁶ G. Sciolla,⁴⁶ S. J. Sekula,⁴⁶ M. Spitznagel,⁴⁶ F. Taylor,⁴⁶ R. K. Yamamoto,⁴⁶ H. Kim,⁴⁷ P. M. Patel,⁴⁷ C. T. Potter,⁴⁷ S. H. Robertson,⁴⁷ A. Lazzaro,⁴⁸ V. Lombardo,⁴⁸ F. Palombo,⁴⁸ J. M. Bauer,⁴⁹ L. Cremaldi,⁴⁹ V. Eschenburg,⁴⁹ R. Godang,⁴⁹ R. Kroeger,⁴⁹ J. Reidy,⁴⁹ D. A. Sanders,⁴⁹ D. J. Summers,⁴⁹ H. W. Zhao,⁴⁹ S. Brunet,⁵⁰ D. Côté,⁵⁰ M. Simard,⁵⁰ P. Taras,⁵⁰ F. B. Viaud,⁵⁰ H. Nicholson,⁵¹ N. Cavallo,⁵² G. De Nardo,⁵² D. del Re,⁵² F. Fabozzi,^{52,‡} C. Gatto,⁵² L. Lista,⁵² D. Monorchio,⁵² D. Picколо,⁵² C. Sciacca,⁵² M. Baak,⁵³ H. Bulten,⁵³ G. Raven,⁵³ H. L. Snoek,⁵³ C. P. Jessop,⁵⁴ J. M. LoSecco,⁵⁴ T. Allmendinger,⁵⁵ G. Benelli,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵ P. D. Jackson,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ T. Pulliam,⁵⁵ A. M. Rahimi,⁵⁵ R. Ter-Antonyan,⁵⁵ Q. K. Wong,⁵⁵ N. L. Blount,⁵⁶ J. Brau,⁵⁶ R. Frey,⁵⁶ O. Igonkina,⁵⁶ M. Lu,⁵⁶ R. Rahmat,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶ J. Strube,⁵⁶ E. Torrence,⁵⁶ F. Galeazzi,⁵⁷ A. Gaz,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ A. Pompili,⁵⁷ M. Posocco,⁵⁷ M. Rotondo,⁵⁷ F. Simonetto,⁵⁷

B. AUBERT *et al.*PHYSICAL REVIEW D **74**, 011106(R) (2006)

- R. Stroili,⁵⁷ C. Voci,⁵⁷ M. Benayoun,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ L. Del Buono,⁵⁸ Ch. de la Vaissière,⁵⁸ O. Hamon,⁵⁸
 B. L. Hartfiel,⁵⁸ M. J. J. John,⁵⁸ Ph. Leruste,⁵⁸ J. Malclès,⁵⁸ J. Ocariz,⁵⁸ L. Roos,⁵⁸ G. Therin,⁵⁸ P. K. Behera,⁵⁹
 L. Gladney,⁵⁹ J. Panetta,⁵⁹ M. Biasini,⁶⁰ R. Covarelli,⁶⁰ M. Pioppi,⁶⁰ C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹
 F. Bucci,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ R. Cenci,⁶¹ F. Forti,⁶¹ M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹
 M. A. Mazur,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ E. Paoloni,⁶¹ G. Rizzo,⁶¹ J. Walsh,⁶¹ M. Haire,⁶² D. Judd,⁶² D. E. Wagoner,⁶²
 J. Biesiada,⁶³ N. Danielson,⁶³ P. Elmer,⁶³ Y. P. Lau,⁶³ C. Lu,⁶³ J. Olsen,⁶³ A. J. S. Smith,⁶³ A. V. Telnov,⁶³ F. Bellini,⁶⁴
 G. Cavoto,⁶⁴ A. D'Orazio,⁶⁴ E. Di Marco,⁶⁴ R. Faccini,⁶⁴ F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ L. Li Gioi,⁶⁴
 M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ G. Piredda,⁶⁴ F. Polci,⁶⁴ F. Safai Tehrani,⁶⁴ C. Vena,⁶⁴ M. Ebert,⁶⁵ H. Schröder,⁶⁵
 R. Waldi,⁶⁵ T. Adye,⁶⁶ N. De Groot,⁶⁶ B. Franek,⁶⁶ E. O. Olaiya,⁶⁶ F. F. Wilson,⁶⁶ S. Emery,⁶⁷ A. Gaidot,⁶⁷ S. F. Ganzhur,⁶⁷
 G. Hamel de Monchenault,⁶⁷ W. Kozanecki,⁶⁷ M. Legendre,⁶⁷ B. Mayer,⁶⁷ G. Vasseur,⁶⁷ Ch. Yèche,⁶⁷ M. Zito,⁶⁷
 W. Park,⁶⁸ M. V. Purohit,⁶⁸ A. W. Weidemann,⁶⁸ J. R. Wilson,⁶⁸ M. T. Allen,⁶⁹ D. Aston,⁶⁹ R. Bartoldus,⁶⁹ P. Bechtle,⁶⁹
 N. Berger,⁶⁹ A. M. Boyarski,⁶⁹ R. Claus,⁶⁹ J. P. Coleman,⁶⁹ M. R. Convery,⁶⁹ M. Cristinziani,⁶⁹ J. C. Dingfelder,⁶⁹
 D. Dong,⁶⁹ J. Dorfan,⁶⁹ G. P. Dubois-Felsmann,⁶⁹ D. Dujmic,⁶⁹ W. Dunwoodie,⁶⁹ R. C. Field,⁶⁹ T. Glanzman,⁶⁹
 S. J. Gowdy,⁶⁹ M. T. Graham,⁶⁹ V. Halyo,⁶⁹ C. Hast,⁶⁹ T. Hryna'ova,⁶⁹ W. R. Innes,⁶⁹ M. H. Kelsey,⁶⁹ P. Kim,⁶⁹
 M. L. Kocian,⁶⁹ D. W. G. S. Leith,⁶⁹ S. Li,⁶⁹ J. Libby,⁶⁹ S. Luitz,⁶⁹ V. Luth,⁶⁹ H. L. Lynch,⁶⁹ D. B. MacFarlane,⁶⁹
 H. Marsiske,⁶⁹ R. Messner,⁶⁹ D. R. Muller,⁶⁹ C. P. O'Grady,⁶⁹ V. E. Ozcan,⁶⁹ A. Perazzo,⁶⁹ M. Perl,⁶⁹ B. N. Ratcliff,⁶⁹
 A. Roodman,⁶⁹ A. A. Salnikov,⁶⁹ R. H. Schindler,⁶⁹ J. Schwiening,⁶⁹ A. Snyder,⁶⁹ J. Stelzer,⁶⁹ D. Su,⁶⁹ M. K. Sullivan,⁶⁹
 K. Suzuki,⁶⁹ S. K. Swain,⁶⁹ J. M. Thompson,⁶⁹ J. Va'vra,⁶⁹ N. van Bakel,⁶⁹ M. Weaver,⁶⁹ A. J. R. Weinstein,⁶⁹
 W. J. Wisniewski,⁶⁹ M. Wittgen,⁶⁹ D. H. Wright,⁶⁹ A. K. Yarritu,⁶⁹ K. Yi,⁶⁹ C. C. Young,⁶⁹ P. R. Burchat,⁷⁰ A. J. Edwards,⁷⁰
 S. A. Majewski,⁷⁰ B. A. Petersen,⁷⁰ C. Roat,⁷⁰ L. Wilden,⁷⁰ S. Ahmed,⁷¹ M. S. Alam,⁷¹ R. Bula,⁷¹ J. A. Ernst,⁷¹ V. Jain,⁷¹
 B. Pan,⁷¹ M. A. Saeed,⁷¹ F. R. Wappler,⁷¹ S. B. Zain,⁷¹ W. Bugg,⁷² M. Krishnamurthy,⁷² S. M. Spanier,⁷² R. Eckmann,⁷³
 J. L. Ritchie,⁷³ A. Satpathy,⁷³ C. J. Schilling,⁷³ R. F. Schwitters,⁷³ J. M. Izen,⁷⁴ I. Kitayama,⁷⁴ X. C. Lou,⁷⁴ S. Ye,⁷⁴
 F. Bianchi,⁷⁵ F. Gallo,⁷⁵ D. Gamba,⁷⁵ M. Bomben,⁷⁶ L. Bosisio,⁷⁶ C. Cartaro,⁷⁶ F. Cossutti,⁷⁶ G. Della Ricca,⁷⁶
 S. Dittongo,⁷⁶ S. Grancagnolo,⁷⁶ L. Lanceri,⁷⁶ L. Vitale,⁷⁶ V. Azzolini,⁷⁷ F. Martinez-Vidal,⁷⁷ Sw. Banerjee,⁷⁸
 B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸
 J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰
 A. M. Eichenbaum,⁸⁰ K. T. Flood,⁸⁰ J. J. Hollar,⁸⁰ J. R. Johnson,⁸⁰ P. E. Kutter,⁸⁰ H. Li,⁸⁰ R. Liu,⁸⁰ B. Mellado,⁸⁰
 A. Mihalyi,⁸⁰ A. K. Mohapatra,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ P. Tan,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France²Universitat de Barcelona Fac. Fisica. Dept. ECM Avda Diagonal 647, 6a planta E-08028 Barcelona, Spain³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy⁴Institute of High Energy Physics, Beijing 100039, China⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany⁹University of Bristol, Bristol BS8 1TL, United Kingdom¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia¹³University of California at Irvine, Irvine, California 92697, USA¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA¹⁵University of California at Riverside, Riverside, California 92521, USA¹⁶University of California at San Diego, La Jolla, California 92093, USA¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA¹⁹California Institute of Technology, Pasadena, California 91125, USA²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA²¹University of Colorado, Boulder, Colorado 80309, USA²²Colorado State University, Fort Collins, Colorado 80523, USA²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

- ²⁵*Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁶*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁷*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*
- ²⁸*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁹*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ³⁰*Harvard University, Cambridge, Massachusetts 02138, USA*
- ³¹*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³²*Imperial College London, London, SW7 2AZ, United Kingdom*
- ³³*University of Iowa, Iowa City, Iowa 52242, USA*
- ³⁴*Iowa State University, Ames, Iowa 50011-3160, USA*
- ³⁵*Johns Hopkins Univ. Dept of Physics & Astronomy 3400 N. Charles Street Baltimore, Maryland 21218, USA*
- ³⁶*Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*
- ³⁷*Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France*
- ³⁸*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
- ³⁹*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ⁴⁰*Queen Mary, University of London, E1 4NS, United Kingdom*
- ⁴¹*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ⁴²*University of Louisville, Louisville, Kentucky 40292, USA*
- ⁴³*University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴⁴*University of Maryland, College Park, Maryland 20742, USA*
- ⁴⁵*University of Massachusetts, Amherst, Massachusetts 01003, USA*
- ⁴⁶*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*
- ⁴⁷*McGill University, Montréal, Québec, Canada H3A 2T8*
- ⁴⁸*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁹*University of Mississippi, University, Mississippi 38677, USA*
- ⁵⁰*Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*
- ⁵¹*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*
- ⁵²*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵³*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵⁴*University of Notre Dame, Notre Dame, Indiana 46556, USA*
- ⁵⁵*Ohio State University, Columbus, Ohio 43210, USA*
- ⁵⁶*University of Oregon, Eugene, Oregon 97403, USA*
- ⁵⁷*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁸*Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France*
- ⁵⁹*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁶⁰*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁶¹*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶²*Prairie View A&M University, Prairie View, Texas 77446, USA*
- ⁶³*Princeton University, Princeton, New Jersey 08544, USA*
- ⁶⁴*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶⁵*Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁶*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁷*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁸*University of South Carolina, Columbia, South Carolina 29208, USA*
- ⁶⁹*Stanford Linear Accelerator Center, Stanford, California 94309, USA*
- ⁷⁰*Stanford University, Stanford, California 94305-4060, USA*
- ⁷¹*State University of New York, Albany, New York 12222, USA*
- ⁷²*University of Tennessee, Knoxville, Tennessee 37996, USA*
- ⁷³*University of Texas at Austin, Austin, Texas 78712, USA*
- ⁷⁴*University of Texas at Dallas, Richardson, Texas 75083, USA*
- ⁷⁵*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*
- ⁷⁶*Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*
- ⁷⁷*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*
- ⁷⁸*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*
- ⁷⁹*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*

^{*}Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France[†]Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy[‡]Also with Università della Basilicata, Potenza, Italy

⁸⁰*University of Wisconsin, Madison, Wisconsin 53706, USA*⁸¹*Yale University, New Haven, Connecticut 06511, USA*

(Received 20 March 2006; published 21 July 2006)

We present measurements of CP -violating asymmetries and branching fractions for the decays $B^+ \rightarrow \omega\pi^+$, $B^+ \rightarrow \omega K^+$, and $B^0 \rightarrow \omega K^0$. The data sample corresponds to $232 \times 10^6 B\bar{B}$ pairs produced by e^+e^- annihilation at the $\Upsilon(4S)$ resonance. For the decay $B^0 \rightarrow \omega K_S^0$, we measure the time-dependent CP -violation parameters $S = 0.51^{+0.35}_{-0.39} \pm 0.02$, and $C = -0.55^{+0.28}_{-0.26} \pm 0.03$. We also measure the branching fractions, in units of 10^{-6} , $\mathcal{B}(B^+ \rightarrow \omega\pi^+) = 6.1 \pm 0.7 \pm 0.4$, $\mathcal{B}(B^+ \rightarrow \omega K^+) = 6.1 \pm 0.6 \pm 0.4$, and $\mathcal{B}(B^0 \rightarrow \omega K^0) = 6.2 \pm 1.0 \pm 0.4$, and charge asymmetries $\mathcal{A}_{ch}(B^+ \rightarrow \omega\pi^+) = -0.01 \pm 0.10 \pm 0.01$ and $\mathcal{A}_{ch}(B^+ \rightarrow \omega K^+) = 0.05 \pm 0.09 \pm 0.01$.

DOI: 10.1103/PhysRevD.74.011106

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Measurements of time-dependent CP asymmetries in B^0 meson decays through a Cabibbo-Kobayashi-Maskawa (CKM) favored $b \rightarrow c\bar{c}s$ amplitude [1,2] have firmly established that CP is not conserved in such decays. The effect, arising from the interference between mixing and decay involving the CP -violating phase $\beta = \arg(-V_{cd}V_{cb}^*/V_{td}V_{tb}^*)$ of the CKM mixing matrix [3], manifests itself as an asymmetry in the time evolution of the $B^0\bar{B}^0$ pair.

Decays to the charmless final states ϕK^0 , $K^+K^-K^0$, $\eta' K^0$, $\pi^0 K^0$, $f_0(980)K^0$, and ωK^0 are all $b \rightarrow q\bar{q}s$ processes dominated by a single penguin (loop) amplitude having the same weak phase β [4]. CKM-suppressed amplitudes and multiple particles in the loop complicate the situation by introducing other weak phases whose contributions are not negligible; see Refs. [5,6] for early quantitative work in addressing the size of these effects. We define ΔS as the difference between the time-dependent CP -violating parameter S (given in detail below) measured in these decays and $S = \sin 2\beta$ measured in charmonium K^0 decays. For the decay $B^0 \rightarrow \omega K^0$, these additional contributions are expected to give $\Delta S \sim 0.1$ [7,8], although this increase may be nullified when final-state interactions are included [8]. A value of ΔS inconsistent with this expectation could be an indication of new physics [9].

We present an improved measurement of the time-dependent CP -violating asymmetry in the decay $B^0 \rightarrow \omega K^0$, previously reported by the Belle Collaboration based on a sample of ~ 30 events [10]. We also measure branching fractions for the decays $B^0 \rightarrow \omega K^0$, $B^+ \rightarrow \omega\pi^+$, and $B^+ \rightarrow \omega K^+$ (charge-conjugate decay modes are implied throughout), and for $B^+ \rightarrow \omega\pi^+$, and $B^+ \rightarrow \omega K^+$, we measure the time-integrated charge asymmetry $\mathcal{A}_{ch} = (\Gamma^- - \Gamma^+)/(\Gamma^- + \Gamma^+)$, where Γ^\pm is the width for these charged decay modes. In the Standard Model \mathcal{A}_{ch} is expected to be consistent with zero within our experimental uncertainty; a nonzero value would indicate direct CP violation in this channel.

The data were collected with the *BABAR* detector [11] at the PEP-II asymmetric e^+e^- collider. An integrated luminosity of 211 fb^{-1} , corresponding to $232 \times 10^6 B\bar{B}$ pairs, was recorded at the $\Upsilon(4S)$ resonance (center-of-mass energy $\sqrt{s} = 10.58 \text{ GeV}$). Charged particles are detected and

their momenta measured by the combination of a silicon vertex tracker (SVT), consisting of five layers of double-sided detectors, and a 40-layer central drift chamber, both operating in a 1.5 T axial magnetic field. Charged-particle identification (PID) is provided by the energy loss in the tracking devices and by the measured Cherenkov angle from an internally reflecting ring-imaging Cherenkov detector (DIRC) covering the central region. A K/π separation of better than 4 standard deviations (σ) is achieved for momenta below $3 \text{ GeV}/c$, decreasing to 2.5σ at the highest momenta in the B decay final states. Photons and electrons are detected by a CsI(Tl) electromagnetic calorimeter.

From a $B^0\bar{B}^0$ pair produced in an $\Upsilon(4S)$ decay, we reconstruct one of the B mesons in the final state $f = \omega K_S^0$, a CP eigenstate with eigenvalue -1 . For the time evolution measurement, we also identify (tag) the flavor (B^0 or \bar{B}^0) and reconstruct the decay vertex of the other B . The asymmetric beam configuration in the laboratory frame provides a boost of $\beta\gamma = 0.56$ to the $\Upsilon(4S)$, which allows the determination of the proper decay time difference $\Delta t \equiv t_f - t_{tag}$ from the vertex separation of the two B meson candidates. Ignoring the Δt resolution (about 0.5 ps), the distribution of Δt is

$$F(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} [1 \mp \Delta w \pm (1 - 2w)(S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t))]. \quad (1)$$

The upper (lower) sign denotes a decay accompanied by a B^0 (\bar{B}^0) tag, τ is the mean B^0 lifetime, Δm_d is the mixing frequency, and the mistag parameters w and Δw are the average and difference, respectively, of the probabilities that a true B^0 (\bar{B}^0) meson is tagged as a \bar{B}^0 (B^0). The parameter C measures direct CP violation. If $C = 0$, then $S = \sin 2\beta + \Delta S$.

The flavor-tagging algorithm [1] has seven mutually exclusive tagging categories of differing purities (including one for untagged events that we retain for yield determinations). The measured analyzing power, defined as efficiency times $(1 - 2w)^2$ summed over all categories, is $(30.5 \pm 0.6)\%$, as determined from a large sample of B -decays to fully reconstructed flavor eigenstates (B_{flav}).

MEASUREMENTS OF CP -VIOLATING ASYMMETRIES ...

We reconstruct a B meson candidate by combining a π^+ , K^+ or K_S^0 with an $\omega \rightarrow \pi^+ \pi^- \pi^0$. We select $K_S^0 \rightarrow \pi^+ \pi^-$ decays by requiring the $\pi^+ \pi^-$ invariant mass to be within 12 MeV of the nominal K^0 mass and by requiring a flight length greater than 3 times its error. We require the primary charged track to have a minimum of six Cherenkov photons in the DIRC. We require the $\pi^+ \pi^- \pi^0$ invariant mass ($m_{3\pi}$) to be between 735 and 825 MeV. Distributions from the data and from Monte Carlo (MC) simulations [12] guide the choice of these selection criteria. We retain regions adequate to characterize the background as well as the signal for those quantities taken subsequently as observables for fitting. We also use in the fit the angle θ_H , defined, in the ω rest frame, as the angle of the direction of the boost from the B rest frame with respect to the normal to the ω decay plane. The quantity $\mathcal{H} \equiv |\cos\theta_H|$ is approximately flat for background and distributed as $\cos^2\theta_H$ for signal.

A B meson candidate is characterized kinematically by the energy-substituted mass $m_{\text{ES}} \equiv \sqrt{(\frac{1}{2}s + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - \mathbf{p}_B^2}$ and the energy difference $\Delta E \equiv E_B^* - \frac{1}{2}\sqrt{s}$, where (E_0, \mathbf{p}_0) and (E_B, \mathbf{p}_B) are four-momenta of the $\Upsilon(4S)$ and the B candidate, respectively, and the asterisk denotes the $\Upsilon(4S)$ rest frame. We require, assuming the $B^+ \rightarrow \omega \pi^+$ hypothesis, $|\Delta E| \leq 0.2$ GeV and $5.25 \leq m_{\text{ES}} \leq 5.29$ GeV.

To reject the dominant background from continuum $e^+ e^- \rightarrow q\bar{q}$ events ($q = u, d, s, c$), we use the angle θ_T between the thrust axis of the B candidate and that of the rest of the tracks and neutral clusters in the event, calculated in the $\Upsilon(4S)$ rest frame. The distribution of $\cos\theta_T$ is sharply peaked near ± 1 for jetlike $q\bar{q}$ pairs and is nearly uniform for the isotropic B decays; we require $|\cos\theta_T| < 0.9$ (0.8 for the charged B decays).

From MC simulations of $B^0 \bar{B}^0$ and $B^+ B^-$ events, we find evidence for a small (0.5%) $B\bar{B}$ background contribution for the charged B decays, so we have added a $B\bar{B}$ component to the fit described below for those channels.

We use an unbinned, multivariate maximum-likelihood fit to extract signal yields and CP -violation parameters. We use the discriminating variables m_{ES} , ΔE , $m_{3\pi}$, \mathcal{H} , and a Fisher discriminant \mathcal{F} [13]. The Fisher discriminant combines five variables: the polar angles with respect to the beam axis in the $\Upsilon(4S)$ frame of the B candidate momentum and of the B thrust axis; the tagging category; and the zeroth and second angular moments of the energy flow, excluding the B candidate, about the B thrust axis [13]. We also use Δt for the $B^0 \rightarrow \omega K_S^0$ decay, while for the charged B decays we use the PID variables T_π and T_K , defined as the number of standard deviations between the measured DIRC Cherenkov angle and that expected for pions and kaons, respectively.

For the $B^0 \rightarrow \omega K_S^0$ decay we define the probability density function (PDF) for each event i , hypothesis j (signal and $q\bar{q}$ background), and tagging category c

PHYSICAL REVIEW D **74**, 011106(R) (2006)

$$\begin{aligned} \mathcal{P}_{j,c}^i &\equiv \mathcal{P}_j(m_{\text{ES}}^i) \mathcal{P}_j(\Delta E^i) \mathcal{P}_j(\mathcal{F}^i) \mathcal{P}_j(m_{3\pi}^i) \mathcal{P}_j(\mathcal{H}^i) \\ &\quad \times \mathcal{P}_j(\Delta t^i, \sigma_{\Delta t}^i, c), \end{aligned} \quad (2)$$

where $\sigma_{\Delta t}^i$ is the error on Δt for event i . We write the extended likelihood function as

$$\mathcal{L} = \prod_c \exp\left(-\sum_j Y_j f_{j,c}\right) \prod_i^{N_c} \left[\sum_j Y_j f_{j,c} \mathcal{P}_{j,c}^i \right], \quad (3)$$

where Y_j is the fit yield of events of species j , $f_{j,c}$ is the fraction of events of species j for each category c , and N_c is the number of events of category c in the sample. We fix $f_{\text{sig},c}$ to $f_{B_{\text{flav}},c}$, the values measured with the large B_{flav} sample [1]. The same likelihood function is used for the charged decays except that the hypothesis j also includes $B\bar{B}$ background, the tagging category is not used and the PDF is slightly different, involving flavor k (primary π^+ or K^+):

$$\mathcal{P}_{jk}^i = \mathcal{P}_j(m_{\text{ES}}^i) \mathcal{P}_j(\Delta E_k^i, T_k^i) \mathcal{P}_j(\mathcal{F}^i) \mathcal{P}_j(m_{3\pi}^i) \mathcal{P}_j(\mathcal{H}^i). \quad (4)$$

The PDF $\mathcal{P}_{\text{sig}}(\Delta t, \sigma_{\Delta t}, c)$, is the convolution of $F(\Delta t; c)$ (Eq. (1)) with the signal resolution function (a sum of three Gaussians) determined from the B_{flav} sample. The other PDF forms are: the sum of two Gaussians for all signal shapes except \mathcal{H} , and the peaking component of the $m_{3\pi}$ background; the sum of three Gaussians for $\mathcal{P}_{q\bar{q}}(\Delta t; c)$; an asymmetric Gaussian with different widths below and above the peak for $\mathcal{P}_j(\mathcal{F})$ (a small “tail” Gaussian is added for $\mathcal{P}_{q\bar{q}}(\mathcal{F})$); Chebyshev functions of second to fourth order for \mathcal{H} signal and the slowly-varying shapes of ΔE , $m_{3\pi}$, and \mathcal{H} backgrounds; and, for $\mathcal{P}_{q\bar{q}}(m_{\text{ES}})$, a phase-space-motivated empirical function [14], with a small Gaussian added for $\mathcal{P}_{B\bar{B}}(m_{\text{ES}})$.

We determine the PDF parameters from simulation for the signal and $B\bar{B}$ background components. We study large control samples of $B \rightarrow D\pi$ decays of similar topology to verify the simulated resolutions in ΔE and m_{ES} , adjusting the PDFs to account for any differences found. For the $q\bar{q}$ background we use $(m_{\text{ES}}, \Delta E)$ sideband data to obtain initial PDF-parameter values but ultimately leave them free to vary in the final fit.

We compute the branching fractions and charge asymmetry from fits performed without Δt or flavor tagging. The free fit parameters are the following: the signal and $q\bar{q}$ background yields (the $B\bar{B}$ yield, if present, is fixed); the three shape parameters of $\mathcal{P}_{q\bar{q}}(\mathcal{F})$; the slope of $\mathcal{P}_{q\bar{q}}(\Delta E)$ and $\mathcal{P}_{q\bar{q}}(m_{3\pi})$; the fraction of the peaking component of $\mathcal{P}_{q\bar{q}}(m_{3\pi})$; ξ [14]; and, for the charged B decays, the signal and background \mathcal{A}_{ch} .

Table I lists the quantities used to determine the branching fraction. Equal production rates of $B^+ B^-$ and $B^0 \bar{B}^0$ pairs have been assumed. Small yield biases are present in the fit, due primarily to unmodeled correlations among the

B. AUBERT *et al.*

TABLE I. Fit sample size, signal yield, estimated yield bias (all in events), estimated purity, detection efficiency, daughter branching fraction product, statistical significance including systematic errors, measured branching fraction, and corrected signal charge asymmetry.

Quantity	$\omega\pi^+$	ωK^+	ωK^0
Events in fit	44175		9145
Signal yield	274 ± 28	266 ± 24	100 ± 15
Yield bias	18	16	8
Purity (%)	34	46	46
Eff. (ϵ , %)	21.8	21.2	23.0
$\prod \mathcal{B}_i$	0.891	0.891	0.307
$\epsilon \times \prod \mathcal{B}_i$ (%)	18.2	17.7	6.4
Significance (σ)	10.8	13.0	8.6
$\mathcal{B}(10^{-6})$	6.1 ± 0.7	6.1 ± 0.6	6.2 ± 1.0
Signal \mathcal{A}_{ch}	-0.01 ± 0.10	0.05 ± 0.09	—

signal PDF parameters. In Table I we include estimates of these biases, evaluated by fitting simulated $q\bar{q}$ experiments drawn from the PDF into which we have embedded the expected number of signal and $B\bar{B}$ background events randomly extracted from the fully simulated MC samples. The estimated purity in Table I is given by the ratio of the signal yield to the effective background plus signal, the latter being defined as the square of the error on the yield.

Figure 1 shows projections onto m_{ES} and ΔE for a subset of the data (including 45–65% of signal events) for which the signal likelihood (computed without the variable plotted) exceeds a threshold that optimizes the sensitivity.




FIG. 1 (color online). The B candidate m_{ES} and ΔE projections for $B^+ \rightarrow \omega\pi^+$ (a, b), $B^+ \rightarrow \omega K^+$ (c, d), and $B^0 \rightarrow \omega K^0$ (e, f) shown for a signal-enhanced subset of the data. Points with error bars represent the data, the solid line the fit function, and the dashed line the background components. Note that the ωK^+ signal in the ΔE plot is displaced from zero since ΔE is defined for the $\omega\pi^+$ hypothesis.

For the time-dependent analysis, we require $|\Delta t| < 20$ ps and $\sigma_{\Delta t} < 2.5$ ps. The free parameters in the fit are the same as for the branching fraction fit plus S , C , the fraction of background events in each tagging category, and the six primary parameters describing the Δt background shape. The parameters τ and Δm_d are fixed to world-average values [15]. Here we find a slightly smaller yield of 95 ± 14 events and $S = 0.51^{+0.35}_{-0.39}$, $C = -0.55^{+0.28}_{-0.26}$. The errors have been scaled by ~ 1.10 to account for a slight underestimate of the fit errors predicted by our simulations when the signal sample size is small. Figure 2 shows the Δt projections and asymmetry of the time-dependent fit with events selected as for Fig. 1.

The major systematic uncertainties affecting the branching fraction measurements include the reconstruction efficiency (0.8% per charged track, 1.5% per photon, and 2.1% per K_S^0) estimated from auxiliary studies. We take one-half of the measured yield bias (3–4%) as a systematic error. The uncertainty due to the signal PDF description is estimated to be $\lesssim 1\%$ in studies where the signal PDF parameters are varied within their estimated errors. The uncertainty due to $B\bar{B}$ background is also estimated to be 1% by variation of the fixed $B\bar{B}$ yield by its estimated uncertainty. The \mathcal{A}_{ch} bias is estimated to be -0.005 ± 0.010 from studies of signal MC, control samples, and calculation of the asymmetry due to particles interacting in the detector. We correct for this bias and assign a systematic uncertainty of 0.01 for \mathcal{A}_{ch} for both $B^+ \rightarrow \omega\pi^+$ and $B^+ \rightarrow \omega K^+$.

For the time-dependent measurements, we estimate systematic uncertainties in S and C due to $B\bar{B}$ background and PDF shape variation (0.01 each), modeling of the signal Δt




FIG. 2 (color online). Projections onto Δt for $B^0 \rightarrow \omega K^0$. Data (points with errors), the fit function (solid line), background component (dashed line), and signal component (dotted line), for events in which the tag meson is (a) B^0 and (b) \bar{B}^0 , and (c) the asymmetry $(N_{B^0} - N_{\bar{B}^0})/(N_{B^0} + N_{\bar{B}^0})$.

MEASUREMENTS OF CP -VIOLATING ASYMMETRIES ...

distribution (0.02), and interference between the CKM-suppressed $\bar{b} \rightarrow \bar{u}cd$ amplitude and the favored $b \rightarrow c\bar{u}d$ amplitude for some tagside B decays [16] (0.02 for C , negligible for S). We also find that the uncertainty due to SVT alignment and position and size of the beam spot are negligible. The B_{flav} sample is used to determine the errors associated with the signal PDF parameters: Δt resolutions, tagging efficiencies, and mistag rates; published measurements [15] are used for τ_B and Δm_d . Summing all systematic errors in quadrature, we obtain 0.02 for S and 0.03 for C .

In conclusion, we have measured the branching fractions and time-integrated charge asymmetry for the decays $B^+ \rightarrow \omega\pi^+$ and $B^+ \rightarrow \omega K^+$ and the branching fraction for $B^0 \rightarrow \omega K^0$. We find $\mathcal{B}(B^+ \rightarrow \omega\pi^+) = (6.1 \pm 0.7 \pm 0.4) \times 10^{-6}$, $\mathcal{B}(B^+ \rightarrow \omega K^+) = (6.1 \pm 0.6 \pm 0.4) \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \omega K^0) = (6.2 \pm 1.0 \pm 0.4) \times 10^{-6}$, $\mathcal{A}_{\text{ch}}(B^+ \rightarrow \omega\pi^+) = -0.01 \pm 0.10 \pm 0.01$, and $\mathcal{A}_{\text{ch}}(B^+ \rightarrow \omega K^+) = 0.05 \pm 0.09 \pm 0.01$, where the first errors are statistical and the second systematic. These results are substantially more precise than earlier measurements [17] and a significant improvement over our previous measurements [18], which they supersede. We also

PHYSICAL REVIEW D **74**, 011106(R) (2006)

measure the time-dependent asymmetry parameters for the decay $B^0 \rightarrow \omega K^0$, $S = 0.51^{+0.35}_{-0.39} \pm 0.02$ and $C = -0.55^{+0.28}_{-0.26} \pm 0.03$, with a precision nearly a factor of 2 better than the previous Belle Collaboration results [10]. If we fix $C = 0$, we find $S = 0.60^{+0.42}_{-0.38}$. This value of S and the world-average value of $\sin 2\beta$ [1,2] yield a value of $\Delta S = 0.12 \pm 0.40$, in good agreement with the expected value near zero.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), Marie Curie EIF (European Union), the A.P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

-
- [1] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 161803 (2005).
 - [2] K. Abe *et al.* (*Belle* Collaboration), Phys. Rev. D **66**, 071102(R) (2002).
 - [3] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
 - [4] Y. Grossman and M. P. Worah, Phys. Lett. B **395**, 241 (1997); D. Atwood and A. Soni, Phys. Lett. B **405**, 150 (1997).
 - [5] Y. Grossman *et al.*, Phys. Rev. D **68**, 015004 (2003).
 - [6] M. Beneke and M. Neubert, Nucl. Phys. **B675**, 333 (2003).
 - [7] M. Beneke, Phys. Lett. B **620**, 143 (2005).
 - [8] H-Y. Cheng, C-K. Chua, and A. Soni, Phys. Rev. D **72**, 014006 (2005).
 - [9] D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
 - [10] K. F. Chen *et al.* (*Belle* Collaboration), Phys. Rev. D **72**, 012004 (2005).
 - [11] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instr. Methods Phys. Res., Sect. A **479**, 1 (2002).
 - [12] The *BABAR* detector Monte Carlo simulation is based on GEANT4: S. Agostinelli *et al.*, Nucl. Instr. Methods Phys. Res., Sect. A **506**, 250 (2003).
 - [13] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **70**, 032006 (2004).
 - [14] $f(x) \propto x\sqrt{1-x^2} \exp[-\xi(1-x^2)]$, with $x \equiv m_{\text{ES}}/E_b$ and ξ a parameter to be fit. See H. Albrecht *et al.* (*ARGUS* Collaboration), Phys. Lett. B **241**, 278 (1990).
 - [15] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
 - [16] Owen Long, Max Baak, Robert N. Cahn, and David Kirkby, Phys. Rev. D **68**, 034010 (2003).
 - [17] C. H. Wang *et al.* (*Belle* Collaboration), Phys. Rev. D **70**, 012001 (2004); C. P. Jessop *et al.* (*CLEO* Collaboration), Phys. Rev. Lett. **85**, 2881 (2000).
 - [18] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **92**, 061801 (2004).