

CMB constraints on the simultaneous variation of the fine structure constant and the electron mass

Kazuhide Ichikawa, Toru Kanzaki, and Masahiro Kawasaki

Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan

(Received 4 March 2006; published 14 July 2006)

We study constraints on time variation of the fine structure constant α from cosmic microwave background (CMB) taking into account simultaneous change in α and the electron mass m_e which might be implied in unification theories. We obtain the constraints $-0.097 < \Delta\alpha/\alpha < 0.034$ at 95% C.L. using WMAP data only, and $-0.042 < \Delta\alpha/\alpha < 0.026$ combining with the constraint on the Hubble parameter by the HST Hubble Key Project. These are improved by 15% compared with constraints assuming only α varies. We discuss other relations between variations in α and m_e but we do not find evidence for varying α .

DOI: [10.1103/PhysRevD.74.023515](https://doi.org/10.1103/PhysRevD.74.023515)

PACS numbers: 98.80.Cq, 98.70.Vc

I. INTRODUCTION

One of fundamental questions in physics is whether or not the physical constants are literally constant. In fact, the physical constants may change in spacetime within the context of some unification theories such as superstring theory and investigating their constancy is an important probe of those theories. In addition to the theoretical possibility, some observations suggest the time variation of the fine structure constant (the coupling constant of electromagnetic interaction) α . Recent constraints including such non-null results are briefly summarized as follows.

Terrestrial limits on α come from atomic clocks [1–7], the Oklo natural fission reactor in Gabon [8] and meteorites. Reference [5] derived the limit on the temporal derivative of α at present as $(-0.3 \pm 2.0) \times 10^{-15} \text{ yr}^{-1}$. Measurements of Sm isotopes in the Oklo provide two bounds on the variation of α as $\Delta\alpha/\alpha = -(0.8 \pm 1.0) \times 10^{-8}$ and $\Delta\alpha/\alpha = (0.88 \pm 0.07) \times 10^{-7}$ [8]. The former result is null, however, the latter is a strong detection that α was larger at $z \sim 0.1$. The meteorite bound obtained by measuring ^{187}Re decay rate is now controversial although varying α is not suggested anyway. Ref. [9] obtained $\Delta\alpha/\alpha = (-8 \pm 8) \times 10^{-7}$, whereas Refs. [10,11] argued that the constraint should be much weaker due to uncertainties in the decay rate modeling.

On the other hand, there are three kinds of celestial probes. One is to use big bang nucleosynthesis (BBN) [12–15] which provides constraints at very high redshifts ($z \sim 10^9\text{--}10^{10}$), for example, $-5.0 \times 10^{-2} < \Delta\alpha/\alpha < 1.0 \times 10^{-2}$ (95% C.L.) [13] or $|\Delta\alpha/\alpha| < 6 \times 10^{-2}$ [16]. The second is from the spectra of high-redshift quasars ($z \sim 1\text{--}3$) [17–23], where there are conflicting results. Refs. [17,18] suggested that α was smaller at $z \geq 1$, $\Delta\alpha/\alpha = (-0.543 \pm 0.116) \times 10^{-5}$ [18]. This result, however, is not supported by other observations [19–23]. For example, Ref. [19] obtained the constraint as $\Delta\alpha/\alpha = (-0.6 \pm 0.6) \times 10^{-6}$ and the others too found no evidence

for varying α . Finally, we can use Cosmic Microwave Background (CMB) to measure α at $z \sim 1100$ [24–27]. Analyses using pre-WMAP data are found in Refs. [26–28]. References [29,30] derive a constraint using the WMAP first-year data, $-0.05 < \Delta\alpha/\alpha < 0.02$ or $-0.06 < \Delta\alpha/\alpha < 0.01$ (95% C.L.), respectively, with or without marginalizing over the running of the spectral index [30].¹

Recently, there have been many studies on constraining the time variation in α which accompanies the variation of the other coupling constants, as would occur rather naturally in unified theories. Under such a framework, BBN has been studied in Refs. [13,35–42], quasar absorption systems in Ref. [43], meteorites in Ref. [9], the Oklo reactor in Ref. [40], and atomic clock experiments in Refs. [43,44]. For example, BBN constraint improves by up to about 2 orders ($-6.0 \times 10^{-4} < \Delta\alpha/\alpha < 1.5 \times 10^{-4}$ [13]) although the factor may vary depending on how they are correlated to α . Hence, it is important to consider the variation of other coupling constants along with α .

In this paper, we investigate constraint on the time variation of α from CMB using the WMAP first-year data. In particular, we consider m_e to vary dependently to the variation in α because, as mentioned above, such might be the case in some unified theory. Since such theories are now under development, how their variations are related to each other can not be predicted. Therefore, we work in a phenomenological way guided by a low energy effective

¹Note that two results suggesting varying α , one of the Oklo results [8] and the quasar observation of Ref. [18] have different signs of $\Delta\alpha$. These results, if correct, can not be explained by homogeneous and monotonically time-varying α . They may indicate that α is not a monotonically varying function of time or as investigated in Refs. [31–33], may suggest a spatial variation of α . In passing, we refer to Ref. [34] for the effects of spatial varying α on CMB.

theory of a string theory and adopt to vary m_e as a power law of α .²

In the next section, we briefly review the recombination process in the early universe and make clear how it depends on α and m_e . In Sec. III, we illustrate their effects on the epoch of recombination and the shape of CMB power spectrum. In Sec. IV, we describe the relations between the variations in α and m_e which we adopt in our analysis. In Sec. V, we present our constraint and we conclude in Sec. VI.

II. α AND m_e DEPENDENCE OF THE RECOMBINATION PROCESS

Nonstandard values of α and m_e modify the CMB angular power spectrum mainly by changing the epoch of recombination. Thus, let us visit briefly the recombination process in the universe and see where those constants appear. We follow the treatment of Ref. [47], which is implemented in the RECFAST code. They have shown that the recombination process is well approximated by the evolutions of three variables: the proton fraction x_p , the singly ionized helium fraction x_{HeII} , and the matter temperature T_M . Their equations are given below. We denote the Boltzmann constant k , the Planck constant h and the speed of light c . In addition to the variables above, $x_p = n_p/n_H$ and $x_{\text{HeII}} = n_{\text{HeII}}/n_H$, we use the electron fraction $x_e = n_e/n_H = x_p + x_{\text{HeII}}$ as an auxiliary variable (note that n_X stands for the number density of species X but n_H is defined as the total hydrogen number density, including both protons and hydrogen atoms). The redshift is denoted by z and the expansion rate by H .

Adopting the three level approximation, the time evolution of the proton fraction of x_p is described by

$$\frac{dx_p}{dz} = \frac{C_H}{H(z)(1+z)} [x_e x_p n_H R_H - \beta_H (1-x_p) e^{-h\nu_H/kT_M}], \quad (1)$$

where $\nu_H = c/(121.5682 \text{ nm})$ is the Ly α frequency and R_H is the case B recombination coefficient for H which is well fitted by

$$R_H = 10^{-19} F \frac{at^b}{1+ct^d} \text{ m}^3 \text{ s}^{-1} \quad (2)$$

with $t = T_M/(10^4 \text{ K})$, $a = 4.309$, $b = -0.6166$, $c = 0.6703$, $d = 0.5300$ [48], and the fudge factor $F = 1.14$ introduced to reproduce the more precise multilevel calculation [47]. β_H is the photoionization coefficient

²References [45,46] studied CMB constraints on the variation of Higgs expectation value whose effect is assumed to appear only in the change in m_e so there is some overlap between their analysis and ours. However, they did not consider α variation at the same time.

$$\beta_H = R_H \left(\frac{2\pi m_e k T_M}{h^2} \right)^{3/2} \exp\left(-\frac{B_{\text{H}2s}}{k T_M}\right), \quad (3)$$

and C_H is the so-called Peebles reduction factor [49]

$$C_H = \frac{[1 + K_H \Lambda_H n_H (1 - x_p)]}{[1 + K_H (\Lambda_H + \beta_H) n_H (1 - x_p)]}, \quad (4)$$

where the binding energy in the 2s energy level is $B_{\text{H}2s} = 3.4 \text{ eV}$, the two-photon decay rate is $\Lambda_H = 8.22458 \text{ s}^{-1}$ and $K_H = c^3/(8\pi\nu_H^3 H)$.

Here, comments on what Eq. (1) means may be in order. The first term in the square brackets in Eq. (1) represents the recombinations to excited states of the atom, ignoring recombination direct to the ground state. The second term represents the rate of ionization from excited states of the atom. The difference of those two terms is the net rate of production of hydrogen atoms when one could ignore the Ly α resonance photons. These photons reduce the rate by the factor C_H , which can be written as the ratio of the net decay rate to the sum of the decay and ionization rates from the $n = 2$ level,

$$C_H = \frac{\Lambda_R + \Lambda_H}{\Lambda_R + \Lambda_H + \beta_H}. \quad (5)$$

We have rewritten Eq. (4) to derive this expression using the decay rate $\Lambda_R \equiv (K_H n_{1s})^{-1}$ allowed by redshifting of Ly α photons out of the line, and using $n_{1s} = n_H - n_p$ which is justified by the far greater occupation number of the hydrogen atom ground state than that of the excited states altogether.

The evolution of the singly ionized helium fraction of x_{HeII} is similarly described by

$$\frac{dx_{\text{HeII}}}{dz} = \frac{C_{\text{He}}}{H(z)(1+z)} [x_{\text{HeII}} x_e n_H R_{\text{HeI}} - \beta_{\text{HeI}} (f_{\text{He}} - x_{\text{HeII}}) e^{-h\nu_{\text{HeI}}/kT_M}], \quad (6)$$

where f_{He} is the total number fraction of helium to hydrogen (using primordial helium mass fraction Y_p , $f_{\text{He}} = Y_p/\{4(1-Y_p)\}$ where we take Y_p to be 0.24), $\nu_{\text{HeI}} = c/(60.1404 \text{ nm})$ is the frequency corresponding to energy between ground state and $2s^1$ state, and R_{HeI} is the case B HeI recombination coefficient for singlets [50]

$$R_{\text{HeI}} = q \left[\sqrt{\frac{T_M}{T_2}} \left(1 + \frac{T_M}{T_2}\right)^{1-p} \left(1 + \frac{T_M}{T_1}\right)^{1+p} \right]^{-1} \text{ m}^3 \text{ s}^{-1}, \quad (7)$$

with $q = 10^{-16.744}$, $p = 0.711$, $T_1 = 10^{5.114} \text{ K}$ and T_2 fixed arbitrary 3K. β_{HeI} is the photoionization coefficient

$$\beta_{\text{HeI}} = R_{\text{HeI}} \left(\frac{2\pi m_e k T_M}{h^2} \right)^{3/2} \exp\left(-\frac{B_{\text{HeI}2s}}{k T_M}\right), \quad (8)$$

$$C_{\text{HeII}} = \frac{[1 + K_{\text{HeI}}\Lambda_{\text{He}}n_{\text{H}}(f_{\text{He}} - x_{\text{HeII}})\exp(\Delta E/kT_M)]}{[1 + K_{\text{HeI}}(\Lambda_{\text{He}} + \beta_{\text{HeI}})n_{\text{H}}(f_{\text{He}} - x_{\text{HeII}})\exp(\Delta E/kT_M)]}, \quad (9)$$

where the binding energy in the 2s energy level is $B_{\text{HeI}2s} = 3.97$ eV and the two-photon decay rate is $\Lambda_{\text{He}} = 51.3$ s $^{-1}$, $K_{\text{HeI}} = c^3/(8\pi\nu_{\text{HeI}}^3H)$ and ΔE is the energy separation between 2s 1 and 2p 1 , $\Delta E/h = c/(58.4334 \text{ nm}) - \nu_{\text{HeI}}$. Note that, contrary to the case with H, the energy separation ΔE between 2s 1 and 2p 1 is so large that we can not neglect it [51].

The matter temperature T_M is evolved as

$$\frac{dT_M}{dz} = \frac{8\sigma_T a_R T_R^4}{3H(z)(1+z)m_e} \frac{x_e}{1+f_{\text{He}}+x_e} (T_M - T_R) + \frac{2T_M}{(1+z)}, \quad (10)$$

where T_R is the radiation temperature, $\sigma_T = 2\alpha^2 h^2/(3\pi m_e^2 c^2)$ is the Thomson cross section, and $a_R = k^4/(120\pi c^3 h^3)$ is the black-body constant. The Compton scattering makes T_R and T_M identical at high redshifts. However, the adiabatic cooling becomes dominant at low redshifts, which leads to the significant difference between T_M and T_R .

Now, we explain how quantities which appear in these equations depend on α and m_e . Two-photon decay rates scale as $\alpha^8 m_e$ [52,53]. Since binding energies scale as $\alpha^2 m_e$, so do ν_{H} and ν_{HeI} , and K_{H} and K_{HeI} scale as $\alpha^{-6} m_e^{-3}$. The remaining task is to investigate how the recombination coefficient R depends on α and m_e . To do this, we follow the treatment of Ref. [25]. The recombination coefficient can be expressed as

$$R = \sum_{n,l}^* [8\pi(2l+1)] \left(\frac{kT_M}{2\pi m_e}\right)^{3/2} \exp\left(\frac{B_n}{kT_M}\right) \times \int_{B_n/kT_M}^{\infty} \frac{\sigma_{nl} y^2 dy}{\exp(y) - 1}, \quad (11)$$

where B_n is the binding energy for n -th excited state and σ_{nl} is the ionization cross section for (n, l) excited state. The asterisk in the upper bound of summation indicates that the sum needs to be regulated, but since this regularization depends only weakly on α and m_e , it can be neglected [54]. The cross section σ_{nl} scales as $\alpha^{-1} m_e^{-2}$ [53]. Altogether,

$$\frac{\partial R(T_M)}{\partial \alpha} = \frac{2}{\alpha} \left(R(T_M) - T_M \frac{\partial R(T_M)}{\partial T_M} \right), \quad (12)$$

$$\frac{\partial R(T_M)}{\partial m_e} = -\frac{1}{m_e} \left(2R(T_M) + T_M \frac{\partial R(T_M)}{\partial T_M} \right). \quad (13)$$

Combining with the fitting formulas (2) and (7), we obtain how R_{H} and R_{HeI} depend on α and m_e .

III. EFFECTS ON THE EPOCH OF RECOMBINATION AND CMB ANGULAR POWER SPECTRUM

We have investigated the equations which describe the process of recombination and how they depend on the coupling constants in the previous section. We incorporate the dependence on α and m_e into the RECFAST code [47] and solve the equations for the ionization fraction x_e as a function of redshift with several different values of α and m_e . The results are shown in Figs. 1 and 2. We have assumed a flat universe and used cosmological parameters $(\omega_b, \omega_m, h) = (0.024, 0.14, 0.72)$, where $\omega_b \equiv \Omega_b h^2$ is the baryon density, $\omega_m \equiv \Omega_m h^2$ is the matter density, h is the Hubble parameter, and Ω denotes the energy density in unit of the critical density. The most important feature is the shift of the epoch of recombination to higher z as α or m_e increases. We can also see this by the rightward shift of the peak of the visibility function shown in Figs. 3 and 4. This is easy to understand because the binding energy B_n scales as $\alpha^2 m_e$ and photons should have higher energy to ionize hydrogens.

In Figs. 5 and 6 we show the power spectrum of the CMB temperature anisotropy for several different values of α and m_e as calculated by the CMBFAST code [55] with the modified RECFAST. We consider a flat Λ CDM universe with a power-law adiabatic primordial fluctuation. The adopted cosmological parameter values are $(\omega_b, \omega_m, h, n_s, \tau) = (0.024, 0.14, 0.72, 0.99, 0.166)$ where τ is the reionization optical depth and n_s is the scalar spectral index. We fix the value of the amplitude of primordial power spectrum. We can see two effects of varying α and m_e in Figs. 5 and 6. Increasing α or m_e shift the peak




FIG. 1 (color online). The ionization fraction x_e as a function of redshift z for no change of α (solid curve), an increase of α by 5% (dashed curve), a decrease of α by 5% (dotted curve).




FIG. 2 (color online). The ionization fraction x_e as a function of redshift z for no change of m_e (solid curve), an increase of m_e by 5% (dashed curve), a decrease of m_e by 5% (dotted curve).

positions to higher values of l and amplify the peak heights.

The peak position shift is understood as follows. Using l_p to denote the position of a peak, $r_\theta(z)$ for the angular diameter distance and $r_s(z)$ for the sound horizon, one can write

$$l_p \sim \frac{r_\theta(z_{ls})}{r_s(z_{ls})}, \quad (14)$$

where z_{ls} is the redshift of the last scattering surface. Increasing α or m_e increases the redshift of the last scattering surface due to the larger binding energy, as in Figs. 3 and 4. The higher z_{ls} in turn corresponds to a smaller sound horizon and a larger angular diameter distance, which lead to a higher value of l_p .




FIG. 3 (color online). The visibility function as a function of conformal time for no change of α (solid curve), an increase of α by 5% (dashed curve), a decrease of α by 5% (dotted curve).




FIG. 4 (color online). The visibility function as a function of conformal time for no change of m_e (solid curve), an increase of m_e by 5% (dashed curve), a decrease of m_e by 5% (dotted curve).

The changes in the peak heights are caused by modifications to the early ISW effect and the diffusion damping. The larger z_{ls} leads to the larger early ISW effect making the first peak higher. To consider the effect beyond the first peak, we focus our attention on the visibility function. The peak of the visibility function moves to a larger redshift since the recombination occurs at higher redshift, when the expansion rate is faster. Hence, the temperature and x_e decreases more rapidly, making the peak width of the visibility function narrower (see Figs. 3 and 4). Since the width of the visibility function corresponds to the damping scale, an increase in α or m_e decreases the effect of damping. This is the reason why the amplitude at larger l increases with increasing α and m_e . Moreover, as seen in Figs. 3 and 4, α changes the visibility function width more than m_e does (quantitatively, an increase of α or m_e by 5%




FIG. 5 (color online). The spectrum of CMB fluctuations for no change of α (solid curve), an increase of α by 5% (dashed curve), a decrease of α by 5% (dotted curve).




FIG. 6 (color online). The spectrum of CMB fluctuations for no change of m_e (solid curve), an increase of m_e by 5% (dashed curve), a decrease of m_e by 5% (dotted curve).

makes the full width at half maximum of the visibility function narrower by 10% or 2%, respectively) because the binding energy which scales as $\alpha^2 m_e$. Thus the damping scale is more sensitive to the change of α than m_e , as appears in Figs. 5 and 6.

Now we discuss the effects of varying α and m_e somewhat more quantitatively using the following four quantities which characterize a shape of CMB power spectrum [56]: the position of the first peak l_1 , the height of the first peak relative to the large angular-scale amplitude evaluated at $l = 10$,

$$H_1 \equiv \left(\frac{\Delta T_{l_1}}{\Delta T_{10}} \right)^2 \quad (15)$$

the ratio of the second peak (l_2) height to the first

$$H_2 \equiv \left(\frac{\Delta T_{l_2}}{\Delta T_{l_1}} \right)^2 \quad (16)$$

the ratio of the third peak (l_3) height to the first

$$H_3 \equiv \left(\frac{\Delta T_{l_3}}{\Delta T_{l_1}} \right)^2 \quad (17)$$

where $(\Delta T_l)^2 \equiv l(l+1)C_l/2\pi$. Note that these four quantities do not depend on overall amplitude. We calculate the response of these four quantities when we vary the parameters ω_b , ω_m , h , τ , n_s , α and m_e . When we vary one parameter, the other parameters are fixed and flatness is always assumed (especially, increasing h means increasing Ω_Λ because $\Omega_\Lambda = 1 - \omega_m/h^2$).

$$\begin{aligned} \Delta l_1 = & 16 \frac{\Delta \omega_b}{\omega_b} - 25 \frac{\Delta \omega_m}{\omega_m} - 47 \frac{\Delta h}{h} + 36 \frac{\Delta n_s}{n_s} \\ & + 290 \frac{\Delta \alpha}{\alpha} + 150 \frac{\Delta m_e}{m_e}, \end{aligned} \quad (18)$$

$$\begin{aligned} \Delta H_1 = & 3.0 \frac{\Delta \omega_b}{\omega_b} - 3.0 \frac{\Delta \omega_m}{\omega_m} - 2.2 \frac{\Delta h}{h} - 1.7 \frac{\Delta \tau}{\tau} \\ & + 18 \frac{\Delta n_s}{n_s} + 3.9 \frac{\Delta \alpha}{\alpha} + 1.4 \frac{\Delta m_e}{m_e}, \end{aligned} \quad (19)$$

$$\begin{aligned} \Delta H_2 = & -0.30 \frac{\Delta \omega_b}{\omega_b} + 0.015 \frac{\Delta \omega_m}{\omega_m} + 0.41 \frac{\Delta n_s}{n_s} \\ & + 0.91 \frac{\Delta \alpha}{\alpha} + 0.30 \frac{\Delta m_e}{m_e}, \end{aligned} \quad (20)$$

$$\begin{aligned} \Delta H_3 = & -0.19 \frac{\Delta \omega_b}{\omega_b} + 0.21 \frac{\Delta \omega_m}{\omega_m} + 0.56 \frac{\Delta n_s}{n_s} \\ & + 0.57 \frac{\Delta \alpha}{\alpha} - 0.019 \frac{\Delta m_e}{m_e}, \end{aligned} \quad (21)$$

and values at the fiducial parameter values are $l_1 = 220$, $H_1 = 6.65$, $H_2 = 0.442$ and $H_3 = 0.449$. Derivatives of l_1 and H_1 with respect to α and m_e are positive as is expected from the considerations above. Furthermore, $\Delta l_1/\Delta \alpha$ and $\Delta l_1/\Delta m_e$ are much larger than the other derivatives of l_1 while $\Delta H_1/\Delta \alpha$ and $\Delta H_1/\Delta m_e$ have relatively similar values to the other derivatives of H_1 . Since such changes are most effectively mimicked by the change in h , it is considered to be the most degenerate parameter with α and m_e . We have seen above that when α or m_e increases, the first peak is enhanced by larger ISW effect and the second or higher peaks are enhanced by smaller diffusion damping. The derivatives of H_2 and H_3 tell us which effect is important. Since $\Delta H_2/\Delta \alpha$ and $\Delta H_3/\Delta \alpha$ are positive and larger than the derivatives with respect to m_e , we see that the effect on the diffusion damping is more significant than that on the early ISW for varying α . They seem to somewhat cancel each other for varying m_e especially regarding H_3 . Such behavior is consistent with the consideration at the end of the previous paragraph, that the diffusion damping is more sensitive to the change in α than m_e .

IV. RELATION BETWEEN VARIATIONS OF α AND m_e

We expect a unified theory can predict the values of the coupling constants, how they are related to each other and how much they vary over a cosmological time scale. In string theory, a candidate for unified theory, there is a dilaton field whose expectation value determines the values of coupling constants. However, since it is not fully formulated at present, we have to assume how α and m_e are related to vary and constrain their variations. To be concrete, following Ref. [35], let us start from considering the low energy action derived from heterotic string theory in the Einstein frame. The action is written as

$$\begin{aligned}
 S = \int d^4x \sqrt{-g} & \left(\frac{1}{2\kappa^2} R - \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \frac{1}{2} D_\mu \phi D^\mu \phi \right. \\
 & - \Omega^{-2} V(\phi) - \bar{\psi} \gamma_\mu D^\mu \psi - \Omega^{-1} m_\psi \bar{\psi} \psi \\
 & \left. - \frac{\alpha'}{16\kappa^2} \Omega^2 F_{\mu\nu} F^{\mu\nu} \right), \quad (22)
 \end{aligned}$$

where Φ is the dilaton field, ϕ is an arbitrary scalar field, and ψ is an arbitrary fermion. D_μ is the gauge covariant derivative corresponding to gauge fields with field strength $F_{\mu\nu}$, $\kappa^2 = 8\pi G$ and $\Omega = e^{-\kappa\Phi/\sqrt{2}}$ is the conformal factor which is used to move from string frame. More concretely, ϕ is the Higgs field and $V(\phi)$ is its potential. The overall factor Ω before the scalar potential means that the Higgs vacuum expectation value $\langle H \rangle$ is independent of the dilaton so it is taken to be constant. $F_{\mu\nu}$ is the gauge field with gauge group including $SU(3) \times SU(2) \times U(1)$. We define its Lagrangian density for the gauge field as $-(1/4g^2)F_{\mu\nu}F^{\mu\nu}$ where g is the unified coupling constant. Compared with equation Eq. (22),

$$\frac{1}{g(M_p)^2} = \frac{\alpha' e^{-\sqrt{2}\kappa\Phi}}{4\kappa^2}, \quad (23)$$

where M_p is the Planck scale. We can calculate the gauge coupling constants at low energy using renormalization group equations. α almost does not run, and hence the α at low energy

$$\alpha \simeq \alpha(M_p) = \frac{g(M_p)^2}{4\pi} = \frac{\kappa^2 e^{\sqrt{2}\kappa\Phi}}{\pi\alpha'}. \quad (24)$$

As for the other gauge coupling constants, variation of the strong coupling constant may affect CMB since its low energy value determines the QCD scale Λ_{QCD} which in turn determines nucleon masses. However, the variation of Λ_{QCD} is related to that of α at low energy can not uniquely be determined from Eq. (22) and especially depends on the details of unification scheme [57]. Therefore, for simplicity, we just assume Λ_{QCD} does not vary. The ψ 's are the ordinary standard model leptons and quarks. As we take $\langle H \rangle = \text{const}$, the Yukawa couplings depend on the dilaton as $e^{\kappa\Phi/\sqrt{2}}$. Therefore the relation between variations of α and m_e is given by

$$\frac{m_e + \Delta m_e}{m_e} = \left(\frac{\alpha + \Delta\alpha}{\alpha} \right)^{1/2}. \quad (25)$$

In this paper, we also consider other possibilities phenomenologically by adopting a power-law relation as

$$\frac{m_e + \Delta m_e}{m_e} = \left(\frac{\alpha + \Delta\alpha}{\alpha} \right)^p, \quad (26)$$

and compute constraints for several values of p . In addition

to the case with changing only α ($p = 0$) and the model described above ($p = 1/2$), we consider cases with $p = 2$ and 4.

V. CONSTRAINTS ON VARYING α AND m_e

We constrain the variation of α in the models described in the previous section using the WMAP first-year data. CMB power spectra are calculated by CMBFAST [55] with RECFAST [47] modified as in Sec. II. The χ^2 is computed for TT and TE data set by the likelihood code supplied by the WMAP team [58–60]. As defined in Sec. III, we consider six cosmological parameters ω_b , ω_m , h , τ , n_s and overall amplitude A in the Λ CDM model assuming the flatness of the universe. We report A in terms of $l(l+1)C_l/2\pi$ at $l = 2$ in unit of μK^2 . In this paper we do not consider gravity waves, running of the spectral index and isocurvature modes. We calculate the minimum χ^2 as a function of α and derive constraints on α . The minimization over six other parameters are performed by iterative applications of the Brent method [61] of the successive parabolic interpolation. More detailed description of this minimization method is found in Ref. [62]. We search for minimum in the region $\tau < 0.3$, which is a prior adopted in Refs. [29,30]. We derive constraints with or without the constraint on the hubble parameter h . When we combine it, we use the Hubble Space Telescope (HST) Hubble Key Project value $h = 0.72 \pm 0.08$ [63] whose error is regarded as gaussian 1σ .

Figure 7 shows our results of χ^2 minimization. It compares varying α only and varying α and m_e with the relation of Eq. (25), respectively, with or without the HST prior. Without the HST prior, we find at 95% C.L. that $-0.107 < \Delta\alpha/\alpha < 0.043$ with changing α only and $-0.097 < \Delta\alpha/\alpha < 0.034$ with the model described in the




FIG. 7 (color online). χ^2 function for variation in α only (solid curves) and α and m_e simultaneously with the relation of Eq. (25) (dashed curves). The HST prior is imposed for the thinner curves.

previous section. Although the best fit α is 4% less than the present value, we find that $\Delta\alpha = 0$ is consistent with the WMAP observation and evidence for varying α is not obtained. The effect of varying m_e simultaneously is found to make the constraint more stringent by 13%. This rather small effect is reasonable since, as is discussed in Sec. III, the effect of m_e on CMB power spectrum is slightly smaller than α , and the relation of Eq. (25) we adopt here does not change m_e much relative to α .

We find that minimum χ^2 is given at $\Delta\alpha/\alpha = -0.04$ with $(\omega_b, \omega_m, h, n_s, \tau, A) = (0.021, 0.132, 0.523, 0.979, 0.146, 942)$ for the case of changing α only, and $\Delta\alpha/\alpha = -0.04$ with $(\omega_b, \omega_m, h, n_s, \tau, A) = (0.020, 0.131, 0.485, 0.979, 0.140, 907)$ for the case of changing α and m_e together. Both cases have notably small values of h . Since h is considered to be the most degenerate parameter with α or m_e as discussed in the end of Sec. III, it is instructive to investigate how constraints tighten when h is limited to higher values such as the HST measurement. From Fig. 7, we obtain, with the HST prior, that $-0.048 < \Delta\alpha/\alpha < 0.032$ with changing α only, and $-0.042 < \Delta\alpha/\alpha < 0.026$ with the model described in the previous section. Compared with no HST prior constraints, they are stringent by about factor of 2 for both cases. Moreover, since low values of h which give good fit with $\Delta\alpha/\alpha \approx -0.04$ are ruled out by the HST prior, the center of allowed region has shifted to larger $\Delta\alpha$.

Here, we comment on the constraint previously obtained by Refs. [29,30] from the WMAP data. As mentioned in Sec. I, they reported the constraint on α to be $-0.06 < \Delta\alpha/\alpha < 0.01$ (95% C.L.). They fixed m_e when varying α and values quoted here is the case with no running for the primordial power spectrum. This constraint seems to have been obtained with marginalization on grid with $0 < \Omega_\Lambda < 0.95$ [29] so it should be compared with our con-




FIG. 8 (color online). χ^2 function for variation in α and m_e with several values of p defined in Eq. (26). The curves are for, from outside to inside, $p = 0, 1/2, 2$ and 4. The HST prior is imposed.

TABLE I. The constraints on α with m_e varies according to various power-law relation Eq. (26) of the index p . The HST prior is imposed.

Power-law index	Constraint (95% C.L.)
$p = 0$	$-0.048 < \Delta\alpha/\alpha < 0.032$
$p = 1/2$	$-0.042 < \Delta\alpha/\alpha < 0.026$
$p = 2$	$-0.031 < \Delta\alpha/\alpha < 0.017$
$p = 4$	$-0.023 < \Delta\alpha/\alpha < 0.011$

straint without the HST prior, $-0.107 < \Delta\alpha/\alpha < 0.043$, which is much weaker than theirs. The difference might be traced to the different analysis method but we could not reproduce their results by our method.

Finally, we investigate the cases in which m_e varies more than α . We consider the models with $p = 2$ and 4 in Eq. (26). We calculate constraints with the HST prior and results are summarized in Fig. 8 and Table I along with $p = 0$ and 1/2 cases. Compared with $p = 0$ (only varying α) case, the constraints become smaller by 40% ($p = 2$) and 60% ($p = 4$). Although those constraints are much smaller than the case with $p = 1/2$, they are still consistent with $\Delta\alpha = 0$.

VI. CONCLUSION

In summary, we have studied a CMB constraint on the time-varying fine structure constant α taking into account simultaneous change of electron mass m_e which might be implied in superstring theories. We have searched sufficiently wide ranges of the cosmological parameters and obtained the WMAP only constraint at 95% C.L. as $-0.097 < \Delta\alpha/\alpha < 0.034$. Combining with the measurement of h by the HST Hubble Key Project, we have obtained more stringent constraint as $-0.042 < \Delta\alpha/\alpha < 0.026$, which improvement is explained by the strong degeneracy between α or m_e and h . These constraints, obtained by adopting the model with Eq. (25) are only slightly tighter than those assuming only α variation: $-0.107 < \Delta\alpha/\alpha < 0.043$ and $-0.048 < \Delta\alpha/\alpha < 0.032$, without and with the HST prior, respectively. This is reasonable since the effect of m_e on CMB power spectrum is similar to that of α (Sec. III) and we adopted to vary m_e milder than α (Eq. (25)).

We have also considered other possibilities for the relation between $\Delta\alpha$ and Δm_e as in the form of Eq. (26) with $p = 2$ and 4. In these cases, the constraints become tighter by roughly a factor of 2. This may not look as drastic as the BBN bounds but CMB bounds are promising and have the advantage that there will be future experiments with higher sensitivities to α as investigated in Refs. [29,30]. In this paper, we do not find evidence of varying α in the CMB data of WMAP. We will wait and see whether future experiments give us more stringent bounds on α or evidence for varying α .

- [1] J. Prestage, R. Tjoelker, and L. Maleki, *Phys. Rev. Lett.* **74**, 3511 (1995).
- [2] S. Bize *et al.*, *Phys. Rev. Lett.* **90**, 150802 (2003).
- [3] H. Marion *et al.*, *Phys. Rev. Lett.* **90**, 150801 (2003).
- [4] M. Fischer *et al.*, *Phys. Rev. Lett.* **92**, 230802 (2004).
- [5] E. Peik *et al.*, *Phys. Rev. Lett.* **93**, 170801 (2004).
- [6] S. Bize *et al.*, physics/0502117.
- [7] E. Peik *et al.*, *Laser Phys.* **15**, 1028 (2005).
- [8] Y. Fujii *et al.*, hep-ph/0205206.
- [9] K. A. Olive, M. Pospelov, Y. Z. Qian, G. Manhes, E. Vangioni-Flam, A. Coc, and M. Casse, *Phys. Rev. D* **69**, 027701 (2004).
- [10] Y. Fujii and A. Iwamoto, *Phys. Rev. Lett.* **91**, 261101 (2003).
- [11] Y. Fujii and A. Iwamoto, *Mod. Phys. Lett. A* **20**, 2417 (2005).
- [12] L. Bergstrom, S. Iguri, and H. Rubinstein, *Phys. Rev. D* **60**, 045005 (1999).
- [13] K. Ichikawa and M. Kawasaki, *Phys. Rev. D* **65**, 123511 (2002).
- [14] K. M. Nollett and R. E. Lopez, *Phys. Rev. D* **66**, 063507 (2002).
- [15] K. Ichikawa and M. Kawasaki, *Phys. Rev. D* **69**, 123506 (2004).
- [16] R. H. Cyburt, B. D. Fields, K. A. Olive, and E. Skillman, *Astropart. Phys.* **23**, 313 (2005).
- [17] J. K. Webb *et al.*, *Phys. Rev. Lett.* **87**, 091301 (2001).
- [18] M. T. Murphy, J. K. Webb, and V. V. Flambaum, *Mon. Not. R. Astron. Soc.* **345**, 609 (2003).
- [19] R. Srianand, H. Chand, P. Petitjean, and B. Aracil, *Phys. Rev. Lett.* **92**, 121302 (2004).
- [20] S. A. Levshakov, M. Centurion, P. Molaro, and S. D'Odorico, *Astron. Astrophys.* **434**, 827 (2005).
- [21] H. Chand, P. Petitjean, R. Srianand, and B. Aracil, *Astron. Astrophys.* **430**, 47 (2005).
- [22] N. Kanekar *et al.*, *Phys. Rev. Lett.* **95**, 261301 (2005).
- [23] S. A. Levshakov, M. Centurion, P. Molaro, S. D'Odorico, D. Reimers, R. Quast, and M. Pollmann, astro-ph/0511765.
- [24] S. Hannestad, *Phys. Rev. D* **60**, 023515 (1999).
- [25] M. Kaplinghat, R. J. Scherrer, and M. S. Turner, *Phys. Rev. D* **60**, 023516 (1999).
- [26] R. A. Battye, R. Crittenden, and J. Weller, *Phys. Rev. D* **63**, 043505 (2001).
- [27] S. Landau, D. Harari, and M. Zaldarriaga, *Phys. Rev. D* **63**, 083505 (2001).
- [28] P. P. Avelino, C. J. A. Martins, G. Rocha, and P. Viana, *Phys. Rev. D* **62**, 123508 (2000).
- [29] C. J. A. Martins, A. Melchiorri, G. Rocha, R. Trotta, P. P. Avelino, and P. Viana, *Phys. Lett. B* **585**, 29 (2004).
- [30] G. Rocha, R. Trotta, C. J. A. Martins, A. Melchiorri, P. P. Avelino, R. Bean, and P. T. P. Viana, *Mon. Not. R. Astron. Soc.* **352**, 20 (2004).
- [31] J. D. Barrow and D. F. Mota, *Classical Quantum Gravity* **20**, 2045 (2003).
- [32] D. F. Mota and J. D. Barrow, *Phys. Lett. B* **581**, 141 (2004).
- [33] D. F. Mota and J. D. Barrow, *Mon. Not. R. Astron. Soc.* **349**, 291 (2004).
- [34] K. Sigurdson, A. Kurylov, and M. Kamionkowski, *Phys. Rev. D* **68**, 103509 (2003).
- [35] B. A. Campbell and K. A. Olive, *Phys. Lett. B* **345**, 429 (1995).
- [36] P. Langacker, G. Segre, and M. J. Strassler, *Phys. Lett. B* **528**, 121 (2002).
- [37] T. Dent and M. Fairbairn, *Nucl. Phys.* **B653**, 256 (2003).
- [38] X. Calmet and H. Fritzsche, *Eur. Phys. J. C* **24**, 639 (2002).
- [39] X. Calmet and H. Fritzsche, *Phys. Lett. B* **540**, 173 (2002).
- [40] K. A. Olive, M. Pospelov, Y. Z. Qian, A. Coc, M. Casse, and E. Vangioni-Flam, *Phys. Rev. D* **66**, 045022 (2002).
- [41] C. M. Muller, G. Schafer, and C. Wetterich, *Phys. Rev. D* **70**, 083504 (2004).
- [42] S. J. Landau, M. E. Mosquera, and H. Vucetich, astro-ph/0411150.
- [43] V. V. Flambaum, D. B. Leinweber, A. W. Thomas, and R. D. Young, *Phys. Rev. D* **69**, 115006 (2004).
- [44] V. V. Flambaum and A. F. Tedesco, *Phys. Rev. C* **73**, 055501 (2006).
- [45] J. Kujat and R. J. Scherrer, *Phys. Rev. D* **62**, 023510 (2000).
- [46] J. J. Yoo and R. J. Scherrer, *Phys. Rev. D* **67**, 043517 (2003).
- [47] S. Seager, D. D. Sasselov, and D. Scott, *Astrophys. J.* **523**, L1 (1999).
- [48] D. Péquignot, P. Petitjean, and C. Boisson, *Astron. Astrophys.* **251**, 680 (1991).
- [49] P. J. E. Peebles, *Principles of Physical Cosmology* (Princeton University Press, New Jersey, 1993).
- [50] D. G. Hummer and D. J. Storey, *Mon. Not. R. Astron. Soc.* **297**, 1073 (1998).
- [51] T. Matsuda, H. Sato, and H. Takeda, *Prog. Theor. Phys.* **42**, 219 (1969).
- [52] G. Breit and E. Teller, *Astrophys. J.* **91**, 215 (1940).
- [53] J. P. Uzan, *Rev. Mod. Phys.* **75**, 403 (2003).
- [54] P. Boschan and P. Biltzinger, *Astron. Astrophys.* **336**, 1 (1998).
- [55] U. Seljak and M. Zaldarriaga, *Astrophys. J.* **469**, 437 (1996).
- [56] W. Hu, M. Fukugita, M. Zaldarriaga, and M. Tegmark, *Astrophys. J.* **549**, 669 (2001).
- [57] M. Dine, Y. Nir, G. Raz, and T. Volansky, *Phys. Rev. D* **67**, 015009 (2003).
- [58] L. Verde *et al.*, *Astrophys. J. Suppl. Ser.* **148**, 195 (2003).
- [59] G. Hinshaw *et al.*, *Astrophys. J. Suppl. Ser.* **148**, 135 (2003).
- [60] A. Kogut *et al.*, *Astrophys. J. Suppl. Ser.* **148**, 161 (2003).
- [61] R. P. Brent, *Algorithms for Minimization Without Derivatives* (Prentice-Hall, Englewood Cliffs, NJ, U.S.A. 1973); see also W. H. Press, B. P. Flannery, S. A. Teukolsky, and W. T. Vetterling, *Numerical Recipes* (Cambridge University Press, New York, 1986).
- [62] K. Ichikawa, M. Fukugita, and M. Kawasaki, *Phys. Rev. D* **71**, 043001 (2005).
- [63] W. L. Freedman *et al.*, *Astrophys. J.* **553**, 47 (2001).