Sign problem via imaginary chemical potential

K. Splittorff

The Niels Bohr Institute, Blegdamsvej 17, DK-2100, Copenhagen Ø, Denmark

B. Svetitsky

School of Physics and Astronomy, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, 69978 Tel Aviv, Israel (Received 19 March 2007; published 28 June 2007)

We calculate an analogue of the average phase factor of the staggered fermion determinant at imaginary chemical potential. Our results from the lattice agree well with the analytical predictions in the microscopic regime for both quenched and phase-quenched QCD. We demonstrate that the average phase factor in the microscopic domain is dominated by the lowest-lying Dirac eigenvalues.

DOI: 10.1103/PhysRevD.75.114504

PACS numbers: 12.38.Gc, 11.30.Rd, 12.39.Fe

I. INTRODUCTION

The numerical study of QCD at nonzero chemical potential μ is hobbled by the complex fermion determinant, which precludes straightforward Monte Carlo calculations based on a real measure. Novel approaches to the problem by means of Taylor expansion [1-4], analytic continuation [5-7], or reweighting [8,9] are consistent in measuring the response to small chemical potential around the temperature of chiral restoration, but they disagree at higher values of μ . Moreover, there will be fundamental technical difficulties at larger μ and larger volumes, and at smaller quark masses and lower temperatures: In all these regimes the fluctuations in the phase of the fermion determinant will become severe [10,11]. Understanding the behavior of the phase of the determinant is thus of crucial importance for the interpretation of lattice data at nonzero chemical potential (see also Ref. [12]).

A common laboratory for studying features of the finitedensity theory is entered by making μ imaginary [13,14]. Here the determinant is real and ordinary Monte Carlo methods can be used. On the face of it, it appears that this would be useless for studying the phase of the determinant since the phase is strictly zero for all configurations. It is perhaps startling, then, to note that one can define and compute an analogue of the average phase factor at imaginary μ [15]. This offers an approach to studying the strength of the sign problem occurring at *real* chemical potential by means of simulations at imaginary μ . The advantages of this method are that one does not have to deal with the sign problem in order to measure its strength, and that one does not have to deal with eigenvalues that have spread out into the complex plane.

The phase factor of the fermion determinant at real chemical potential is given by

$$e^{2i\theta} = \frac{\det(D + \mu\gamma_0 + m)}{\det(D + \mu\gamma_0 + m)^*} = \frac{\det(D + \mu\gamma_0 + m)}{\det(D - \mu\gamma_0 + m)}.$$
 (1)

Here we have used the fact that when μ is real, the complex conjugate determinant is obtained by flipping its sign. The average phase factor at real chemical potential is then

$$\langle e^{2i\theta(\mu)} \rangle = \left\langle \frac{\det(D + \mu \gamma_0 + m)}{\det(D - \mu \gamma_0 + m)} \right\rangle.$$
(2)

As in Ref. [15], we define the average phase factor at imaginary μ by simply substituting $i\mu$ for μ

$$\langle e^{2i\theta(i\mu)}\rangle \equiv \left\langle \frac{\det(D+i\mu\gamma_0+m)}{\det(D-i\mu\gamma_0+m)} \right\rangle,$$
 (3)

where the parameter μ is still real. Both determinants in Eq. (3) are now real. We present in this note a numerical study of the phase factor defined in this manner at imaginary μ . In particular we study the dependence on the chemical potental, the quark mass, and the volume. We work in the microscopic domain of QCD (also called the ϵ regime) [16,17], where the quark mass *m* and the chemical potential are chosen to fulfill

$$|\mu|^2 F_\pi^2 \ll \frac{1}{\sqrt{V}} \quad \text{and} \quad m\Sigma \ll \frac{1}{\sqrt{V}},$$
 (4)

while the Euclidean volume $V = L^3/T$ (where T is the temperature) is taken large

$$V\Lambda_{\rm OCD}^4 \gg 1.$$
 (5)

Here $\Sigma = \langle \bar{\psi}\psi \rangle$ is the chiral condensate and F_{π} is the pion decay constant.¹ Formulas for the average phase factor have recently been derived in this regime [15,18].

The values of Σ and F_{π} on the lattices we use were calculated from a two-point spectral correlation function in Refs. [19,20]. Given these values, we can make parameterfree comparisons of the numerical measurements of the average phase factor with the analytical predictions. In all cases we study the agreement is within the statistical errors. This confirmation of the analytic predictions at imaginary μ gives support to the analytic results derived for real μ where direct lattice tests are harder to obtain.

¹In fact we take $|\mu|^2 F_{\pi}^2 \sim 1/V = T/L^3$, which implies that $\mu/T \ll 1$ when *L* is large. Thus the periodicity in the imaginary chemical potential noted by Roberge and Weiss [13] will never appear in the microscopic domain.

K. SPLITTORFF AND B. SVETITSKY

Our results also show that the average phase factor in the microscopic domain is dominated by the lowest-lying eigenvalues of the Dirac operator.

Below we will recall the theoretical predictions in the microscopic domain and their analytic properties as functions of the chemical potential. Then we will compare these to the lattice data.

II. ANALYTICAL FORMULAS

Our numerical results are based on data obtained in a quenched ensemble and in an ensemble with dynamical fermions given an imaginary isospin chemical potential. Here we present the microscopic formulas obtained for these two cases.

Below we work with standard unimproved staggered fermions at a coupling where the chiral symmetry breaking pattern is identical to that of the continuum theory in the sector of zero topological charge (see e.g. Ref. [21]). We therefore present the analytical predictions in the trivial topological sector.

A. Quenched theory

The average phase factor at real chemical potential in the quenched theory is given by [15,18]

$$\langle e^{2i\theta(\mu)} \rangle_{N_{f}=0} = 1 - 4\hat{\mu}^{2}I_{0}(\hat{m})K_{0}(\hat{m}) - \frac{e^{-2\hat{\mu}^{2}}}{4\hat{\mu}^{2}}e^{-(\hat{m}^{2}/8\hat{\mu}^{2})} \\ \times \int_{\hat{m}}^{\infty} dxxe^{-(x^{2}/8\hat{\mu}^{2})}K_{0}\left(\frac{x\hat{m}}{4\hat{\mu}^{2}}\right) \\ \times \left[I_{0}(x)\hat{m}I_{1}(\hat{m}) - xI_{1}(x)I_{0}(\hat{m})\right],$$
(6)

where we have defined microscopic variables via

$$\hat{m} \equiv m\Sigma V$$
 and $\hat{\mu} \equiv \mu F_{\pi} \sqrt{V}$. (7)

The formula (6) cannot be continued to imaginary μ because of the essential singularity of the last term at $\mu = 0$. The nonanalyticity has its origin in the inverse determinant of the non-Hermitian Dirac operator in Eq. (2)—it is due to the eigenvalues with real part larger than the quark mass [15,22]. For purely imaginary μ the eigenvalues remain on the imaginary axis and are thus always inside the quark mass. The nonanalytic term, therefore, is not expected to appear at imaginary μ . Indeed, a direct calculation at imaginary μ gives [15]

$$\langle e^{2i\theta(i\mu)} \rangle_{N_f=0} = 1 + 4\hat{\mu}^2 I_0(\hat{m}) K_0(\hat{m}).$$
 (8)

As expected the analytic continuation of this result from imaginary values of the chemical potential back to real values gives just the first two terms in Eq. (6). For real μ the eigenvalue density outside the quark mass is highly suppressed if $\mu \leq m_{\pi}/2$. Consequently the first two terms in Eq. (6) are dominant when $\mu \leq m_{\pi}/2$ (see Fig. 1). The measurement of the average phase factor at imaginary



FIG. 1 (color online). The quenched average phase factor as a function of $(2\mu/m_{\pi})^2$ for $\hat{m} = 3$. The average phase factor from $\mu^2 < 0$, Eq. (8), gives the solid line when continued via $i\mu \rightarrow \mu$. This line agrees with the direct result for real μ , Eq. (6), when $\mu \leq m_{\pi}/2$. For larger values of μ the essential singularity in Eq. (6) at $\mu = 0$ becomes important and the analytic continuation fails to reproduce the true result. The same occurs in unquenched QCD.

values of μ is therefore indicative of the strength of the sign problem for real $\mu \leq m_{\pi}/2$.

B. Two dynamical flavors with isospin chemical potential

In this theory, also known as the phase-quenched theory, there are two flavors of fermion with *opposite* values of chemical potential $\pm \mu$. The average phase of the singleflavor determinant is given (for real μ) by

$$\langle e^{2i\theta(\mu)} \rangle_{1+1^*} = \frac{\langle \frac{\det(D+\mu\gamma_0+m)}{\det(D+\mu\gamma_0+m)^*} | \det(D+\mu\gamma_0+m) |^2 \rangle}{\langle |\det(D+\mu\gamma_0+m) |^2 \rangle},$$
(9)

where the averages on the right-hand side are quenched averages. This simplifies to

$$\langle e^{2i\theta(\mu)}\rangle_{1+1^*} = \frac{\langle \det(D+\mu\gamma_0+m)^2\rangle}{\langle |\det(D+\mu\gamma_0+m)|^2\rangle} = \frac{Z_{N_f=2}}{Z_{1+1^*}},$$

a ratio of partition functions in ensembles with dynamical fermions. These can be evaluated exactly in the microscopic regime [17,23] to obtain

$$\langle e^{2i\theta(\mu)} \rangle_{1+1^*} = \frac{I_0^2(\hat{m}) - I_1^2(\hat{m})}{2e^{2\hat{\mu}^2} \int_0^1 dt t e^{-2\hat{\mu}^2 t^2} I_0^2(\hat{m}t)}$$
(10)

for real μ , with scaling variables as defined in Eq. (7) (note that the numerator, which is $Z_{N_f=2}$, does not depend on μ). In this case the average phase factor is analytic at $\mu = 0$,

SIGN PROBLEM VIA IMAGINARY CHEMICAL POTENTIAL

because the inverse determinant in the numerator of Eq. (9) has been canceled. The average phase factor therefore is

$$\langle e^{2i\theta(i\mu)} \rangle_{1+1^*} = \frac{I_0^2(\hat{m}) - I_1^2(\hat{m})}{2e^{-2\hat{\mu}^2} \int_0^1 dt t e^{2\hat{\mu}^2 t^2} I_0^2(\hat{m}t)}$$
(11)

for imaginary chemical potential $i\mu$.

III. NUMERICAL DATA

In the lattice theory, we work with standard, unimproved staggered fermions and introduce the chemical potential using the Hasenfratz-Karsch prescription [24]. As in the continuum, each operator $D \pm i\mu\gamma_0$ is anti-Hermitian, and anticommutes with γ_5 . Each operator's eigenvalues therefore come in pairs of opposite sign on the imaginary axis. For each gauge field configuration we measure the two sets of eigenvalues defined by

$$[D(A) + i\mu\gamma_0]\psi_+^{(j)} = i\lambda_+^{(j)}\psi_+^{(j)}, \qquad (12)$$

$$[D(A) - i\mu\gamma_0]\psi_{-}^{(j)} = i\lambda_{-}^{(j)}\psi_{-}^{(j)}.$$
 (13)

Thus $\det(D \pm i\mu\gamma_0 + m) = \prod_j (i\lambda_{\pm}^{(j)} + m)$. Combining positive with negative λ 's, the average phase factor is given by

$$\langle e^{2i\theta(i\mu)}\rangle = \left\langle \prod_{j=1}^{\infty} \frac{\lambda_{+}^{(j)2} + m^2}{\lambda_{-}^{(j)2} + m^2} \right\rangle \equiv \langle \Phi_{+-}\rangle \qquad (14)$$

$$= \left\langle \prod_{j=1}^{\infty} \frac{\lambda_{-}^{(j)2} + m^2}{\lambda_{+}^{(j)2} + m^2} \right\rangle \equiv \langle \Phi_{-+} \rangle, \qquad (15)$$

where the products run over positive λ 's only. The equality of $\langle \Phi_{+-} \rangle$ and $\langle \Phi_{-+} \rangle$ follows from charge conjugation symmetry, which exchanges λ_+ with λ_- and is true in both the quenched and the phase-quenched theory (but not in the unquenched theory). With finite statistics, the two estimators Φ_{+-} and Φ_{-+} give distinct measurements of the average phase.

A. Quenched theory

Our quenched eigenvalue data are a combination of data used in Ref. [19] with new data. We simulated the SU(3) gauge theory with the plaquette action at $\beta = 5.7$ on lattices with 8⁴ and 12⁴ sites. For each gauge configuration we calculated the smallest 24 pairs of positive eigenvalues $\lambda_{+}^{(j)}$.

On the smaller lattice, we chose two values of the chemical potential: $a\mu = 0.01$ and $a\mu = 0.1$. In Ref. [19] we determined F_{π} and Σ , giving the scaled values $\hat{\mu} = 0.159$ and $\hat{\mu} = 1.59$, respectively; also $\Sigma V/a = 1039$, which permits converting the lattice masses *am* into scaled masses \hat{m} . With these values we



FIG. 2 (color online). The average phase factor calculated via Eq. (14) (circles) and via Eq. (15) (crosses) from the lowest 24 positive eigenvalues of the staggered Dirac operator on an 8⁴ lattice, for fixed chemical potential $a\mu = 0.01$ as a function of the scaled quark mass; 4000 configurations. The full line is the analytical microscopic prediction using the parameters determined in Ref. [19].

can make a parameter-free comparison between the analytical prediction (8) and the measurement of the (truncated) average phase factor. The result for $a\mu = 0.01$, based on data from Ref. [19], is shown in Fig. 2. The agreement is well within the statistical error bars.

The value $a\mu = 0.01$ was chosen in Ref. [19] to ensure that the scaling variable $\hat{\mu}$ was roughly 1/10, since this facilitates the determination of F_{π} . At such small values of μ the shift of the eigenvalues due to the imaginary chemical potential is small compared to the average level spacing.² This was our motivation in running a new quenched simulation in order to calculate eigenvalues with the larger value $a\mu = 0.1$. The result is shown in Fig. 3; again the agreement with Eq. (8) is good. Note that in this case $\mu L = 0.8$ which is pushing the limit of the microscopic domain.

Since *m* is not a dynamical parameter of the quenched simulation, the data points for different values of *m* are highly correlated, and the variation among them is systematic, not statistical. In effect, *m* determines which eigenvalues contribute the most to the average phase in Eqs. (14) and (15). When *m* is small, all eigenvalues contribute to the phase; the smallest eigenvalues cause large fluctuations, while the large eigenvalues show but little dependence on μ and thus contribute little to the ratio. When *m* is large it eliminates the effect of small eigenvalues on the phase.

²For a real μ of this order the width of the strip of eigenvalues in the complex plane is much smaller than the average level spacing along the imaginary axis [23,25,26].



FIG. 3 (color online). As in Fig. 2 but with $a\mu = 0.1$; 6068 configurations.

Therefore as *m* approaches the largest of the calculated eigenvalues there is a systematic deviation from the analytic curve. We illustrate this by cutting the number of calculated eigenvalues to 10. Figure 4 shows the deviation from the analytic curve as $m\Sigma V$ approaches $\lambda^{(10)}\Sigma V \sim 30$.

For the larger lattice, with 12^4 sites, we use data from Ref. [19] with $a\mu = 0.002$. In Ref. [19] we determined that $\hat{\mu} = \mu F_{\pi} \sqrt{V} = 0.159$ and $\Sigma V/a = 5259$. Again we can compare the analytical prediction (8) to the measurement of the truncated average phase factor without free parameters, as shown in Fig. 5. The agreement is clear, although statistically weaker than for the smaller lattice.



FIG. 4 (color online). Same eigenvalue data as in Fig. 2, but retaining only 10 eigenvalues. As the quark mass approaches the largest retained eigenvalue ($\lambda^{(10)}\Sigma V \sim 30$) we observe that the error band drops below the analytic curve (thin line).



FIG. 5 (color online). As in Fig. 2, but for the 12^4 lattice; 2000 configurations.

B. Two dynamical flavors

For the two-flavor theory with imaginary isospin chemical potential we use eigenvalue data calculated in the long runs of Ref. [20], which should be consulted for numerical details. The lattice theory is based on the plaquette action and unimproved staggered fermions, simulated at $\beta = 4.2$ with volume 6⁴. We have data for two masses, am = 0.002and 0.005, and the imaginary chemical potential coupled to isospin is $a\mu = 0.0055$. A determination [20] of aF_{π} gives $\hat{\mu} = 0.1338$.

For the smaller quark mass, am = 0.002, we calculated [20] that $\hat{m} = m\Sigma V = 3.318$. From the lowest 24 eigenvalues we measure

$$\langle \Phi_{+-} \rangle_{1+1^*} = 1.0092 \pm 0.0020,$$
 (16)

$$\langle \Phi_{-+} \rangle_{1+1^*} = 1.0082 \pm 0.0020.$$
 (17)

The analytical prediction for the average phase factor (11) is

$$\langle e^{2i\theta(i\mu)} \rangle_{1+1^*} = 1.0094,$$
 (18)

in agreement with the numerical averages.

For the larger quark mass (ma = 0.005) we have $m\Sigma V = 8.295$. From the lowest 24 eigenvalues we compute

$$\langle \Phi_{+-} \rangle_{1+1^*} = 1.0036 \pm 0.0030$$
 (19)

$$\langle \Phi_{-+} \rangle_{1+1^*} = 1.0032 \pm 0.0030.$$
 (20)

The analytical formula (11) gives

$$\langle e^{2i\theta(i\mu)} \rangle_{1+1^*} = 1.0041.$$
 (21)

Here also we see satisfactory agreement with theory, although the deviation from unity is not really significant.

IV. CONCLUSIONS

We have computed an analogue of the complex phase factor of the fermion determinant at imaginary values of the chemical potential. The results have been used as a parameter-free test of the predictions from the microscopic domain of QCD. In the quenched as well as in the dynamical cases studied the numerical and analytical results are in agreement. Furthermore, we have demonstrated that the phase factor in the microscopic domain is dominated by the low-lying Dirac eigenvalues. We hope that this first numerical study of the phase factor at imaginary chemical potential will encourage studies also outside the microscopic domain.

ACKNOWLEDGMENTS

We wish to thank P. H. Damgaard, P. de Forcrand, and J. J. M. Verbaarschot for discussions. This work was supported by the Israel Science Foundation under Grant No. 173/05 (B.S.) and by the Carlsberg Foundation (K.S.). B.S. thanks the Niels Bohr Institute for its hospitality.

- S. Choe *et al.*, Phys. Rev. D **65**, 054501 (2002); I. Pushkina *et al.* (QCD-TARO Collaboration), Phys. Lett. B **609**, 265 (2005).
- [2] C. R. Allton *et al.*, Phys. Rev. D 66, 074507 (2002); 68, 014507 (2003); 71, 054508 (2005).
- [3] R. V. Gavai and S. Gupta, Phys. Rev. D 68, 034506 (2003);
 71, 114014 (2005).
- [4] C. Bernard et al., Proc. Sci., LAT2006 (2006) 139.
- [5] M. D'Elia and M. P. Lombardo, Phys. Rev. D 67, 014505 (2003); 70, 074509 (2004).
- [6] P. de Forcrand and O. Philipsen, Nucl. Phys. B642, 290 (2002); B673, 170 (2003); J. High Energy Phys. 01 (2007) 077.
- [7] V. Azcoiti, G. Di Carlo, A. Galante, and V. Laliena, J. High Energy Phys. 12 (2004) 010; Nucl. Phys. B723, 77 (2005).
- [8] I. M. Barbour *et al.*, Phys. Rev. D **56**, 7063 (1997), and references therein.
- Z. Fodor and S. D. Katz, Phys. Lett. B 534, 87 (2002);
 J. High Energy Phys. 03 (2002) 014; 04 (2004) 050;
 Z. Fodor, S. D. Katz, and C. Schmidt, J. High Energy Phys. 03 (2007) 121.
- [10] M. Golterman, Y. Shamir, and B. Svetitsky, Phys. Rev. D 74, 071501 (2006); B. Svetitsky, Y. Shamir, and M. Golterman, Proc. Sci., LAT2006 (2006) 148.
- [11] K. Splittorff, Proc. Sci., LAT2006 (2006) 023.

- [12] S. Ejiri, Phys. Rev. D 69, 094506 (2004); 73, 054502 (2006).
- [13] A. Roberge and N. Weiss, Nucl. Phys. B275, 734 (1986).
- [14] M. Alford, A. Kapustin, and F. Wilczek, Phys. Rev. D 59, 054502 (1999).
- [15] K. Splittorff and J.J.M. Verbaarschot, arXiv:hep-lat/ 0702011 [Phys. Rev. D (to be published)].
- [16] J. Gasser and H. Leutwyler, Phys. Lett. B 188, 477 (1987).
- [17] H. Leutwyler and A. Smilga, Phys. Rev. D 46, 5607 (1992).
- [18] K. Splittorff and J. J. M. Verbaarschot, Phys. Rev. Lett. 98, 031601 (2007).
- [19] P.H. Damgaard, U.M. Heller, K. Splittorff, and B. Svetitsky, Phys. Rev. D 72, 091501 (2005).
- [20] P. H. Damgaard, U. M. Heller, K. Splittorff, B. Svetitsky, and D. Toublan, Phys. Rev. D 73, 074023 (2006).
- [21] P.H. Damgaard, U.M. Heller, R. Niclasen, and K. Rummukainen, Phys. Rev. D 61, 014501 (1999).
- [22] K. Splittorff and J. J. M. Verbaarschot, Nucl. Phys. B757, 259 (2006).
- [23] K. Splittorff and J. J. M. Verbaarschot, Nucl. Phys. B683, 467 (2004).
- [24] P. Hasenfratz and F. Karsch, Phys. Lett. B 125, 308 (1983).
- [25] J.C. Osborn, Phys. Rev. Lett. 93, 222001 (2004).
- [26] G. Akemann, J.C. Osborn, K. Splittorff, and J.J.M. Verbaarschot, Nucl. Phys. B712, 287 (2005).