

QCD sum rule analysis of the rare radiative $B_c \rightarrow D_s^* \gamma$ decayK. Azizi^{1,*} and V. Bashiry^{2,†}¹*Physics Department, Middle East Technical University, 06531 Ankara, Turkey*²*Institute for Studies in Theoretical Physics and Mathematics (IPM), P.O. Box 19395-5531, Tehran, Iran*

(Received 20 August 2007; published 10 December 2007)

In this work, the radiative $B_c \rightarrow D_s^* \gamma$ decay is investigated in the framework of QCD sum rules. The transition form factors responsible for the decay are calculated. The total branching ratio for this decay is estimated to be in the order of 10^{-5} , so this decay can be measurable at LHC in the near future.

DOI: [10.1103/PhysRevD.76.114007](https://doi.org/10.1103/PhysRevD.76.114007)

PACS numbers: 13.20.He, 11.55.Hx, 13.40.Hq

I. INTRODUCTION

The heavy pseudoscalar meson B_c contains two heavy quarks of different flavor. This meson has been discovered in 1998 via the decay mode $B_c \rightarrow J/\psi l^\pm \nu$ in 1.8 TeV $p\bar{p}$ collisions, using the CDF detector at the Fermi Lab [1]. The B_c meson constitutes a very rich laboratory for studying various decay channels. There are three classes of decays of B_c meson, namely, b -quark decay (when c is spectator), c -quark decay (when b is spectator), and the weak annihilation channels. Because of two heavy quark contents, the B_c decay channels are expected to be very rich in comparison with other B mesons, so investigation of this meson is essential from both theoretical and experimental points of view. The B_c meson decays provide windows for reliable determination of the Cabibbo-Kobayashi-Maskawa matrix element V_{cb} and can shed light on new physics beyond the standard model.

At LHC with the luminosity values of $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ and $\sqrt{s} = 14 \text{ TeV}$, the number of B_c^\pm events is expected to be about 10^8 – 10^{10} per year, so there is a high probability to study not only some rare B_c decays, but also CP violation, T violation, and polarization asymmetries. Some possible channels are $B_c \rightarrow l\bar{\nu}\gamma$, $B_c \rightarrow \rho^+\gamma$, $B_c \rightarrow K^{*+}\gamma$ and $B_c \rightarrow B_u^* l^+ l^-$, $B_c \rightarrow B_u^* \gamma$, which have been studied in the frame of light-cone QCD and three point QCD sum rules [2–5]. A large set of exclusive non-leptonic and semileptonic decays of the B_c meson, which has been studied within a relativistic constituent quark model, can be found in [6]. Another possible decay channel of B_c is $B_c^- \rightarrow \eta' \ell^- \bar{\nu}$ decay, which is studied both for decay rate and lepton polarization asymmetry [7]. We analyzed the radiative $B_c \rightarrow D_s^* \gamma$ decay by using QCD sum rules method. Using our calculations at the end, we also analyze the $B_c \rightarrow D^* \gamma$ decay by making the necessary changes.

The main quantities in analyzing $B_c \rightarrow D_s^* \gamma$ decay are the form factors. For the calculation of form factors, relevant to this transition, we need some nonperturbative approaches. Among the nonperturbative approaches, the

QCD sum rules method has received special attention, because this approach is based on QCD Lagrangian. This method has been successfully applied to a wide variety of problems (for a review see [2–4,8–13]).

The $B_c \rightarrow D_s^* \gamma$ decay occurs via flavor changing neutral current (FCNC) transition ($b \rightarrow s\gamma$) and weak annihilation channels. The b -quark decay (electromagnetic penguin) for $B_c \rightarrow B_u^* \gamma$ has been calculated in [5] (for more details about the electromagnetic penguin diagram see also [14]). We repeated the similar calculations for our problem and found that the corresponding branching ratio contribution was less important than that of the weak annihilation channel. Note that, the $B_c \rightarrow D_s^* \gamma$ decay has been investigated in the perturbative QCD (PQCD) approach [15], relativistic independent quark model (RIQM) [16], and in the framework of perturbative QCD in standard model [SM (PQCD)], multiscale walking technicolor (MWTM), and topcolor assisted MWTM (TAMWTM) models [17]. They found also that the contribution of the weak annihilation was more important than that of the electromagnetic penguin in PQCD, RIQM, and TAMWTM. In addition, $B_c \rightarrow D^* \gamma$ decay has also been investigated in the relativistic independent quark model (RIQM) [16].

The paper is organized as follows: In Sec. II we construct the transition amplitude for the weak annihilation channel in terms of four relevant form factors, where a photon can be radiated from B_c or D_s^* . Two of the relevant form factors for this decay are calculated in [2] in the framework of the light-cone QCD sum rules. In Sec. III, we calculate the remaining two form factors, when a photon is radiated from the D_s^* meson also in the light-cone QCD. In Sec. IV, we calculate the transition form factors for the electromagnetic penguin in the framework of the three point QCD sum rules method. Finally, in Sec. V numerical analysis, discussion, and comparison of our results to those of the other approaches are forwarded and the conclusion is presented.

II. TRANSITION AMPLITUDE OF THE WEAK ANNIHILATION FOR THE RADIATIVE $B_c \rightarrow D_s^* \gamma$ DECAY

In this section, we concentrated on the main points for obtaining the matrix elements of the radiative decay

*kazizi@newton.physics.metu.edu.tr

†bashiry@ipm.ir

$B_c \rightarrow D_s^* \gamma$ along the lines similar to [18]. The weak annihilation mechanism for this decay is shown in Fig. 1. The transition amplitude for this decay can be written as

$$M(B_c \rightarrow D_s^* \gamma) = \frac{G_F}{\sqrt{2}} V_{cb} V_{cs}^* \langle D_s^*(p) \gamma(q) | (\bar{s} \Gamma_\nu c) (\bar{c} \Gamma^\nu b) | B_c(p+q) \rangle, \quad (1)$$

where $\Gamma_\nu = \gamma_\nu(1 - \gamma_5)$ and p , q , and $p+q$ are the momentum of D_s^* , photon, and B_c , respectively. Using factorization hypothesis, the matrix element in Eq. (1) can be written in the following form:

$$\begin{aligned} & \langle D_s^*(p) \gamma(q) | (\bar{s} \Gamma_\nu c) (\bar{c} \Gamma^\nu b) | B_c(p+q) \rangle \\ &= -e \varepsilon^\mu \varepsilon^{(D_s^*)\nu} f_{D_s^*} m_{D_s^*} T_{\mu\nu}^{(B_c)} - ie \varepsilon^\mu (p+q)^\nu f_{B_c} T_{\mu\nu}^{(D_s^*)}, \end{aligned} \quad (2)$$

where the covariant decomposition of hadronic matrix elements $T_{\mu\nu}^{(B_c)}$ and $T_{\mu\nu}^{(D_s^*)}$ are responsible for the emission of photon from initial and final states, $f_{D_s^*}$, f_{B_c} are the leptonic decay constants of D_s^* and B_c mesons, respectively, and ε^μ and $\varepsilon^{(D_s^*)\nu}$ are the polarization vectors of a photon and D_s^* meson. The covariant decompositions of hadronic matrix elements are defined by the following two-point correlation functions:

$$T_{\mu\nu}^{(B_c)}(p, q) = i \int d^4x e^{iqx} \langle 0 | T \{ j_\mu^{\text{em}} \bar{c} \Gamma_\nu b(0) \} | B_c(p+q) \rangle \quad (3)$$

and

$$T_{\mu\nu}^{(D_s^*)}(p, q) = i \int d^4x e^{iqx} \langle D_s^*(p) | T \{ j_\mu^{\text{em}} \bar{s} \Gamma_\nu c(0) \} | 0 \rangle, \quad (4)$$

where j^{em} stands for electromagnetic current. Our aim is to construct the $T_{\mu\nu}^{(B_c)}$ and $T_{\mu\nu}^{(D_s^*)}$ in terms of form factors and other physical quantities. Let us first focus on $T_{\mu\nu}^{(B_c)}$. This quantity can be written in terms of two independent 4-momenta p and q in general as follows:

$$\begin{aligned} T_{\mu\nu}^{(B_c)}(p, q) &= g_{\mu\nu} a + p_\mu q_\nu b + q_\mu p_\nu c + p_\mu p_\nu d \\ &+ q_\mu q_\nu e + \varepsilon_{\nu\mu\lambda\sigma} p^\lambda q^\sigma F_V^{(B_c)}, \end{aligned} \quad (5)$$

where a , b , c , d , e , and $F_V^{(B_c)}$ are invariant amplitudes. Applying the Ward identity for electromagnetic current to Eq. (5) and using the fact that for a real photon $q^2 = 0$, we rewrite Eq. (5) in the following form:

$$\begin{aligned} T_{\mu\nu}^{(B_c)}(p, q) &= (g_{\mu\nu}(p \cdot q) - p_\mu q_\nu) i F_A^{(B_c)} + g_{\mu\nu}(p \cdot q) \alpha \\ &+ p_\mu q_\nu \beta + q_\mu q_\nu c + i \frac{p_\mu p_\nu}{p \cdot q} f_{B_c} + q_\mu q_\nu e \\ &+ \varepsilon_{\nu\mu\lambda\sigma} p^\lambda q^\sigma F_V^{(B_c)}, \end{aligned} \quad (6)$$

where $F_A^{(B_c)}$, α , and β are the new invariant amplitudes. To obtain the relation between α and β , we compare Eqs. (5) and (6), which leads to

$$\begin{aligned} g_{\mu\nu} a + p_\mu q_\nu b &= (g_{\mu\nu}(p \cdot q) - p_\mu q_\nu) i F_A^{(B_c)} \\ &+ g_{\mu\nu}(p \cdot q) \alpha + p_\mu q_\nu \beta \end{aligned} \quad (7)$$

by multiplying both sides of Eq. (7) with q^μ and using $a + (p \cdot q)b = i f_{B_c}$, we get the following relation between α and β , which is called the Ward identity:

$$\alpha + \beta = \frac{i f_{B_c}}{(p \cdot q)}. \quad (8)$$

From Eq. (8), it is clear that α and β can take different choices. Within the scope of the present work, in parallel with [18], we set $\beta = 0$ and $\alpha = \frac{i f_{B_c}}{(p \cdot q)}$. Substituting the values of α and β , we obtain

$$\begin{aligned} T_{\mu\nu}^{(B_c)}(p, q) &= (g_{\mu\nu}(p \cdot q) - p_\mu q_\nu) i F_A^{(B_c)} + i f_{B_c} g_{\mu\nu} \\ &+ q_\mu q_\nu c + i \frac{p_\mu p_\nu}{p \cdot q} f_{B_c} + q_\mu q_\nu e \\ &+ \varepsilon_{\nu\mu\lambda\sigma} p^\lambda q^\sigma F_V^{(B_c)}. \end{aligned} \quad (9)$$


Using $\varepsilon \cdot q = 0$, $\varepsilon^{(D_s^*)} \cdot p = 0$ and Eq. (9), we obtain the following expression for the first term in Eq. (2) in terms of two form factors ($F_A^{(B_c)}$, $F_V^{(B_c)}$):

$$\begin{aligned} e \varepsilon^\mu \varepsilon^{(D_s^*)\nu} f_{D_s^*} m_{D_s^*} T_{\mu\nu}^{(B_c)} &= e f_{D_s^*} m_{D_s^*} \{ [(\varepsilon \cdot \varepsilon^{(D_s^*)})(p \cdot q) \\ &- (\varepsilon \cdot p)(\varepsilon^{(D_s^*)} \cdot q)] i F_A^{(B_c)} \\ &+ i f_{B_c} (\varepsilon \cdot \varepsilon^{(D_s^*)}) \\ &+ \varepsilon_{\nu\mu\lambda\sigma} \varepsilon^{(D_s^*)\nu} \varepsilon^\mu p^\lambda q^\sigma F_V^{(B_c)} \}. \end{aligned} \quad (10)$$

Omitting the details for the calculation $T_{\mu\nu}^{(D_s^*)}$, we get the following result for the second term in Eq. (2):

$$\begin{aligned} ie \varepsilon^\mu (p+q)^\nu f_{B_c} T_{\mu\nu}^{(D_s^*)} &= i e f_{B_c} \{ [(\varepsilon \cdot \varepsilon^{(D_s^*)})(p \cdot q) \\ &- (\varepsilon \cdot p)(\varepsilon^{(D_s^*)} \cdot q)] i F_A^{(D_s^*)} \\ &+ f_{D_s^*} m_{D_s^*} (\varepsilon \cdot \varepsilon^{(D_s^*)}) \\ &+ \varepsilon_{\nu\mu\lambda\sigma} \varepsilon^{(D_s^*)\nu} \varepsilon^\mu p^\lambda q^\sigma F_V^{(D_s^*)} \}, \end{aligned} \quad (11)$$

where $F_A^{(D_s^*)}$ and $F_V^{(D_s^*)}$ are two form factors of D_s^* . Now, we can write the transition amplitude for the radiative $B_c \rightarrow D_s^* \gamma$ decay in terms of four form factors $F_A^{(B_c)}$, $F_V^{(B_c)}$, $F_A^{(D_s^*)}$, and $F_V^{(D_s^*)}$ as follows:


 FIG. 1. The weak annihilation mechanism for $B_c \rightarrow D_s^* \gamma$.

$$\begin{aligned}
 M(B_c \rightarrow D_s^* \gamma) = & e \frac{G_F}{\sqrt{2}} V_{cb} V_{cs}^* [-f_{D_s^*} m_{D_s^*} \{[(\varepsilon \cdot \varepsilon^{(D_s^*)})(p \cdot q) - (\varepsilon \cdot p)(\varepsilon^{(D_s^*)} \cdot q)] i F_A^{(B_c)} \\
 & + i f_{B_c} (\varepsilon \cdot \varepsilon^{(D_s^*)}) + \varepsilon_{\nu\mu\lambda\sigma} \varepsilon^{(D_s^*)\nu} \varepsilon^\mu p^\lambda q^\sigma F_V^{(B_c)}\} - i f_{B_c} \{[(\varepsilon \cdot \varepsilon^{(D_s^*)})(p \cdot q) - (\varepsilon \cdot p)(\varepsilon^{(D_s^*)} \cdot q)] i F_A^{(D_s^*)} \\
 & + f_{D_s^*} m_{D_s^*} (\varepsilon \cdot \varepsilon^{(D_s^*)}) + \varepsilon_{\nu\mu\lambda\sigma} \varepsilon^{(D_s^*)\nu} \varepsilon^\mu p^\lambda q^\sigma F_V^{(D_s^*)}\}]. \quad (12)
 \end{aligned}$$

The form factors $F_A^{(B_c)}$ and $F_V^{(B_c)}$ corresponding to the emission of the photon from b - and c -quark [see Figs. 1(i) and 1(ii)], are calculated in [2]. Therefore, we will concentrate on the calculation of the form factors $F_A^{(D_s^*)}$ and $F_V^{(D_s^*)}$ [see Figs. 1(iii) and 1(iv)].

III. LIGHT-CONE QCD SUM RULES FOR THE FORM FACTORS $F_A^{(D_s^*)}$ AND $F_V^{(D_s^*)}$

Based on the general idea of the QCD sum rules method, we will calculate the transition form factors by equating the representation of a suitable correlator in hadronic and quark-gluon languages. For this aim, we consider the following correlation function:

$$\begin{aligned}
 \Pi_{\mu\nu}(p, q) = & i \int d^4x e^{iqx} \langle \gamma(q) | T \{ \bar{c}(x) \gamma_\mu (1 - \gamma_5) s(x) \bar{s}(0) \\
 & \times \gamma_\nu c(0) \} | 0 \rangle. \quad (13)
 \end{aligned}$$

Here, q and p are the momentum values of photon and D_s^* , respectively, and $Q = p + q$ is the transferred momentum. Now, we insert the hadronic state $D_s^*(p)$ to Eq. (13). This can be rewritten as

$$\Pi_{\mu\nu}(p, q) = \frac{\langle \gamma(q) | \bar{c} \gamma_\mu (1 - \gamma_5) s | D_s^*(p) \rangle \langle D_s^*(p) | \bar{s} \gamma_\nu c | 0 \rangle}{m_{D_s^*}^2 - p^2}. \quad (14)$$

The second term in Eq. (14) by definition is

$$\langle D_s^*(p) | \bar{s} \gamma_\nu c | 0 \rangle = f_{D_s^*} m_{D_s^*} \varepsilon_\nu^{(D_s^*)}. \quad (15)$$

From Figs. 1(iii) and 1(iv) and considering the parity, Lorentz and gauge invariances, we can write the matrix element for the emission of the photon from D_s^* meson as

$$\langle \gamma(q) | \bar{c} \gamma_\mu (1 - \gamma_5) s | D_s^*(p) \rangle = e \left\{ i \varepsilon_{\mu\alpha\beta\sigma} \varepsilon^\alpha \varepsilon^{(D_s^*)\beta} q^\sigma \frac{F_V^{(D_s^*)}(Q^2)}{m_{D_s^*}^2} + [\varepsilon_\mu (\varepsilon^{(D_s^*)} \cdot q) - (\varepsilon \cdot \varepsilon^{(D_s^*)}) q_\mu] \frac{F_A^{(D_s^*)}(Q^2)}{m_{D_s^*}^2} \right\}. \quad (16)$$

Substituting Eqs. (15) and (16) to (14), we have

$$\Pi_{\mu\nu}(p, q) = \frac{ef_{D_s^*}m_{D_s^*}}{m_{D_s^*}^2 - p^2} \left\{ i\varepsilon_{\mu\alpha\beta\sigma}\varepsilon^\alpha\varepsilon_\nu^{(D_s^*)}\varepsilon^{(D_s^*)\beta}q^\sigma \frac{F_V^{(D_s^*)}(Q^2)}{m_{D_s^*}^2} + [\varepsilon_\mu\varepsilon_\nu^{(D_s^*)}(\varepsilon^{(D_s^*)} \cdot q) - (\varepsilon \cdot \varepsilon^{(D_s^*)})\varepsilon_\nu^{(D_s^*)}q_\mu] \frac{F_A^{(D_s^*)}(Q^2)}{m_{D_s^*}^2} \right\}. \quad (17)$$

The summation over polarization of the D_s^* meson is performed by using

$$\varepsilon_\nu^{(D_s^*)}\varepsilon_\beta^{(D_s^*)} = -g_{\nu\beta} + \frac{P_\nu P_\beta}{m_{D_s^*}^2}. \quad (18)$$

After performing the standard calculations for the phenomenological part, we get

$$\Pi_{\mu\nu}(p, q) = \frac{ef_{D_s^*}m_{D_s^*}}{m_{D_s^*}^2 - p^2} \left\{ i\varepsilon_{\mu\nu\alpha\sigma}\varepsilon^\alpha q^\sigma \frac{F_V^{(D_s^*)}(Q^2)}{m_{D_s^*}^2} + [q_\mu\varepsilon_\nu - \varepsilon_\mu q_\nu] \frac{F_A^{(D_s^*)}(Q^2)}{m_{D_s^*}^2} \right\}. \quad (19)$$

The theoretical part (QCD side) of the correlator is calculated by means of operator product expansion (OPE) up to operators having dimension $d = 5$ in deep Euclidean space, where both p^2 and Q^2 are large and negative. It is determined by the bare-loop [Figs. 2(a) and 2(b)] and the power corrections from the operators

with $d = 3$, $\langle\bar{\psi}\psi\rangle$, $d = 4$, $m_s\langle\bar{\psi}\psi\rangle$, $d = 5$, $m_0^2\langle\bar{\psi}\psi\rangle$ [Figs. 2(c)–2(e)] and the photon interaction with a soft quark line [Fig. 2(f)]. In calculating the bare-loop and nonperturbative correction contributions, we first write the Lorentz decomposition of the correlator as

$$\Pi_{\mu\nu}(p, q) = i\varepsilon_{\mu\nu\alpha\sigma}\varepsilon^\alpha q^\sigma \Pi_1 + [q_\mu\varepsilon_\nu - \varepsilon_\mu q_\nu] \Pi_2 \quad (20)$$

and the dispersion representation (Cutkosky method) for the coefficients of corresponding Lorentz structures appearing in the $\Pi_{\mu\nu}(p, q)$ as follows:

$$\Pi_{1,2}(p, q) = \int ds \frac{\rho_{1,2}(s, p^2)}{s - Q^2} + \text{subtraction terms}, \quad (21)$$

where $\rho_{1,2}(s, p^2)$ are spectral density corresponding to two structures in $\Pi_{\mu\nu}(p, q)$ and subtraction terms stand for corrections. To calculate $\rho_{1,2}(s, p^2)$, we consider diagrams in Figs. 2(a) and 2(b). For instance, for the contribution of diagram (a) we get




FIG. 2. diagrams for bare-loop [(a), (b)] power corrections from the operators with 3, 4, and 5 dimensions [(c), (d), (e)] and propagation of the soft quark in electromagnetic field (f).

$$T_{\mu\nu} = eN_c Q_s \int \frac{d^4k}{(2\pi)^4} \left\{ \text{Tr} \left[\frac{i(\not{k} + m_c)}{k^2 - m_c^2} \gamma_\mu (1 - \gamma_5) \frac{i(\not{Q} + \not{k} + m_s)}{(Q+k)^2 - m_s^2} \not{\epsilon} \frac{i(\not{p} + \not{k} + m_s)}{(p+k)^2 - m_s^2} \gamma_\nu \right] \right\}. \quad (22)$$

With the help of the above equations, we obtain the following expressions corresponding to the coefficients of the structures $i\varepsilon_{\mu\nu\alpha\sigma}\varepsilon^\alpha q^\sigma$ and $[q_\mu\varepsilon_\nu - \varepsilon_\mu q_\nu]$:

$$T_1 = \frac{-eN_c Q_s}{4\pi^2} \left\{ \int_0^1 dx x \int_0^1 dy [m_c m_s - m_s^2 x \bar{y} + p^2(X_1 \bar{y} + X_3) - (p \cdot q)(X_2 \bar{y}^2 - 4x^2 \bar{x}y + 2x^2 \bar{x}y^2 - 4x \bar{x}y)] \int_0^\infty d\alpha e^{-\alpha\Delta} \right\},$$

$$T_2 = \frac{eN_c Q_s}{16\pi^2} \left\{ \int_0^1 dx x \int_0^1 dy [m_c m_s - m_s^2 x \bar{y} + p^2(X_1 \bar{y} - X_3) - (p \cdot q)X_3(2\bar{y} + 4\bar{x}y)] \int_0^\infty d\alpha e^{-\alpha\Delta} \right\}, \quad (23)$$

where $\bar{x}(\bar{y}) = 1 - x(y)$, $\Delta = -p^2 x \bar{x} y - Q^2 x \bar{x} \bar{y} + m_c^2 \bar{x} + m_s^2 x$, $X_1 = x - 2x^2 + x^3$, $X_2 = x^2 - 2x^3$, and $X_3 = -x + x^2$. In this calculation, we have also used the exponential representation for the denominator as

$$\frac{1}{\Delta^n} = \frac{1}{(n-1)!} \int_0^\infty d\alpha \alpha^{n-1} e^{-\alpha\Delta}. \quad (24)$$

Next, we apply the double Borel operator $\hat{B}(M_1^2)\hat{B}(M_2^2)$ on $T_{(1,2)}$ and we get

$$\tilde{T}_1 = \frac{-eN_c Q_s}{4\pi^2} \frac{\sigma_1 \sigma_2}{\sigma_1 + \sigma_2} \int_0^1 dx \frac{1}{\bar{x}} e^{(1/x\bar{x})(m_c^2 \bar{x} + m_s^2 x)(\sigma_1 + \sigma_2)}$$

$$\times \left[m_c m_s - m_s^2 x \frac{\sigma_2}{\sigma_1 + \sigma_2} + p^2 \left((x^3 - 2x^2) \frac{\sigma_2}{\sigma_1 + \sigma_2} + x^2 - x \frac{\sigma_1}{\sigma_1 + \sigma_2} \right) \right.$$

$$\left. - (p \cdot q) \left(x^2 - 2x^3 + (3x^2 - 4x^3) \frac{\sigma_1^2}{(\sigma_1 + \sigma_2)^2} + (8x^3 - 2x^2 - 4x) \frac{\sigma_1}{\sigma_1 + \sigma_2} \right) \right],$$

$$\tilde{T}_2 = \frac{eN_c Q_s}{16\pi^2} \frac{\sigma_1 \sigma_2}{\sigma_1 + \sigma_2} \int_0^1 dx \frac{1}{\bar{x}} e^{(1/x\bar{x})(m_c^2 \bar{x} + m_s^2 x)(\sigma_1 + \sigma_2)} \quad (25)$$

$$\times \left[m_c m_s - m_s^2 x \frac{\sigma_2}{\sigma_1 + \sigma_2} + p^2 \left((x + x^3 - 2x^2) \frac{\sigma_2}{\sigma_1 + \sigma_2} - x^2 + x \right) - (p \cdot q) (-2x + 2x^2) \frac{\sigma_2}{\sigma_1 + \sigma_2} \right.$$

$$\left. - 2x^3 \frac{\sigma_2^2}{(\sigma_1 + \sigma_2)^2} + (8x^2 - 4x^3 - 4x) \frac{\sigma_1}{\sigma_1 + \sigma_2} + (2x^3 - 2x^2) \frac{\sigma_1^2}{(\sigma_1 + \sigma_2)^2} \right],$$

where $\sigma_1 = \frac{1}{M_1^2}$ and $\sigma_2 = \frac{1}{M_2^2}$.

In deriving Eq. (25), we use the definition

$$\hat{B}(M^2)e^{-\alpha p^2} = \delta(1 - \alpha M^2). \quad (26)$$

For the determination of the spectral density, we apply the Borel transformations to \tilde{T}_1 and \tilde{T}_2 [19] and we obtain

$$\varrho_{1,2}(s, t, p^2) = \frac{1}{st} \hat{B}\left(\frac{1}{s}, \sigma_1\right) \hat{B}\left(\frac{1}{t}, \sigma_2\right) \frac{\tilde{T}_{1,2}}{\sigma_1 \sigma_2}. \quad (27)$$

In this step, we use the following relations:

$$\sigma^n e^{-\alpha\sigma} = \left(-\frac{d}{d\alpha}\right)^n e^{-\alpha\sigma}, \quad (28)$$

$$\rho_{1,2}(s, p^2) = \int dt \frac{\varrho_{1,2}(s, t, p^2)}{t - p^2}, \quad (29)$$

and

$$\hat{B}\left(\frac{1}{s}, \sigma_1\right) \hat{B}\left(\frac{1}{t}, \sigma_2\right) e^{-\alpha(\sigma_1 + \sigma_2)} = \delta\left(1 - \frac{\alpha}{s}\right) \delta\left(1 - \frac{\alpha}{t}\right). \quad (30)$$

Then, we get the following expressions for the two spectral densities, as follows:

$$\begin{aligned}
\rho_1(s, p^2) &= \frac{eN_c Q_s}{4\pi^2} \frac{1}{(s-p^2)^3} \int_{x_0}^{x_1} dx \frac{1}{\bar{x}} \left\{ (m_c m_s - m_s^2 x)(s-p^2)^2 + p^2 \left[(x^3 - x^2)(s-p^2)^2 - x \left(p^2 + \frac{m_c^2}{x} + \frac{m_s^2}{\bar{x}} \right) (s-p^2) \right] \right. \\
&\quad - \frac{1}{2} (m_s^2 - p^2) \left[(x^2 - 2x^3)(s-p^2)^2 - 2(3x^2 - 4x^3) \left(p^2 + \frac{m_c^2}{x} + \frac{m_s^2}{\bar{x}} \right)^2 \right. \\
&\quad \left. \left. + (8x^3 - 2x^2 - 4x) \left(p^2 + \frac{m_c^2}{x} + \frac{m_s^2}{\bar{x}} \right) (s-p^2) \right] \right\}, \\
\rho_2(s, p^2) &= \frac{eN_c Q_s}{16\pi^2} \frac{1}{(s-p^2)^3} \int_{x_0}^{x_1} dx \frac{1}{\bar{x}} \left\{ [m_c m_s - m_s^2 x + p^2(2x - 3x^2 + x^3)](s-p^2)^2 \right. \\
&\quad - \frac{1}{2} (m_s^2 - p^2) \left[(-2x + 2x^2 - 4x^3)(s-p^2)^2 + (8x^2 - 4x^3 - 4x) \left(p^2 + \frac{m_c^2}{x} + \frac{m_s^2}{\bar{x}} \right) (s-p^2) \right. \\
&\quad \left. \left. - 2(2x^3 - 2x^2) \left(p^2 + \frac{m_c^2}{x} + \frac{m_s^2}{\bar{x}} \right)^2 \right] \right\}, \tag{31}
\end{aligned}$$

where the integration region is determined by the following inequality:

$$sx\bar{x} - (m_c^2 \bar{x} + m_s^2 x) \geq 0. \tag{32}$$

Similar to the above calculations for diagram (a), we have repeated the entire calculations for diagram (b). Finally, we get the following results for the spectral densities:

$$\begin{aligned}
\rho_1(s, p^2) &= \frac{eN_c}{48\pi^2} \frac{1}{(s-p^2)^3} \left\{ Q_s \left[\lambda \{ 3(-3 + 5\alpha - 5\beta)p^6 + (28 + \alpha(-103 + \alpha(17 + 4\alpha)) + 52\beta - \alpha(61 + 8\alpha))\beta \right. \right. \\
&\quad + 4(10 + \alpha)\beta^2 p^4 s - 3(-2 + 16\alpha^3 + \beta - 31\beta^2 - \alpha^2(75 + 32\beta) + \alpha(39 + 2\beta(41 + 8\beta)) + 7\lambda^2)p^2 s^2 \\
&\quad + \alpha(1 - 86\alpha^2 + \beta - 86\beta^2 + \alpha(67 + 172\beta) + 18\lambda^2)s^3 \} + 6\{p^6 + 2m_c m_s (s-p^2)^2 \\
&\quad + (2 - 11\alpha + 4\beta)p^4 s(-1 + 26\alpha^2 + 2\beta^2 - 2\alpha(7 + 11\beta))p^2 s^2 - \alpha(1 + 18\alpha^2 + 2\beta(1 + \beta) \\
&\quad - 4\alpha(4 + 5\beta))s^3 \} \ln \frac{1 + \alpha - \beta - \lambda}{1 + \alpha - \beta + \lambda} \left. \right] + Q_c \left[\lambda \{ 3(-3 + 5\beta - 5\alpha)p^6 + (28 + \beta(-103 + \beta(17 + 4\beta)) \right. \\
&\quad + 52\alpha - \beta(61 + 8\beta))\alpha + 4(10 + \beta)\alpha^2 p^4 s - 3(-2 + 16\beta^3 + \alpha - 31\alpha^2 - \beta^2(75 + 32\alpha) \\
&\quad + \beta(39 + 2\alpha(41 + 8\alpha)) + 7\lambda^2)p^2 s^2 + \beta(1 - 86\beta^2 + \alpha - 86\alpha^2 + \beta(67 + 172\alpha) + 18\lambda^2)s^3 \} \\
&\quad + 6\{p^6 + 2m_c m_s (s-p^2)^2 + (2 - 11\beta + 4\alpha)p^4 s(-1 + 26\beta^2 + 2\alpha^2 - 2\beta(7 + 11\alpha))p^2 s^2 \\
&\quad - \beta(1 + 18\beta^2 + 2\alpha(1 + \alpha) - 4\beta(4 + 5\alpha))s^3 \} \ln \frac{1 + \beta - \alpha - \lambda}{1 + \beta - \alpha + \lambda} \left. \right] \right\}, \\
\rho_2(s, p^2) &= \frac{eN_c}{96\pi^2} \frac{1}{(s-p^2)^3} \left\{ Q_s \left[\lambda \left\{ -(29 + 2\alpha^2 + \beta(17 + 2\beta) - \alpha(19 + 4\beta))p^6 + (4\alpha^3 + \alpha(5 + \beta)(5 + 4\beta) \right. \right. \right. \\
&\quad - \alpha^2(23 + 8\beta) + 6(8 + \beta - \beta^2))p^4 s + \left(-23 + 2\alpha^3 - 5\beta(1 + 4\beta) + \alpha(-9 + 2\beta(21 + \beta)) \right. \\
&\quad \left. \left. + \alpha^2 \left(-30 - 4\beta + \frac{48}{\lambda^2 - (1 + \alpha - \beta)^2} \right) \right) p^2 s^2 + \alpha(13 + \alpha(-5 + 22\alpha) + \beta - 44\alpha\beta + 22\beta^2)s^3 \right\} \\
&\quad + 6\{-2p^6 + m_c m_s (s-p^2)^2 + (4 + 5\alpha)p^4 s - 2(1 + \alpha + 4\alpha^2 - 2\alpha\beta)p^2 s^2 + \alpha(1 + 4\alpha(\alpha - \beta))s^3 \} \\
&\quad \times \ln \frac{1 - \alpha + \beta - \lambda}{1 - \alpha + \beta + \lambda} \left. \right] + Q_c \left[\lambda \left\{ -(29 + 2\beta^2 + \alpha(17 + 2\alpha) - \beta(19 + 4\alpha))p^6 + (4\beta^3 + \beta(5 + \alpha)(5 + 4\alpha) \right. \right. \\
&\quad - \beta^2(23 + 8\alpha) + 6(8 + \alpha - \alpha^2))p^4 s + \left(-23 + 2\beta^3 - 5\alpha(1 + 4\alpha) + \beta(-9 + 2\alpha(21 + \alpha)) \right. \\
&\quad \left. \left. + \beta^2 \left(-30 - 4\alpha + \frac{48}{\lambda^2 - (1 + \beta - \alpha)^2} \right) \right) p^2 s^2 + \beta(13 + \beta(-5 + 22\beta) + \alpha - 44\beta\alpha + 22\alpha^2)s^3 \right\} \\
&\quad + 6\{-2p^6 + m_c m_s (s-p^2)^2 + (4 + 5\beta)p^4 s - 2(1 + \beta + 4\beta^2 - 2\beta\alpha)p^2 s^2 + \beta(1 + 4\beta(\beta - \alpha))s^3 \} \\
&\quad \times \ln \frac{1 - \beta + \alpha - \lambda}{1 - \beta + \alpha + \lambda} \left. \right] \right\}. \tag{33}
\end{aligned}$$

where $\alpha = \frac{m_s^2}{s}$ and $\lambda = \sqrt{1 + \alpha^2 + \beta^2 - 2\alpha - 2\beta - 2\alpha\beta}$.

The next step is to calculate contributions coming from the power correction terms. After standard but lengthy calculations for the contributions of the diagrams (c), (d), (e), we get

$$\begin{aligned}\Pi_1(p, q)^{(c,d,e)} &= \frac{m_c}{r'r} \langle \bar{s}s \rangle - \frac{m_s}{2} \langle \bar{s}s \rangle \left[\frac{m_c^2}{r'^2 r} + \frac{1}{r'r} + \frac{m_c^2}{r' r^2} \right] + \frac{m_s^2}{2} \langle \bar{s}s \rangle \left[\frac{2m_c^3}{r'^3 r} + \frac{2m_c^3}{r'^2 r^2} + \frac{m_c}{r'^2 r} + \frac{m_c}{r' r^2} + \frac{2m_c^3}{r' r^3} \right] \\ &\quad - \frac{m_0^2}{4} \langle \bar{s}s \rangle \left[\frac{2m_c^3}{r'^3 r} + \frac{2m_c^3}{r'^2 r^2} + \frac{2m_c}{3r'^2 r} + \frac{m_c}{r' r^2} + \frac{2m_c^3}{r' r^3} \right], \\ \Pi_2(p, q)^{(c,d,e)} &= \frac{-m_c}{r'r} \langle \bar{s}s \rangle + \frac{m_s}{2} \langle \bar{s}s \rangle \left[\frac{m_c^2}{r'^2 r} + \frac{m_c^2}{r' r^2} \right] - \frac{m_s^2}{2} \langle \bar{s}s \rangle \left[\frac{2m_c^3}{r'^3 r} + \frac{2m_c^3}{r'^2 r^2} + \frac{m_c}{r'^2 r} + \frac{m_c}{r' r^2} + \frac{2m_c^3}{r' r^3} \right] \\ &\quad + \frac{m_0^2}{4} \langle \bar{s}s \rangle \left[\frac{2m_c^3}{r'^3 r} + \frac{2m_c^3}{r'^2 r^2} + \frac{2m_c}{3r'^2 r} + \frac{m_c}{r' r^2} + \frac{2m_c^3}{r' r^3} \right],\end{aligned}\tag{34}$$

where $r^2 = p^2 - m_c^2$ and $r'^2 = Q^2 - m_c^2$. Finally, we calculate the contribution of diagram (f). For the calculation of this diagram corresponding to the propagation of the soft quark in the external electromagnetic field, we use the light-cone expansion for the nonlocal operators. After contracting the c -quark lines in

$$\begin{aligned}\Pi_{\mu\nu}(p, q) &= i \int d^4x e^{iQx} \langle \gamma(q) | T \{ \bar{s}(0) \gamma_\mu c(0) \bar{c}(x) \\ &\quad \times \gamma_\nu (1 - \gamma_5) s(x) \} | 0 \rangle,\end{aligned}\tag{35}$$

we obtain

$$\begin{aligned}\Pi_{\mu\nu}(p, q) &= i \int d^4x \frac{d^4k}{(2\pi)^2} \frac{e^{i(Q-k)x}}{m_c^2 - k^2} \langle \gamma(q) | \bar{s} \gamma_\mu (\not{k} + m_c) \\ &\quad \times \gamma_\nu (1 - \gamma_5) s \rangle | 0 \rangle.\end{aligned}\tag{36}$$

To calculate the matrix element appearing in the above equation, we use the following identities:

$$\begin{aligned}\gamma_\mu \gamma_\nu &= g_{\mu\nu} + i\sigma_{\mu\nu}, \\ \gamma_\mu \gamma_\nu \gamma_5 &= g_{\mu\nu} \gamma_5 - \frac{i}{2} \varepsilon_{\mu\nu\alpha\sigma} \sigma_{\alpha\sigma},\end{aligned}\tag{37}$$

$$\gamma_\mu \gamma_\alpha \gamma_\nu = g_{\mu\alpha} \gamma_\nu + g_{\alpha\nu} \gamma_\mu - g_{\mu\nu} \gamma_\alpha + i\varepsilon_{\mu\nu\alpha\sigma} \gamma_\sigma \gamma_5$$

and photon distribution amplitudes (DA's) for twist 2, 3, and 4 [20,21]:

$$\begin{aligned}\langle \gamma(q) | \bar{s} \gamma_\nu s | 0 \rangle &= -\frac{Q_s}{2} f_{3\gamma} \int_0^1 du \bar{\psi}^{(V)}(u) x^\theta F_{\theta\nu}(ux) \\ \langle \gamma(q) | \bar{s} \gamma_\alpha \gamma_5 s | 0 \rangle &= -\frac{iQ_s}{4} f_{3\gamma} \int_0^1 du \psi^{(A)}(u) x^\theta \tilde{F}_{\theta\alpha}(ux) \\ \langle \gamma(q) | \bar{s} \sigma_{\alpha\beta} s | 0 \rangle &= Q_s \langle \bar{s}s \rangle \int_0^1 du \phi(u) F_{\alpha\beta}(ux) \\ &\quad + \frac{Q_s \langle \bar{s}s \rangle}{16} \int_0^1 du x^2 \mathcal{A}(u) F_{\alpha\beta}(ux) \\ &\quad + \frac{Q_s \langle \bar{s}s \rangle}{8} \int_0^1 du \mathcal{B}(u) \\ &\quad \times x^\rho (x_\beta F_{\alpha\rho}(ux) - x_\alpha F_{\beta\rho}(ux)),\end{aligned}\tag{38}$$

where $F_{\mu\nu}$ is the field strength tensor of the electromagnetic field, which is defined by

$$\begin{aligned}F_{\mu\nu}(x) &= i(\varepsilon_{\nu\mu} q_\mu - \varepsilon_{\mu\nu} q_\nu) e^{iqx} \\ \tilde{F}_{\mu\nu}(x) &= \frac{1}{2} \varepsilon_{\mu\nu\alpha\beta} F_{\alpha\beta}(x).\end{aligned}\tag{39}$$

The asymptotic expression for the photon wave function $\phi(u)$ in terms of magnetic susceptibility of the quark condensate, $\chi(\mu)$, at a renormalization scale ($\mu = 1 \text{ GeV}^2$) is defined by

$$\phi(u) = \chi(\mu) u(1-u).\tag{40}$$

Other functions used in Eq. (38) are defined by [20,21]

$$\begin{aligned}\bar{\psi}^{(V)}(u) &= -20u(1-u)(2u-1) + \frac{15}{16}(\omega_\gamma^A - 3\omega_\gamma^V) \\ &\quad \times u(1-u)(2u-1)(7(2u-1)^2 - 3), \\ \psi^{(A)}(u) &= (1 - (2u-1)^2)(5(2u-1)^2 - 1) \frac{5}{2} \\ &\quad \times \left(1 + \frac{9}{16} \omega_\gamma^V - \frac{3}{16} \omega_\gamma^A \right), \\ \mathcal{A}(u) &= 40u(1-u)(3k - k^+ + 1) + 8(\xi_2^+ - 3\xi_2) \\ &\quad \times [u(1-u)(2 + 13u(1-u)) \\ &\quad + 2u^3(10 - 15u + 6u^2) \ln u \\ &\quad + 2(1-u)^3(10 - 15(1-u) + 6(1-u^2)) \\ &\quad \times \ln(1-u)], \\ \mathcal{B}(u) &= 40 \int_0^u d\alpha (4 - \alpha)(1 + 3k^+) \left[\frac{-1}{2} + \frac{3}{2}(2\alpha - 1)^2 \right],\end{aligned}\tag{41}$$

where k, k^+, ξ_2, ξ_2^+ , and $f_{3\gamma}$ are constants (see [20,21]). Using the above relations in Eq. (36), we obtain

$$\begin{aligned}
\Pi_{\mu\nu}(p, q) = & - \int d^4x \frac{d^4k}{(2\pi)^2} \frac{e^{i(p-k)x}}{m_c^2 - k^2} \left\{ \frac{-Q_s}{2} f_{3\gamma} k_\mu \int_0^1 du \bar{\psi}^{(V)}(u) x^\sigma F_{\sigma\nu} - \frac{Q_s}{2} f_{3\gamma} k_\nu \int_0^1 du \bar{\psi}^{(V)}(u) x^\sigma F_{\sigma\mu} \right. \\
& - \frac{Q_s}{2} f_{3\gamma} g_{\mu\nu} k_\alpha \int_0^1 du \bar{\psi}^{(V)}(u) x^\sigma F_{\sigma\alpha} + \frac{Q_s}{4} f_{3\gamma} k_\alpha \int_0^1 du \psi^{(A)}(u) [-x^\alpha F_{\mu\nu} + x_\mu F_{\alpha\nu} + x_\nu F_{\mu\alpha}] \\
& + \frac{iQ_s}{8} f_{3\gamma} [k_\mu \varepsilon_{\theta\nu\eta\lambda} + k_\nu \varepsilon_{\theta\mu\eta\lambda} - g_{\mu\nu} k_\alpha \varepsilon_{\theta\alpha\eta\lambda}] \int_0^1 du \psi^{(A)}(u) x^\theta F_{\eta\lambda} \\
& + \frac{iQ_s}{2} f_{3\gamma} k_\alpha \varepsilon_{\mu\nu\alpha\sigma} \int_0^1 du \bar{\psi}^{(V)}(u) x^\theta F_{\theta\sigma} + im_c \left[Q_s \langle \bar{s}s \rangle \int_0^1 du \phi(u) F_{\mu\nu}(ux) + \frac{Q_s \langle \bar{s}s \rangle}{16} \int_0^1 du x^2 \mathcal{A}(u) F_{\mu\nu}(ux) \right. \\
& + \frac{Q_s \langle \bar{s}s \rangle}{8} \int_0^1 du \mathcal{B}(u) x^\sigma (x_\nu F_{\mu\sigma}(ux) - x_\mu F_{\nu\sigma}(ux)) \left. \right] \\
& + \frac{m_c}{2} Q_s \langle \bar{s}s \rangle \varepsilon_{\mu\nu\alpha\sigma} \left[\int_0^1 du \phi(u) F_{\alpha\sigma}(ux) + \frac{1}{16} \int_0^1 du x^2 \mathcal{A}(u) F_{\alpha\sigma}(ux) \right. \\
& \left. + \frac{1}{8} \int_0^1 du \mathcal{B}(u) x^\theta (x_\sigma F_{\alpha\theta}(ux) - x_\alpha F_{\sigma\theta}(ux)) \right] \left. \right\}. \tag{42}
\end{aligned}$$

After performing integration over x and k , the following results corresponding to the coefficients of two invariant structures $i\varepsilon_{\mu\nu\alpha\sigma}\varepsilon^\alpha q^\sigma$ and $[q_\mu \varepsilon_\nu - \varepsilon_\mu q_\nu]$ are obtained as follows:

$$\begin{aligned}
\Pi_1(p, q)^{(f)} = & \frac{m_c Q_s \langle \bar{s}s \rangle}{2} \left[\int_0^1 du \phi(u) \frac{1}{p^2 - m_c^2} - \frac{1}{16} \int_0^1 du \mathcal{A}(u) \left(\frac{10}{(p^2 - m_c^2)^2} + \frac{8m_c^2}{(p^2 - m_c^2)^3} \right) \right], \\
\Pi_2(p, q)^{(f)} = & \frac{-Q_s}{4} f_{3\gamma} \int_0^1 du \psi^{(A)}(u) \left(\frac{1}{p^2 - m_c^2} + \frac{2m_c^2}{(p^2 - m_c^2)^2} \right) \\
& - m_c Q_s \langle \bar{s}s \rangle \left[\int_0^1 du \phi(u) \frac{1}{p^2 - m_c^2} - \frac{1}{16} \int_0^1 du \mathcal{A}(u) \left(\frac{10}{(p^2 - m_c^2)^2} + \frac{8m_c^2}{(p^2 - m_c^2)^3} \right) \right]. \tag{43}
\end{aligned}$$

These results are the final results of the QCD part (OPE expression) of the correlator. The next step is to equate Eqs. (19) and (20) (the physical or phenomenological side of the correlation function) and perform the Borel transformation, with respect to the momentum of the D_s^* meson ($p^2 \rightarrow M_B^2$), in order to suppress the contributions of higher states and continuum. We obtain the following sum rules for the transition form factors, namely:

$$\begin{aligned}
F_{V,A}^{(D_s^*)}(Q^2) = & \frac{m_{D_s^*}}{f_{D_s^*}} e^{(m_{D_s^*}/M_B^2)} \hat{B} \left\{ \int_{(m_c+m_s)^2}^{s_0} ds \frac{\rho_{1,2}(s, p^2)}{s - Q^2} \right. \\
& \left. + \Pi_{1,2}^{c+d+e+f} \right\}, \tag{44}
\end{aligned}$$

where V and A correspond to 1 and 2 in the right-hand side, respectively. In Eq. (44), in order to subtract the contributions of the higher states and the continuum, quark-hadron duality assumption is used, i.e. it is assumed that

$$\rho^{\text{higher states}}(s) = \rho^{\text{OPE}}(s) \theta(s - s_0). \tag{45}$$

In the calculations, the following rule for the Borel transformation is used:

$$\hat{B} \frac{1}{(p^2 - s)^n} = (-1)^n \frac{e^{(-s/M_B^2)}}{\Gamma(n)(M_B^2)^{n-1}}. \tag{46}$$

IV. QCD SUM RULES FOR THE FORM FACTORS INDUCED BY ELECTROMAGNETIC PENGUIN

The effective Hamiltonian for the $b \rightarrow s\gamma$ transition can be written as follows:

$$\begin{aligned}
H = & - \frac{G_F e}{4\pi^2 \sqrt{2}} V_{ib} V_{is}^* C_7(\mu) \bar{s} \sigma_{\mu\nu} \\
& \times \left[m_b \frac{1 + \gamma_5}{2} + m_s \frac{1 - \gamma_5}{2} \right] b F^{\mu\nu}. \tag{47}
\end{aligned}$$

In order to obtain the transition amplitude, we need to calculate the following matrix element:

$$\langle D_s^* | \bar{s} \sigma_{\mu\nu} (1 \pm \gamma_5) q^\nu b | B_c \rangle. \tag{48}$$

At $q^2 = 0$, we can write this matrix element in terms of the two gauge invariant form factors $T_1(0)$ and $T_2(0)$:

$$\begin{aligned}
\langle D_s^*(p, \varepsilon^{(D_s^*)}) | \bar{s} \sigma_{\mu\nu} q^\nu b | B_c(Q) \rangle \\
= & i \varepsilon_{\mu\alpha\beta\lambda} \varepsilon^{(D_s^*)\alpha} p^\beta Q^\lambda T_1(0), \\
\langle D_s^*(p, \varepsilon^{(D_s^*)}) | \bar{s} \sigma_{\mu\nu} q^\nu \gamma_5 b | B_c(Q) \rangle \\
= & [(m_{B_c}^2 - m_{D_s^*}^2) \varepsilon_\mu^{(D_s^*)} - (\varepsilon^{(D_s^*)} \cdot q)(p + Q)_\mu] T_2(0). \tag{49}
\end{aligned}$$

Using the relation

$$\sigma_{\mu\nu}\gamma_5 = -\frac{i}{2}\varepsilon_{\mu\nu\alpha\beta}\sigma^{\alpha\beta}, \quad (50)$$

one can immediately obtain that $T_2(0) = \frac{1}{2}T_1(0)$. Then, we need to calculate only the form factor $T_1(0)$. For this aim, we define the following three point correlation function:

$$\begin{aligned} \Pi_{\mu\alpha} = & - \int d^4x d^4y e^{i(Qx - py)} \langle 0 | T \{ \bar{c}(y) \gamma_\alpha s(y) \bar{s}(0) \\ & \times \sigma_{\mu\nu} q^\nu b(0) \bar{c}(x) i \gamma_5 b(x) \} | 0 \rangle, \end{aligned} \quad (51)$$

where $\bar{c}\gamma_\alpha s$ and $\bar{c}i\gamma_5 b$ are the interpolating currents of D_s^* and B_c mesons, respectively.

After inserting the hadrons full set with quantum numbers of corresponding interpolating currents (see also [13]), we obtain the following expression for the phenomenological part of the correlation function:

$$\begin{aligned} \Pi_{\mu\alpha} = & i \frac{f_{B_c} m_{B_c}^2}{(m_b + m_c)} \frac{f_{D_s^*} m_{D_s^*}}{(p^2 - m_{D_s^*}^2)(Q^2 - m_{B_c}^2)} \\ & \times \varepsilon_{\mu\alpha\beta\lambda} p^\beta Q^\lambda T_1(0) + \text{excited states.} \end{aligned} \quad (52)$$

For the calculation of the QCD part, we write the Lorentz structure in the above correlator as

$$\Pi_{\mu\alpha} = i\varepsilon_{\mu\alpha\beta\lambda} p^\beta Q^\lambda \Pi(p^2, Q^2), \quad (53)$$

where

$$\begin{aligned} \Pi^{\text{per}}(p^2, Q^2) = & -\frac{1}{(2\pi)^2} \int d\tilde{s} ds' \frac{\rho^{\text{per}}(p^2, Q^2)}{(\tilde{s} - Q^2)(s' - p^2)} \\ & + \text{subtraction terms.} \end{aligned} \quad (54)$$

The standard calculations lead to the following result for the perturbative part (bare-loop diagram):

$$\rho^{\text{per}}(s', \tilde{s}) = 4N_C [m_b m_c (A_1 + A_2 + I_0) - m_b^2 A_1 - 2A_3], \quad (55)$$

where

$$\begin{aligned} A_1 = & \frac{2I_0}{(\tilde{s} - s')^2} \left[s'(\tilde{s} + m_c^2 - m_b^2) - \frac{1}{2}(\tilde{s} + s')(s' + m_c^2) \right], \\ A_2 = & \frac{2I_0}{(\tilde{s} - s')^2} \left[\frac{1}{2}(\tilde{s} + s')(m_b^2 - m_c^2 - \tilde{s}) + \tilde{s}(s' + m_c^2) \right], \\ A_3 = & I_0 \frac{m_b^2 [m_b^2 s' + (m_c^2 - s')(\tilde{s} - s')]}{2(\tilde{s} - s')^2}, \\ I_0 = & -\frac{1}{4(\tilde{s} - s')}. \end{aligned} \quad (56)$$




FIG. 3. Feynman diagrams for gluon corrections.

The integration regions over \tilde{s} and s' are obtained from the following inequalities:

$$m_c^2 \leq s' \leq s'_0, \quad s' - \frac{s' m_b^2}{m_c^2 - s'} \leq \tilde{s} \leq \tilde{s}_0. \quad (57)$$

The quark condensate terms give zero contribution after applying the double Borel transformation, with respect to the p^2 ($p^2 \rightarrow M_2^2$) and Q^2 ($Q^2 \rightarrow M_1^2$). Only the gluon condensates can contribute to the form factor. Figure 3 shows such types of diagrams. After lengthy calculations for the gluon condensates contribution and equating the phenomenological and QCD parts and applying double Borel transformation with respect to the p^2 and Q^2 , we find the following expression for the form factor $T_1(0)$:

$$T_1(0) = -\frac{1}{(2\pi)^2} \frac{(m_b + m_c)}{f_{B_c} m_{B_c}^2 f_{D_s^*} m_{D_s^*}} \times \left[\int d\tilde{s} ds' \rho^{\text{per}}(s', \tilde{s}) e^{-(\tilde{s}/M_1^2)} e^{-(s'/M_2^2)} + M_1^2 M_2^2 \left\langle \frac{\alpha_s}{\pi} G^2 \right\rangle C_{G^2} \right], \quad (58)$$

where C_{G^2} is the Wilson coefficient of the gluon condensate and we thus have (see Fig. 3)

$$C_{G^2} = C_{G^2}^a + C_{G^2}^b + C_{G^2}^c + C_{G^2}^d + C_{G^2}^e + C_{G^2}^f. \quad (59)$$

The explicit expressions for $C_{G^2}^i$ are given below as follows:

$$C_{G^2}^a = 96m_b \{ [m_c(I_0[1, 3, 1] + m_b^2 I_0[1, 4, 1] + I_1[1, 3, 1] + m_b^2 I_1[1, 4, 1] + I_2[1, 3, 1] - m_b^2 I_2[1, 4, 1])] + m_b(I_1[1, 3, 1] + m_b^2 I_1[1, 4, 1] + 2I_3[1, 4, 1]) \}, \quad (60)$$

$$C_{G^2}^b = 16\{2I_0[1, 1, 2] + 2m_b m_c I_0[1, 1, 3] + 2I_0[2, 1, 1] + 3m_b m_c I_0[2, 1, 2] + 4m_c^2 I_0[2, 1, 2] + 4m_b m_c^3 I_0[2, 1, 3] + 2m_b m_c I_0[3, 1, 1] + 2m_c^2 I_0[3, 1, 1] + 6m_b m_c^3 I_0[3, 1, 2] + 2m_c^4 I_0[3, 1, 2] + 2m_b m_c^5 I_0[3, 1, 3] + 2I_1[1, 1, 2] - 2m_b^2 I_1[1, 1, 3] + 2m_b m_c I_1[1, 1, 3] + 2I_1[2, 1, 1] - m_b^2 I_1[2, 1, 2] + m_b m_c I_1[2, 1, 2] + 4m_c^2 I_1[2, 1, 2] - 4m_b^2 m_c^2 I_1[2, 1, 3] + 4m_b m_c^3 I_1[2, 1, 3] - 2m_b^2 I_1[3, 1, 1] + 2m_c^2 I_1[3, 1, 1] - 6m_b^2 m_c^2 I_1[3, 1, 2] + 4m_b m_c^3 I_1[3, 1, 2] + 2m_c^4 I_1[3, 1, 2] - 2m_b^2 m_c^4 I_1[3, 1, 3] + 2m_b m_c^5 I_1[3, 1, 3] + 2m_b m_c I_2[1, 1, 3] + I_2[2, 1, 1] + m_b m_c I_2[2, 1, 2] + 4m_b m_c^3 I_2[2, 1, 3] - 4m_c^2 I_2[3, 1, 1] + 4m_b m_c^3 I_2[3, 1, 2] + 2m_b m_c^5 I_2[3, 1, 3] - 4I_3[1, 1, 3] - 4I_3[2, 1, 2] - 8m_c^2 I_3[2, 1, 3] - 8I_3[3, 1, 1] - 16m_c^2 I_3[3, 1, 2] - 4m_c^4 I_3[3, 1, 3]\} - 32M_2^2 \frac{d}{dM_2^2} \{M_2^2 [I_0[2, 1, 2] + 2m_b m_c I_0[2, 1, 3] + 2m_b m_c I_0[3, 1, 2] + m_c^2 I_0[3, 1, 2] + 2m_b m_c^3 I_0[3, 1, 3] + I_1[2, 1, 2] - 2m_b^2 I_1[2, 1, 3] + 2m_b m_c I_1[2, 1, 3] - 2m_b^2 I_1[3, 1, 2] + m_b m_c I_1[3, 1, 2] + m_c^2 I_1[3, 1, 2] - 2m_b^2 m_c^2 I_1[3, 1, 3] + 2m_b m_c^3 I_1[3, 1, 3] - I_2[2, 1, 2] + 2m_b m_c I_2[2, 1, 3] - 2I_2[3, 1, 1] + m_b m_c I_2[3, 1, 2] - m_c^2 I_2[3, 1, 2] + 2m_b m_c^3 I_2[3, 1, 3] - 4I_3[2, 1, 3] - 6I_3[3, 1, 2] - 4m_c^2 I_3[3, 1, 3]\} - 32M_2^4 \left(\frac{d^2}{dM_2^2}\right)^2 \{M_2^4 [m_b^2 I_1[3, 1, 3] + 2I_2[3, 1, 2] - m_b m_c (I_0[3, 1, 3] + I_1[3, 1, 3] + I_2[3, 1, 3]) + 2I_3[3, 1, 3]\}, \quad (61)$$

$$C_{G^2}^c = 96m_c \{ -(m_b^2 m_c I_1[4, 1, 1]) + m_b (I_0[3, 1, 1] + m_c^2 I_0[4, 1, 1] + I_1[3, 1, 1] + m_c^2 I_1[4, 1, 1] + I_2[3, 1, 1] + m_c^2 I_2[4, 1, 1]) - 2m_c I_3[4, 1, 1] \}, \quad (62)$$

$$C_{G^2}^d = -32m_b m_c \{ I_0[2, 1, 2] + I_0[3, 1, 1] + m_c^2 I_0[3, 1, 2] + I_1[2, 1, 2] + m_c^2 I_1[3, 1, 2] + I_2[2, 1, 2] + m_c^2 I_2[3, 1, 2] - 4I_3[3, 1, 2] - M_2^2 \frac{d}{dM_2^2} [M_2^2 (I_0[3, 1, 2] + I_1[3, 1, 2] + I_2[3, 1, 2])] \} + 16\{m_b^2 I_1[2, 1, 2] + I_2[2, 1, 1] - m_b m_c (I_0[2, 1, 2] + I_1[2, 1, 2] + I_2[2, 1, 2]) + 4I_3[2, 1, 2]\}, \quad (63)$$

$$\begin{aligned}
 C_{G^2}^e = & 16\{2I_0[1, 1, 2] + I_0[1, 2, 1] + 2m_b^2 I_0[1, 2, 2] + m_b m_c I_0[1, 2, 2] + m_c^2 I_0[2, 1, 2] + m_b m_c I_0[2, 2, 1] + m_c^2 I_0[2, 2, 1] \\
 & + m_b^2 m_c^2 I_0[2, 2, 2] + m_b m_c^3 I_0[2, 2, 2] + I_1[1, 1, 2] + I_1[1, 2, 1] + m_b^2 I_1[1, 2, 2] + m_b m_c I_1[1, 2, 2] - I_1[2, 1, 1] \\
 & - 2m_b m_c I_1[2, 1, 2] - m_b^2 I_1[2, 2, 1] + m_c^2 I_1[2, 2, 1] - 2m_b^3 m_c I_1[2, 2, 2] + m_b m_c^3 I_1[2, 2, 2] + I_2[1, 1, 2] \\
 & + I_2[1, 2, 1] + m_b^2 I_2[1, 2, 2] + m_b m_c I_2[1, 2, 2] - 2m_b m_c I_2[2, 2, 1] + m_c^2 I_2[2, 2, 1] + m_b m_c^3 I_2[2, 2, 2] \\
 & - 2I_3[2, 1, 2] - 4I_3[2, 2, 1] - 2m_b^2 I_3[2, 2, 2] - 4m_b m_c I_3[2, 2, 2]\} - 16M_1^2 \frac{d}{dM_1^2} \{M_1^2 [I_2[2, 2, 1] + 2I_3[2, 2, 2]]\} \\
 & + 16M_2^2 \frac{d}{dM_2^2} \{M_2^2 [-m_b m_c I_0[2, 2, 2] + I_1[2, 1, 2] + m_b^2 I_1[2, 2, 2] - m_b m_c I_1[2, 2, 2] + I_2[2, 1, 2] + I_2[2, 2, 1] \\
 & + m_b^2 I_2[2, 2, 2] - m_b m_c I_2[2, 2, 2] + 2I_3[2, 2, 2]]\} + 16\{I_1[1, 1, 2] + m_b^2 I_1[1, 2, 2] + I_2[1, 2, 1] \\
 & - m_b m_c (I_0[1, 2, 2] + I_1[1, 2, 2] + I_2[1, 2, 2]) + 2I_3[1, 2, 2]\}, \tag{64}
 \end{aligned}$$

$$\begin{aligned}
 C_{G^2}^f = & 16\{2I_0[1, 2, 1] + 2I_0[2, 1, 1] + 2m_b^2 I_0[2, 2, 1] + 6m_b m_c I_0[2, 2, 1] + 2m_c^2 I_0[2, 2, 1] + 2I_1[1, 2, 1] - 5I_1[2, 1, 1] \\
 & - 5m_c^2 I_1[2, 2, 1] + 6m_b m_c I_1[2, 2, 1] + 2m_c^2 I_1[2, 2, 1] + 2I_2[1, 2, 1] - I_2[2, 1, 1] - m_b^2 I_2[2, 2, 1] \\
 & + 6m_b m_c I_2[2, 2, 1] + 2m_c^2 I_2[2, 2, 1] - 14I_3[2, 2, 1]\} - 32M_1^2 \frac{d}{dM_1^2} \{M_1^2 [I_0[2, 2, 1] + I_1[2, 2, 1] + I_2[2, 2, 1]]\} \tag{65}
 \end{aligned}$$

and for the explicit form of the $I_i[a, b, c]$, we obtain

$$\begin{aligned}
 I_0[a, b, c] &= \frac{(-1)^{a+b+c}}{16\pi^2 \Gamma(a)\Gamma(b)\Gamma(c)} (M_1^2)^{2-a-b} (M_2^2)^{2-a-c} \mathcal{U}_0(a+b+c-4, 1-c-b), \\
 I_1[a, b, c] &= \frac{(-1)^{a+b+c+1}}{16\pi^2 \Gamma(a)\Gamma(b)\Gamma(c)} (M_1^2)^{2-a-b} (M_2^2)^{3-a-c} \mathcal{U}_0(a+b+c-5, 1-c-b), \\
 I_2[a, b, c] &= \frac{(-1)^{a+b+c+1}}{16\pi^2 \Gamma(a)\Gamma(b)\Gamma(c)} (M_1^2)^{3-a-b} (M_2^2)^{2-a-c} \mathcal{U}_0(a+b+c-5, 1-c-b), \\
 I_3[a, b, c] &= \frac{(-1)^{a+b+c+1}}{32\pi^2 \Gamma(a)\Gamma(b)\Gamma(c)} (M_1^2)^{3-a-b} (M_2^2)^{3-a-c} \mathcal{U}_0(a+b+c-6, 2-c-b). \tag{66}
 \end{aligned}$$

The function $\mathcal{U}_0(i, j)$, also, is given by

$$\begin{aligned}
 \mathcal{U}_0(i, j) &= \int_0^\infty dy (y + M_1^2 + M_2^2)^i y^j \\
 &\times \exp\left[-\frac{B_{-1}}{y} - B_0 - B_1 y\right], \tag{67}
 \end{aligned}$$

where

$$\begin{aligned}
 B_{-1} &= \frac{m_b^2}{M_1^2} \left[M_1^2 + M_2^2 \right], \\
 B_0 &= \frac{1}{M_1^2 M_2^2} \left[M_1^2 m_c^2 + M_2^2 (m_c^2 + m_b^2) \right], \\
 B_1 &= \frac{m_c^2}{M_1^2 M_2^2}. \tag{68}
 \end{aligned}$$

V. NUMERICAL ANALYSIS

In this section, we present our numerical analysis for the form factors. From the sum rule expressions of these form

factors, we see that the condensates, leptonic decay constants of B_c and D_s^* mesons, continuum thresholds s_0 , \tilde{s}_0 , and s'_0 , the relevant parameters in photon DA's and Borel parameters M_B^2 , M_1^2 , and M_2^2 are the main input parameters. In further numerical analysis, we choose the value of the condensates at a fixed renormalization scale of about 1 GeV. The values of the condensates are [22]: $\langle \bar{\psi}\psi \rangle_{\mu=1 \text{ GeV}} = -(240 \pm 10 \text{ MeV})^3$, $\langle \bar{s}s \rangle = (0.8 \pm 0.2) \langle \bar{\psi}\psi \rangle$ and $m_0^2 = 0.8 \text{ GeV}^2$. The quark and mesons masses are taken to be $m_c(\mu = m_c) = 1.275 \pm 0.015 \text{ GeV}$, $m_s(1 \text{ GeV}) \simeq 142 \text{ MeV}$ [23], $m_b = (4.7 \pm 0.1) \text{ GeV}$ [22], $m_{D_s^*} = 2.112 \text{ GeV}$, and $m_{B_c} = 6.258 \text{ GeV}$. For the values of the leptonic decay constants of B_c and D_s^* mesons, we use the results obtained from the two-point QCD analysis: $f_{B_c} = 0.35 \text{ GeV}$ [24–26] and $f_{D_s^*} = 266 \pm 32 \text{ MeV}$ [27]. The relevant parameters in photon DA's are taken to be $\chi = 3.15 \pm 0.3 \text{ GeV}^{-2}$, $\kappa = 0.2$, $\kappa^+ = 0$, $\zeta_1 = 0.4$, $\zeta_1^+ = 0$, $\zeta_2 = 0.3$, $\zeta_2^+ = 0$, $f_{3\gamma} = -(4 \pm 2) \times 10^{-3} \text{ GeV}^2$, $w_\gamma^A = -2.1 \pm 1.0$, and $w_\gamma^V = 3.8 \pm 1.8$ [20,21,28]. The threshold parameters are also

determined from the two-point QCD sum rules: $s_0 = 8 \text{ GeV}^2$, $\tilde{s}_0 = 45 \text{ GeV}^2$, $s'_0 = 8 \text{ GeV}^2$ [2,27,29]. The Borel parameters M_B^2 , M_1^2 , and M_2^2 are auxiliary quantities and therefore the results of physical quantities should not depend on them. In the QCD sum rule method, OPE is truncated at finite order, leaving a residual dependence on the Borel parameters. For this reason, the working regions for the Borel parameters should be chosen such that in these regions the form factors are practically independent of them. The working regions for the Borel parameters M_B^2 , M_1^2 , and M_2^2 can be determined on the condition that, on the one side, the continuum contribution should be small, and on the other side, the contribution of the operator with the highest dimension should be small. As a result of the above-mentioned requirements, the working regions for this transition are obtained to be

$$\begin{aligned} 4 \text{ GeV}^2 < M_B^2 < 10 \text{ GeV}^2, \\ 10 \text{ GeV}^2 \leq M_1^2 \leq 25 \text{ GeV}^2, \\ 4 \text{ GeV}^2 \leq M_2^2 \leq 10 \text{ GeV}^2. \end{aligned} \quad (69)$$

Now, by calculating the total decay widths and taking $|V_{cs}| = 0.957 \pm 0.017$, $|V_{cb}| = 0.0416 \pm 0.0006$, $|V_{tb}| = 0.77_{-0.24}^{+0.18}$, $|V_{ts}| = (40.6 \pm 2.7) \times 10^{-3}$ [30], $\langle \frac{\alpha_s}{\pi} G^2 \rangle = 0.012 \text{ GeV}^4$ [29], $C_7(\mu = m_c) = -0.0068 - 0.02i$ [5], and $\tau_{B_c} = 0.52 \times 10^{-12} \text{ s}$ [31], we obtain the numerical results of the electromagnetic penguin (EP), weak annihilation (WA), and total branching ratios for this decay as follows:

$$\begin{aligned} \mathbf{B}^{(\text{EP})}(B_c \rightarrow D_s^* \gamma) &= 3.468 \times 10^{-6} \\ \mathbf{B}^{(\text{WA})}(B_c \rightarrow D_s^* \gamma) &= 1.557 \times 10^{-5} \\ \mathbf{B}^{(\text{total})}(B_c \rightarrow D_s^* \gamma) &= 2.462 \times 10^{-5}. \end{aligned} \quad (70)$$

From the above results, we see that the weak annihilation contribution to the total branching ratio is about 4.48 times greater than that of the electromagnetic penguin diagram. Here, it is observed that the difference between the total branching ratio with the sum of the weak annihilation and electromagnetic penguin branching ratios comes from the cross term in total decay width. Also our result for the total branching ratio shows that the $B_c \rightarrow D_s^* \gamma$ decay can be measured at LHC.

Now, we compare our results of the $B_c \rightarrow D_s^* \gamma$ to the results of the perturbative QCD [15], relativistic independent quark model [16], SM (PQCD) [17], multiscale walking technicolor (MWTTCM) [17], and topcolor assisted MWTTCM (TAMWTTCM) [17] for $\tau_{B_c} = 0.52 \times 10^{-12} \text{ s}$ as shown in Table I.

Looking at this table, it is seen that there is a good agreement between the present study and the PQCD [15], in order of magnitude for the total branching ratio. However, our result is approximately 1 order of magnitude less than that of the RIQM and MWTTCM. Also, it is 1 order of magnitude and 2 orders of magnitude greater than that of the TAMWTTCM and SM (PQCD) [17], respectively. The ratio of $\mathbf{B}^{\text{WA}}/\mathbf{B}^{\text{EP}}$ in the present work, PQCD [15], RIQM, SM (PQCD) [17], TAMWTTCM, MWTTCM are 4.48, 1.34, 1.9, 3.4, 1.23, and 0.01, respectively. As a result of the above discussions, we can say that in the QCD sum rules (present study), relativistic independent quark model, perturbative QCD and TAMWTTCM approaches, the weak annihilation contribution to the total branching ratio dominates the contribution coming from the electromagnetic penguin diagram, but this is not true only for the MWTTCM approach. The presence of the pseudo-Goldstone bosons in the MWTTCM leads to a discrepancy between this model and the other two models in [17] (for more details see [17]) and a part of inconsistency in the results of the different methods may be related to the different magnitudes of the input parameters, getting from different references; e.g. we use $m_c(\mu = m_c) = 1.275 \pm 0.015 \text{ GeV}$ for the c -quark masses while the authors of [17] use $m_c = 1.6 \text{ GeV}$ and also to the nature of the methods and their accuracy.

In this step, for the analysis of $B_c \rightarrow D^* \gamma$, in the entire calculations we replace the s quark with the d quark. Making $m_{D_s^*} \rightarrow m_{D^*}$, $f_{D_s^*} \rightarrow f_{D^*}$, $V_{ts} \rightarrow V_{td}$, $V_{cs} \rightarrow V_{cd}$ changes and taking $m_{D^*} = 2.010 \text{ GeV}$, $f_{D^*} = 0.23 \pm 0.02 \text{ GeV}$ [32], $V_{cd} = 0.230 \pm 0.011$, $V_{td} = (7.4 \pm 0.8) \times 10^{-3}$ [30], and $m_d = 5 \text{ MeV}$, we obtain the numerical results as below:

$$\begin{aligned} \mathbf{B}^{(\text{EP})}(B_c \rightarrow D^* \gamma) &= 1.151 \times 10^{-7} \\ \mathbf{B}^{(\text{WA})}(B_c \rightarrow D^* \gamma) &= 2.162 \times 10^{-6} \\ \mathbf{B}^{(\text{total})}(B_c \rightarrow D^* \gamma) &= 2.786 \times 10^{-6}. \end{aligned} \quad (71)$$

TABLE I. Comparison of the branching ratio for $B_c \rightarrow D_s^* \gamma$ decay based on the $\tau_{B_c} = 0.52 \times 10^{-12} \text{ s}$.

	$\mathbf{B}^{\text{EP}}(B_c \rightarrow D_s^* \gamma)$	$\mathbf{B}^{\text{WA}}(B_c \rightarrow D_s^* \gamma)$	$\mathbf{B}^{\text{total}}(B_c \rightarrow D_s^* \gamma)$
Present study	3.468×10^{-6}	1.557×10^{-5}	2.462×10^{-5}
PQCD	3.70×10^{-6}	4.94×10^{-6}	1.14×10^{-5}
RIQM	2.40×10^{-5}	4.51×10^{-5}	1.39×10^{-4}
MWTTCM	$(0.68-3.42) \times 10^{-4}$	$(0.74-0.81) \times 10^{-6}$	$(0.74-3.57) \times 10^{-4}$
TAMWTTCM	$(5.18-7.23) \times 10^{-7}$	$(7.24-8.13) \times 10^{-7}$	$(1.78-9.95) \times 10^{-6}$
SM (PQCD)	1.73×10^{-7}	5.89×10^{-7}	7.83×10^{-7}

TABLE II. Comparison of the branching ratio for $B_c \rightarrow D^* \gamma$ decay based on the $\tau_{B_c} = 0.52 \times 10^{-12} s$.

	$\mathbf{B}^{\text{EP}}(B_c \rightarrow D^* \gamma)$	$\mathbf{B}^{\text{WA}}(B_c \rightarrow D^* \gamma)$	$\mathbf{B}^{\text{total}}(B_c \rightarrow D^* \gamma)$
Present study	1.151×10^{-7}	2.162×10^{-6}	2.786×10^{-6}
RIQM [16]	5.70×10^{-7}	1.33×10^{-6}	3.64×10^{-6}

These results also enhance the importance of the weak annihilation contribution to the total branching ratio in comparing with the electromagnetic penguin diagram ones for the $B_c \rightarrow D^* \gamma$. Finally, we compare our results to the relativistic independent quark model (RIQM) [16] for $\tau_{B_c} = 0.52 \times 10^{-12} s$ in Table II.

From the Table II, a good agreement is also seen in the order of magnitude between the present study and the relativistic independent quark model.

In conclusion, the present study concentrated on the radiative $B_c \rightarrow D_s^* \gamma$ and $B_c \rightarrow D^* \gamma$ decays in the frame-

work of QCD sum rules. The form factors responsible for these decays were calculated. The branching ratio for this decays were estimated. The results show that the $B_c \rightarrow D_s^* \gamma$ case can be measured at LHC in the near future.

ACKNOWLEDGMENTS

One of the authors (K.A.) would like to thank TUBITAK, Turkish Scientific and Research Council, for their partial support. Also, V.B. would like to thank theory group of CERN for their hospitality.

-
- [1] F. Abe *et al.* (CDF Collaboration), Phys. Rev. D **58**, 112004 (1998).
- [2] T.M. Aliev and M. Savci, Phys. Lett. B **434**, 358 (1998).
- [3] T.M. Aliev and M. Savci, J. Phys. G **24**, 2223 (1998).
- [4] T.M. Aliev and M. Savci, Eur. Phys. J. C **47**, 413 (2006).
- [5] T.M. Aliev and M. Savci, Phys. Lett. B **480**, 97 (2000).
- [6] M. A. Ivanov, J. G. Korner, and P. Santorelli, Phys. Rev. D **73**, 054024 (2006).
- [7] V. Bashiry, Eur. Phys. J. C **47**, 423 (2006).
- [8] P. Colangelo and A. Khodjamirian, in *At the Frontier of Particle Physics/Handbook of QCD*, edited by M. Shifman (World Scientific, Singapore, 2001), Vol. 3, p. 1495.
- [9] P. Ball, V.M. Braun, and H.G. Dosch, Phys. Rev. D **44**, 3567 (1991).
- [10] P. Ball, Phys. Rev. D **48**, 3190 (1993).
- [11] A. A. Ovchinnikov and V. A. Slobodenyuk, Z. Phys. C **44**, 433 (1989); V. N. Baier and A. Grozin, Z. Phys. C **47**, 669 (1990).
- [12] A. A. Ovchinnikov, Sov. J. Nucl. Phys. **50**, 519 (1989).
- [13] T.M. Aliev, K. Azizi, and A. Ozpineci, Eur. Phys. J. C **51**, 593 (2007).
- [14] V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, Nucl. Phys. **B569**, 473 (2000).
- [15] D. S. Du, X. Li, and Y. Yang, Phys. Lett. B **380**, 193 (1996).
- [16] N. Barik, Sk. Naimuddin, S. Kar, and P. C. Dash, Phys. Rev. D **63**, 014024 (2001).
- [17] Gongru Lu, Chongxing Yue, Yigang Cao, Zhaohua Xiong, and Zhenjun Xiao, Phys. Rev. D **54**, 5647 (1996).
- [18] Alexander Khodjamirian and Daniel Wyler, in *Sergei Matinian Festschrift*, From Integrable Models to Gauge Theories, edited by V. Gurzadyan and A. Sedrakyan (World Scientific, Singapore, 2002).
- [19] V. A. Nesterenko and A. V. Radyushkin, Sov. J. Nucl. Phys. **39**, 811 (1984).
- [20] J. Rohrwild, Phys. Rev. D **75**, 074025 (2007).
- [21] P. Ball, V.M. Braun, and N. Kivel, Nucl. Phys. **B649**, 263 (2003).
- [22] B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, 232 (2006).
- [23] Ming Qiu Huang, Phys. Rev. D **69**, 114015 (2004).
- [24] P. Colangelo, G. Nardulli, and N. Paver, Z. Phys. C **57**, 43 (1993).
- [25] V. V. Kiselev and A. V. Tkabladze, Phys. Rev. D **48**, 5208 (1993).
- [26] T.M. Aliev and O. Yilmaz, Nuovo Cimento A **105**, 827 (1992).
- [27] P. Colangelo, F. De Fazio, and A. Ozpineci, Phys. Rev. D **72**, 074004 (2005).
- [28] I. I. Balitsky, V.M. Braun, and A. V. Kolesnichenko, Nucl. Phys. **B312**, 509 (1989).
- [29] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. **B147**, 385 (1979).
- [30] A. Ceccucci *et al.* (Particle Data Group), J. Phys. G **33**, 139 (2006).
- [31] M. Beneke and G. Buchalla, Phys. Rev. D **53**, 4991 (1996).
- [32] K. C. Bowler, L. Del Debbio, J. M. Flynn, G. N. Lacagnina, V. I. Lesk, C. M. Maynard, and D. G. Richards, Nucl. Phys. **B619**, 507 (2001).