

Naturalness and the neutrino matrix

J. Sayre¹ and S. Wiesenfeldt^{1,2}

¹*Department of Physics, University of Illinois at Urbana-Champaign, 1110 West Green Street, Urbana, Illinois 61801, USA*

²*Institut für Theoretische Teilchenphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*

(Received 14 November 2007; published 13 March 2008)

The observed pattern of neutrino mass splittings and mixing angles indicates that their family structure is significantly different from that of the charged fermions. We investigate the implications of these data for the fermion mass matrices in grand-unified theories with a type-I seesaw mechanism. We show that, with simple assumptions, naturalness leads to a strongly hierarchical Majorana mass matrix for heavy right-handed neutrinos and a partially cascade form for the Dirac neutrino matrix. We consider various model building scenarios which could alter this conclusion, and discuss their consequences for the construction of a natural model. We find that including partially lopsided matrices can aid us in generating a satisfying model.

DOI: [10.1103/PhysRevD.77.053005](https://doi.org/10.1103/PhysRevD.77.053005)

PACS numbers: 14.60.Pq, 12.10.Dm

I. INTRODUCTION

The measurement of neutrino mass splittings and mixing angles [1,2] has provided a new window into physics beyond the standard model. The fact that the hierarchy between at least one pair of the neutrinos is weak and that two leptonic mixing angles are large, in contrast to the strongly hierarchical masses of quarks and charged leptons and small Cabibbo-Kobayashi-Maskawa (CKM) mixing, was initially surprising. It leads us to surmise that neutrino masses arise through a somewhat different mechanism than the quark and charged lepton masses. Thus, the relation between the charged fermion and neutrino observables is not necessarily obvious. In fact, we have such a mechanism in the form of the type-I seesaw [3], which can naturally yield neutrino masses in the range indicated by experiment. Moreover, the physical light neutrino mass matrix is a product of more fundamental matrices. This fact can potentially explain the differences between the mixing angles and mass hierarchies of the charged fermion and neutrino sectors.

The seesaw mechanism arises naturally within a grand-unified theory (GUT) such as SO(10) [4], where each generation of standard model fermions is unified into the 16-dimensional spinor representation, together with the right-handed neutrinos. The breaking of $B - L$ (where B and L denote baryon and lepton number, respectively), which is a subgroup of SO(10), automatically gives rise to Majorana masses for the singlet neutrinos, and then to the seesaw mechanism. Indeed, the neutrino data have encouraged GUT model building [5,6].

Although GUTs provide a natural framework for massive neutrinos and, combined with family symmetries or textures, have allowed for a number of successful models of quark masses and mixing, it has proven difficult to incorporate neutrinos in a completely satisfactory manner. In this paper, we reconsider neutrino masses and mixings under the guidance of naturalness. That is, rather than focusing on a particular theoretical structure and modify-

ing it as necessary to obtain the best fit to the data, we will try to minimize the dependence on specific model assumptions and work up from the experimental data to see where it naturally leads us. In particular, we will show that the construction of a natural, unified picture of all standard model fermion masses and mixing angles imposes non-trivial constraints on the structure of both sectors.

In this framework, we are interested only in the orders of magnitude of various parameters and, in pursuing natural solutions, we seek to avoid unnatural cancellations, i.e., that terms of a given order must cancel to produce a term of lower order. It may be possible to arrange such cancellations in a technically natural way via a judicious choice of symmetries, but this is by no means trivial. Furthermore, an exact symmetry is a strong assumption to make, given the current uncertainty in the neutrino data. We will instead adopt naturalness as described above, seeking to constrain the approximate structure of our theory without *ad hoc* symmetries. Ideally, this structure can serve as a guide for developing well-motivated symmetries upon which an ultimately satisfying theory can be built.

Of course, one must make some assumptions based on previous successes to make progress and, in this capacity, we will focus on the SO(10) models with small representations [7,8]. This scenario will serve as a concrete example; however, much of the analysis could be adapted to SO(10) models with large representations and/or type-II seesaw mechanisms, as well as to other unifying groups.

This paper is organized as follows: We start by introducing our theoretical framework in Sec. II and reviewing the experimental data in Sec. III. In Sec. IV we derive natural constraints on the neutrino mass matrices. Since the fermion mass matrices are related by the GUT symmetry, we study the implications of quark mixing in Sec. V. In Sec. VI we show how mass matrices consistent with our constraints can be generated via family symmetries, and we investigate how well they can fit the charged fermion masses. In SO(10) models with small representations, the neutrino Dirac mass matrix can receive additional contributions

via couplings to a second up-type Higgs doublet, present in the $B - L$ breaking Higgs field. We consider this possibility in Sec. VII, supplemented by the appendix. The remaining sections are devoted to two cases which generalize beyond our initial assumptions. These involve models wherein otherwise negligible leptonic rotations play an important role in neutrino mixing, either due to a lopsided structure in some mass matrices (Sec. VIII), or to a particular form for the effective neutrino matrix (Sec. IX). We conclude in Sec. X.

II. GENERAL STRUCTURE OF THEORY

The standard model fermions are found in three copies of the spinor representation $\mathbf{16}_i$.¹ We will make use of the small representations $\mathbf{10}_H$, $\mathbf{45}_H$, $\mathbf{16}_H$, $\mathbf{16}'_H$, $\overline{\mathbf{16}}_H$, and potentially $\overline{\mathbf{16}}'_H$ to break the GUT symmetry and to generate fermion masses. Several authors have used this framework to build interesting models [7,8].

The SO(10) symmetry is broken to the standard model by GUT scale vacuum expectation values (vevs), one in the SU(5) singlet direction of $\mathbf{16}_H$ and $\overline{\mathbf{16}}_H$, denoted ν , and $\langle \mathbf{45}_H \rangle$ along the $B - L$ direction. The electroweak symmetry is broken when weak doublets in $\mathbf{10}_H$ acquire vevs. It is also possible that the doublets in $\mathbf{16}'_H$ and $\overline{\mathbf{16}}'_H$ acquire weak scale vevs, in which case the light Higgs doublets are a mixture of weak doublets from the vector and spinor representations [9]. We will assume for now that $\overline{\mathbf{16}}'_H$ does not acquire a weak vev.

Charged fermion masses are generated via several operators: the renormalizable operator $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H$, which contributes to all Dirac mass matrices for the standard model fermions; the higher-dimensional operator $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H \mathbf{45}_H$, which differentiates the quark mass matrices from the lepton matrices due to their differing charges under $B - L$; and $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}'_H$, which contributes only to down quark and charged lepton mass matrices. The operator $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H$ is symmetric in generation space while $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H \mathbf{45}_H$ is antisymmetric ($\mathbf{16}_i$ and $\mathbf{16}_j$ are contracted as a $\mathbf{120}$, for $\langle \mathbf{45} \rangle \propto B - L$ this is the only contraction that contributes to the mass matrices). The operator $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}'_H$ may be symmetric or asymmetric, depending on how the fields are contracted.

With this set of operators, the Dirac neutrino matrix M_D receives contributions from the operators $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H$ and $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H \mathbf{45}_H$, and we expect it to be somewhat similar to the up quark matrix, i.e., to have a similarly strong hierarchy of mass eigenstates from the first to the third generation. For the up quarks this is approximately 5 orders of magnitude. Although the neutrino hierarchy can be somewhat weaker due to factors of 3 coming from the $B - L$ direction vev of the $\mathbf{45}_H$, one would still expect roughly a

10^{-4} ratio between the lightest and heaviest Dirac matrix eigenvalues.

We define the orientation of M_D as $\nu^i M_D^{ij} N^j$, where N is the standard model singlet. Then we can parametrize the Dirac matrix as

$$M_D \equiv L_D D_D R_D^\dagger. \quad (1)$$

Here and throughout the paper the matrices M are dimensionless and the largest eigenvalue is normalized to 1. Since we are primarily concerned with interfamily relations this causes no problems, but one should bear in mind that there is an overall scale associated with all mass matrices. In the above case, the dimensionful Dirac mass operator is $u \nu M_D N$, where u is the mass of the largest eigenvalue. Similarly, throughout the paper L and R will signify unitary matrices defined by the diagonalization equations

$$L^\dagger M M^\dagger L = R^\dagger M^\dagger M R = D^2 \equiv \text{diag}(\eta^2, \epsilon^2, 1), \quad (2)$$

where $\eta, \epsilon, 1$ are the normalized eigenvalues of M .

In general, L_D and R_D are arbitrary unitary matrices and D_D is a diagonal matrix of the eigenvalues of M_D ; however, we expect the eigenvalues to be strongly hierarchical. This hierarchy will be naturally generated if we posit the forms

$$\begin{aligned} D_D &\equiv \text{diag}(\eta, \epsilon, 1), \\ L_D &\sim \begin{pmatrix} 1 & \mu' \sqrt{\frac{\eta}{\epsilon}} & \nu' \sqrt{\eta} \\ \mu' \sqrt{\frac{\eta}{\epsilon}} & 1 & \rho' \sqrt{\epsilon} \\ \nu' \sqrt{\eta} & \rho' \sqrt{\epsilon} & 1 \end{pmatrix}, \\ R_D &\sim \begin{pmatrix} 1 & \mu \sqrt{\frac{\eta}{\epsilon}} & \nu \sqrt{\eta} \\ \mu \sqrt{\frac{\eta}{\epsilon}} & 1 & \rho \sqrt{\epsilon} \\ \nu \sqrt{\eta} & \rho \sqrt{\epsilon} & 1 \end{pmatrix}. \end{aligned} \quad (3)$$

We expect $\eta \ll \epsilon \ll 1$. Based on the quark hierarchy we may estimate their approximate size as $\eta \sim 10^{-4}$ and $\epsilon \sim 10^{-2}$, but most of the analysis does not depend on this assumption.

L_D and R_D are unitary matrices and the parametrizations above should be read as giving the orders of magnitude only of the various entries. The parameters μ, ν, ρ and their primed counterparts are generally expected to be less than or equal to order one. If they were significantly larger, various entries would need to cancel to preserve the smaller eigenvalues. Thus $\mu, \nu, \rho \sim 1$ is the minimal requirement for naturalness in the absence of an exact symmetry relating the Yukawa couplings. This is known as a geometrical hierarchy pattern [10]. It corresponds to the following form for M_D :

¹The subscripts i, j will be used to indicate generations while Higgs fields will be denoted with a subscript H .

$$M_D \sim \begin{pmatrix} \leq \eta & \sqrt{\eta\epsilon} & \sqrt{\eta} \\ \sqrt{\eta\epsilon} & \leq \epsilon & \sqrt{\epsilon} \\ \sqrt{\eta} & \sqrt{\epsilon} & 1 \end{pmatrix}. \quad (4)$$

The central feature of such a matrix is that the off-diagonal entries play a dominant or codominant role in determining the two smaller eigenvalues. A geometric hierarchy can be easily obtained with a U(1) symmetry via the Froggatt-Nielsen mechanism [11].

On the other hand, μ , ν , and ρ may be arbitrarily smaller without endangering the eigenvalue hierarchy. In this case the diagonal entries in M_D become dominant and must be correspondingly close to the eigenvalues. We will refer to this possibility as a subgeometric hierarchy. With three generations it is, of course, possible to have a mixed case which is partially geometric and partially subgeometric.

There is one exception to these naturalness considerations, which occurs if M_D is highly asymmetric, i.e., if $(M_D)_{ji}$ and $(M_D)_{ij}$ are of different orders for some i and j . However, if it arises only from $\mathbf{16}_i\mathbf{16}_j\mathbf{10}_H$ and $\mathbf{16}_i\mathbf{16}_j\mathbf{10}_H\mathbf{45}_H$, we would not expect this; these operators give symmetric and antisymmetric contributions, respectively, which would have to be arranged to cancel in a seemingly unnatural way. Thus we generally expect L_D and R_D to have similar values for their parameters, i.e., $\mu \sim \mu'$, $\nu \sim \nu'$ and $\rho \sim \rho'$.

To implement the type-I seesaw, we need a matrix for the heavy neutrinos: $N^i M_R^{ij} N^j$. Such a coupling may arise from $\frac{1}{m}(M_R)_{ij}\mathbf{16}_i\mathbf{16}_j\overline{\mathbf{16}}_H\overline{\mathbf{16}}_H$ when $\overline{\mathbf{16}}_H$ acquires its GUT scale vev v . This nonrenormalizable operator is suppressed by some mass m , which is by default the Planck scale but which in practice may be somewhat less, depending on the origin of the effective operator. The seesaw formula then gives

$$M_\nu \simeq -M_D M_R^{-1} (M_D)^T. \quad (5)$$

As discussed above, M_D , M_ν and M_R are dimensionless. The massive parameter which sets the scale for the neutrinos is $u^2 m / v^2$. For $u \sim 100$ GeV, $v \sim 10^{16}$ GeV, and $m \sim m_{\text{Pl}} \sim 10^{18}$ GeV, this comes out to be 0.1 eV, consistent with the range indicated by experiment.

$$V_{\text{PMNS}} \equiv L_e^\dagger L_\nu = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \times \text{diag}(e^{i\alpha_1/2}, e^{i\alpha_2/2}, 1). \quad (7)$$

For concreteness, we will assume the tribimaximal solution which sets the mixing angles $\theta_{12} = 30^\circ$, $\theta_{13} = 0^\circ$, $\theta_{23} = 45^\circ$ [16],

$$V_{\text{PMNS}} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{3}} & 0 \\ -\sqrt{\frac{1}{6}} & \sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \\ \sqrt{\frac{1}{6}} & -\sqrt{\frac{1}{3}} & \sqrt{\frac{1}{2}} \end{pmatrix}, \quad (8)$$

We stress that the discussion above depends very little on the assumption of small representations or the vevs used to do symmetry breaking. One may, for example, use $\langle \mathbf{45}_H \rangle$ proportional to the hypercharge generator or use a $\mathbf{54}_H$ in place of the $\mathbf{45}_H$ to accomplish the breaking from SU(5) to the standard model [12]. Alternatively, we could have used the large representation approach with $\mathbf{10}_H$, $\mathbf{120}_H$, and $\mathbf{126}_H$, which many authors have used for model building [13]. In any case, we still expect a hierarchy in the quark and charged lepton mass matrices. Because of SO(10) relations, this hierarchy should manifest itself in the Dirac neutrino matrix as well and the same naturalness considerations apply.

III. EXPERIMENTAL CONSTRAINTS

The detection of neutrino oscillation is successfully explained by massive neutrinos with nontrivial mixing. We know two mass squared splittings among the neutrinos and two mixing angles of the leptonic mixing matrix, with a limit on the third for the physical light neutrinos [2],

$$\begin{aligned} \tan^2 \theta_{12} &= 0.45 \pm 0.05; \\ \Delta m_{\text{sol}}^2 &= (8.0 \pm 0.3) \times 10^{-5} \text{ eV}^2; \\ \sin^2 2\theta_{23} &= 1.02 \pm 0.04; \\ \Delta m_{\text{atm}}^2 &= (2.5 \pm 0.2) \times 10^{-3} \text{ eV}^2; \\ \sin^2 2\theta_{13} &= 0 \pm 0.05. \end{aligned} \quad (6)$$

Additionally, cosmological considerations place a limit on the total mass of the neutrinos [14], along with limits from tritium beta decay and neutrinoless double beta decay on the electron neutrino [1,2,15]. These experimental results constrain the total mass of the light neutrinos to be less than or of the order of 1 eV. Our discussion does not depend on the exact number since the masses are degenerate in this limit. The bound will only become important to our analysis if it approaches the atmospheric mass splitting.

The mixing is characterized by the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix, a unitary matrix parametrized by three angles and three phases,

neglecting phases. This is in some sense an extreme solution consistent with the data. Given the several seemingly disparate factors which influence the angles, it seems highly unlikely that any model will predict exactly zero for θ_{13} , or exactly maximal atmospheric mixing, unless carefully designed to do so [6]. Therefore it may well be that experiments eventually favor a less striking set of angles. Furthermore, in a detailed model one would also need to carefully consider renormalization, which can have

a significant effect on the mixing angles and mass splittings [17].² We do not address these effects in further detail in this paper because they make little difference in our analysis. We are only looking at relative orders of magnitude of masses and mixing angles. Because of its simple structure, we will use the tribimaximal solution as an experimental input. The critical facts we need are the existence of two large neutrino mixing angles and a relatively weak neutrino mass hierarchy, both of which will remain true despite renormalization effects.

We will assume for now that the tribimaximal structure is generated essentially in the neutrino sector; given the

charged lepton hierarchy, we usually expect relatively small rotations in L_e compared to the large PMNS entries. Since we are only concerned with orders of magnitude, we will (for now) neglect the charged lepton component. As with the geometric hierarchy discussed in Sec. II, there is one exception to this rule associated with a highly asymmetric structure, this time in the charged lepton matrix. Such a lopsided matrix can introduce large rotations, as shown in the Albright-Barr model [7]. This case will be discussed further in Sec. VIII.

The neutrino mass matrix will be diagonalized by the tribimaximal rotations if it has the form

$$M_\nu = V_{\text{PMNS}} D_\nu V_{\text{PMNS}}^T \propto \begin{pmatrix} \left(m_1 + \frac{1}{2}m_2\right) & -\frac{1}{2}(m_1 - m_2) & \frac{1}{2}(m_1 - m_2) \\ -\frac{1}{2}(m_1 - m_2) & \frac{1}{2}\left(\frac{1}{2}m_1 + m_2 + \frac{3}{2}m_3\right) & -\frac{1}{2}\left(\frac{1}{2}m_1 + m_2 - \frac{3}{2}m_3\right) \\ \frac{1}{2}(m_1 - m_2) & -\frac{1}{2}\left(\frac{1}{2}m_1 + m_2 - \frac{3}{2}m_3\right) & \frac{1}{2}\left(\frac{1}{2}m_1 + m_2 + \frac{3}{2}m_3\right) \end{pmatrix}, \quad (9)$$

i.e., $L_\nu = R_\nu = V_{\text{PMNS}}$. The m 's are the physical neutrino masses with an arbitrary phase for m_1 and m_2 . Since we know the two mass squared differences, we may rewrite these in terms of a single mass,

$$m_1 = e^{i\phi_1}|m_1|, \quad m_2 = e^{i\phi_2}\sqrt{|m_1|^2 + \Delta_{\text{sol}}^2}, \quad (10)$$

$$m_3 = \sqrt{|m_1|^2 + \Delta_{\text{sol}}^2 \pm \Delta_{\text{atm}}^2},$$

where we have introduced the notation $\Delta \equiv \sqrt{\Delta m^2}$. The \pm in the definition of m_3 represents the choice of normal (+) or inverted (-) hierarchy. We take the phase factors $e^{i\phi_{1,2}}$ to be ± 1 so that there are just a few choices of relative positive or negative to make. Since we are only concerned with orders of magnitude and this will give the extrema, this should not limit the analysis. Then it is simple to scan through the allowed range of m_1 . By doing this, one can observe the patterns of relative order in the neutrino entries which are consistent with experiment. The potentially interesting possibilities are

(1)

$$M_\nu \sim \begin{pmatrix} \lambda & \lambda & \lambda \\ \lambda & 1 & 1 \\ \lambda & 1 & 1 \end{pmatrix},$$

corresponding to $m_1 \ll m_2 \simeq \Delta_{\text{sol}}$, normal hierarchy.

(2)

$$M_\nu \sim \begin{pmatrix} 0 & \lambda & \lambda \\ \lambda & 1 & 1 \\ \lambda & 1 & 1 \end{pmatrix},$$

²For example, a bimaximal mixing scenario ($\theta_{12}, \theta_{23} = 45^\circ$, $\theta_{13} = 0$) at the GUT scale can produce weak scale mixing angles consistent with the data quoted above [17].

corresponding to $2m_1 \simeq m_2 \simeq \frac{2}{\sqrt{3}}\Delta_{\text{sol}}$, $\phi_2 - \phi_1 = \pi$, normal hierarchy.

(3)

$$M_\nu \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix},$$

corresponding to $\Delta_{\text{sol}}(\Delta_{\text{atm}}) \lesssim m_1 \simeq m_2 \lesssim \Delta_{\text{atm}}(\sqrt{2}\Delta_{\text{atm}})$, $\phi_2 - \phi_1 = 0$, normal (inverted) hierarchy.

(4)

$$M_\nu \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

corresponding to degenerate masses, $\phi_2 = 0$, $\phi_1 = 0$.

(5)

$$M_\nu \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

corresponding to degenerate masses, $\phi_2 = \pi$, $\phi_1 = \pi$.

(6)

$$M_\nu \sim \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix},$$

corresponding to degenerate masses, $\phi_2 - \phi_1 = \pi$. Here $\lambda \equiv \frac{\Delta_{\text{sol}}}{\Delta_{\text{atm}}} \simeq 0.2$ and 0 should be read as at least a few orders of magnitude smaller than 1. Any other possibilities should be roughly an interpolation between those listed and we do not expect them to lead to significant deviations from the results following.

The cases with nondegenerate masses, namely, the first through third above, violate the geometrical hierarchy naturalness limit discussed in Sec. II. In each case the democratic 2-3 block generically leads to two large eigenvalues of order 1 and one large mixing angle. Then the couplings of the first generation give a naive estimate for the third eigenvalue of λ , λ^2 , and 1 for the first, second, and third cases, respectively. This is not compatible with the eigenvalue ranges listed above, so some unexpected cancellations would have to take place. Moreover, these cases are more compatible with a small θ_{12} due to the smallness of all off-diagonal first generation entries. The fourth and fifth cases naturally lead to degenerate eigenvalues as listed but imply unnatural precision to account for the large mixing angles.

In short, hierarchical neutrino masses are unexpected in conjunction with large mixing angles, and large mixing angles naturally proceed from large off-diagonal entries in the effective mass matrix. Thus, case 6 above is the most natural simple assumption to account for the experimental data; it is known as a democratic mass matrix [18].

We can also consider evidence from neutrinoless double beta decay experiments. A positive signal would confirm the Majorana nature of neutrinos and lend credence to seesaw models. The experimental status is controversial: After the Heidelberg-Moscow collaboration set the limit $|m_{ee}| = |(M_\nu)_{11}| < 0.35h$ eV, where h denotes the uncertainty of the nuclear matrix element [2,15], a subset of the collaboration claimed evidence for a signal [19]. Depending on the value of h , this signal points at quasidegenerate neutrino masses in the range 0.1–0.9 eV [2]. This result clearly requires confirmation from current and future experiments. If confirmed, the hierarchical scenarios would be ruled out, consistent with our conclusions from naturalness. However, since this claim is still controversial, we will not rule out the hierarchical scenarios in our analysis.

We note that for an inverted hierarchy with $m_2 \approx \frac{3}{2\sqrt{2}} \Delta_{\text{atm}}$, we could have

$$M_\nu \sim \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}, \quad (11)$$

depending on the phases $\phi_{1,2}$. These should be thought of as special subcases of case 6. As will be shown in the next section, these possibilities will only add additional modeling constraints compared to case 6 without additional explanatory power, so they are not particularly interesting in this context. Bearing these caveats in mind we shall, however, consider some cases besides 6 because they may relax other naturalness constraints.

IV. MODELING

Now we will do a little rearranging of the seesaw formula in terms of the eigenvalues and unitary matrix decomposition of M_D :

$$R_D^\dagger M_R^{-1} R_D^* = D_D^{-1} L_D^\dagger M_\nu L_D^* D_D^{-1}. \quad (12)$$

Applying this to the sixth and henceforth canonical case above, we get

$$R_D^\dagger M_R^{-1} R_D^* \sim \begin{pmatrix} \frac{1}{\eta^2} & \frac{1}{\eta\epsilon} & \frac{1}{\eta} \\ \frac{1}{\eta\epsilon} & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{1}{\eta} & \frac{1}{\epsilon} & 1 \end{pmatrix}, \quad (13)$$

where we have kept only the leading terms. The salient point is that, with the assumption $\mu', \nu', \rho' \leq 1$, the L_D rotations (and similarly the charged lepton rotations) cannot change the orders of the entries. From this we see the apparent double hierarchy for M_R : its eigenvalues naturally scale as $\eta^2, \epsilon^2, 1$ compared to $\eta, \epsilon, 1$ for M_D .

Most of the other cases are similar and retain at least a $\frac{1}{\eta^2}$ ratio between the first and third eigenvalues. For the cases where M_ν has entries less than order one, the unitary rotations can contribute significantly, in particular, they can “fill in” the zero entries, but they cannot make any entries larger than order unity in $L_D^\dagger M_\nu L_D^*$.

There are two cases which may differ importantly from the others. Case 1 in Sec. III is interesting since it yields

$$R_D^\dagger M_R^{-1} R_D^* \sim \begin{pmatrix} \frac{\lambda}{\eta^2} & \frac{1}{\eta\epsilon}(\lambda + \mu'\frac{\eta}{\epsilon}) & \frac{1}{\eta}(\lambda + \mu'\frac{\eta}{\epsilon}) \\ \frac{1}{\eta\epsilon}(\lambda + \mu'\frac{\eta}{\epsilon}) & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{1}{\eta}(\lambda + \mu'\frac{\eta}{\epsilon}) & \frac{1}{\epsilon} & 1 \end{pmatrix}. \quad (14)$$

Similarly, for the second case we get

$$R_D^\dagger M_R^{-1} R_D^* \sim \begin{pmatrix} \frac{1}{\eta^{3/2}}(\lambda + \mu'\frac{\eta}{\epsilon})(\frac{\mu'}{\sqrt{\epsilon}} + \nu') & \frac{1}{\eta\epsilon}(\lambda + \mu'\frac{\eta}{\epsilon}) & \frac{1}{\eta}(\lambda + \mu'\frac{\eta}{\epsilon}) \\ \frac{1}{\eta\epsilon}(\lambda + \mu'\frac{\eta}{\epsilon}) & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{1}{\eta}(\lambda + \mu'\frac{\eta}{\epsilon}) & \frac{1}{\epsilon} & 1 \end{pmatrix}. \quad (15)$$

In these cases we see that we have mitigated the largest ratio of entries from $\frac{1}{\eta^2}$ to a smaller value, although said ratio remains significantly larger than $\frac{1}{\eta}$.

Let us now consider the effects of the matrix R_D on the canonical case. We will show that, under the current assumptions, one can put additional constraints on μ, ν and ρ . To begin, we parametrize the inverse heavy neutrino matrix

$$M_R^{-1} \equiv \begin{pmatrix} A & B & C \\ B & D & E \\ C & E & F \end{pmatrix} \quad (16)$$

and evaluate both Eq. (12) and

$$M_R^{-1} = R_D D_D^{-1} L_D^\dagger M_\nu L_D^* D_D^{-1} R_D^T, \quad (17)$$

which is just another rearrangement of the seesaw formula. Keeping only potentially leading terms, we find

$$\begin{aligned} A &\simeq \frac{1}{\eta^2}, \\ B &\simeq \frac{\mu}{\eta^{3/2}\epsilon^{1/2}} + \frac{1}{\eta\epsilon}, \\ C &\simeq \frac{\nu}{\eta^{3/2}} + \frac{\rho}{\eta\epsilon^{1/2}} + \frac{\mu\nu}{\epsilon^{3/2}} + \frac{1}{\eta}, \\ D &\simeq \frac{\mu^2}{\eta\epsilon} + \frac{2\mu}{\eta^{1/2}\epsilon^{3/2}} + \frac{2\rho\mu}{\eta^{1/2}} + \frac{1}{\epsilon^2}, \\ E &\simeq \frac{\mu\nu}{\eta\epsilon^{1/2}} + \frac{\nu + \rho\mu}{\eta^{1/2}\epsilon} + \frac{\mu}{\eta^{1/2}\epsilon^{1/2}} + \frac{\rho\nu\epsilon^{1/2}}{\eta^{1/2}} + \frac{\rho}{\epsilon^{3/2}} + \frac{1}{\epsilon}, \\ F &\simeq \frac{\nu^2}{\eta} + \frac{2\rho\nu}{\eta^{1/2}\epsilon^{1/2}} + \frac{2\nu}{\eta^{1/2}} + \frac{\rho^2}{\epsilon} + \frac{\rho}{\epsilon^{1/2}} + 1. \end{aligned} \quad (18)$$

Now, with a little consideration, one can see that each entry should only be as big as the rightmost term. This is because Eq. (12) must still be satisfied looking only at the order of the terms. For example, we can look at the equation for the (12) entry of Eq. (13) in terms of A through F and μ , ν , and ρ via Eqs. (3) and (16). This comes out to be

$$\begin{aligned} A\mu\sqrt{\frac{\eta}{\epsilon}} + B + C(\rho\sqrt{\epsilon} + \mu\nu\frac{\eta}{\sqrt{\epsilon}}) + D\mu\sqrt{\frac{\eta}{\epsilon}} \\ + E(\mu\rho + \nu)\sqrt{\eta} + F\rho\nu\sqrt{\eta\epsilon} \sim \frac{1}{\eta\epsilon}. \end{aligned} \quad (19)$$

B appears in this equation with a coefficient of order 1, thus any solution to the set of conditions in Eqs. (18) with $B > \frac{1}{\eta\epsilon}$ will apparently not satisfy Eq. (19).³ This is a naturalness condition. One can, of course, numerically satisfy both equations but it requires a cancellation between two terms to at least an order of magnitude. If we want to avoid the need for a symmetry precisely relating various parameters, the only natural solution is to set $B \sim \frac{1}{\eta\epsilon}$.⁴

³Here the important number is actually the ratio $B/F \sim 1/(\eta\epsilon)$. Using the conventions above we find $F \sim 1$, but there is an overall numerical factor which we omit because it can be absorbed into the dimensionful vevs.

⁴Technically, it could be smaller since Eq. (12) depends on experimental numbers. Thus in Eq. (19), it may cancel the theoretical parameter term $\mu/(\eta^{3/2}\sqrt{\epsilon})$ without fine-tuning as long as it is consistent with the experimentally allowed range. At any rate, it would only make the hierarchy stronger since $A \sim \frac{1}{\eta^2}$ regardless.

Applying the same analysis to the rest of Eqs. (18), we come to the conclusion that

$$M_R^{-1} \sim D_D^{-1} L_D^\dagger M_\nu L_D^* D_D^{-1}, \quad (20)$$

or that the hierarchy of M_R^{-1} could be even stronger, regardless of R_D . Then we must impose constraints on the mixing parameters in Eqs. (18) so that the parameters $B - F$ do not become too large:

$$\mu \lesssim \sqrt{\frac{\eta}{\epsilon}}, \quad \nu \lesssim \sqrt{\eta}, \quad \rho \lesssim \sqrt{\epsilon}. \quad (21)$$

For $\eta \sim 10^{-4}$ and $\epsilon \sim 10^{-2}$, this corresponds to $\mu, \rho \lesssim 10^{-1}$ and $\nu \lesssim 10^{-2}$.

If we take the minimum required suppression and apply it to μ' , ν' , and ρ' as well, we get the cascade hierarchy pattern [10,20] for the Dirac matrix,

$$M_D \sim \begin{pmatrix} \eta & \eta & \eta \\ \eta & \epsilon & \epsilon \\ \eta & \epsilon & 1 \end{pmatrix}. \quad (22)$$

For any hierarchical texture of $R_D^\dagger M_R^{-1} R_D^*$ we will find that M_R^{-1} generally retains the same hierarchy. Intuitively, this is because R_D will tend to smear out any hierarchy in M_R^{-1} ; the larger entries will be rotated into the smaller. The hierarchy would only be sharpened if there were a very precise relation between R_D and M_R^{-1} , which we have no reason to expect. So in general, if $R_D^\dagger M_R^{-1} R_D^*$ has a hierarchy of entries, M_R^{-1} should have at least as strong a hierarchy. Conversely, to maintain a strong hierarchy in $R_D^\dagger M_R^{-1} R_D^*$, the unitary rotations cannot be too far from diagonal, a fact reflected in the constraints on μ , ν and ρ .

For the other possible textures of M_ν with one or two suppressed entries, we mostly find equal or stronger constraints on μ , ν , and ρ . For example, if the (23) and (32) entries of M_ν are small so that the corresponding entries in $D_D^{-1} L_D^\dagger M_\nu L_D^* D_D^{-1}$ are much less than $\frac{1}{\epsilon}$, then we also require $E \ll \frac{1}{\epsilon}$. This in turn imposes stronger constraints on the mixing parameters. This is the situation for cases 3–5 as well as the special subcases of 6 mentioned in Sec. III.

It is interesting that the constraints on μ , ν , and ρ remain valid even if we take the first case of the list,

$$M_\nu \sim \begin{pmatrix} \lambda & \lambda & \lambda \\ \lambda & 1 & 1 \\ \lambda & 1 & 1 \end{pmatrix}, \quad R_D^\dagger M_R^{-1} R_D^* \sim \begin{pmatrix} \frac{\lambda}{\eta^2} & \frac{\lambda}{\eta\epsilon} & \frac{\lambda}{\eta} \\ \frac{\lambda}{\eta\epsilon} & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{\lambda}{\eta} & \frac{1}{\epsilon} & 1 \end{pmatrix}. \quad (23)$$

This is because we retain the strong hierarchy along the first column and row, as well as in the (23) block, whose entries remain less than or equal in order to the first generation entries.

The one exceptional case is the other form noted before, case 2. This leads one to the conclusion

$$M_R^{-1} \sim \begin{pmatrix} \frac{1}{\eta^{3/2}}(\lambda + \mu' \frac{\eta}{\epsilon}) (\frac{\mu'}{\sqrt{\epsilon}} + \nu') & \frac{1}{\eta \epsilon} (\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}}) & \frac{1}{\eta} (\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}}) \\ \frac{1}{\eta \epsilon} (\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}}) & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{1}{\eta} (\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}}) & \frac{1}{\epsilon} & 1 \end{pmatrix}, \quad (24)$$

and the naturalness conditions

$$\mu \simeq \frac{\sqrt{\eta}}{\sqrt{\epsilon} \lambda} \sim 1, \quad \nu \simeq \frac{\sqrt{\eta}}{\lambda} \sim 0.1, \quad \rho \leq \sqrt{\epsilon} \sim 0.1. \quad (25)$$

So in this case we are not as constrained as the cascade pattern but still more constrained than the geometric pattern; only the constraint on ρ remains the same. This makes sense since, in this case, we have a relatively

weak hierarchy in the first row and column compared to the canonical case. Therefore, we find weaker constraints on the rotation parameters for the first generation.

In general then, we are led to both a double (or at least enhanced) hierarchy for M_R and a cascade (or subgeometrical) pattern for M_D in a simple type-I scenario. Other authors have come to similar conclusions following from the assumption of hierarchical Yukawa matrices [21,22].

V. CKM CONSTRAINTS

The Dirac mass matrices of quarks and leptons are related by SO(10) and possibly family symmetries. Thus, we should also consider the size of the unitary rotations in the up and down quark mass matrices, which are measurable through the CKM matrix, $V_{\text{CKM}} \equiv L_u^\dagger L_d$. The experimental CKM values are [1]

$$V_{\text{CKM}} = \begin{pmatrix} 1 & 0.226 \pm 0.002 & [4.3 \pm 0.3] \times 10^{-3} \\ 0.23 \pm 0.01 & 1 & [4.2 \pm 0.06] \times 10^{-2} \\ [7.4 \pm 0.8] \times 10^{-3} & 3.5 \times 10^{-2} & 1 \end{pmatrix}. \quad (26)$$

If we suppose for the moment a geometric pattern for both the up and down quark matrices, then the predicted CKM matrix is

$$V_{\text{CKM}} \simeq \begin{pmatrix} 1 & \sqrt{\frac{m_d}{m_s}} - \sqrt{\frac{m_u}{m_c}} + \sqrt{\frac{m_u m_s}{m_t m_b}} & \sqrt{\frac{m_d}{m_b}} - \sqrt{\frac{m_u}{m_t}} - \sqrt{\frac{m_u m_s}{m_c m_b}} \\ -\sqrt{\frac{m_d}{m_s}} + \sqrt{\frac{m_u}{m_c}} - \sqrt{\frac{m_c m_d}{m_t m_b}} & 1 & \sqrt{\frac{m_s}{m_b}} - \sqrt{\frac{m_c}{m_t}} + \sqrt{\frac{m_u m_d}{m_c m_b}} \\ -\sqrt{\frac{m_d}{m_b}} + \sqrt{\frac{m_u}{m_t}} - \sqrt{\frac{m_c m_d}{m_t m_s}} & -\sqrt{\frac{m_s}{m_b}} + \sqrt{\frac{m_c}{m_t}} - \sqrt{\frac{m_c m_d}{m_t m_s}} & 1 \end{pmatrix} \\ \simeq \begin{pmatrix} 1 & 0.23 - 0.06 + 4 \times 10^{-4} & 0.03 - 0.003 - 0.008 \\ -0.23 + 0.06 - 0.001 & 1 & 0.14 - 0.04 + 0.002 \\ -0.03 + 0.003 - 0.01 & -0.14 + 0.04 - 7 \times 10^{-4} & 1 \end{pmatrix}. \quad (27)$$

A few features are striking. One is that the geometric ratio $\sqrt{\frac{m_d}{m_s}} - \sqrt{\frac{m_u}{m_c}}$ nicely reproduces the experimental value for the first-second generation mixing [23]. The dominant term comes from the down quark mixing, while the contribution from the up quark mixing is significantly too small to account for the mixing by itself. Secondly, the down quark contribution to the first-third mixing is too large by roughly an order of magnitude. Lastly, the down quark contribution to the second-third generation mixing is also too large by roughly a factor of 3. So the geometric hierarchy does a good job for the Cabibbo angle but gives too much mixing with the third generation.

This result is consistent with a partially cascade structure in M_d and M_u .⁵ The relatively large Cabibbo angle

indicates that the down quark matrix should be close to geometrical in the 1-2 block. However, it will fit the data better if it is cascadelike in the third generation. If the same were true of M_D , we would be consistent with the second case from Sec. III. On the other hand, since $\mathbf{16}_i, \mathbf{16}_j, \mathbf{16}_H, \mathbf{16}_H$ only contributes to the down quark and charged lepton matrices, the neutrino matrix could remain completely cascadelike without conflict.

VI. IMPLEMENTING THE CASCADE HIERARCHY

Since we argue that a cascade texture is theoretically desirable, we will investigate how it can be generated. We will make use of the Froggatt-Nielsen mechanism [11] and consider a global $U(1) \times \mathbb{Z}_2 \times \mathbb{Z}'_2$ symmetry. We introduce three SO(10) singlets ϕ_i . The flavor symmetry is broken spontaneously at a high scale m by vevs of the singlet fields, which we expect to be all of the same order, $\langle \phi \rangle$. The symmetry breaking is assumed to be transmitted to quarks and leptons through interactions with heavy particles so that the Yukawa couplings are constructed out of

⁵Since the largest terms come from the down quark sector, the CKM values are also consistent with a geometric hierarchy in M_u . Given SO(10) relations and possible family symmetries, the simplest assumption is that M_u has a similar hierarchy structure to M_d .

powers of $\zeta \equiv \langle \phi \rangle / m$ with a texture dictated by the family symmetry.

We assign the following charges:

Field	$\mathbf{16}_1$	$\mathbf{16}_2$	$\mathbf{16}_3$	$\mathbf{10}_H$	ϕ_1	ϕ_2	ϕ_3
U(1)	2	1	0	0	-1	0	0
\mathbb{Z}_2	-	-	+	+	+	-	+
\mathbb{Z}'_2	-	+	+	+	+	+	-

Then the operator $M_{10}^{ij} \mathbf{16}_i \mathbf{16}_j \mathbf{10}_H$ originates from $\Phi^{ij} \mathbf{16}_i \mathbf{16}_j \mathbf{10}_H$, where Φ represents the higher-dimensional couplings,

$$\Phi = \begin{pmatrix} \frac{1}{m^4} (\phi_1)^4 & \frac{1}{m^4} (\phi_1)^3 \phi_3 & \frac{1}{m^4} (\phi_1)^2 \phi_2 \phi_3 \\ \frac{1}{m^4} (\phi_1)^3 \phi_3 & \frac{1}{m^2} (\phi_1)^2 & \frac{1}{m^2} \phi_1 \phi_2 \\ \frac{1}{m^4} (\phi_1)^2 \phi_2 \phi_3 & \frac{1}{m^2} \phi_1 \phi_2 & 1 \end{pmatrix}, \quad (28)$$

so that

$$M_{10} \sim \begin{pmatrix} \zeta^4 & \zeta^4 & \zeta^4 \\ \zeta^4 & \zeta^2 & \zeta^2 \\ \zeta^4 & \zeta^2 & 1 \end{pmatrix}. \quad (29)$$

This is the cascade form of Eq. (21) with $\eta = \zeta^4$ and $\epsilon = \zeta^2$. The same pattern can easily be reproduced in the other operators which contribute to fermion masses. Note that in the absence of the \mathbb{Z}_2 symmetries we would have generated a geometric hierarchy.

We must also consider whether a cascade hierarchy can naturally accommodate the fermion masses in a unified theory. Restricting ourselves to two generations, the operators discussed in Sec. II contribute to the (normalized) mass matrices as follows:

$$\begin{aligned} M_u &= \begin{pmatrix} \alpha' & \alpha + \beta \\ \alpha - \beta & 1 \end{pmatrix}, \\ M_D &= \begin{pmatrix} \alpha' & \alpha - 3\beta \\ \alpha + 3\beta & 1 \end{pmatrix}, \\ M_d &= \begin{pmatrix} \alpha' + \gamma' & \alpha + \beta + \gamma \\ \alpha - \beta + \gamma & 1 \end{pmatrix}, \\ M_e &= \begin{pmatrix} \alpha' + \gamma' & \alpha - 3\beta + \gamma \\ \alpha + 3\beta + \gamma & 1 \end{pmatrix}. \end{aligned} \quad (30)$$

Here, the terms α and α' parametrize the operator $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H$. The parameter β derives from $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H \mathbf{45}_H$, while γ and γ' characterize $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}'_H$. Looking at the determinants, we calculate the mass ratios:

$$\begin{aligned} \frac{m_c}{m_t} &\simeq |\alpha' - \alpha^2 + \beta^2|, \\ \epsilon &\simeq |\alpha' - \alpha^2 + 9\beta^2|, \\ \frac{m_s}{m_b} &\simeq |\alpha' + \gamma' + \beta^2 - (\alpha + \gamma)^2|, \\ \frac{m_\mu}{m_\tau} &\simeq |\alpha' + \gamma' + 9\beta^2 - (\alpha + \gamma)^2|. \end{aligned} \quad (31)$$

As expected, β accounts for the difference of down quark and charged fermion masses,

$$8\beta^2 = \frac{m_\mu}{m_\tau} \mp \frac{m_s}{m_b} \simeq \begin{cases} 4 \times 10^{-2} & - \\ 8 \times 10^{-2} & + \end{cases} \quad (32)$$

where we used $(m_\mu/m_\tau)_{\text{GUT}} \simeq 0.06$ and $(m_s/m_b)_{\text{GUT}} \simeq 0.02$. Since we wish to minimize off-diagonal terms in a cascadelike matrix, we will use the smaller value for β ,⁶

$$\beta \simeq 7 \times 10^{-2}. \quad (33)$$

Then we obtain

$$\epsilon = \frac{m_c}{m_t} + 8\beta^2 \simeq 7 \times 10^{-2}, \quad (34)$$

with $(m_c/m_t)_{\text{GUT}} \simeq 0.03$.

In order to have a cascade form for M_D , we require $\alpha' \sim \alpha \pm 3\beta \sim \epsilon$. Since $3\beta \simeq 0.2$, this implies $\alpha' \sim 0.1$, independent of α . This value of α' can be consistent with the value of ϵ in Eq. (34), but it needs to cancel significantly with α^2 to ensure a suitably small value for m_c/m_t . Conversely, m_c/m_t implies $\alpha' \lesssim 10^{-2}$, which leads to a geometric hierarchy in M_D . Since we have been trying to avoid requiring the cancellation of theoretical parameters, this simple cascade ansatz is problematic.

One particularly attractive way out of this dilemma is to consider the possibility that M_D , but not M_u , receives additional contributions, e.g., via particular higher-dimensional operators. If such an operator gave a contribution to the (22) element of M_D of order $\epsilon \sim 0.1$, α' could be made sufficiently small. We consider such a scenario in the following section.

VII. NEW CONTRIBUTIONS TO M_D

In Sec. IV we saw that the observed pattern of neutrino masses and mixings leads us to an enhanced hierarchy for M_R , compared to M_D . One should note, however, that while M_D is related to the observed quark and charged lepton hierarchies by SO(10) and any family symmetries, it is not directly observed. In particular, one may include another operator, $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}'_H$. As noted above, the weak doublet in $\mathbf{16}'_H$ can acquire a weak scale vev u' such that this operator potentially contributes to the up quark and neutrino masses. However, it can be constructed to contribute only to the Dirac neutrino matrix. In this case we expect $u' < u$, since u is required to generate a large top quark mass and the sum of the squares of weak scale vevs must equal $(246 \text{ GeV})^2$.

A simple possibility for generating this operator is to integrate out SO(10) singlets, \mathbf{S} , at some scale above the relevant GUT scale vevs. For this purpose we can propose the operators

⁶The larger value, $\beta \simeq 0.1$, leads to $\epsilon \simeq 0.1$.

$$\bar{M}_{ij} \mathbf{16}_i \bar{\mathbf{16}}_H \mathbf{S}_j + \bar{M}'_{ij} \mathbf{16}_i \bar{\mathbf{16}}'_H \mathbf{S}_j + m_s (M_S)_{ij} \mathbf{S}_i \mathbf{S}_j. \quad (35)$$

We assume at least three singlets to guarantee that all three right-handed neutrinos become heavy. As usual, we define m_s to have units of mass so that M_S is dimensionless with entries of order 1 or smaller, and similarly we normalize \bar{M} and \bar{M}' in Eq. (38). In the following analysis we assume that all the \mathbf{S} singlets are integrated out to generate an effective Majorana mass for the N 's. To compute this via a straightforward seesaw mechanism, we will work in the basis where M_S is diagonal and impose the conditions

$$(M_S)_{ii} > \frac{v}{m_s} \quad (36)$$

for all i .

The mass matrix for the electrically neutral particles reads⁷

$$\begin{pmatrix} \nu & N & S \end{pmatrix} \begin{pmatrix} 0 & \frac{1}{2} u M_D & \frac{1}{2} u' \bar{M}' \\ \frac{1}{2} u M_D^T & 0 & \frac{1}{2} v \bar{M} \\ \frac{1}{2} u' \bar{M}'^T & \frac{1}{2} v \bar{M}^T & m_s M_S \end{pmatrix} \begin{pmatrix} \nu \\ N \\ S \end{pmatrix}. \quad (37)$$

As derived in the appendix, the light neutrino mass matrix is then given by

$$\begin{aligned} M_\nu &\simeq M_D (\bar{M}^{-1})^T M_S \bar{M}^{-1} M_D^T \\ &\quad - \frac{x}{2} [\bar{M}' \bar{M}^{-1} M_D^T + M_D (\bar{M}' \bar{M}^{-1})^T], \\ x &\equiv \frac{u' v}{u m_s} \ll 1. \end{aligned} \quad (38)$$

The mass of the heaviest neutrino is of order $u^2 m_s / v^2$. It is crucial that, in the final formula, M_D appears in all terms, i.e., terms quadratic in $\bar{M}' \bar{M}^{-1}$ have not appeared.

Let us study the effect of the new contributions. We parametrize the various matrices as follows:

$$\begin{aligned} M_R^{-1} &= (\bar{M}^{-1})^T M_S \bar{M}^{-1} \equiv \begin{pmatrix} A & B & C \\ B & D & E \\ C & E & F \end{pmatrix}, \\ \bar{M}' \bar{M}^{-1} &= \begin{pmatrix} a & b & c \\ b' & d & e \\ c' & e' & f \end{pmatrix} \end{aligned} \quad (39)$$

(note that the matrix $\bar{M}' \bar{M}^{-1}$ is generally not symmetric), and

$$\begin{aligned} M_D^{-1} M_\nu (M_D^{-1})^T &= M_R^{-1} - \frac{x}{2} [M_D^{-1} \bar{M}' \bar{M}^{-1} \\ &\quad + (\bar{M}' \bar{M}^{-1})^T (M_D^{-1})^T] \\ &= \begin{pmatrix} A' & B' & C' \\ B' & D' & E' \\ C' & E' & F' \end{pmatrix}. \end{aligned} \quad (40)$$

⁷Barr calls this scenario a type-III seesaw mechanism [24]; however, it can also be understood as a product of two type-I mechanisms.

The last matrix, with primed capital letters, is the total effective matrix which takes the place of M_R^{-1} in Sec. IV. The unprimed capital letters parametrize the familiar heavy neutrino matrix and the lower case letters parametrize the new terms. Before proceeding to consider the effects of these new terms, we note that M_R can easily acquire a double hierarchy if it is generated by integrating out heavy singlets, as described above. If \bar{M} has a hierarchy comparable to M_D and M_S is roughly democratic, a double hierarchy occurs naturally.

We can write the total effective parameters in terms of these old and new components and perform the same analysis on the total effective matrix ($A' - F'$) as we did on the simple type-I parameters ($A - F$) in Sec. IV. Then we obtain the following set of equations:

$$\begin{aligned} A' &\simeq A + \left[\frac{a}{\eta} + b' \frac{\mu'}{\sqrt{\eta \epsilon}} + c' \frac{\nu'}{\sqrt{\eta}} \right] x \\ B' &\simeq B + \frac{1}{2} \left[a \frac{\mu}{\sqrt{\eta \epsilon}} + \frac{b}{\eta} + \frac{b'}{\epsilon} + c' \frac{\rho'}{\sqrt{\epsilon}} + d \frac{\mu'}{\sqrt{\eta \epsilon}} + e' \frac{\nu'}{\sqrt{\eta}} \right] x \\ C' &\simeq C + \frac{1}{2} \left[a \frac{\nu}{\sqrt{\eta}} + b' \frac{\rho}{\sqrt{\epsilon}} + c' + \frac{c}{\eta} + e \frac{\mu'}{\sqrt{\eta \epsilon}} + f \frac{\nu'}{\sqrt{\eta}} \right] x \\ D' &\simeq D + \left[b \frac{\mu}{\sqrt{\eta \epsilon}} + \frac{d}{\epsilon} + e' \frac{\rho'}{\sqrt{\epsilon}} \right] x \\ E' &\simeq E + \frac{1}{2} \left[b \frac{\nu}{\sqrt{\eta}} + c \frac{\mu}{\sqrt{\eta \epsilon}} + d \frac{\rho}{\sqrt{\epsilon}} + \frac{e}{\epsilon} + e' + f \frac{\rho'}{\sqrt{\epsilon}} \right] x \\ F' &\simeq F + \left[c \frac{\nu}{\sqrt{\eta}} + e \frac{\rho}{\sqrt{\epsilon}} + f \right] x. \end{aligned} \quad (41)$$

In these equations we have kept only the leading terms. In doing so, we make use of the important fact that the constraints on μ , ν , and ρ still apply. They follow from consideration of the experimental data and the geometric constraints on M_D only.⁸

Although these equations still appear somewhat complicated, the requirement that we fit the same hierarchy of orders as imposed in Eqs. (18) can only be satisfied in a few ways. In general, the new terms give us new parameters which could play a role in a precision fit to the data, but they will not affect the conclusions of this paper unless they dominate over the old terms. Let us consider the canonical case, which implies

$$\begin{pmatrix} A' & B' & C' \\ B' & D' & E' \\ C' & E' & F' \end{pmatrix} \propto \begin{pmatrix} \frac{1}{\eta^2} & \frac{1}{\eta \epsilon} & \frac{1}{\eta} \\ \frac{1}{\eta \epsilon} & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{1}{\eta} & \frac{1}{\epsilon} & 1 \end{pmatrix}. \quad (42)$$

Examining Eq. (41), this puts some initial constraints on our new parameters. For example,

⁸This would not be the case if there were new terms in the effective total matrix which did not involve M_D .

$$\frac{bx}{\eta} \leq B' \sim \frac{F'}{\eta\epsilon}. \quad (43)$$

These constraints may be summarized in matrix form:

$$\begin{pmatrix} a & b & c \\ b' & d & e \\ c' & e' & f \end{pmatrix} \lesssim \begin{pmatrix} \frac{1}{\eta} & \frac{1}{\epsilon} & 1 \\ \frac{1}{\eta} & \frac{1}{\epsilon} & 1 \\ \frac{1}{\eta} & \frac{1}{\epsilon} & 1 \end{pmatrix} \frac{F'}{x}. \quad (44)$$

Taking these restrictions into account, we conclude that to satisfy $\frac{A'}{F'} \sim \frac{1}{\eta^2}$ we must have

$$A \sim F' \frac{1}{\eta^2} \quad \text{or} \quad a \sim F' \frac{1}{\eta x}. \quad (45)$$

The latter case is initially appealing because one can apparently trade the strong double hierarchy constraint on M_R for a weaker standard hierarchy in $\bar{M}'\bar{M}^{-1}$ if the term involving a dominates.

This turns out not to be feasible. Recall that \bar{M} and \bar{M}' have all entries of order 1 or less. Thus, if $a = F' \frac{1}{\eta x}$, there exists some i and some $n \geq 1$ for which

$$\bar{M}'_{i1} = \frac{1}{n} \quad \text{and} \quad \bar{M}'_{i1} = \frac{nF'}{\eta x} \quad (46)$$

[cf. Eq. (39)]. Then the assumption that the a term dominates over A gives us the inequality

$$\begin{aligned} \frac{1}{\eta^2} \sim \frac{ax}{\eta F'} &\geq \frac{A}{F'} = \frac{1}{F'} [(\bar{M}^{-1})^T M_S \bar{M}^{-1}]_{11} \\ &\geq \left(\frac{n}{\eta x}\right)^2 (M_S)_{ii} F', \end{aligned} \quad (47)$$

from which we obtain $(M_S)_{ii} F' \leq \frac{x^2}{n^2}$. Applying the seesaw constraint on M_S and inserting the definition of x gives us

$$\frac{v}{m_s} F' \leq (M_S)_{ii} F' \leq \frac{u^2 v^2}{n^2 u^2 m_s^2}. \quad (48)$$

This requires F' to be too small, that is,

$$1 \lesssim [(\bar{M}^{-1})^T M_S \bar{M}^{-1}]_{33} = F \leq F' \leq \frac{u^2 v}{n^2 u^2 m_s}. \quad (49)$$

Since $v \ll m_s$ and $u' \lesssim u$, this condition cannot be satisfied. Thus the additional contributions cannot dominate over the type-I contributions or change the need for a double hierarchy.

One can instead look at case 1 from Sec. III. If the new terms dominate in the largest ratio, which is still A'/F' , this implies $\bar{M}'_{i1} = \lambda \frac{nF'}{\eta x}$. Proceeding as in the canonical case above, one finds $\lambda (M_S)_{ii} F' \leq \frac{x^2}{n^2}$. Since we require $F' \geq 1$ this is only possible if

$$x^2 \geq (M_S)_{ii} \lambda n^2 \geq \frac{v}{m_s} \lambda n^2, \quad (50)$$

or equivalently,

$$\lambda \leq \frac{u^2 v}{n^2 u^2 m_s} \ll 1. \quad (51)$$

This is a very marginal case since we are relying on $v \ll m_s$ to use the seesaw formula as a valid approximation and $\lambda \sim 0.2$.

If we proceed nonetheless, then we impose the conditions on B' :

$$\frac{\lambda}{\eta\epsilon} \sim \frac{B'}{F'} \geq \frac{B}{F'} \sim \frac{1}{F'} \sum_k \bar{M}_{k1}^{-1} \bar{M}_{k2}^{-1}, \quad (52)$$

which implies the constraint $\bar{M}_{i2}^{-1} \leq x/[n\epsilon(M_S)_{ii}]$. Now we turn to $D' \sim \frac{1}{\epsilon^2}$. By similar reasoning as in the canonical case it can be shown that D must dominate to satisfy D' of the appropriate magnitude, due to the suppression of the new terms by x . Then

$$\frac{1}{\epsilon^2} \sim \frac{D'}{F'} \geq \frac{D}{F'} \sim \frac{(M_S)_{jj}}{F'} (\bar{M}_{j2}^{-1})^2, \quad (53)$$

for some $j \neq i$, which gives us the condition

$$\bar{M}_{j2}^{-1} \sim \frac{1}{\epsilon} \sqrt{\frac{F'}{(M_S)_{jj}}}. \quad (54)$$

This in turn implies

$$\bar{M}_{j1}^{-1} \leq \frac{\lambda}{\eta} \sqrt{\frac{F'}{(M_S)_{jj}}}, \quad (55)$$

so as not to violate the bound on b . We find then that the new contributions can technically dominate in the (11) entry but the type-I terms remain comparable and dominate in other entries, still exhibiting a strong hierarchy compared to M_D .

The related case 2, with $(M_\nu)_{11} \ll \lambda$, is, not surprisingly, similar. One finds that the new term a can dominate if $\lambda \sqrt{\frac{v}{\epsilon}} \leq \frac{u^2 v}{n^2 u^2 m_s}$, which provides somewhat more room for consistency with the seesaw approximation. The constraints on the matrices are

$$\begin{aligned} \bar{M}_{i1}^{-1} &\simeq \frac{\lambda n F'}{x \sqrt{\eta \epsilon}}, & \bar{M}_{i2}^{-1} &\leq \frac{x}{n \sqrt{\eta \epsilon} (M_S)_{ii}}, \\ \bar{M}_{j2}^{-1} &\simeq \frac{1}{\epsilon} \sqrt{\frac{F'}{(M_S)_{jj}}}, & \bar{M}_{j1}^{-1} &\leq \frac{\lambda}{\eta} \sqrt{\frac{F'}{(M_S)_{jj}}}. \end{aligned} \quad (56)$$

In both cases the new terms can dominate in some entries, but the type-I terms remain important and retain a strong, albeit not quite double, hierarchy. We note that this is due largely to the structure of the theory: if M_R is a dimension-five operator generated by integrating out singlets, then a hierarchy in \bar{M} similar to that in M_D naturally leads to a doubled hierarchy in M_R . Because of the suppression of the new terms by v/m_s , M_R will always play an important role.

It is interesting that even with the $\mathbf{16}_i \mathbf{16}_j \overline{\mathbf{16}}_H \overline{\mathbf{16}}_H'$ operator only contributing to the neutrino sector, we still derive the cascade constraints. Although this operator only contributes to the Dirac neutrino matrix, the constraints apply to the operators which generate the up quark matrix. This follows from the precise relations between the higher dimensional operators induced by their common origin. These relations result in M_D appearing in all terms of the formula for M_ν . As a consequence of the persistent cascade constraints, we cannot use the new terms to solve the mass splitting problems discussed in Sec. VI.

If one treats $\mathbf{16}_i \mathbf{16}_j \overline{\mathbf{16}}_H \overline{\mathbf{16}}_H'$ and $\mathbf{16}_i \mathbf{16}_j \overline{\mathbf{16}}_H \overline{\mathbf{16}}_H$ as independent, it is possible to relax said constraints. That is, in the discussion above both operators depend on the coupling $\overline{M}_{ij} \mathbf{16}_i \overline{\mathbf{16}}_H \mathbf{S}_j$ and are therefore related. If we allow them to vary arbitrarily, then the modified seesaw formula in Eq. (38) would have additional terms which did not involve M_D . In effect, we would be adding new terms to the Dirac neutrino matrix which could strongly alter its hierarchy compared to the quarks and charged leptons. If this resulted in a relatively weak Dirac neutrino hierarchy, M_R would have a correspondingly weakened hierarchy and the mixing parameter constraints would also weaken. However, as shown above, this is not necessarily the case when one begins with a more complete theory.

In general, if one can weaken the Dirac neutrino hierarchy without upsetting the charged fermion hierarchies, the requirement of a double hierarchy in M_R and a cascade hierarchy in M_D becomes less restrictive, since they are specified relative to the eigenvalue hierarchy of M_D . One possibility for doing so may be to introduce a vectorlike fourth generation of down quarks and leptons at the GUT scale. This can relate M_D to the down quark hierarchy such that the hierarchy of M_R is similar to that of the up quarks [25].

VIII. LOPSIDED MODELS

Thus far, we have not allowed for any cancellations between terms in our equations, in keeping with our aim to eliminate unnatural models. There are, however, two scenarios where one must be more careful. These are cases where unitary rotations play a very significant role either due to large rotations or small entries in the neutrino matrix.

In this section we will consider the first type of these cases, lopsided models, wherein the operator $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}_H'$ is constructed so as to contribute in a highly asymmetrical way to the down quark and charged lepton mass matrices [7].⁹ These lopsided matrices can yield a natural hierarchy while violating the geometric pattern limit discussed above. Lopsidedness results in large off-

diagonal terms in the unitary rotations on one side of the matrix but not both.

To illustrate these features we will restrict ourselves to two generations first. The following table summarizes the three natural cases we have discussed for a generic matrix M with eigenvalues ϵ and 1, which is diagonalized by the unitary rotation matrices L and R .

Hierarchy	M	L	R
Geometric	$\begin{pmatrix} \epsilon & \sqrt{\epsilon} \\ \sqrt{\epsilon} & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & \sqrt{\epsilon} \\ \sqrt{\epsilon} & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & \sqrt{\epsilon} \\ \sqrt{\epsilon} & 1 \end{pmatrix}$
Cascade	$\begin{pmatrix} \epsilon & \epsilon \\ \epsilon & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & \epsilon \\ \epsilon & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & \epsilon \\ \epsilon & 1 \end{pmatrix}$
Lopsided	$\begin{pmatrix} \epsilon & \epsilon \\ 1 & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & \epsilon \\ \epsilon & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

For both the geometric and cascade cases L and R are similar to each other. As expected, the off-diagonal entries of L and R for the cascade case are smaller than in the geometric case.

The lopsided case, being highly asymmetric, leads to very different rotation matrices on the left and right. We see that to generate large mixing on one side, i.e., R with all entries of the same order, we are led to L being closer to diagonal than in the geometric case. Rather, it is similar to the cascade rotation matrices. So in this simple case, to preserve naturalness, there is a trade off between the left and right sides. If one side's unitary rotation violates the geometric naturalness bound, the other's is concomitantly constrained to be closer to unity.

To take potentially large mixing in the charged lepton sector into account, we have to reevaluate our seesaw formula. In Eq. (12), we neglected the rotations from the charged lepton sector, parametrized by L_e [cf. Eq. (7)]. To include them we rewrite the formula as

$$R_D^\dagger M_R^{-1} R_D^* = D_D^{-1} V_0 M'_\nu V_0^T D_D^{-1}, \quad (57)$$

where

$$M'_\nu \equiv V_{\text{PMNS}} D_\nu V_{\text{PMNS}}^T = L_e^\dagger M_\nu L_e^*, \quad V_0 \equiv L_D^\dagger L_e. \quad (58)$$

M'_ν is the light neutrino mass matrix in the basis where the charged leptons are diagonal. It can have the same forms as discussed in Sec. III for M_ν . With the substitutions $M_\nu \rightarrow M'_\nu$ and $L_D \rightarrow V_0$, the equations used above are unaltered.

The crucial difference is that the assumed form of L_D in Eq. (3) does not necessarily apply to V_0 in the lopsided case. Since V_0 contains off-diagonal entries of order one, we may arrange for terms of equal order to cancel each other in $V_0 M'_\nu V_0^T$. This is not fine-tuning because we are, in effect, canceling an experimental term with a theoretical one, rather than canceling two theoretical parameters against each other. To put it another way, we are simply using a theoretical term to generate an experimental parameter of the same order. The result is that we may be able

⁹We will not discuss the origin of these lopsided matrices, which, e.g., can be due to family symmetries [7].

to have a form for $R_D^\dagger M_R^{-1} R_D^*$ which does not have such a strong hierarchy, and which in turn may not imply the restrictive cascade form for M_D . In such a scenario, some or all of the large mixing in V_{PMNS} comes from charged lepton unitary rotations.

To examine the lopsided case further we must see what can be said about the matrix V_0 . Again, we can look at the CKM matrix for possible constraints. It can tell us about the potential lopsidedness in the down quark and charged lepton mass matrices. The operator we are using to generate lopsidedness contributes to M_e as the transpose of its contribution to M_d , as is familiar from SU(5) models.¹⁰ Hence, large rotations in L_e would coincide with large rotations in R_d and vice versa. We now see that the experimental values are consistent with either a cascade structure or a lopsided structure for the down quark mass matrix in the third generation couplings.

One might hope that the relatively large 1-2 mixing, which is consistent with a geometric hierarchy in the down quark matrix (cf. Sec. V), would constrain the 1-2 mixing in R_d . This, however, turns out not to be the case. We can construct a matrix with all the desired features and generically large right-handed mixing, e.g.,

$$\begin{aligned}
 M_d &\sim \begin{pmatrix} \frac{m_d}{m_b} & \frac{\sqrt{m_d m_s}}{m_b} & \frac{m_d}{m_b} \\ \frac{\sqrt{m_d m_s}}{m_b} & \frac{m_s}{m_b} & \frac{m_s}{m_b} \\ 1 & 1 & 1 \end{pmatrix}, \\
 L_d &\sim \begin{pmatrix} 1 & \sqrt{\frac{m_d}{m_s}} & \frac{\sqrt{m_d m_s}}{m_b} \\ \sqrt{\frac{m_d}{m_s}} & 1 & \frac{m_s}{m_b} \\ \frac{\sqrt{m_d m_s}}{m_b} & \frac{m_s}{m_b} & 1 \end{pmatrix}, \\
 R_d &\sim \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}.
 \end{aligned} \tag{59}$$

Thus, although the CKM matrix is highly suggestive of either a partially cascade or lopsided form for the down quark mass matrix, it is difficult to constrain the form of R_d and its counterpart L_e in the latter case.

On the other hand, we note that since V_{PMNS} has a small value for the (13) entry, naturalness requires that at least one entry in the column L_e^{i1} be correspondingly small. This suggests that we can rule out the extreme lopsided case shown above.

Lopsidedness also modifies the eigenvalue fitting we did in Sec. VI. Let us consider the case where $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}'_H$ is lopsided and assume that it contributes to only one off-diagonal entry in M_d and M_e in a significant way. Then the mass matrices in Eq. (30) are modified to

¹⁰This is simply due to the fact that $\mathbf{16}_H$ breaks SO(10) to SU(5), so $\mathbf{16}_i \mathbf{16}_j \mathbf{16}_H \mathbf{16}'_H$ is basically an SU(5) Yukawa operator for down quarks and charged fermions, suppressed by ν/M .

$$\begin{aligned}
 M_d &= \begin{pmatrix} \alpha' + \gamma' & \alpha + \beta \\ \alpha - \beta + \gamma & 1 \end{pmatrix}, \\
 M_e &= \begin{pmatrix} \alpha' + \gamma' & \alpha - 3\beta + \gamma \\ \alpha + 3\beta & 1 \end{pmatrix},
 \end{aligned} \tag{60}$$

with the corresponding eigenvalues

$$\begin{aligned}
 \frac{m_s}{m_b} &= \left| \frac{\alpha' + \gamma' + \beta^2 - \alpha^2 - \gamma(\alpha + \beta)}{1 + \gamma^2} \right|, \\
 \frac{m_\mu}{m_\tau} &= \left| \frac{\alpha' + \gamma' + 9\beta^2 - \alpha^2 - \gamma(\alpha + 3\beta)}{1 + \gamma^2} \right|.
 \end{aligned} \tag{61}$$

This yields

$$\frac{m_\mu}{m_\tau} - \frac{m_s}{m_b} = \frac{2\beta(4\beta - \gamma)}{1 + \gamma^2} \xrightarrow{\gamma \sim 1} \beta \sim 4 \times 10^{-2}, \tag{62}$$

which is only a slight improvement over the symmetric case, a cascade structure in M_D is still inconsistent with the charged fermion hierarchies. Thus, in the absence of additional contributions to the mass matrices, it seems we must rely on large, lopsided mixing between the second and third generations to alleviate the need for a cascade structure in the 2-3 block of M_D .

IX. SMALL ENTRIES AND MIXING

Aside from lopsided matrices, there is another scenario in which V_0 can play an important role. We saw in Sec. IV that the entries of the first row and column of $R_D^\dagger M_R^{-1} R_D^*$ are smaller in the cases 1 and 2. If we allow cancellations between these entries of order λ and the mixing parameter $\mu' \sqrt{\frac{\eta}{\epsilon}}$, we might expect some qualitatively different results. In this more general case, we use μ' , ν' and ρ' to parametrize V_0 rather than L_D . Including the effects of L_e , we no longer have the symmetry constraints $(\mu, \nu, \rho) \sim (\mu', \nu', \rho')$.

We consider case 1:

$$\begin{aligned}
 R_D^\dagger M_R^{-1} R_D^* &\sim \begin{pmatrix} \frac{\lambda}{\eta^2} & \frac{1}{\eta \epsilon} \left(\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}} \right) & \frac{1}{\eta} \left(\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}} \right) \\ \frac{1}{\eta \epsilon} \left(\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}} \right) & \frac{1}{\epsilon^2} & \frac{1}{\epsilon} \\ \frac{1}{\eta} \left(\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}} \right) & \frac{1}{\epsilon} & 1 \end{pmatrix}.
 \end{aligned}$$

Here, although the unitary rotations remain relatively close to unity, the rotation parameter $\mu' \sqrt{\frac{\eta}{\epsilon}}$ may be large enough to cancel the experimental term λ . Such cancellation is only possible if $\mu' \sim 1$.¹¹ Under geometrical constraints,

¹¹Since μ' includes contributions from L_e , its coefficient $\sqrt{\eta/\epsilon}$ should be $\sqrt{m_e/m_\mu}$ if the charged lepton ratio is larger. However, since $\sqrt{m_e/m_\mu} \sim 0.1 \sim \sqrt{\eta/\epsilon}$ under our assumptions, we keep our familiar notation.

the (11) entry will be $\frac{\lambda}{\eta^2}$, while the (12) entry could be much smaller than $\frac{\lambda}{\eta\epsilon}$. One can proceed to analyze the mixing parameters in R_D as in Sec. IV. Because of the relatively large (11) entry, one finds that the constraints

$$\mu \lesssim \sqrt{\frac{\eta}{\epsilon}} \left(1 + \frac{\mu'}{\lambda} \sqrt{\frac{\eta}{\epsilon}}\right), \quad \nu \lesssim \sqrt{\nu'} \left(1 + \frac{\mu'}{\lambda} \sqrt{\frac{\eta}{\epsilon}}\right), \quad \rho \lesssim \sqrt{\epsilon} \quad (63)$$

are required to preserve the small (12) and (13) entries. Since this requires $\mu \ll \mu' \sim 1$, the charged lepton rotations would have to be significantly larger than those from the Dirac neutrino matrix, at least for the 1-2 mixing. This would suggest an approximately geometric structure in the charged lepton matrix and a Dirac neutrino matrix with very small first generation mixing. Thus, the Dirac neutrino matrix would have a more restricted form than the cascade hierarchy derived for the simpler case without cancellations. Unless some additional information prompts us to favor these textures for M_ν , M_D and M_e , there is no compelling reason to further pursue this route.

In the second case, where $(M_\nu)_{11} \sim 0$, we find that the constraints are the same as those listed in Eq. (25), i.e., the same as we found for this case without allowing for cancellations. These results hold because, regardless of how small $\lambda + \mu' \sqrt{\frac{\eta}{\epsilon}}$ may be, we retain the same relative hierarchy between the first generation entries and the same hierarchy in the 2-3 block, cf. Eq. (15).

We conclude that these potential cancellations have little effect on our previous considerations.

X. OUTLOOK

Barring cancellations or additional flavor symmetries, the observed pattern of neutrino mass splittings and mixing angles leads us to two related propositions for simple model building in the general context of a grand-unified theory with type-I seesaw mechanism. The first is a double hierarchy, with respect to the hierarchy of the Dirac matrix, M_D , in the effective heavy neutrino matrix M_R . The second, contingent upon the first, is a cascade structure in M_D , or a texture which is even closer to diagonal. These conclusions follow only from the structure of the type-I seesaw formula, together with the observation that the experimental neutrino data most naturally arise from an approximately democratic effective light neutrino matrix. If the neutrino masses obey a normal hierarchy, i.e., $m_1 \lesssim m_2 \sim \sqrt{\Delta m_{\text{sol}}^2} \ll m_3 \sim \sqrt{\Delta m_{\text{atm}}^2}$, it is possible to relax these constraints, but it remains true that M_R should have an enhanced hierarchy and M_D should have a subgeometrical structure. Moreover, in this case some approximate symmetry must exist to generate a second large mixing angle and a hierarchy consistent with experiment.

These conclusions are rather general and not restricted to the specific model with small representations outlined in

Sec. II. They hold for hierarchical, symmetric matrices, up to factors of order one. In light of the quark and charged lepton mass hierarchies, it is natural for M_D to be hierarchical. In particular, this matrix is closely related to the up quark matrix in many GUT models. Family symmetries will also tend to engender such relations. In Sec. VII we showed that even adding an operator which ostensibly only contributes to the Dirac neutrino matrix does not necessarily relax our conclusions.

Can we implement these textures in a complete model? We discussed a scenario with a $U(1) \times \mathbb{Z}_2 \times \mathbb{Z}_2$ flavor symmetry, where we generated a cascade structure for the Dirac matrices through the Froggatt-Nielsen mechanism. A double hierarchy in M_R is natural if it is an effective operator generated by integrating out singlets coupled to $\mathbf{16}_i \mathbf{16}_H$, where this coupling has an eigenvalue hierarchy similar to that in M_D (cf. Ref. [21]). However, this structure led to problems in the quark sector. We have seen that the relatively large Cabibbo angle implies that the down quark matrix is not purely cascadelike, although a cascade structure in the third generation is supported. This does not necessarily conflict with a fully cascade pattern in M_D , but it requires a somewhat more complicated picture than the simple model described above. Furthermore, in our specific model, we rely on the antisymmetric operator $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H \mathbf{45}_H$ to differentiate the down quark and charged lepton matrices. This implies that its contributions cannot be too small. Since it also contributes to the up quark and neutrino matrices, it becomes difficult to reconcile a cascade structure in these matrices with the strong up quark hierarchy in a natural way.

Lopsided models may provide us a way out of these potential difficulties. Compared with a cascade pattern, they are equally compatible with the CKM matrix. For the purposes of mass fitting, lopsidedness slightly relaxes the need for large off-diagonal contributions from $\mathbf{16}_i \mathbf{16}_j \mathbf{10}_H \mathbf{45}_H$. More importantly, a lopsided charged lepton matrix introduces large rotations which contribute to the PMNS matrix. If these are primarily responsible for one or both of the large mixing angles, it is possible to reduce the pull towards a double hierarchy in M_R . This in turn can relax the constraints that lead us to a cascade structure for M_D and so for M_u . Exactly how much lopsidedness can obviate the need for a double hierarchy remains an open question. The atmospheric mass splitting remains small compared to the quark mass splittings, irrespective of the origin of the large mixing angles. This will tend to require an enhanced hierarchy in at least part of M_R . Additionally, while it is technically possible that most or all of the PMNS structure comes from charged lepton rotations, we must ask how much can be done in a natural way. For example, as discussed at the end of Sec. VIII, a small value for θ_{13} precludes generically large mixing from lopsidedness in all generations.

This brings us to the nature and origin of θ_{13} in general, which we have not addressed in detail in this paper. We

chose to leave this an open question in light of the current uncertainty in the size of θ_{13} ; only an upper bound is known. While it is clear that the solar and atmospheric mixing angles are large compared with those in the quark sector, θ_{13} may or may not be comparatively small. Actually, the experimental upper limit, approximately 10° , is of the same order as the Cabibbo angle. This is large enough that its smallness compared to the other neutrino angles may be explained by normal fluctuations of order one parameters without violating our sense of naturalness [6]. However, if θ_{13} is significantly closer to zero we should seek some more robust explanation. For the forms of M_ν listed in Sec. III, this would require a symmetry closely relating various matrix elements. Another possibility arises for partially lopsided matrices: if one large mixing angle arises from the charged lepton sector and the other from M_ν , then it is natural to preserve a small third angle. Clearly, it is important to determine the order of θ_{13} .

In summary, a combination of partially lopsided and partially cascade matrices, in conjunction with an enhanced hierarchy in M_R , seems to be the most natural route to explain the generic features of the quark and lepton data in a grand-unified model. The details of a complete model remain to be worked out, but our conclusions follow from a fairly general framework. It will be interesting to see if a workable model can be obtained with relatively simple family symmetries and what consequences there might be for experimental predictions.

ACKNOWLEDGMENTS

We would like to thank S. Willenbrock for useful comments on the manuscript. This work was supported in part by the U.S. Department of Energy under Contract No. DE-FG02-91ER40677, as well as the Sonderforschungsbereich Transregio 9 Computergestützte Theoretische Teilchenphysik of the Deutsche Forschungsgemeinschaft.

APPENDIX: DERIVATION OF EXPANDED SEESAW FORMULA

In this appendix, we derive the extended seesaw formula, given in Eq. (38). As mentioned in Sec. VII, it is crucial that terms quadratic in $\bar{M}'\bar{M}^{-1}$ do not appear. The

formula was originally derived in Ref. [24] through a slightly different calculation.

As displayed in Eq. (35), we propose the operators

$$W_S = \bar{M}_{ij} \mathbf{16}_i \bar{\mathbf{16}}_H \mathbf{S}_j + \bar{M}'_{ij} \mathbf{16}_i \bar{\mathbf{16}}'_H \mathbf{S}_j + m_s (M_S)_{ij} \mathbf{S}_i \mathbf{S}_j.$$

We integrate out the singlet fields, \mathbf{S} , by taking a partial derivative and setting it equal to zero,

$$\frac{\partial W_S}{\partial \mathbf{S}_j} \equiv 0; \tag{A1}$$

$$\mathbf{S}_i = -\frac{1}{2m_s} [\bar{M}_{ij} \mathbf{16}_i \bar{\mathbf{16}}_H + \bar{M}'_{ij} \mathbf{16}_i \bar{\mathbf{16}}'_H] (M_S^{-1})_{jk}.$$

Plugging this into our initial equation yields

$$W_S^{\text{eff}} = -\frac{1}{4m_s} \mathbf{16}_i [(\bar{M} M_S^{-1} \bar{M}^T)_{ij} \bar{\mathbf{16}}_H \bar{\mathbf{16}}_H + (\bar{M}' M_S^{-1} \bar{M}'^T)_{ij} \bar{\mathbf{16}}'_H \bar{\mathbf{16}}'_H + 2(\bar{M}' M_S^{-1} \bar{M}'^T)_{ij} \bar{\mathbf{16}}_H \bar{\mathbf{16}}'_H] \mathbf{16}_j. \tag{A2}$$

Now we let the Higgs fields acquire their GUT and weak scale vevs and we include the Dirac term $\nu M_D N$, where ν and N are the left- and right-handed neutrinos, respectively. Suppressing the generation indices, we obtain

$$W_N = -\frac{1}{4m_s} [\nu^2 N (\bar{M} M_S^{-1} \bar{M}^T) N + u^2 \nu (\bar{M}' M_S^{-1} \bar{M}'^T) \nu + 2u' \nu \nu (\bar{M}' M_S^{-1} \bar{M}'^T) N] + u \nu M_D N. \tag{A3}$$

We extremize with respect to N and find

$$\frac{\partial W_N}{\partial N} \equiv 0; \tag{A4}$$

$$N = \left[\frac{um_s}{v^2} M_D (\bar{M}^{-1})^T M_S \bar{M}^{-1} - \frac{u'}{v} \bar{M}' \bar{M}^{-1} \right] \nu.$$

Inserting this into the last equation and performing a little algebra gives the amended seesaw formula

$$W_\nu^{\text{eff}} \simeq \left\{ M_D (\bar{M}^{-1})^T M_S \bar{M}^{-1} M_D^T - \frac{1}{2} \frac{u'v}{um_s} [\bar{M}' \bar{M}^{-1} M_D^T + M_D (\bar{M}' \bar{M}^{-1})^T] \right\} \frac{u^2 m_s}{v^2}. \tag{A5}$$

[1] W. M. Yao *et al.* (Particle Data Group), *J. Phys. G* **33**, 1 (2006).
 [2] A. Strumia and F. Vissani, arXiv:hep-ph/0606054.
 [3] P. Minkowski, *Phys. Lett.* **67B**, 421 (1977).
 [4] H. Georgi, in *Particles and Fields*, edited by C. Carlson, AIP Conf. Proc. No. 23 (AIP, New York, 1975), p. 575; H. Fritzsch and P. Minkowski, *Ann. Phys. (N.Y.)* **93**, 193

(1975).
 [5] R. N. Mohapatra and A. Y. Smirnov, *Annu. Rev. Nucl. Part. Sci.* **56**, 569 (2006); G. Altarelli, arXiv:0705.0860.
 [6] G. Altarelli and F. Feruglio, *New J. Phys.* **6**, 106 (2004).
 [7] C. H. Albright and S. M. Barr, *Phys. Rev. D* **62**, 093008 (2000); **64**, 073010 (2001).
 [8] See, e.g., K. S. Babu, J. C. Pati, and F. Wilczek, *Nucl.*

- Phys. **B566**, 33 (2000); R. Dermisek and S. Raby, Phys. Rev. D **62**, 015007 (2000); T. Blazek, S. Raby, and K. Tobe, Phys. Rev. D **62**, 055001 (2000); J. Sayre and S. Wiesenfeldt, Phys. Lett. B **637**, 295 (2006).
- [9] K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **74**, 2418 (1995).
- [10] I. Dorsner and S. M. Barr, Nucl. Phys. **B617**, 493 (2001).
- [11] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. **B147**, 277 (1979).
- [12] G. Anderson, S. Raby, S. Dimopoulos, L. J. Hall, and G. D. Starkman, Phys. Rev. D **49**, 3660 (1994); S. Wiesenfeldt and S. Willenbrock, Phys. Lett. B **661**, 268 (2008).
- [13] For a recent brief review, see A. Melfo, AIP Conf. Proc. **917** 252 (2007).
- [14] J. Lesgourgues and S. Pastor, Phys. Rep. **429**, 307 (2006); S. Hannestad and G. G. Raffelt, J. Cosmol. Astropart. Phys. **11** (2006) 016.
- [15] L. Baudis *et al.*, Phys. Rev. Lett. **83**, 41 (1999); H. V. Klapdor-Kleingrothaus *et al.*, Eur. Phys. J. A **12**, 147 (2001).
- [16] P. F. Harrison, D. H. Perkins, and W. G. Scott, Phys. Lett. B **530**, 167 (2002).
- [17] S. Antusch, J. Kersten, M. Lindner, M. Ratz, and M. A. Schmidt, J. High Energy Phys. **03** (2005) 024.
- [18] H. Harari, H. Haut, and J. Weyers, Phys. Lett. **78B**, 459 (1978); L. Lavoura, Phys. Lett. B **228**, 245 (1989); H. Fritzsch and J. Plankl, Phys. Lett. B **237**, 451 (1990).
- [19] H. V. Klapdor-Kleingrothaus and I. V. Krivosheina, Mod. Phys. Lett. A **21**, 1547 (2006).
- [20] G. Altarelli, F. Feruglio, and I. Masina, Phys. Lett. B **472**, 382 (2000).
- [21] R. Dermisek, Phys. Rev. D **70**, 073016 (2004).
- [22] J. A. Casas, A. Ibarra, and F. Jimenez-Alburquerque, J. High Energy Phys. **04** (2007) 064.
- [23] H. Fritzsch, Phys. Lett. **70B**, 436 (1977).
- [24] S. M. Barr, Phys. Rev. Lett. **92**, 101601 (2004).
- [25] Y. Nomura and T. Yanagida, Phys. Rev. D **59**, 017303 (1998); T. Asaka, Phys. Lett. B **562**, 291 (2003); W. Buchmuller, L. Covi, D. Emmanuel-Costa, and S. Wiesenfeldt, J. High Energy Phys. **12** (2007) 030.