

Magnetic moments of heavy Ξ_Q baryons in light cone QCD sum rules

T. M. Aliev,^{*} K. Azizi,⁺ and A. Ozpineci[‡]

Physics Department, Middle East Technical University, 06531, Ankara, Turkey

(Received 31 March 2008; published 9 June 2008)

The magnetic moments of heavy Ξ_Q baryons containing a single charm or bottom quark are calculated in the framework of the light cone QCD sum rules method. A comparison of our results with the predictions of other approaches, such as relativistic and nonrelativistic quark models, the hypercentral model, chiral perturbation theory, and soliton and skyrmion models, is presented.

DOI: [10.1103/PhysRevD.77.114006](https://doi.org/10.1103/PhysRevD.77.114006)

PACS numbers: 11.55.Hx, 13.40.Em, 14.20.Lq, 14.20.Mr

I. INTRODUCTION

During the last few years, exciting experimental results were obtained in the baryon sector containing a single b quark. The CDF Collaboration observed the states Σ_b^{\pm} and $\Sigma_b^{* \pm}$ [1], while both D0 [2] and CDF [3] Collaborations have seen Ξ_b . Recently, *BABAR* Collaboration reported the discovery of Ω_c^* with mass splitting $m_{\Omega_c^*} - m_{\Omega_c} = (70.8 \pm 1.0 \pm 1.1)$ MeV [4].

The masses of the heavy baryons have been studied in the framework of various phenomenological models [5–13] and also in the framework of the QCD sum rules method [14–26]. Along with their masses, another static parameter of the heavy baryons is their magnetic moment. Study of the magnetic moments can give valuable information about the internal structures of hadrons.

The magnetic moments of heavy baryons have been studied in the framework of different methods. In [27,28] the magnetic moments of charmed baryons are calculated within the naive quark model. In [29,30], magnetic moments of charmed and bottom baryons are analyzed in the quark model, and in [31] heavy baryon magnetic moments are studied in the bound state approach. Magnetic moments of heavy baryons are calculated in the relativistic three-quark model [32], the hypercentral model [33], the chiral perturbation model [34], the soliton model [35], the skyrmion model [36], and the nonrelativistic constituent quark model with light and strange $\bar{q}q$ pairs [37]. In [38] the magnetic moments of Σ_c and Λ_c baryons are calculated with the QCD sum rules method in an external electromagnetic field. In [39,40], the light cone QCD sum rules method is applied to study the magnetic moments of the Λ_Q ($Q = c, b$) and $\Sigma_Q \Lambda_Q$ transition magnetic moments (more about this method can be found in [41–44] and references therein).

The aim of the present work is to calculate the magnetic moments of the Ξ_b baryons that were recently observed by D0 and CDF Collaborations within the light cone QCD sum rules framework. The plan of the paper is as follows.

In Sec. II, using the general form of the baryon current, the light cone QCD sum rules for Ξ_b and Ξ_c baryons are calculated. In Sec. III we present our numerical calculations on the Ξ_b and Ξ_c baryons. In this section we also present a comparison of our results with the predictions of other approaches.

II. LIGHT CONE QCD SUM RULES FOR THE Ξ_Q MAGNETIC MOMENTS

In order to calculate the magnetic moments of Ξ_Q ($Q = b, c$) in the framework of the light cone QCD sum rules, we need the expression for the interpolating current of Ξ_Q . To construct it, we follow [11]; i.e. we assume that the strange and light quarks in Ξ_Q are in a relative spin zero state (scalar or pseudoscalar diquarks). Therefore, the most general current without derivatives and with the quantum numbers of Ξ_Q can be constructed from the combination of the aforementioned scalar or pseudoscalar diquarks in the following way:

$$\eta_Q = \varepsilon_{abc}[(q^{aT} C s^b) \gamma_5 + \beta(q^{aT} C \gamma_5 s^b)] Q^c. \quad (1)$$

Here a, b , and c are color indices; C is the charge conjugation operator; $Q = b$ or c ; $q = u$ or d ; and β is an arbitrary parameter. Having the explicit expression for the interpolating current, our next task is to construct light cone QCD sum rules for the magnetic moments of Ξ_Q baryons. They are constructed from the following correlation function:

$$\Pi(p, q) = i \int d^4 x e^{ipx} \langle \gamma | T \{ \eta_Q(x) \bar{\eta}_Q(0) \} | 0 \rangle. \quad (2)$$

The calculation of the phenomenological side at the hadronic level proceeds by inserting into the correlation function a complete set of hadronic states with the quantum numbers of Ξ_Q . We get

$$\begin{aligned} \Pi &= \sum_i \frac{\langle 0 | \eta_Q | \Xi_{Q_i}(p_2) \rangle \langle \Xi_{Q_i}(p_2) | \\ &| \Xi_{Q_i}(p_1) \rangle_\gamma \frac{\langle \Xi_{Q_i}(p_1) | \bar{\eta}_Q | 0 \rangle}{p_1^2 - m_{\Xi_Q}^2}. \end{aligned} \quad (3)$$

Isolating the ground state's contributions, Eq. (3) can be

^{*}taliev@metu.edu.tr

⁺e146342@metu.edu.tr

[‡]ozpineci@metu.edu.tr

written as

$$\begin{aligned} \Pi &= \frac{\langle 0 | \eta_Q | \Xi_Q(p_2) \rangle \langle \Xi_Q(p_2) | \Xi_Q(p_1) \rangle_\gamma}{p_2^2 - m_{\Xi_Q}^2} \\ &\times \frac{\langle \Xi_Q(p_1) | \bar{\eta}_Q | 0 \rangle}{p_1^2 - m_{\Xi_Q}^2} \\ &+ \sum_{h_i} \frac{\langle 0 | \eta_{Qi} | h_i(p_2) \rangle \langle h_i(p_2) | h_i(p_1) \rangle_\gamma}{p_2^2 - m_{h_i}^2} \\ &\times \frac{\langle h_i(p_1) | \bar{\eta}_Q | 0 \rangle}{p_1^2 - m_{h_i}^2}, \end{aligned} \quad (4)$$

where $p_1 = p + q$, $p_2 = p$, and q is the photon momentum. The second term in Eq. (4) describes the higher resonances and continuum contributions. The coupling of the interpolating current with the baryons Ξ_Q is determined as

$$\langle 0 | \eta_Q | \Xi_Q(p) \rangle = \lambda_Q u_{\Xi_Q}(p), \quad (5)$$

where $u_{\Xi_Q}(p)$ is a spinor describing the baryon Ξ_Q with four-momentum p , and λ_Q is the corresponding residue.

The last step for obtaining the expression for the physical part of the correlator function is to write down the matrix element $\langle \Xi_Q(p_2) | \Xi_Q(p_1) \rangle_\gamma$ in terms of the form factors. Using Lorentz covariance, this matrix element can be written as

$$\begin{aligned} &\langle \Xi_Q(p_1) | \Xi_Q(p_2) \rangle_\gamma \\ &= \varepsilon^\mu \bar{u}_{\Xi_Q}(p_1) \left[f_1 \gamma_\mu - i \frac{\sigma_{\mu\alpha} q_\alpha}{2m_{\Xi_Q}} f_2 \right] u_{\Xi_Q}(p_2) \\ &= \bar{u}_{\Xi_Q}(p_1) \left[(f_1 + f_2) \gamma_\mu + \frac{(p_1 + p_2)_\mu}{2m_{\Xi_Q}} f_2 \right] u_{\Xi_Q}(p_2) \varepsilon^\mu, \end{aligned} \quad (6)$$

where $f_1(q^2)$ and $f_2(q^2)$ are the form factors and ε^μ is the photon polarization vector.

For the calculation of the Ξ_Q magnetic moments, only the values of the form factors at $q^2 = 0$ are needed because the photon is real in our problem. Using Eqs. (4)–(6) for the physical part of the correlator and summing over the spins of initial and final Ξ_Q baryons, the correlation function becomes

$$\begin{aligned} \Pi &= -\lambda_Q^2 \varepsilon^\mu \frac{\not{p}_2 + m_{\Xi_Q}}{p_2^2 - m_{\Xi_Q}^2} \left[(f_1 + f_2) \gamma_\mu + \frac{(p_1 + p_2)_\mu}{2m_{\Xi_Q}} f_2 \right] \\ &\times \frac{\not{p}_1 + m_{\Xi_Q}}{p_1^2 - m_{\Xi_Q}^2}. \end{aligned} \quad (7)$$

From this expression, we see that there are various structures which can be chosen for studying the magnetic moments of Ξ_Q . In the present work, following [45], we choose the structure $\not{p}_2 \not{\not{q}}$ which contains the magnetic form factor $f_1 + f_2$, and at $q^2 = 0$ it gives the magnetic moment of Ξ_Q in units of $e\hbar/2m_{\Xi_Q}$. Choosing this struc-

ture in the physical part of the correlator, for the magnetic moments of Ξ_Q we obtain

$$\Pi = -\lambda_Q^2 \frac{1}{p_1^2 - m_{\Xi_Q}^2} \mu_{\Xi_Q} \frac{1}{p_2^2 - m_{\Xi_Q}^2}, \quad (8)$$

where $\mu_{\Xi_Q} = (f_1 + f_2)|_{q^2=0}$ are the magnetic moments of Ξ_Q in units of $e\hbar/2m_{\Xi_Q}$.

In order to calculate the magnetic moments of Ξ_Q baryons, the expression of the theoretical part of the correlation function is needed. After simple calculations for the theoretical part of the correlation function in QCD, we obtain

$$\begin{aligned} \Pi &= -i \varepsilon_{abc} \varepsilon_{a'b'c'} \int d^4x e^{ipx} \langle \gamma(q) | \{ \gamma_5 S_Q^{cc'} \gamma_5 \text{Tr}(S_q^{ba'} S_s^{ab'}) \\ &+ \beta \gamma_5 S_Q^{cc'} \text{Tr}(S_q^{ba'} \gamma_5 S_s^{ab'}) + \beta S_Q^{cc'} \gamma_5 \text{Tr}(\gamma_5 S_q^{ba'} S_s^{ab'}) \\ &+ \beta^2 S_Q^{cc'} \text{Tr}(\gamma_5 S_s^{ab'} \gamma_5 S_q^{ba'}) \} | 0 \rangle \end{aligned} \quad (9)$$

where $S'_i = CS_i^T C$, C and T are the charge conjugation and transposition operators, respectively, and S_Q and $S_{q(s)}$ are the heavy and light (strange) quark propagators.

The correlation function from the QCD part receives three different contributions: (a) perturbative contributions, (b) nonperturbative contributions where a photon is emitted from the freely propagating quark (in other words, at short distances), (c) nonperturbative contributions where a photon is radiated at long distances. To obtain the expression for the contribution from the emission of a photon at short distances, the following procedure can be used: Each one of the quarks can emit the photon, and hence each term in Eq. (9) corresponds to three terms in which the propagator of the photon emitting the quark is replaced by

$$S_{\alpha\beta}^{ab} \Rightarrow -\frac{1}{2} \left\{ \int d^4y F^{\mu\nu} y_\nu S^{\text{free}}(x-y) \gamma_\mu S^{\text{free}}(y) \right\}_{\alpha\beta}^{ab}, \quad (10)$$

where the Fock-Schwinger gauge, $x_\mu A^\mu(x) = 0$, has been used. Note that the explicit expressions of the free light and heavy quark propagators in the x representation are

$$\begin{aligned} S_q^{\text{free}} &= \frac{i\not{x}}{2\pi^2 x^4} - \frac{m_q}{4\pi^2 x^2}, \\ S_Q^{\text{free}} &= \frac{m_Q^2}{4\pi^2} \frac{K_1(m_Q \sqrt{-x^2})}{\sqrt{-x^2}} - i \frac{m_Q^2 \not{x}}{4\pi^2 x^2} K_2(m_Q \sqrt{-x^2}), \end{aligned} \quad (11)$$

where K_i are Bessel functions, $m_{u,d} = 0$, and $m_s \neq 0$. The expression for the nonperturbative contributions to the correlation function can be obtained from Eq. (9) by replacing one of the light quark propagators by

$$S_{\alpha\beta}^{ab} \rightarrow -\frac{1}{4} \bar{q}^a \Gamma_j q^b (\Gamma_j)_{\alpha\beta}, \quad (12)$$

where $\Gamma_j = \{1, \gamma_5, \gamma_\alpha, i\gamma_5 \gamma_\alpha, \sigma_{\alpha\beta}/\sqrt{2}\}$ and the sum over Γ_j is implied; the other two propagators are the full propa-

gators involving both perturbative and nonperturbative contributions. In order to calculate the correlation function from the QCD side, we need the explicit expressions of the heavy and light quark propagators in the presence of an external field.

The light cone expansion of the propagator in an external field is obtained in [43]. It receives contributions from

various $\bar{q}Gq$, $\bar{q}GGq$, $\bar{q}q\bar{q}q$ nonlocal operators, where G is the gluon field strength tensor. In this work, we consider operators with only one gluon field and contributions coming from three-particle nonlocal operators, and neglect terms with two gluons $\bar{q}GGq$ and four quarks $\bar{q}q\bar{q}q$ [44]. In this approximation the expressions for the heavy and light quark propagators are

$$iS_Q(x) = iS_Q^{\text{free}}(x) - ig_s \int \frac{d^4k}{(2\pi)^4} e^{-ikx} \int_0^1 dv \left[\frac{\not{k} + m_Q}{(m_Q^2 - k^2)^2} G^{\mu\nu}(vx) \sigma_{\mu\nu} + \frac{1}{m_Q^2 - k^2} vx^\mu G^{\mu\nu} \gamma_\nu \right],$$

$$S_q(x) = S_q^{\text{free}}(x) - \frac{m_q}{4\pi^2 x^2} - \frac{\langle \bar{q}q \rangle}{12} \left(1 - i \frac{m_q}{4} \not{x} \right) - \frac{x^2}{192} m_0^2 \langle \bar{q}q \rangle \left(1 - i \frac{m_q}{6} \not{x} \right) - ig_s \int_0^1 du \left[\frac{\not{x}}{16\pi^2 x^2} G_{\mu\nu}(ux) \sigma_{\mu\nu} \right. \quad (13)$$

$$\left. - ux^\mu G_{\mu\nu}(ux) \gamma^\nu \frac{i}{4\pi^2 x^2} - i \frac{m_q}{32\pi^2} G_{\mu\nu} \sigma^{\mu\nu} \left(\ln \left(\frac{-x^2 \Lambda^2}{4} \right) + 2\gamma_E \right) \right],$$

where Λ is the energy cutoff separating perturbative and nonperturbative domains.

In order to calculate the theoretical part, from Eqs. (9)–(13) it follows that the matrix elements of nonlocal operators $\bar{q}\Gamma_i q$ between the photon and vacuum states are needed, i.e. $\langle \gamma(q) | \bar{q}(x_1) \Gamma_i q(x_2) | 0 \rangle$. These matrix elements can be expanded near the light cone $x^2 = 0$ in terms of the photon distribution amplitudes [46].

$$\begin{aligned} \langle \gamma(q) | \bar{q}(x) \sigma_{\mu\nu} q(0) | 0 \rangle &= -ie_q \bar{q}q (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int_0^1 du e^{i\bar{u}qx} \left(\chi \varphi_\gamma(u) + \frac{x^2}{16} \mathbb{A}(u) \right) - \frac{i}{2(qx)} e_q \langle \bar{q}q \rangle \left[x_\nu \left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \right. \\ &\quad \left. - x_\mu \left(\varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \right] \int_0^1 du e^{i\bar{u}qx} h_\gamma(u) \langle \gamma(q) | \bar{q}(x) \gamma_\mu q(0) | 0 \rangle \\ &= e_q f_{3\gamma} \left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \int_0^1 du e^{i\bar{u}qx} \psi^v(u) \langle \gamma(q) | \bar{q}(x) \gamma_\mu \gamma_5 q(0) | 0 \rangle \\ &= -\frac{1}{4} e_q f_{3\gamma} \varepsilon_{\mu\nu\alpha\beta} \varepsilon^\nu q^\alpha x^\beta \int_0^1 du e^{i\bar{u}qx} \psi^a(u) \langle \gamma(q) | \bar{q}(x) g_s G_{\mu\nu}(vx) q(0) | 0 \rangle \\ &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{S}(\alpha_i) \langle \gamma(q) | \bar{q}(x) g_s \tilde{G}_{\mu\nu} i\gamma_5(vx) q(0) | 0 \rangle \\ &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \tilde{\mathcal{S}}(\alpha_i) \langle \gamma(q) | \bar{q}(x) g_s \tilde{G}_{\mu\nu}(vx) \gamma_\alpha \gamma_5 q(0) | 0 \rangle \\ &= e_q f_{3\gamma} q_\alpha (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{A}(\alpha_i) \langle \gamma(q) | \bar{q}(x) g_s G_{\mu\nu}(vx) i\gamma_\alpha q(0) | 0 \rangle \\ &= e_q f_{3\gamma} q_\alpha (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{V}(\alpha_i) \langle \gamma(q) | \bar{q}(x) \sigma_{\alpha\beta} g_s G_{\mu\nu}(vx) q(0) | 0 \rangle \\ &= e_q \langle \bar{q}q \rangle \left\{ \left[\left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \left(g_{\alpha\nu} - \frac{1}{qx} (q_\alpha x_\nu + q_\nu x_\alpha) \right) q_\beta - \left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \left(g_{\beta\nu} - \frac{1}{qx} (q_\beta x_\nu + q_\nu x_\beta) \right) q_\alpha \right. \right. \\ &\quad \left. \left. - \left(\varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \left(g_{\alpha\mu} - \frac{1}{qx} (q_\alpha x_\mu + q_\mu x_\alpha) \right) q_\beta + \left(\varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \left(g_{\beta\mu} - \frac{1}{qx} (q_\beta x_\mu + q_\mu x_\beta) \right) q_\alpha \right] \right. \\ &\quad \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_1(\alpha_i) + \left[\left(\varepsilon_\alpha - q_\alpha \frac{\varepsilon x}{qx} \right) \left(g_{\mu\beta} - \frac{1}{qx} (q_\mu x_\beta + q_\beta x_\mu) \right) q_\nu - \left(\varepsilon_\alpha - q_\alpha \frac{\varepsilon x}{qx} \right) \right. \\ &\quad \times \left(g_{\nu\beta} - \frac{1}{qx} (q_\nu x_\beta + q_\beta x_\nu) \right) q_\mu - \left(\varepsilon_\beta - q_\beta \frac{\varepsilon x}{qx} \right) \left(g_{\mu\alpha} - \frac{1}{qx} (q_\mu x_\alpha + q_\alpha x_\mu) \right) q_\nu + \left(\varepsilon_\beta - q_\beta \frac{\varepsilon x}{qx} \right) \\ &\quad \times \left. \left(g_{\nu\alpha} - \frac{1}{qx} (q_\nu x_\alpha + q_\alpha x_\nu) \right) q_\mu \right] \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_2(\alpha_i) + \frac{1}{qx} (q_\mu x_\nu - q_\nu x_\mu) (\varepsilon_\alpha q_\beta - \varepsilon_\beta q_\alpha) \\ &\quad \times \left. \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_3(\alpha_i) + \frac{1}{qx} (q_\alpha x_\beta - q_\beta x_\alpha) (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_4(\alpha_i) \right\} \quad (14) \end{aligned}$$

where χ is the magnetic susceptibility of the quarks, $\varphi_\gamma(u)$ is the leading twist-2 photon distribution amplitude (DA), $\psi^\nu(u)$, $\psi^a(u)$, \mathcal{A} , and \mathcal{V} are the twist-3 DA's, and $h_\gamma(u)$, \mathbb{A} , \mathcal{T}_i ($i = 1, 2, 3, 4$) are the twist-4 DA's, respectively. The explicit expressions of DA's are presented in the next section.

The theoretical part of the correlation function can be obtained in terms of QCD parameters by substituting photon DA's and expressions for heavy and light quark

propagators into Eq. (9). Sum rules for the Ξ_Q magnetic moments are obtained by equating two representations of the correlation function. The higher states and continuum contributions are modeled using hadron-quark duality. Applying double Borel transformations on the variables $p_1^2 = (p + q)^2$ and $p_2^2 = p^2$ on both sides of the correlator, for the Ξ_Q baryon magnetic moments we get

$$\begin{aligned}
-\lambda_Q^2(\beta)\mu_{\Xi_Q} e^{-m_{\Xi_Q}^2/M_B^2} &= \int_{m_Q^2}^{s_0} e^{-s/M_B^2} \rho(s) ds + \frac{(\beta^2 - 1)e^{-m_Q^2/M_B^2}}{288\pi^2} \{\gamma_E(6e_Q + e_s)m_s m_0^2 \langle \bar{q}q \rangle\} - \frac{(\beta^2 - 1)e^{-m_Q^2/M_B^2} m_Q^2}{72M_B^4} \\
&\times \{(e_s + e_q)m_0^2 \langle \bar{s}s \rangle \langle \bar{q}q \rangle \eta_1\} + \frac{e^{-m_Q^2/M_B^2} m_Q^2}{432M_B^4} \{(\beta^2 - 1)m_0^2 \langle \bar{s}s \rangle \langle \bar{q}q \rangle [36e_Q + (e_s + e_q)\mathbb{A}(u_0)] \\
&+ (\beta^2 + 1)f_{3\gamma} e_q m_s m_0^2 \langle \bar{s}s \rangle (\eta_2 + \psi^a(u_0))\} + \frac{e^{-m_Q^2/M_B^2}}{72} \{(1 - \beta^2) \langle \bar{s}s \rangle \langle \bar{q}q \rangle [12e_Q - (e_s + e_q) \\
&\times [\eta_1 - m_0^2 \chi_i \varphi_\gamma(u_0)]] - 3e_q(\beta^2 + 1)f_{3\gamma} m_s \langle \bar{s}s \rangle \eta_2\} - \frac{(\beta^2 - 1)e^{-m_Q^2/M_B^2} M_B^2 m_s}{96\pi^2 m_Q^2} \\
&\times [(6e_Q + e_s)\gamma_E m_0^2 \langle \bar{q}q \rangle] - \frac{e^{-m_Q^2/M_B^2} m_s}{288\pi^2} \left(3(\beta^2 - 1)e_Q m_0^2 \langle \bar{q}q \rangle \left[-3 + 2\gamma_E + 2 \ln \left[\frac{\Lambda^2}{m_Q^2} \right] \right] \right. \\
&\left. + e_q m_0^2 \langle \bar{s}s \rangle (1 + \beta^2) + e_s m_0^2 \langle \bar{q}q \rangle \left[(1 - \beta^2) \left(\gamma_E + \ln \left[\frac{\Lambda^2}{m_Q^2} \right] \right) \right] \right) + \frac{9e_Q m_Q^2 m_s}{144\pi^2} \{ \langle \bar{s}s \rangle (1 + \beta^2) \\
&+ 2\langle \bar{q}q \rangle (1 - \beta^2) \}, \tag{15}
\end{aligned}$$

where $M_B^2 = \frac{M_1^2 M_2^2}{M_1^2 + M_2^2}$ and $u_0 = \frac{M_1^2}{M_1^2 + M_2^2}$. Since the masses of the initial and final baryons are the same, we will set $M_1^2 = M_2^2$ and $u_0 = 1/2$. The functions appearing in Eq. (15) are defined as

$$\begin{aligned}
\eta_1 &= \int \mathcal{D}\alpha_i \int_0^1 dv \mathcal{S}(\alpha_i) \delta(\alpha_q + v\alpha_g - u_0), \\
\eta_2 &= \int \mathcal{D}\alpha_i \int_0^1 dv \mathcal{V}(\alpha_i) \delta'(\alpha_q + v\alpha_g - u_0), \\
\rho(s) &= \frac{(\beta^2 - 1)}{144\pi^2 M_B^2} \left\{ m_0^2 (6e_Q + e_s) m_s \langle \bar{q}q \rangle \ln \left(\frac{-m_Q^2 + s}{\Lambda^2} \right) \right\} + \frac{3(1 + \beta^2)e_Q m_Q^4}{64\pi^4} \left\{ \frac{13}{2} + \psi_{10} - \frac{1}{6}\psi_{20} - \frac{1}{6}\psi_{30} \right. \\
&+ \left[\psi_{10} + \frac{3}{2} \right] \ln \left(\frac{s}{m_Q^2} \right) + \frac{1}{6}\psi_{41} \left. \right\} + \frac{(1 - \beta^2)m_s \langle \bar{q}q \rangle \psi_{10}}{48\pi^2 m_Q^2} \left\{ 2e_q m_Q^2 \eta_1 - (e_s + e_q) m_0^2 \left[8 + \ln \left(\frac{s - m_Q^2}{\Lambda^2} \right) \right] \right\} \\
&- \frac{m_s}{288\pi^2 m_Q^2} \left\{ (\beta^2 - 1) m_0^2 (e_s + 6e_Q) \langle \bar{q}q \rangle \left[3 \ln \left(\frac{s - m_Q^2}{\Lambda^2} \right) - \left(4\gamma_E + \ln \left[\frac{\Lambda^2}{m_Q^2} \right] \right) \right] + 6e_Q [3m_Q^2 \{ (1 + \beta^2) \langle \bar{s}s \rangle \right. \right. \\
&+ 2(1 - \beta^2) \langle \bar{q}q \rangle \} \psi_{10}] \left. \right\} + \frac{m_Q^2}{576\pi^2} \left((e_s + 12e_Q) \left\{ \frac{(\beta^2 - 1)}{m_Q^4} m_0^2 m_s \langle \bar{q}q \rangle \left[-(1 + \gamma_E)(\psi_{22} + 2\psi_{12}) - \psi_{02} - \psi_{32} \right. \right. \right. \\
&- \psi_{22} - \frac{\gamma_E}{2}\psi_{20} + 3(2\psi_{32} + 3\psi_{22} + \psi_{02}) \ln \left(\frac{s - m_Q^2}{\Lambda^2} \right) - \ln \left(\frac{\Lambda^2}{m_Q^2} \right) \left. \right] \left. \right\} + 12e_q \left\{ \frac{2}{m_Q^2} (\beta^2 - 1) m_s \langle \bar{q}q \rangle \eta_1 \psi_{21} \right. \\
&\left. + (1 + \beta^2) f_{3\gamma} \eta_2 \left[\psi_{21} - \psi_{10} + \frac{1}{2}\psi_{20} + \frac{1}{2}\psi_{00} + \ln \left(\frac{m_Q^2}{s} \right) \right] \right\}, \tag{16}
\end{aligned}$$

where

$$\int \mathcal{D}\alpha_i = \int_0^1 d\alpha_{\bar{q}} \int_0^1 d\alpha_q \int_0^1 d\alpha_g \delta(1 - \alpha_{\bar{q}} - \alpha_q - \alpha_g), \tag{17}$$

and the functions ψ_{nm} are defined as

$$\psi_{nm} = \frac{(s - m_Q^2)^n}{s^m (m_Q^2)^{n-m}}. \quad (18)$$

Note that the contributions of the terms $\sim \langle G^2 \rangle$ are also calculated, but their numerical values are very small, and

$$\begin{aligned} \lambda_Q^2(\beta) = & e^{m_{\Xi_Q}^2/M_B^2} \left(\int_{m_Q^2}^{s_0} ds e^{-s/M_B^2} \left\{ \frac{(1 + \beta^2)m_Q^4}{2^9 \pi^4} \left[(1 - x^2) \left(\frac{1}{x^2} - \frac{8}{x} + 1 \right) - 12 \ln x \right] \right. \right. \\ & + \frac{m_s}{2^4 \pi^2} (1 - x^2) \left[(1 - \beta^2) \langle \bar{q}q \rangle + \frac{(1 + \beta^2) \langle \bar{s}s \rangle}{2} \right] + \frac{(1 + \beta^2) \langle g^2 G^2 \rangle}{2^{10} 3 \pi^4} (1 - x)(1 + 5x) \left. \right\} \\ & + \frac{m_s}{2^5 \pi^2} \left\{ \frac{(1 + \beta^2)m_0^2 \langle \bar{s}s \rangle}{6} e^{-m_Q^2/M_B^2} - (1 - \beta^2)m_0^2 \langle \bar{q}q \rangle \left[e^{-m_Q^2/M_B^2} + \int_0^1 d\alpha (1 - \alpha) e^{-m_Q^2/(1-\alpha)M_B^2} \right] \right. \\ & \left. \left. - \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{6} (1 - \beta^2) e^{-m_Q^2/M_B^2} \right\}, \right. \end{aligned} \quad (19)$$

where $x = m_Q^2/s$.

III. NUMERICAL ANALYSIS

The present section is devoted to the numerical analysis of the magnetic moments of Ξ_Q baryons. The values of the input parameters, appearing in the sum rules expression for magnetic moments, are $\langle \bar{u}u \rangle(1 \text{ GeV}) = \langle \bar{d}d \rangle(1 \text{ GeV}) = -(0.243)^3 \text{ GeV}^3$, $\langle \bar{s}s \rangle(1 \text{ GeV}) = 0.8 \langle \bar{u}u \rangle(1 \text{ GeV})$, $m_0^2(1 \text{ GeV}) = 0.8 \text{ GeV}^2$ [47], $\Lambda = 300 \text{ MeV}$, and $f_{3\gamma} = -0.0039 \text{ GeV}^2$ [46]. The value of the magnetic susceptibility, $\chi(1 \text{ GeV}) = -3.15 \pm 0.3 \text{ GeV}^{-2}$, was obtained by a combination of the local duality approach and QCD sum

therefore, for customary in Eq. (15), these terms are omitted. From Eq. (15) it follows that, for the determination of the Ξ_Q baryon magnetic moments, we need to know the residue λ_Q . The residue can be obtained from the two-point sum rules and is calculated in [25]. For the current given in Eq. (1), it takes the following form:

rules [46]. Recently, from the analysis of radiative heavy meson decay for $\chi(1 \text{ GeV}) = -(2.85 \pm 0.5) \text{ GeV}^{-2}$ was obtained [48], which is in good agreement with the instanton liquid model prediction [49], but slightly below the QCD sum rules prediction [46]. Note that, first, the magnetic susceptibility in the framework of QCD sum rules is calculated in [50], and it is obtained that $\chi(1 \text{ GeV}) = -4.4 \text{ GeV}^{-2}$. In the numerical analysis, we have used all three values of χ existing in the literature and obtained that the values of the magnetic moments of Ξ_Q baryons are practically insensitive to the value of χ . The photon DA's entering the sum rules for the magnetic moments of Ξ_Q are calculated in [46], and their expressions are

$$\begin{aligned} \varphi_\gamma(u) &= 6u\bar{u}(1 + \varphi_2(\mu)C_2^{3/2}(u - \bar{u})), \\ \psi^v(u) &= 3(3(2u - 1)^2 - 1) + \frac{3}{64}(15w_\gamma^v - 5w_\gamma^A)(3 - 30(2u - 1)^2 + 35(2u - 1)^4), \\ \psi^a(u) &= (1 - (2u - 1)^2)(5(2u - 1)^2 - 1) \frac{5}{2} \left(1 + \frac{9}{16}w_\gamma^v - \frac{3}{16}w_\gamma^A \right), \quad \mathcal{A}(\alpha_i) = 360\alpha_q\alpha_{\bar{q}}\alpha_g^2 \left(1 + w_\gamma^A \frac{1}{2}(7\alpha_g - 3) \right), \\ \mathcal{V}(\alpha_i) &= 540w_\gamma^v(\alpha_q - \alpha_{\bar{q}})\alpha_q\alpha_{\bar{q}}\alpha_g^2, \quad h_\gamma(u) = -10(1 + 2\kappa^+)C_2^{1/2}(u - \bar{u}), \\ \mathbb{A}(u) &= 40u^2\bar{u}^2(3\kappa - \kappa^+ + 1) + 8(\zeta_2^+ - 3\zeta_2^-)[u\bar{u}(2 + 13u\bar{u}) + 2u^3(10 - 15u + 6u^2)\ln(u) \\ & \quad + 2\bar{u}^3(10 - 15\bar{u} + 6\bar{u}^2)\ln(\bar{u})], \\ \mathcal{T}_1(\alpha_i) &= -120(3\zeta_2 + \zeta_2^+)(\alpha_{\bar{q}} - \alpha_q)\alpha_{\bar{q}}\alpha_q\alpha_g, \\ \mathcal{T}_2(\alpha_i) &= 30\alpha_g^2(\alpha_{\bar{q}} - \alpha_q)((\kappa - \kappa^+) + (\zeta_1 - \zeta_1^+)(1 - 2\alpha_g) + \zeta_2(3 - 4\alpha_g)), \\ \mathcal{T}_3(\alpha_i) &= -120(3\zeta_2 - \zeta_2^+)(\alpha_{\bar{q}} - \alpha_q)\alpha_{\bar{q}}\alpha_q\alpha_g, \\ \mathcal{T}_4(\alpha_i) &= 30\alpha_g^2(\alpha_{\bar{q}} - \alpha_q)((\kappa + \kappa^+) + (\zeta_1 + \zeta_1^+)(1 - 2\alpha_g) + \zeta_2(3 - 4\alpha_g)), \\ \mathcal{S}(\alpha_i) &= 30\alpha_g^2\{(\kappa + \kappa^+)(1 - \alpha_g) + (\zeta_1 + \zeta_1^+)(1 - \alpha_g)(1 - 2\alpha_g) + \zeta_2[3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g)]\}, \\ \tilde{\mathcal{S}}(\alpha_i) &= -30\alpha_g^2\{(\kappa - \kappa^+)(1 - \alpha_g) + (\zeta_1 - \zeta_1^+)(1 - \alpha_g)(1 - 2\alpha_g) + \zeta_2[3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g)]\}. \end{aligned} \quad (20)$$

The constants appearing in the wave functions are given as [46] $\varphi_2(1 \text{ GeV}) = 0$, $w_y^V = 3.8 \pm 1.8$, $w_y^A = -2.1 \pm 1.0$, $\kappa = 0.2$, $\kappa^+ = 0$, $\zeta_1 = 0.4$, $\zeta_2 = 0.3$, $\zeta_1^+ = 0$, and $\zeta_2^+ = 0$.

From the explicit expressions of the magnetic moments of Ξ_Q baryons, it follows that it contains three auxiliary parameters: Borel mass squared M_B^2 , continuum threshold s_0 , and β , which enters the expression of the interpolating current for Ξ_Q . The physical quantity, magnetic moment μ_{Ξ_Q} , should be independent of these auxiliary parameters. In other words, we should find the “working regions” of these auxiliary parameters, where the magnetic moments are independent of them.

The value of the continuum threshold is fixed from the analysis of the two-point sum rules, where the mass and residue λ_{Ξ_Q} of the Ξ_Q baryons are determined [25], which leads to the value $s_0 = 6.5^2 \text{ GeV}^2$ for Ξ_b and $s_0 = 3.0^2 \text{ GeV}^2$ for Ξ_c . If we choose the value $s_0 = 6.4^2 \text{ GeV}^2$ for Ξ_b and $s_0 = 8 \text{ GeV}^2$ for Ξ_c , the results remain practically unchanged. Next, we try to find the working region of M_B^2 where μ_{Ξ_Q} are independent of it at fixed values of β and the above-mentioned values of s_0 . The upper bound of M_B^2 is obtained, requiring that the continuum contribution should be less than the contribution of the first resonance. The lower bound of M_B^2 is determined by requiring that the highest power of $1/M_B^2$ be less than 30% of the highest power of M_B^2 . These two conditions are both satisfied in the region $15 \text{ GeV}^2 \leq M_B^2 \leq 20 \text{ GeV}^2$ and $5 \text{ GeV}^2 \leq M_B^2 \leq 8 \text{ GeV}^2$ for Ξ_b and Ξ_c , respectively.

In Figs. 1 and 2, we depict the dependence of $\mu_{\Xi_b^0}$ and $\mu_{\Xi_b^-}$ on M_B^2 at fixed values of β and $s_0 = 6.5^2 \text{ GeV}^2$. In Figs. 3 and 4, we present the dependence of $\mu_{\Xi_c^0}$ and $\mu_{\Xi_c^+}$ on M_B^2 at fixed values of β and $s_0 = 3.0^2 \text{ GeV}^2$. From these figures, we see that the values of the magnetic mo-




FIG. 1. The dependence of the magnetic moment $\mu_{\Xi_b^0}$ on M_B^2 at $s_0 = 6.5^2 \text{ GeV}^2$ and $\beta = \pm 5, -1$.




FIG. 2. The same as Fig. 1 but for $\mu_{\Xi_b^-}$.




FIG. 3. The same as Fig. 1 but for $\mu_{\Xi_c^0}$ and at $s_0 = 3.0^2 \text{ GeV}^2$.




FIG. 4. The same as Fig. 3 but for $\mu_{\Xi_c^+}$.




FIG. 5. The dependence of the magnetic moment $\mu_{\Xi_b^0}$ on $\cos\theta$ at $s_0 = 6.5^2 \text{ GeV}^2$ and for $M_B^2 = 15 \text{ GeV}^2$ and $M_B^2 = 20 \text{ GeV}^2$.




FIG. 6. The same as Fig. 5 but for $\mu_{\Xi_b^-}$.




FIG. 7. The same as Fig. 5 but for $\mu_{\Xi_c^0}$ and $s_0 = 3.0^2 \text{ GeV}^2$ and for $M_B^2 = 5 \text{ GeV}^2$ and $M_B^2 = 8 \text{ GeV}^2$.




FIG. 8. The same as Fig. 7 but for $\mu_{\Xi_c^+}$.

ments of Ξ_b and Ξ_c exhibit good stability when M_B^2 varies in the region $15 \text{ GeV}^2 \leq M_B^2 \leq 20 \text{ GeV}^2$ and $5 \text{ GeV}^2 \leq M_B^2 \leq 8 \text{ GeV}^2$, respectively. The last step of our analysis is the determination of the working region for the auxiliary parameter β . For this aim, in Figs. 5–8 we present the dependence of the magnetic moments of Ξ_Q baryons on $\cos\theta$ where $\tan\theta = \beta$, using the values of M_B^2 from the working region which we already determined and at fixed values of s_0 .

From these figures we obtained that the prediction of the magnetic moment μ_{Ξ_b} (μ_{Ξ_c}) is practically independent of the value of the auxiliary parameter β . From all these analysis we deduce the final results for the magnetic moments in Table I for $\chi = -3.15 \text{ GeV}^2$. Comparison of our results on the magnetic moments of Ξ_Q baryons with predictions of other approaches, as we already noted, is also presented in Table I.

We see that within errors our predictions on the magnetic moments are in good agreement with the quark model predictions. Our results on the magnetic moments of Ξ_c are also close to the predictions of the other approaches, except the prediction of [33] on $\mu_{\Xi_c^0}$.

In summary, the magnetic moments of Ξ_Q baryons, which were discovered recently (more precisely, Ξ_b was discovered), are calculated in the framework of light cone

TABLE I. Results for the magnetic moments of Ξ_Q baryons in different approaches.

	$\mu_{\Xi_b^0}$	$\mu_{\Xi_b^-}$	$\mu_{\Xi_c^0}$	$\mu_{\Xi_c^+}$
Our results	-0.045 ± 0.005	-0.08 ± 0.02	0.35 ± 0.05	0.50 ± 0.05
RQM [32]	-0.06	-0.06	0.39	0.41
NQM [32]	-0.06	-0.06	0.37	0.37
[33]	$-1.02 \div -1.06$	$0.45 \div 0.48$
[34]	0.32	0.42
[35]	0.38	0.38
[36]	0.28	0.28
[37]	$0.28 \div 0.34$	$0.39 \div 0.46$

QCD sum rules. Our results on magnetic moments are close to the predictions of the other approaches existing in the literature.

ACKNOWLEDGMENTS

Two of the authors (K. A. and A. O.) would like to thank TUBITAK, Turkish Scientific and Research Council, for

their partial financial support through Project No. 106T333. One of the authors (A. O.) would like to thank TUBA for funds provided through the GEBIP program.

-
- [1] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **99**, 202001 (2007).
- [2] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **99**, 052001 (2007).
- [3] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **99**, 052002 (2007).
- [4] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **97**, 232001 (2006).
- [5] S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
- [6] R. Roncaglia, D. B. Lichtenberg, and E. Predazzi, Phys. Rev. D **52**, 1722 (1995).
- [7] E. Jenkins, Phys. Rev. D **54**, 4515 (1996).
- [8] N. Mathur, R. Lewis, and R. M. Woloshyn, Phys. Rev. D **66**, 014502 (2002).
- [9] D. Ebert, R. N. Faustov, and V. O. Galkin, Phys. Rev. D **72**, 034026 (2005).
- [10] M. Karliner and H. J. Lipkin, arXiv:hep-ph/0307343; arXiv:hep-ph/0611306.
- [11] M. Karliner, B. Kereu-Zura, H. J. Lipkin, and J. L. Rosner, arXiv:0706.2163.
- [12] J. L. Rosner, Phys. Rev. D **75**, 013009 (2007).
- [13] M. Karliner and H. J. Lipkin, Phys. Lett. B **575**, 249 (2003).
- [14] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. **B147**, 385 (1979).
- [15] E. Bagan, M. Chabab, H. G. Dosch, and S. Narison, Phys. Lett. B **278**, 367 (1992); **287**, 176 (1992); **301**, 243 (1993).
- [16] C. S. Navarra and M. Nielsen, Phys. Lett. B **443**, 285 (1998).
- [17] E. V. Shuryak, Nucl. Phys. **B198**, 83 (1982).
- [18] A. G. Grozin and O. I. Yakovlev, Phys. Lett. B **285**, 254 (1992).
- [19] Y. B. Dai, C. S. Huang, C. Liu, and C. D. Lu, Phys. Lett. B **371**, 99 (1996).
- [20] D. W. Wang, M. Q. Huang, and C. Z. Li, Phys. Rev. D **65**, 094036 (2002).
- [21] S. L. Zhu, Phys. Rev. D **61**, 114019 (2000).
- [22] C. S. Huang, A. L. Zhang, and S. L. Zhu, Phys. Lett. B **492**, 288 (2000).
- [23] D. W. Wang and M. Q. Huang, Phys. Rev. D **68**, 034019 (2003).
- [24] Z. G. Wang, Eur. Phys. J. C **54**, 231 (2008).
- [25] F. O. Duraes and M. Nielsen, Phys. Lett. B **658**, 40 (2007).
- [26] X. Liu, H. X. Chen, Y. R. Liu, A. Hosaka, and S. L. Zhu, Phys. Rev. D **77**, 014031 (2008).
- [27] A. L. Choudhury and V. Joshi, Phys. Rev. D **13**, 3115 (1976).
- [28] D. B. Lichtenberg, Phys. Rev. D **15**, 345 (1977).
- [29] L. Y. Glozman and D. O. Riska, Nucl. Phys. **A603**, 326 (1996).
- [30] B. Julia-Diaz and D. O. Riska, Nucl. Phys. **A739**, 69 (2004).
- [31] S. Scholl and H. Weigel, Nucl. Phys. **A735**, 163 (2004).
- [32] A. Faessler *et al.*, Phys. Rev. D **73**, 094013 (2006).
- [33] B. Patel, A. K. Rai, and P. C. Vinodkumar, arXiv:0803.0221.
- [34] M. Savage, Phys. Lett. B **326**, 303 (1994).
- [35] D. O. Riska, Nucl. Instrum. Methods Phys. Res., Sect. B **119**, 259 (1996).
- [36] Y. Oh, D. P. Min, M. Rho, and N. N. Scoccola, Nucl. Phys. **A534**, 493 (1991).
- [37] C. S. An, Nucl. Phys. **A797**, 131 (2007); **A801**, 82(E) (2008).
- [38] S. L. Zhu, W. Y. Hwang, and Z. S. Yang, Phys. Rev. D **56**, 7273 (1997).
- [39] T. M. Aliev, A. Ozpineci, and M. Savci, Phys. Rev. D **65**, 096004 (2002).
- [40] T. M. Aliev, A. Ozpineci, and M. Savci, Phys. Rev. D **65**, 056008 (2002).
- [41] P. Colangelo and A. Khodjamirian, in *At the Frontier of Particle Physics/Handbook of QCD*, edited by M. Shifman (World Scientific, Singapore, 2001), Vol. 3, p. 1495.
- [42] V. M. Braun, arXiv:hep-ph/9801222.
- [43] I. I. Balitsky and V. M. Braun, Nucl. Phys. **B311**, 541 (1989).
- [44] V. M. Braun and I. E. Filyanov, Z. Phys. C **48**, 239 (1990).
- [45] T. M. Aliev, A. Ozpineci, and M. Savci, Phys. Rev. D **66**, 016002 (2002); **67**, 039901(E) (2003).
- [46] P. Ball, V. M. Braun, and N. Kivel, Nucl. Phys. **B649**, 263 (2003).
- [47] V. M. Belyaev and B. L. Ioffe, JETP **56**, 493 (1982).
- [48] J. Rohrwild, J. High Energy Phys. **09** (2007) 073.
- [49] A. E. Dorokhov, Eur. Phys. J. C **42**, 309 (2005).
- [50] V. M. Belyaev and I. I. Kogan, Yad. Fiz. **40**, 1035 (1984).