

Noncommutative corrections to the Robertson-Walker metric

S. Fabi, B. Harms, and A. Stern

Department of Physics, University of Alabama, Tuscaloosa, Alabama 35487, USA

(Received 20 August 2008; published 23 September 2008)

Upon applying Chamseddine's noncommutative deformation of gravity, we obtain the leading order noncommutative corrections to the Robertson-Walker metric tensor. We get an isotropic inhomogeneous metric tensor for a certain choice of the noncommutativity parameters. Moreover, the singularity of the commutative metric at $t = 0$ is replaced by a more involved space-time structure in the noncommutative theory. In a toy model we construct a scenario where there is no singularity at $t = 0$ at leading order in the noncommutativity parameter. Although singularities may still be present for nonzero t , they need not be the source of all timelike geodesics and the result resembles a bouncing cosmology.

DOI: [10.1103/PhysRevD.78.065037](https://doi.org/10.1103/PhysRevD.78.065037)

PACS numbers: 11.10.Nx, 02.40.Gh, 04.20.-q

I. INTRODUCTION

Noncommutative deformations of general relativity offer the promise of modeling effects of quantum gravity. A number of different deformations have been given [1–4]. The approach of Aschieri *et al.* [2] has the advantage of preserving the full diffeomorphism symmetry of the commutative theory. As it is technically rather involved, it so far however has not been very convenient for practical applications. An older approach of Chamseddine [1] is based on the noncommutative analogue $SO(4, 1)$ gauge theory and uses the Seiberg-Witten map [5]. It makes contact with general relativity using a Wigner-Inönü contraction. Ideally, one should look for solutions to a noncommutative deformation of the field equations and map back to the commutative theory in order to obtain a physical interpretation. This procedure could be easily carried out in the case of $U(1)$ gauge theory in order to obtain noncommutative corrections to the Coulomb solution [6]. However in the case of gravity, a deformation of Einstein equations which is covariant under a noncommutative version of local Lorentz transformations remains obscure within the $SO(4, 1)$ gauge theory approach. An alternative procedure has been adopted recently to obtain noncommutative corrections to black holes [7,8]. (See also [9–13].) There, rather than solving some noncommutative analogue of the Einstein equations subject to the appropriate boundary conditions, one maps the known black hole solutions of general relativity to the noncommutative theory. One then defines a noncommutative analogue of the metric tensor in order to give a physical interpretation of the results. As is typical with noncommutative gravity, the leading order corrections are second order in noncommutativity parameters [14].

Cosmology offers another possible realm of application of noncommutativity. Previous studies have led to corrections to the cosmic microwave background radiation [15], and noncommutative scalar fields have been coupled to the Robertson-Walker metric tensor in order to study effects on inflation [16–18]. Noncommutativity could also potentially resolve the big bang singularity. With this in mind we apply

the procedure discussed above to obtain leading order corrections to the Robertson-Walker metric tensor. We get an isotropic inhomogeneous metric tensor (with respect to one world line) after making a specific choice of the noncommutativity parameters. Isotropic inhomogeneous cosmologies have been studied previously [19], and some specific models have been proposed in order to explain the cosmological acceleration [20–26]. For an arbitrary expansion, the second order corrections to the Robertson-Walker metric tensor which we obtain are rather involved. They simplify considerably for the special case of a linear expansion which allows for an analysis at small time t (associated with the noncommutativity scale). In this toy model we can construct a scenario where the noncommutative metric tensor is everywhere well defined at $t = 0$ to leading order in the noncommutativity scale. New singularities do appear at nonzero t in this case, but these singularities are not the source of all timelike geodesics. Instead, geodesics can be extended through the $t = 0$ time slice, and range from $t \rightarrow -\infty$ to $t \rightarrow +\infty$. The noncommutative metric tensor is invariant under $t \rightarrow -t$ and describes a bouncing universe.

This article is organized as follows. We review the gauge theory formalism for gravity in Sec. II and the noncommutative generalization obtained by Chamseddine in Sec. III. There we introduce a recursion relation found recently in [27] for the second order potentials. It is employed in obtaining the leading noncommutative corrections to the Robertson-Walker metric in Sec. IV. There we analyze the resulting space-time structure near $t = 0$ for the case of a linear expansion. We briefly remark on a slightly more realistic expansion associated with a flat radiation dominated universe in Sec. V.

II. COMMUTATIVE THEORY

The gauge theory formalism for gravity [28] is expressed in terms of spin connection and vierbein one-forms, $\omega^{ab} = -\omega^{ba}$ and e^a , respectively. $a, b, \dots = 0, 1, 2, 3$ are Lorentz indices which are raised and lowered with the flat metric tensor $\eta = \text{diag}(-1, 1, 1, 1)$, while the

space-time metric is

$$g_{\mu\nu} = e^a{}_\mu e^b{}_\nu \eta_{ab}. \quad (2.1)$$

Infinitesimal variations of ω^{ab} and e^a induced from local $ISO(3, 1)$ transformations are given by

$$\begin{aligned} \delta\omega^{ab} &= d\lambda^{ab} + [\omega, \lambda]^{ab} \\ \delta e^a &= d\rho^a + \omega^a{}_c \rho^c - \lambda^a{}_c e^c, \end{aligned} \quad (2.2)$$

for infinitesimal parameters $\lambda^{ab} = -\lambda^{ba}$ and ρ^a , and where $[\omega, \lambda]^{ab} = \omega^a{}_c \lambda^{cb} - \lambda^a{}_c \omega^{cb}$. The spin curvature and torsion two-forms, $R^{ab} = -R^{ba}$ and T^a , respectively, are constructed from ω^{ab} and e^a according to

$$R^{ab} = d\omega^{ab} + \omega^a{}_c \wedge \omega^{cb} \quad T^a = de^a + \omega^a{}_b \wedge e^b, \quad (2.3)$$

and they satisfy the Bianchi identities

$$\begin{aligned} dR^{ab} &= R^a{}_c \wedge \omega^{cb} - R^b{}_c \wedge \omega^{ca} \\ dT^a &= R^a{}_b \wedge e^b - \omega^a{}_c \wedge T^c. \end{aligned} \quad (2.4)$$

The field action

$$S = \frac{1}{4} \int \epsilon_{abcd} R^{ab} \wedge e^c \wedge e^d, \quad (2.5)$$

describing pure gravity is invariant under local Lorentz transformations (and the full set of local Poincaré transformations when the torsion vanishes). The field equations obtained from arbitrary variations of ω^{ab} and e^a are

$$T^{[a} \wedge e^{b]} = 0 \quad (2.6)$$

$$R^{[ab} \wedge e^{c]} = 0, \quad (2.7)$$

where the brackets indicate antisymmetrization of indices. Provided that the vierbeins have an inverse, (2.6) implies a vanishing torsion, while (2.7) implies a vanishing Ricci curvature $\mathcal{R}_{\mu\nu} = \mathcal{R}_{\mu\sigma\nu}{}^\sigma$, where the Riemann curvature $\mathcal{R}_{\mu\nu\rho}{}^\lambda$ is given in terms of the spin curvature by

$$\mathcal{R}_{\mu\nu\rho}{}^\lambda = -R_{\mu\nu}^{ab} e_{b\rho} [e^{-1}]^\lambda{}_a, \quad (2.8)$$

where $e^b{}_\rho [e^{-1}]^\rho{}_a = \delta_a^b$.

The above $ISO(3, 1)$ gauge theory is obtained from a Wigner-Inönü contraction of $SO(4, 1)$ gauge theory. Denote the potential one-forms and the infinitesimal gauge parameters of $SO(4, 1)$ gauge theory by $A^{AB} = -A^{BA}$ and $\Lambda^{AB} = -\Lambda^{BA}$, respectively, with indices $A, B, \dots = 0, 1, 2, 3, 4$ which are raised and lowered with the metric

tensor $\text{diag}(-1, 1, 1, 1, 1)$. An $SO(4, 1)$ gauge variation is given by

$$\delta A^{AB} = D\Lambda^{AB} = d\Lambda^{AB} + [A, \Lambda]^{AB}, \quad (2.9)$$

where $[A, \Lambda]^{AB} = A^A{}_C \Lambda^{CB} - \Lambda^A{}_C A^{CB}$, and the curvature two-forms $F^{AB} = -F^{BA}$ are

$$F^{AB} = dA^{AB} + A^A{}_C \wedge A^{CB}. \quad (2.10)$$

The contraction to $ISO(3, 1)$ gauge theory is obtained by setting

$$\begin{aligned} \Lambda^{ab} &= \lambda^{ab} & \Lambda^{a4} &= \kappa \rho^a & A^{ab} &= \omega^{ab} \\ A^{a4} &= \kappa e^a & F^{ab} &= R^{ab} & F^{a4} &= \kappa T^a, \end{aligned} \quad (2.11)$$

and taking the limit $\kappa \rightarrow 0$.

III. NONCOMMUTATIVE THEORY

The noncommutative generalization for gauge theories based on nonunitary groups was obtained in [29,30]. For the case of $SO(4, 1)$ gauge theory, denote by \hat{A}^{AB} and $\hat{\Lambda}^{AB}$, respectively, the noncommutative analogues of the $SO(4, 1)$ connection one-forms and infinitesimal gauge parameters. The noncommutative analogue of (2.9) is given by

$$\delta \hat{A}^{AB} = D_\star \hat{\Lambda}^{AB} = d\hat{\Lambda}^{AB} + [\hat{A}, \hat{\Lambda}]_\star^{AB}, \quad (3.1)$$

where $[\hat{A}, \hat{\Lambda}]_\star^{AB} = \hat{A}^A{}_C \star \hat{\Lambda}^{CB} - \hat{\Lambda}^A{}_C \star \hat{A}^{CB}$, and the \star denotes the Groenewold-Moyal star product. Acting between two functions the latter is given by

$$\star = \exp\left\{\frac{i}{2} \Theta^{\mu\nu} \overleftarrow{\partial}_\mu \overrightarrow{\partial}_\nu\right\}, \quad (3.2)$$

where $\Theta^{\mu\nu} = -\Theta^{\nu\mu}$ are constant matrix elements denoting the noncommutativity parameters and $\overleftarrow{\partial}_\mu$ and $\overrightarrow{\partial}_\mu$ are left and right derivatives, respectively, with respect to some coordinates x^μ . The noncommutative analogue of the $SO(4, 1)$ curvature two-form is

$$\hat{F}^{AB} = d\hat{A}^{AB} + \hat{A}^A{}_C \star \hat{\Lambda}^{CB}. \quad (3.3)$$

\star denotes an exterior product where the usual pointwise product between components of the forms is replaced by the Groenewold-Moyal star product. The noncommutative spin connection, vierbein, curvature, and torsion forms, denoted, respectively, by $\hat{\omega}^{ab}$, \hat{e}^a , \hat{R}^{ab} , and \hat{T}^a , can be extracted from \hat{A}^{AB} as in the commutative case, i.e.,

$$\begin{aligned} \hat{A}^{ab} &= \hat{\omega}^{ab} & \hat{A}^{a4} &= \kappa \hat{e}^a & \hat{F}^{ab} &= \hat{R}^{ab} \\ \hat{F}^{a4} &= \kappa \hat{T}^a, & & & & \text{as } \kappa \rightarrow 0. \end{aligned} \quad (3.4)$$

It is known [29,30] that \hat{A} , \hat{F} , and $\hat{\Lambda}$, unlike their commutative analogues, are not valued in the $SO(4, 1)$ Lie algebra, since $(\hat{A}^{AB}, \hat{\Lambda}^{AB}) \rightarrow (-\hat{A}^{BA}, -\hat{\Lambda}^{BA})$ is not an isomorphism of the gauge algebra (3.1). Moreover, \hat{A}^{AB} , \hat{F}^{AB} , and $\hat{\Lambda}^{AB}$ cannot be restricted to real-valued forms, although one can impose anti-Hermiticity

$$\begin{aligned} (\hat{A}^{AB})^* &= -\hat{A}^{BA} & (\hat{F}^{AB})^* &= -\hat{F}^{BA} \\ (\hat{\Lambda}^{AB})^* &= -\hat{\Lambda}^{BA}, \end{aligned} \quad (3.5)$$

and the diagonal components are purely imaginary. It was observed in [30] that, if one enlarges the domain of \hat{A}^{AB} , \hat{F}^{AB} , and $\hat{\Lambda}^{AB}$ to the product of the space-time manifold (with local coordinates x^μ) with the space of all noncommutativity parameters $\Theta^{\mu\nu}$, then the following conditions can be imposed consistent with the gauge algebra:

$$\begin{aligned} \hat{A}^{AB}(x, \Theta) &= -\hat{A}^{BA}(x, -\Theta) \\ \hat{F}^{AB}(x, \Theta) &= -\hat{F}^{BA}(x, -\Theta) \\ \hat{\Lambda}^{AB}(x, \Theta) &= -\hat{\Lambda}^{BA}(x, -\Theta). \end{aligned} \quad (3.6)$$

$\hat{A}^{AB}(x, \Theta)$, $\hat{F}^{AB}(x, \Theta)$, and $\hat{\Lambda}^{AB}(x, \Theta)$ can be expanded in terms of a power series in $\Theta^{\mu\nu}$:

$$\begin{aligned} \hat{A}_\mu^{AB}(x, \Theta) &= A_\mu^{AB}(x) + A_{\mu(1)}^{AB}(x) + A_{\mu(2)}^{AB}(x) + \dots \\ \hat{F}_{\mu\nu}^{AB}(x, \Theta) &= F_{\mu\nu}^{AB}(x) + F_{\mu\nu(1)}^{AB}(x) + F_{\mu\nu(2)}^{AB}(x) + \dots \\ \hat{\Lambda}^{AB}(x, \Theta) &= \Lambda^{AB}(x) + \Lambda_{(1)}^{AB}(x) + \Lambda_{(2)}^{AB}(x) + \dots, \end{aligned} \quad (3.7)$$

where the (n) subscript indicates the n th order in $\Theta^{\mu\nu}$,

$$M^{AB}(x) = M_{\rho_1\sigma_1\rho_2\sigma_2\dots\rho_n\sigma_n}^{AB}(x) \Theta^{\rho_1\sigma_1} \Theta^{\rho_2\sigma_2} \dots \Theta^{\rho_n\sigma_n}. \quad (3.8)$$

Then (3.6) implies that the coefficients $M_{\rho_1\sigma_1\rho_2\sigma_2\dots\rho_n\sigma_n}^{AB}(x)$ are (anti)symmetric under interchange of the A and B indices for n odd (even). Equation (3.5) then implies that the coefficients are imaginary (real) for n odd (even).

The power series (3.7) have been defined using the Seiberg-Witten map from the commutative gauge theory [29,30]:

$$\hat{A}_\mu = \hat{A}_\mu(A) \quad \hat{F}_{\mu\nu} = \hat{F}_{\mu\nu}(A) \quad \hat{\Lambda} = \hat{\Lambda}(A, \Lambda), \quad (3.9)$$

where A , F , and Λ again denote the commutative potentials, curvatures, and infinitesimal gauge parameters, respectively. Since the latter are valued in the $SO(4, 1)$ Lie

algebra, this puts restrictions on the allowable \hat{A} , \hat{F} , and $\hat{\Lambda}$. The Seiberg-Witten map [5] then defines the space $\hat{\mathcal{A}}$ of allowable noncommutative potentials \hat{A} . The map is required to satisfy

$$\begin{aligned} \hat{A}_\mu(A + \partial\Lambda + [A, \Lambda]) - \hat{A}_\mu(A) \\ = \partial_\mu \hat{\Lambda}(\Lambda, A) + [\hat{A}_\mu(A), \hat{\Lambda}(\Lambda, A)]_\star, \end{aligned} \quad (3.10)$$

for infinitesimal Λ . The zeroth order in the expansion (3.7) agrees with the commutative theory. Up to homogeneous terms, the first order expressions for the noncommutative potentials and infinitesimal gauge parameters are

$$\begin{aligned} A_\mu &= -\frac{i}{4} \Theta^{\rho\sigma} \{A_\rho, \partial_\sigma A_\mu + F_{\sigma\mu}\} \\ \Lambda &= -\frac{i}{4} \Theta^{\rho\sigma} \{A_\rho, \partial_\sigma \Lambda\}, \end{aligned} \quad (3.11)$$

where the parentheses denote the anticommutator $\{A, B\}^{AB} = A^AC B_C^B + B^AC A_C^B$. Recently, a relatively simple recursion relation was found for the higher order potentials and gauge parameters [27]. At second order one gets

$$\begin{aligned} A_\mu &= -\frac{i}{8} \Theta^{\rho\sigma} (\{A_\rho, \partial_\sigma A_\mu + F_{\sigma\mu}\} + \{A_\rho, \partial_\sigma A_\mu + F_{\sigma\mu}\}_{(1)}) \\ &\quad + \{A_\rho, \partial_\sigma A_\mu + F_{\sigma\mu}\}_{\star(1)} \\ \Lambda &= -\frac{i}{8} \Theta^{\rho\sigma} (\{A_\rho, \partial_\sigma \Lambda\} + \{A_\rho, \partial_\sigma \Lambda\}_{(1)} + \{A_\rho, \partial_\sigma \Lambda\}_{\star(1)}), \end{aligned} \quad (3.12)$$

where the subscript $\star_{(n)}$ on the bracket indicates the n th order term in the Θ expansion of the star-anticommutator $\{A, B\}_\star^{AB} = A^AC \star B_C^B + B^AC \star A_C^B$.

Using (3.4), one next defines noncommutative vierbeins and spin connections through a power series expansion in Θ [1]:

$$\begin{aligned} \hat{e}_\mu^a(x, \Theta) &= e_\mu^a(x) + e_{\mu(1)}^a(x) + e_{\mu(2)}^a(x) + \dots \\ \hat{\omega}_\mu^{ab}(x, \Theta) &= \omega_\mu^{ab}(x) + \omega_{\mu(1)}^{ab}(x) + \omega_{\mu(2)}^{ab}(x) + \dots, \end{aligned} \quad (3.13)$$

which in turn is defined through the Seiberg-Witten map of the potential one-forms

$$\hat{e} = \hat{e}(e, \omega) \quad \hat{\omega} = \hat{\omega}(e, \omega). \quad (3.14)$$

The zeroth order again agrees with the commutative theory, while for the first and second orders one gets

$$e_{\mu}^a = -\frac{i}{4} \Theta^{\rho\sigma} ([\omega_{\rho}]^a_b (\partial_{\sigma} e_{\mu}^b + T_{\sigma\mu}^b) + (\partial_{\sigma} \omega_{\mu} + R_{\sigma\mu})^a_b e_{\rho}^b) \quad (3.15)$$

$$\omega_{\mu}^{ab} = -\frac{i}{4} \Theta^{\rho\sigma} \{\omega_{\rho}, \partial_{\sigma} \omega_{\mu} + R_{\sigma\mu}\}^{ab},$$

and

$$\begin{aligned} e_{\mu}^a &= -\frac{i}{8} \Theta^{\rho\sigma} ([\omega_{\rho}]^a_b (\partial_{\sigma} e_{\mu}^b + T_{\sigma\mu}^b) + [\omega_{\rho}]^a_b (\partial_{\sigma} e_{\mu}^b + T_{\sigma\mu}^b) + [\omega_{\rho}]^a_b \star_{(1)} (\partial_{\sigma} e_{\mu}^b + T_{\sigma\mu}^b) + (\partial_{\sigma} \omega_{\mu} + R_{\sigma\mu})^a_b e_{\rho}^b + (\partial_{\sigma} \omega_{\mu} + R_{\sigma\mu})^a_b e_{\rho}^b + (\partial_{\sigma} \omega_{\mu} + R_{\sigma\mu})^a_b \star_{(1)} e_{\rho}^b) \\ \omega_{\mu}^{ab} &= -\frac{i}{8} \Theta^{\rho\sigma} (\{\omega_{\rho}, \partial_{\sigma} \omega_{\mu} + R_{\sigma\mu}\}^{ab} + \{\omega_{\rho}, \partial_{\sigma} \omega_{\mu} + R_{\sigma\mu}\}^{ab}_{\star_{(1)}}), \end{aligned} \quad (3.16)$$

where the first order corrections to the curvature and torsion are defined as

$$\begin{aligned} R_{\mu\nu}^{ab} &= \partial_{\mu} \omega_{\nu}^{ab} - \partial_{\nu} \omega_{\mu}^{ab} + [\omega_{\mu}, \omega_{\nu}]^{ab} + [\omega_{\mu}, \omega_{\nu}]^{ab} + [\omega_{\mu}, \omega_{\nu}]_{\star_{(1)}}^{ab} \\ T_{\mu\nu}^a &= \partial_{\mu} e_{\nu}^a + [\omega_{\mu}]^a_b e_{\nu}^b + [\omega_{\mu}]^a_b e_{\nu}^b + [\omega_{\mu}]^a_b \star_{(1)} e_{\nu}^b - (\mu \rightleftharpoons \nu). \end{aligned} \quad (3.17)$$

For the discussion below we follow [7,8] and specialize to the case of zero torsion in the commutative theory; i.e.,

$$T_{\mu\nu}^a = 0. \quad (3.18)$$

Furthermore, in order to make a physical interpretation of the noncommutative vierbeins we define the real symmetric noncommutative version of the metric tensor according to

$$\hat{g}_{\mu\nu} = \frac{1}{2} \eta_{ab} (\hat{e}_{\mu}^a \star \hat{e}_{\nu}^{b*} + \hat{e}_{\nu}^b \star \hat{e}_{\mu}^{a*}). \quad (3.19)$$

IV. ROBERTSON-WALKER METRIC

We now apply the above formalism to the case of the Robertson-Walker metric. Starting with the usual expression for the Robertson-Walker invariant measure

$$ds^2 = -dt^2 + a(t)^2 \left(\frac{dr^2}{1-kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right), \quad (4.1)$$

where $a(t)$ is the scale factor, one can assign vierbein one-forms according to

$$\begin{aligned} e^0 &= dt & e^1 &= \frac{a(t)dr}{\sqrt{1-kr^2}} & e^2 &= a(t)r d\theta \\ e^3 &= a(t)r \sin\theta d\phi. \end{aligned} \quad (4.2)$$

The torsion vanishes upon choosing the following for the spin connection one-forms:

$$\begin{aligned} \omega^{01} &= \chi dr & \omega^{02} &= \dot{a} r d\theta & \omega^{03} &= \dot{a} r \sin\theta d\phi \\ \omega^{12} &= -\sqrt{1-kr^2} d\theta & \omega^{31} &= \sqrt{1-kr^2} \sin\theta d\phi \\ \omega^{23} &= -\cos\theta d\phi, \end{aligned} \quad (4.3)$$

where the dot denotes differentiation with respect to t . To determine χ one can compute the curvature scalar $\mathcal{R} = \mathcal{R}_{\mu\nu}{}^{\mu\nu}$ using (2.8), and compare with the known result for the Robertson-Walker metric; i.e.,

$$\mathcal{R} = 6 \left(\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 + \frac{k}{a^2} \right). \quad (4.4)$$

They agree for

$$\chi = \frac{\dot{a}}{\sqrt{1-kr^2}}. \quad (4.5)$$

Next we compute the second order noncommutative corrections to the metric tensor. For simplicity we set all components of $\Theta^{\mu\nu}$ equal to zero except for

$$\Theta^{tr} = -\Theta^{rt} = \Theta. \quad (4.6)$$

This choice leads to an isotropic inhomogeneous space-time. Up to second order in Θ , we find the following noncommutative vierbein one-forms after substituting into (3.13), (3.14), (3.15), and (3.16):

$$\begin{aligned} \hat{e}^0 &= dt + \frac{i\Theta}{4} \frac{\dot{a}^2 + 2a\ddot{a}}{1-kr^2} dr - \frac{5\Theta^2(\ddot{a}^2 + \dot{a}a^{(3)})}{32(1-kr^2)} dt \\ &\quad + \frac{r\Theta^2 k(9\dot{a}\ddot{a} - 2aa^{(3)})}{16(1-kr^2)^2} dr \\ \hat{e}^1 &= \frac{adr}{\sqrt{1-kr^2}} + \frac{i\Theta\dot{a}dt}{4\sqrt{1-kr^2}} - \frac{ir\Theta k\dot{a}dr}{4(1-kr^2)^{3/2}} \\ &\quad - \frac{3\Theta^2(3\ddot{a}\dot{a}^2 + 4aa^{(3)}\dot{a} + 4a\ddot{a}^2)}{32(1-kr^2)^{3/2}} dr \\ \hat{e}^2 &= \Phi d\theta & \hat{e}^3 &= \Phi \sin\theta d\phi, \end{aligned} \quad (4.7)$$

where

$$\Phi = ar - \frac{i\Theta}{4} \dot{a} - \frac{r\Theta^2(8a\ddot{a}^2 + (9\dot{a}^2 + 4k)\ddot{a} + 4a\dot{a}a^{(3)})}{32(1-kr^2)}, \quad (4.8)$$

and $a^{(3)}$ denotes the third time derivative of a . Only one off diagonal element of $\hat{g}_{\mu\nu}$ (3.19) results in these coordinates:

$$\begin{aligned}\hat{g}_{tt} &= -1 + \frac{\Theta^2(6\ddot{a}^2 + 5\dot{a}a^{(3)})}{16(1-kr^2)} + O(\Theta^4) \\ \hat{g}_{rr} &= \frac{a^2}{1-kr^2} - \frac{\Theta^2}{16(1-kr^2)^3} \\ &\quad \times ((1-kr^2)(\dot{a}^4 + 13a\ddot{a}\dot{a}^2 + 12a^2a^{(3)}\dot{a} + 16(a\ddot{a})^2) \\ &\quad + k(3kr^2 + 4)\dot{a}^2 + 4a\ddot{a}k(kr^2 + 1)) + O(\Theta^4) \\ \hat{g}_{\theta\theta} &= r^2a^2 + \frac{\Theta^2}{16} \left(-\frac{a(8a\ddot{a}^2 + (9\dot{a}^2 + 4k)\ddot{a} + 4a\dot{a}a^{(3)})r^2}{1-kr^2} \right. \\ &\quad \left. + 5\dot{a}^2 + 4a\ddot{a} \right) + O(\Theta^4) \\ \hat{g}_{\phi\phi} &= \sin^2\theta\hat{g}_{\theta\theta} \quad \hat{g}_{tr} = -\frac{r\Theta^2k\dot{a}\ddot{a}}{2(1-kr^2)^2} + O(\Theta^4). \quad (4.9)\end{aligned}$$

When interpreted as a metric tensor $\hat{g}_{\mu\nu}$ describes an inhomogeneous isotropic space-time with respect to the world line at $r = 0$.¹ We note that there are no second order corrections when the scale factor is a constant.

We wish to examine $\hat{g}_{\mu\nu}$ for small t (which we define later). As a general analysis with arbitrary scale factor is quite involved, we shall limit the discussion to a toy model. The simplest nontrivial example is the case of $a(t) = vt$, associated with a linear expansion in the commutative theory.² Here we can construct a scenario where there is no singularity at $t = 0$ to second order in Θ . We first note that the case of $a(t) = vt$ implies that the off diagonal matrix element \hat{g}_{tr} vanishes at second order and that the diagonal elements are invariant under $t \rightarrow -t$. If in the noncommutative theory we define the analogue of the invariant measure according to $d\hat{s}^2 = \hat{g}_{\mu\nu}dx^\mu dx^\nu$, it here has the form

$$\begin{aligned}d\hat{s}^2 &= -dt^2 + \frac{a_r(t,r)^2 dr^2}{1-kr^2} + r^2 a_\Omega(t,r)^2 (d\theta^2 + \sin^2\theta d\phi^2) \\ &\quad + O(\Theta^4), \quad (4.10)\end{aligned}$$

$$\mathcal{R} = \mathcal{R}_{\mu\nu}{}^{\mu\nu} = \frac{6(v^2 + k)}{t^2 v^2} - \frac{\Theta^2(k(v^4 + 8kv^2 + 7k^2)r^6 - (v^4 + 26kv^2 + 2k^2)r^4 + (11v^2 + 4k)r^2 + 5)}{8r^4 t^4 v^2 (1 - r^2 k)^2} + O(\Theta^4). \quad (4.15)$$

It is well behaved everywhere on the surfaces defined by (4.14) except at the spatial origin. It follows from the

¹This is not the case for generic $\Theta^{\mu\nu}$.

²If one further restricts $k = -v^2$, then the commutative theory corresponds to the Milne universe. In this case, all components of the Riemann curvature vanish and the commutative metric can be mapped into a region of Minkowski space using $(t, r) \rightarrow (t' = t\sqrt{1 + v^2 r^2}, r' = vtr)$.

where

$$\begin{aligned}a_r(t, r)^2 &= a(t)^2 - \frac{\Theta^2 v^2}{16} \left(\frac{v^2}{1-kr^2} + \frac{k(3kr^2 + 4)}{(1-kr^2)^2} \right) \\ a_\Omega(t, r)^2 &= a(t)^2 + \frac{5\Theta^2 v^2}{16r^2}. \quad (4.11)\end{aligned}$$

The second order correction to $a_\Omega(t, r)$ renders $\hat{g}_{\theta\theta}$ and $\hat{g}_{\phi\phi}$ nonsingular at $t = 0$. The second order correction to $a_r(t, r)^2$ is everywhere negative when

$$-\frac{v^2}{4} < k \leq 0, \quad (4.12)$$

which means that \hat{g}_{rr} is also everywhere nonsingular at $t = 0$. Thus when (4.12) holds, the leading corrections imply that there is no singularity at $t = 0$. Instead, the noncommutative metric tensor is everywhere well defined on the $t = 0$ time slice, which has a three-dimensional Minkowski signature $(-1, 1, 1)$. The same result applies for

$$k > 0, \quad 0 \leq r^2 < \frac{1}{k}. \quad (4.13)$$

(The metric tensor is ill defined at $r^2 = 1/k$ for the case of $k > 0$.) On the other hand, the noncommutative metric tensor is singular for these two cases when

$$t^2 = \frac{\Theta^2}{16} \left(\frac{v^2}{1-kr^2} + \frac{k(3kr^2 + 4)}{(1-kr^2)^2} \right), \quad (4.14)$$

the solutions of which define two disconnected surfaces, associated with positive and negative values for t . (For the choice of a dimensionful radial coordinate, Θ^{-1} , v^2 , and k have units of $1/\text{length}^2$.) One can compute the scalar curvature in order to determine whether the surfaces contain real (as opposed to coordinate) singularities. Treated as a space-time metric tensor, $\hat{g}_{\mu\nu}$ leads to the following (commutative) space-time scalar curvature:³

second order analysis that there are (at least) two singular points on the space-time manifold,

$$(t, r) = \left(\pm \frac{\Theta}{4} \sqrt{v^2 + 4k}, 0 \right), \quad (4.16)$$

³Alternatively, one can define a noncommutative analogue of the scalar curvature, as is done in [8]; however, its geometrical meaning is not obvious.

which go to the big bang singularity when $\Theta \rightarrow 0$.⁴ [Equation (4.16) can be used to define “small t ” in this case.] Unlike the big bang singularity, the two singular points in (4.16) are not the source of all timelike geodesics when $\Theta \neq 0$. To see this we next look at the geodesic equations. Call $u^\mu = \frac{dx^\mu}{d\sigma}$ where σ parametrizes the geodesic. Because of rotational invariance, we can consistently set $u^\theta = u^\phi = 0$. The geodesic equations for u^t and u^r then read

$$\begin{aligned} \frac{du^t}{d\sigma} &= -\frac{t(vu^r)^2}{1-kr^2} + O(\Theta^4) \\ \frac{du^r}{d\sigma} &= -\frac{kr(u^r)^2}{1-kr^2} - \frac{2u^t u^r}{t} \\ &\quad - \frac{\Theta^2}{16t^3(kr^2-1)^3} \{rtk((1-kr^2)v^2 \\ &\quad + k(3kr^2+11))(u^r)^2 - 2(1-kr^2)((1-kr^2)v^2 \\ &\quad + k(3kr^2+4))u^t u^r\} + O(\Theta^4). \end{aligned} \quad (4.17)$$

The comoving world lines $u^t = 1$, $u^r = 0$ of the commutative theory are unaffected by the second order corrections in Θ . Consequently, all of them, except for the central one at $r = 0$ which intersects the singular points (4.16), can be extended through the $t = 0$ time slice, and range from $t \rightarrow$

$-\infty$ to $t \rightarrow +\infty$. Therefore, although cosmic singularities are still present at leading order in Θ , they are no longer the source of all timelike geodesics. This result also holds when (4.12) or (4.13) are no longer true, as is the case with the Milne universe. Then there are singularities on the $t = 0$ time slice on the surface of a sphere of radius⁵

$$r = \sqrt{\frac{v^2 + 4k}{k(v^2 - 3k)}}, \quad (4.18)$$

but they are not the source of all timelike geodesics.

V. CONCLUDING REMARKS

It is of course of interest to go beyond the toy model considered in the previous section and consider more realistic functions for the scale parameter. Unfortunately, the analysis then becomes quite a bit more involved, and so we only briefly comment on a couple of examples here.

For the example of $a(t) = Ct^{1/2}$, which is standardly associated with a flat radiation dominated universe, the noncommutative metric tensor (4.9) is no longer diagonal in the coordinates (t, r, θ, ϕ) unless $k = 0$. From (4.9) we can compute the volume form for this case:

$$\det \hat{g}_{\mu\nu} = -\left(t^3 - \frac{\Theta^2(83r^2(1-kr^2)C^2 - 4t(-5k^2r^4 + 4kr^2 + 2))}{256r^2(1-kr^2)^2}\right) \frac{C^6 r^4 \sin^2 \theta}{1-kr^2}. \quad (5.1)$$

It is well behaved at $t = 0$ for $k \leq 0$ and $k > 0$, $0 \leq r^2 < \frac{1}{k}$, except for the origin $r = 0$. The origin appears to be a singularity in space-time from the expression for the space-time scalar curvature which in this case is

$$\begin{aligned} \mathcal{R} &= \mathcal{R}_{\mu\nu}{}^{\mu\nu} \\ &= \frac{6k}{C^2 t} + \frac{(933C^4(r^2k-1)r^4 - 4C^2t(82k^2r^4 - 284kr^2 + 21)r^2 + 16t^2(5k^3r^6 - 22k^2r^4 - 4kr^2 - 1))\Theta^2}{512C^2r^4t^5(1-kr^2)^2} + O(\Theta^4). \end{aligned} \quad (5.2)$$

More generally, upon setting the parenthesis in (5.1) equal to zero, one now gets a cubic equation in t , defining surfaces where the noncommutative metric tensor is singular. Equation (5.2) may be employed to determine whether or not points on these surfaces are coordinate singularities. The geodesic equations for u^t and u^r now have Θ^2 terms proportional to $(u^t)^2$, and so unlike in the previous case, the comoving world lines $u^t = 1$, $u^r = 0$ of the commutative theory are not geodesics of the noncommutative metric due to Θ^2 corrections.

⁴The scalar curvature given in (4.15) is still singular at $t = 0$. However, this is due to the truncation of the expansion in Θ . The exact expression for the scalar curvature which follows from the second order corrected metric tensor is well defined at $t = 0$ for (4.12) or (4.13).

Of course it is also of interest to consider the example of $a(t) = Ct^{2/3}$, which is standardly associated with a matter dominated universe. One can then try to perform spatial averages of the second order correction in this case in order to obtain the best fit for the map of the luminosity distance of the supernova versus the redshift along the lines of [20–26]. Since we are not required to make any particular choice for $\Theta^{\mu\nu}$, as in (4.6), a reasonable fit may be possible.

Lastly, we note that there are limitations to the procedure of Refs. [7,8] and adopted in this article. It is reasonable to expect that there could be other second order noncommutative gravitational effects which are not readily accessible from this procedure, such as:

⁵The argument given in the previous footnote suggests that these are coordinate singularities.

- (1) The scale factor $a(t)$ may receive corrections due to some noncommutative aspects of the matter contribution to the Einstein equations.
- (2) The geodesic equations for test particles can pick up noncommutative corrections.

The procedure of Refs. [7,8] was to map known solutions of general relativity to the noncommutative theory, using the Seiberg-Witten map, and then to define a noncommutative analogue of the metric tensor in order to give a physical interpretation of the results. On the other hand, the computation of both corrections (1) and (2). would require detailed knowledge of the noncommutative gravi-

tational field equations. [Concerning (2), the noncommutative particle equations of motion can presumably be obtained by taking some sort of particle limit of the noncommutative energy momentum conservation law which follows from the field equations, in a manner similar to what is done in the commutative theory.] However, as stated in the Introduction, a deformation of Einstein equations consistent with the noncommutative analogue of local Lorentz covariance is problematic in this approach. In order to know whether or not corrections such as (1) and (2) can influence physical predictions it is therefore important to pursue a better understanding of the noncommutative dynamics.

-
- [1] A. H. Chamseddine, Phys. Lett. B **504**, 33 (2001).
 - [2] P. Aschieri, C. Blohmann, M. Dimitrijevic, F. Meyer, P. Schupp, and J. Wess, Classical Quantum Gravity **22**, 3511 (2005).
 - [3] X. Calmet and A. Kobakhidze, Phys. Rev. D **72**, 045010 (2005); **74**, 047702 (2006).
 - [4] R. Banerjee, P. Mukherjee, and S. Samanta, Phys. Rev. D **75**, 125020 (2007).
 - [5] N. Seiberg and E. Witten, J. High Energy Phys. **09** (1999) 032.
 - [6] A. Stern, Phys. Rev. Lett. **100**, 061601 (2008); Phys. Rev. D **78**, 065006 (2008).
 - [7] M. Chaichian, A. Tureanu, and G. Zet, Phys. Lett. B **660**, 573 (2008); M. Chaichian, M.R. Setare, A. Tureanu, and G. Zet, J. High Energy Phys. **04** (2008) 064.
 - [8] P. Mukherjee and A. Saha, Phys. Rev. D **77**, 064014 (2008).
 - [9] B. P. Dolan, K. S. Gupta, and A. Stern, Classical Quantum Gravity **24**, 1647 (2007).
 - [10] A. Kobakhidze, arXiv:0712.0642.
 - [11] R. Banerjee, B. R. Majhi, and S. Samanta, Phys. Rev. D **77**, 124035 (2008); R. Banerjee, B. R. Majhi, and S. K. Modak, arXiv:0802.2176.
 - [12] M. Buric and J. Madore, arXiv:0807.0960.
 - [13] P. Nicolini, arXiv:0807.1939.
 - [14] P. Mukherjee and A. Saha, Phys. Rev. D **74**, 027702 (2006).
 - [15] E. Akofer, A. P. Balachandran, S. G. Jo, A. Joseph, and B. A. Qureshi, J. High Energy Phys. **05** (2008) 092; E. Akofer, A. P. Balachandran, A. Joseph, L. Pekowsky, and B. A. Qureshi, arXiv:0806.2458.
 - [16] F. Lizzi, G. Mangano, G. Miele, and M. Peloso, J. High Energy Phys. **06** (2002) 049.
 - [17] R. Brandenberger and P. M. Ho, Phys. Rev. D **66**, 023517 (2002).
 - [18] O. Bertolami and L. Guisado, Phys. Rev. D **67**, 025001 (2003).
 - [19] A. Krasinski, *Inhomogeneous Cosmological Models* (Cambridge University Press, Cambridge, England, 1997).
 - [20] S. Rasanen, J. Cosmol. Astropart. Phys. **02** (2004) 003.
 - [21] E. W. Kolb, S. Matarrese, A. Notari, and A. Riotto, Phys. Rev. D **71**, 023524 (2005).
 - [22] T. Buchert, Classical Quantum Gravity **22**, L113 (2005).
 - [23] Y. Nambu and M. Tanimoto, arXiv:gr-qc/0507057.
 - [24] A. E. Romano, Phys. Rev. D **75**, 043509 (2007).
 - [25] K. Enqvist, Gen. Relativ. Gravit. **40**, 451 (2008).
 - [26] I. Jakacka and J. Stelmach, Classical Quantum Gravity **18**, 2643 (2001).
 - [27] K. Ulker and B. Yapiskan, Phys. Rev. D **77**, 065006 (2008).
 - [28] For a recent review, see A. H. Chamseddine, Int. J. Geom. Methods Mod. Phys. **3**, 149 (2006).
 - [29] B. Jurco, S. Schraml, P. Schupp, and J. Wess, Eur. Phys. J. C **17**, 521 (2000).
 - [30] L. Bonora, M. Schnabl, M. M. Sheikh-Jabbari, and A. Tomasiello, Nucl. Phys. **B589**, 461 (2000).