

**Erratum: Branching ratios and polarization in $B \rightarrow VV, VA, AA$ decays
[Phys. Rev. D **78**, 094001 (2008)]**

Hai-Yang Cheng and Kwei-Chou Yang
(Received 26 January 2009; published 10 February 2009)

DOI: [10.1103/PhysRevD.79.039903](https://doi.org/10.1103/PhysRevD.79.039903)

PACS numbers: 13.25.Hw, 12.38.Bx, 14.40.-n, 99.10.Cd

For Eq. (3.2), the LCDA $\Phi^{M,h}$ for $h = \pm$ should read Φ_{\pm}^M for $i = 1, 2, 3, 4, 9, 10$ and Φ_{\mp}^M for $i = 5, 7$. In Table IV there are several typos in the experimental results of f_L and theoretical predictions of f_{\perp} for the decay modes $\bar{B}^0 \rightarrow \rho^0 \omega$, $B^- \rightarrow K^{*-} \omega$, and $\bar{B}^0 \rightarrow \bar{K}^{*0} \omega$. Only those three modes are shown in the following table.

| Decay | \mathcal{B} | | f_L | | f_{\perp} | |
|---|-----------------------------|--------|------------------------|------|------------------------|------|
| | Theory | Expt | Theory | Expt | Theory | Expt |
| $\bar{B}^0 \rightarrow \rho^0 \omega$ | $0.1^{+0.1+0.4}_{-0.1-0.0}$ | <1.5 | $0.55^{+0.47}_{-0.29}$ | | $0.22^{+0.16}_{-0.23}$ | |
| $B^- \rightarrow K^{*-} \omega$ | $3.5^{+0.4+3.0}_{-0.4-1.7}$ | <3.4 | $0.66^{+0.32}_{-0.38}$ | | $0.17^{+0.20}_{-0.17}$ | |
| $\bar{B}^0 \rightarrow \bar{K}^{*0} \omega$ | $3.0^{+0.5+2.9}_{-0.4-1.8}$ | <2.7 | $0.57^{+0.44}_{-0.46}$ | | $0.21^{+0.25}_{-0.22}$ | |