

Comment on “Two-photon decay width of the sigma meson”

Eef van Beveren*

Centro de Física Teórica, Departamento de Física, Universidade de Coimbra, P-3004-516 Coimbra, Portugal

Frieder Kleefeld[†] and George Rupp[‡]

Centro de Física das Interações Fundamentais, Instituto Superior Técnico, Technical University of Lisbon, P-1049-001 Lisboa Codex, Portugal

Michael D. Scadron[§]

Department of Physics, University of Arizona, Tucson, Arizona 85721, USA

(Received 12 November 2008; published 21 May 2009)

We comment on a recent paper by Giacosa, Gutsche, and Lyubovitskij [Phys. Rev. D **77**, 034007 (2008)], in which it is argued that a quarkonium interpretation of the σ meson should give rise to a much smaller two-photon decay width than commonly assumed. The reason for this claimed discrepancy is a term in the transition amplitude, necessary for gauge invariance, which allegedly is often omitted in the literature, including the work of the present authors. Here we show their claims to be incorrect by demonstrating, in the context of the quark-level linear σ model, that the recently extracted experimental values are compatible with a $q\bar{q}$ assignment for the σ , provided that meson loops are taken into account as well.

DOI: 10.1103/PhysRevD.79.098501

PACS numbers: 12.39.Ki, 13.25.Jx, 13.30.Eg, 13.40.Hq

I. INTRODUCTION

In a recent paper [1], Giacosa, Gutsche, and Lyubovitskij (GGL) studied the two-photon decay width of the σ meson, alias $f_0(600)$ [2,3], based on the presupposition that it is a $q\bar{q}$ state. They employed two simple perturbative sigma models, one purely local, comprising σ , π , quark and antiquark fields, and the other nonlocal, with only σ , q , and \bar{q} , besides an extended covariant vertex function. The principal result of their work was that, in contrast with what is generally assumed, a $q\bar{q}$ assignment for the σ should lead to a width $\Gamma_{\sigma\rightarrow\gamma\gamma}$ much smaller—probably even below 1 keV—than the recently reported value of (4.1 ± 0.3) keV resulting from an analysis by Pennington [4] (also see Ref. [5]), as well as the 3 prior values given in the 2006 [2] and 2008 [3] PDG tables. Therefore, GGL concluded that, if the large experimental $\gamma\gamma$ width is confirmed, a quarkonium interpretation of the σ is not favored, “*contrary to usual belief*.” As an explanation for their very small $\Gamma_{\sigma\rightarrow\gamma\gamma}$ prediction, GGL argued that a term in the quark-triangle loop diagram, necessary for gauge invariance, largely cancels the lead term, thus resulting in a small total amplitude. Moreover, GGL claimed that the former term is “*often neglected*,” including in previous work of ours and our coauthors [6–9].

In this comment, we shall show that GGL are mistaken on several points. First of all, we have not unduly neglected any term in the evaluation of the quark-triangle diagram in

Refs. [6–9]. When we disregarded the term in question, this was fully justified, since the term was zero or negligible. Second, the small $\Gamma_{\sigma\rightarrow\gamma\gamma}$ value obtained by GGL is a consequence of a very low σ mass, in combination with a relatively large constituent quark mass, at least in the local case. For the nonlocal Lagrangian, their tiny $\Gamma_{\sigma\rightarrow\gamma\gamma}$ value is rather an indication for the inadequacy of the Lagrangian itself. Third, we demonstrate, by explicit calculation, how important meson-loop contributions are, which is in principle admitted by GGL, but not concretized.

In Sec. II of this comment, we study in detail the two-photon width of the σ meson, in the context of the quark-level linear σ model (QLL σ M) [10], showing that a good agreement with data is achieved. In Sec. III we present our conclusions.

II. TWO-PHOTON WIDTH OF THE σ IN THE QLL σ M

Given the scalar amplitude structure [7,8,11] $\mathcal{M}\epsilon_\nu(k')\epsilon_\mu(k)(g^{\mu\nu}k' \cdot k - k'^\mu k^\nu)$, the rate for the decay of a scalar meson S into two photons reads

$$\Gamma(S \rightarrow \gamma\gamma) = \frac{m_S^3 |\mathcal{M}_{S\rightarrow\gamma\gamma}|^2}{64\pi}. \quad (1)$$

If one assumes, as GGL do, that the σ is a scalar $q\bar{q}$ state, then the principal contribution to the amplitude $\mathcal{M}_{\sigma\rightarrow\gamma\gamma}$ comes from the up and down quark-triangle diagrams (see e.g. Fig. 1 in Ref. [1]), yielding (with $N_c = 3$)

$$\mathcal{M}_{\sigma\rightarrow\gamma\gamma}^{n\bar{n}} = \frac{5\alpha}{3\pi f_\pi} 2\xi_n [2 + (1 - 4\xi_n)I(\xi_n)], \quad (2)$$

*eef@teor.fis.uc.pt

†kleefeld@cfif.ist.utl.pt

‡george@ist.utl.pt

§scadron@physics.arizona.edu

where $\alpha = e^2/4\pi$, $\xi_n = m_n^2/m_\sigma^2$ (n stands for u or d), and $I(\xi)$ is the triangle loop integral given by

$$I(\xi) \begin{cases} = \frac{\pi^2}{2} - 2\log^2\left[\sqrt{\frac{1}{4\xi}} + \sqrt{\frac{1}{4\xi} - 1}\right] + 2\pi i \log\left[\sqrt{\frac{1}{4\xi}} + \sqrt{\frac{1}{4\xi} - 1}\right] & (\xi \leq 0.25), \\ = 2\arcsin^2\left[\sqrt{\frac{1}{4\xi}}\right] & (\xi \geq 0.25). \end{cases} \quad (3)$$

These Eqs. (2) and (3) exactly correspond to Eqs. (2) and (4) in Ref. [1], with the proviso that GGL defined the σ - \bar{q} - q coupling in their Lagrangian as $g_\sigma/\sqrt{2}$ instead of our QLL σ M coupling g , the latter being related to f_π above via the Goldberger-Treiman relation $m_q = f_\pi g$ [6–9]. Note that we use here, just like GGL in their local approach as well as in the Nambu-Jona-Lasinio (NJL) model [12], a point σ - q - \bar{q} coupling, which is also done for the different two-meson couplings of the σ described below. However, in the QLL σ M these couplings should not be understood as perturbative parameters, since all couplings are self-consistently interrelated via dynamical generation and loop shrinking [10].

Ignoring for the moment possible meson-loop contributions as well as an $s\bar{s}$ component in the σ , we can use Eq. (2) to calculate $\Gamma_{\sigma \rightarrow \gamma\gamma}$, for different σ and quark masses. Also, we can check what the importance is of the term involving $I(\xi)$. However, let us first deal with the allegation by GGL that we had erroneously neglected the $I(\xi)$ term in previous work. Well, in Ref. [6] we simply worked in the, perfectly well-defined, NJL limit ($m_\sigma = 2m_q$) of the QLL σ M, in which the term in question vanishes identically, using quite reasonable σ and quark masses of 630 MeV and 315 MeV, respectively. The resulting $\Gamma_{\sigma \rightarrow \gamma\gamma}$, ignoring meson-loops, would then be ≈ 2.2 keV. But accounting for an estimate of the pion-loop contribution as well yielded the prediction of ≈ 3.8 keV [6], in good agreement with experiment, then and now. In Ref. [9], Eq. (101), again the NJL limit of the QLL σ M was used, but now also including an estimate for the kaon loop, besides the pion loop, leading to a slightly smaller result, but still very much larger than any of GGL’s predictions (also see Ref. [13]). Finally, in Refs. [7,8] $\Gamma_{\sigma \rightarrow \gamma\gamma}$ was not even considered, thus making the critique by GGL completely void. Moreover, note that in Ref. [7] we did use the full expressions of Eqs. (2) and (3) above when necessary, namely, in the detailed discussion of the $f_0(1370) \rightarrow \gamma\gamma$ width. Only when *estimating* the $f_0(980) \rightarrow \gamma\gamma$ width, and a possible $n\bar{n}$ admixture in the latter scalar resonance, did we use an approximate expression in Refs. [7,8], with totally immaterial consequences in view of the small scalar mixing angle.

Let us now carry out a more detailed analysis of $\Gamma_{\sigma \rightarrow \gamma\gamma}$ in a QLL σ M setting, employing Eqs. (2) and (3). Working beyond the chiral limit (CL), we may take the NJL value $m_\sigma = 675$ MeV for $m_n = 337.5$ MeV [14], where m_n stands for the nonstrange (up or down) quark mass. Still neglecting $n\bar{n}$ - $s\bar{s}$ mixing and meson loops, this gives $\Gamma_{\sigma \rightarrow \gamma\gamma}^{qq} \approx 2.7$ keV. Also away from the NJL limit, corre-

sponding to a mass $m_\sigma \approx 666$ MeV [14], the latter width gets reduced to ≈ 2.4 keV. If we now moreover allow for the admixture of a small $s\bar{s}$ component in the σ , with a nonstrange-strange mixing angle of, say, -10° [14], then we get $\Gamma_{\sigma \rightarrow \gamma\gamma}^{qq} \approx 2.5$ keV, for the often used [9] QLL σ M quark masses $m_n = 337.5$ MeV and $m_s = 486$ MeV. Note that this $s\bar{s}$ component, with amplitude

$$\mathcal{M}_{\sigma \rightarrow \gamma\gamma}^{s\bar{s}} = \frac{\sqrt{2}\alpha g}{3\pi m_s} 2\xi_s [2 + (1 - 4\xi_s)I(\xi_s)], \quad (4)$$

contributes with a weight factor of only $\sqrt{2}\alpha m_n/3\pi f_\pi m_s$ (using the GT relation $m_n = f_\pi g$), as compared to $5\alpha/3\pi f_\pi$ from Eq. (2) in the $n\bar{n}$ case, since the charge of a strange quark is $-1/3$ [7].

Next we are going to add meson-loop contributions as well. Now, in the framework of the QLL σ M, loops with charged mesons that couple to the σ include those with pions and kaons, as well as those with the scalar mesons $\kappa(800)$ and $a_0(980)$. The expression for a gauge-invariant meson-loop contribution to the two-photon amplitude mainly differs from the quark triangle in Eq. (2) because of the presence of a seagull graph (see e.g. Ref. [11], first paper), yielding a total amplitude

$$\mathcal{M}_{\sigma \rightarrow \gamma\gamma}^{MM} = -\frac{2g'\alpha}{\pi m_M^2} \left[-\frac{1}{2} + \xi I(\xi) \right], \quad \xi = \frac{m_M^2}{m_\sigma^2}, \quad (5)$$

where the minus sign stems from the opposite statistics with respect to the quark-loop case, and g' is the cubic QLL σ M meson coupling. For the charged-meson loops pertinent to the σ , we shall need the three-meson couplings [7,9,10]

$$\begin{aligned} g_{\sigma n\bar{n}, \pi\pi} &= \frac{\cos^2(\phi_S)m_\sigma^2 + \sin^2(\phi_S)m_{f_0(980)}^2 - m_{\pi^\pm}^2}{2f_\pi}, \\ g_{\sigma s\bar{s}, \pi\pi} &= 0, \\ g_{\sigma n\bar{n}, KK} &= \frac{\cos^2(\phi_S)m_\sigma^2 + \sin^2(\phi_S)m_{f_0(980)}^2 - m_{K^\pm}^2}{2f_K}, \\ g_{\sigma s\bar{s}, KK} &= \frac{\sin^2(\phi_S)m_\sigma^2 + \cos^2(\phi_S)m_{f_0(980)}^2 - m_{K^\pm}^2}{\sqrt{2}f_K}, \\ g_{\sigma n\bar{n}, \kappa\kappa} &= \frac{\cos^2(\phi_S)m_\sigma^2 + \sin^2(\phi_S)m_{f_0(980)}^2 - m_\kappa^2}{2(f_\pi - f_K)}, \\ g_{\sigma s\bar{s}, \kappa\kappa} &= \frac{\sin^2(\phi_S)m_\sigma^2 + \cos^2(\phi_S)m_{f_0(980)}^2 - m_\kappa^2}{\sqrt{2}(f_K - f_\pi)}, \\ g_{\sigma n\bar{n}, a_0a_0} &= 3g_{\sigma n\bar{n}, \pi\pi}, \quad g_{\sigma s\bar{s}, a_0a_0} = 0, \end{aligned} \quad (6)$$

where ϕ_S is the scalar mixing angle, and $f_K = f_\pi(m_s/m_n + 1)/2 \approx 1.22f_\pi$. The cubic coupling of the physical σ meson to these four meson-meson channels is then given by

$$g'_{\sigma,MM} = \cos(\phi_S)g_{\sigma n\bar{n},MM} - \sin(\phi_S)g_{\sigma s\bar{s},MM}. \quad (7)$$

Note that we neglect here small OZI-violating corrections to the QLL σ M three-meson couplings, just as in previous work of ours [7]. Such contributions will be included in a forthcoming study.

Now we are in a position to do a complete calculation of $\Gamma_{\sigma \rightarrow \gamma\gamma}$, with both quark and meson loops accounted for. Note that the imaginary part of $I(\xi)$, as given by the $\xi < 0.25$ case in Eq. (3), will be included for the pion-loop amplitude. If we choose again a scalar mixing angle of -10° and take $m_\kappa = 800$ MeV, we obtain a total two-photon width

$$\Gamma_{\sigma \rightarrow \gamma\gamma}^{q\bar{q}+MM} \approx 3.5 \text{ keV}. \quad (8)$$

This rate corresponds to a total amplitude modulus $|\mathcal{M}| = 4.88 \times 10^{-2} \text{ GeV}^{-1}$, which can be decomposed in terms of the partial quark- and meson-loop amplitudes

$$\begin{aligned} \Re \mathcal{M}_{n\bar{n}} &= 4.01 \times 10^{-2} \text{ GeV}^{-1}, \\ \Re \mathcal{M}_{s\bar{s}} &= 1.08 \times 10^{-2} \text{ GeV}^{-1}, \\ \mathcal{M}_{\pi\pi} &= (1.19 - i1.03) \times 10^{-2} \text{ GeV}^{-1}, \\ \mathcal{M}_{KK} &= -1.82 \times 10^{-3} \text{ GeV}^{-1}, \\ \mathcal{M}_{\kappa\kappa} &= -2.05 \times 10^{-3} \text{ GeV}^{-1}, \\ \mathcal{M}_{a_0a_0} &= -1.50 \times 10^{-3} \text{ GeV}^{-1}. \end{aligned} \quad (9)$$

Note that here the relative sign between quark and meson loops has already been included. Also observe that the kaon, κ , and $a_0(980)$ loops reduce the contribution of the pion loop, so that the net effect of the meson loops on the two-photon width is about +40%. Changing the scalar mixing angle to e.g. $\phi_S = -18^\circ$ [9] only reduces the total two-photon width by about 3%. So the predictions, for a wide range of reasonable mixing angles, are fully compatible with the corresponding PDG [2] data, and also not at odds with Pennington's recent results [4,5].

In contrast, the sensitivity of $\Gamma_{\sigma \rightarrow \gamma\gamma}$ to the σ mass is much stronger, which is obvious from Eq. (1), relating width and amplitude via m_σ cubed. This can also be seen in Fig. 2 of the paper [1] by GGL themselves, where e.g. an m_σ of 650 MeV, with $m_q = 350$ MeV, would yield a $\Gamma_{\sigma \rightarrow \gamma\gamma}^{q\bar{q}}$ of roughly 2.5 keV, so in agreement with the values 2.4–2.5 keV found above. However, by taking a very small m_σ of 440 MeV, as GGL choose to do, one obtains a much smaller $\Gamma_{\sigma \rightarrow \gamma\gamma}$, even when meson loops are included. For instance, if we assume the σ to be purely $n\bar{n}$ and take $m_q = 250$ MeV, $\Gamma_{\sigma \rightarrow \gamma\gamma}$ becomes ≈ 0.7 keV, even with the 3 meson-loop contributions included, which should be compared to GGL's value of 0.54 keV (see Table I of Ref. [1])

for the pure $q\bar{q}$ case. Neglecting in this scenario the term proportional to $I(\xi)$ would indeed increase our result of ≈ 0.7 keV to ≈ 1.4 keV, but this is of course an error we have not and will not make.

At this point, we also take exception at GGL's claim “...the results for $\Gamma_{\sigma \rightarrow \gamma\gamma}$ at a fixed pole mass of $M_\sigma = 440$ MeV as favored by recent theoretical and experimental works [16, 20],” where their Ref. 20 is our Ref. [2], i.e., the 2006 PDG Review of Particle Physics. It is simply false to state that the PDG favors a σ pole mass of 440 MeV. The truth is that the PDG listings mention “(400–1200) – i (250–500) OUR ESTIMATE,” for the $f_0(600)$ T -matrix pole (i.e., S -matrix pole) as a function of \sqrt{s} . On the other hand, the theoretical papers referred to by GGL include the Roy-equation analysis by Caprini, Colangelo, and Leutwyler [15], which indeed found 441 MeV for the real part of the σ S -matrix pole, besides an imaginary part of 272 MeV. However, it is a common mistake to confuse the real part of the pole with the “mass” of a broad resonance, especially when the resonance is certainly not of a pure Breit-Wigner (BW) type, like e.g. the σ , which is strongly distorted due to the $\pi\pi$ threshold and the Adler zero not far below [16]. Notice that, in the latter analysis, the mass of the σ at which the $\pi\pi$ phase shift passes through 90° —by definition the K -matrix pole—lies at 926 MeV. This does not mean that this *is* the σ mass, but just demonstrates the difficulty of assigning *any* specific mass to a broad non-BW resonance. Anyhow, our above choice of 666 MeV, in the context of the QLL σ M, is surely more reasonable than naively taking the real part of a pole that is already significantly lower than the “world average” [2,17] of σ poles.

To conclude this section, we note that the $Z = 0$ compositeness condition, discussed by GGL in the context of their nonlocal Lagrangian, is manifestly satisfied in the—nonperturbative and self-consistent—QLL σ M, provided $\xi = m_q^2/m_\sigma^2 \leq 0.25$, with g_σ not depending on m_σ , up to corrections of the order of 3% beyond the CL [9].

III. CONCLUSIONS

In the present comment we have shown that GGL incorrectly referred to and criticized our previous papers on the subject. Moreover, we have demonstrated, via an explicit and detailed calculation in the context of the QLL σ M, that the reported experimental values of $\Gamma_{\sigma \rightarrow \gamma\gamma}$ give quantitative support to a $q\bar{q}$ interpretation of the σ meson, provided that one uses a reasonable σ mass and also includes meson-loop contributions, besides the quark loop considered by GGL.

Finally, let us comment on the nonlocal Lagrangian employed by GGL besides the local one. Their justification was “However, the local approach is no longer applicable for values of M_σ close to threshold, as will be evident from the discussion of the next section.” Well, as already mentioned above, the QLL σ M is a *local* renormalizable field

theory, exactly satisfying the $Z = 0$ compositeness condition close to—but below—threshold, due to its nonperturbative and self-consistent formulation [10]. This condition can be rigorously described in both the QLL σ M and the NJL model, in terms of a log-divergent gap equation [18]. The latter can also be expressed via a four-dimensional ultraviolet cutoff Λ , resulting in a value $\Lambda \approx 2.3m_q$. For a nonstrange quark mass of 337.5 MeV, this gives $\Lambda \approx 750$ MeV, which is an energy scale that clearly separates the “elementary” σ from e.g. the “composite” ρ meson. For further details, we refer to Ref. [18].

In contrast, GGL prefer a nonlocal σ -model Lagrangian for the calculation of $\Gamma_{\sigma \rightarrow \gamma\gamma}$ in a $q\bar{q}$ scenario. In view of the

numerical results of the latter model, which produces even tinier values for $\Gamma_{\sigma \rightarrow \gamma\gamma}$ than their local approach, we are led to conclude that nature rather disfavors the nonlocal realization of chiral symmetry as proposed by GGL than a $q\bar{q}$ interpretation of the σ meson.

ACKNOWLEDGMENTS

This work was supported by the *Fundação para a Ciência e a Tecnologia* of the *Ministério da Ciência, Tecnologia e Ensino Superior* of Portugal, under Contract No. POCI/FP/81913/2007.

-
- [1] F. Giacosa, T. Gutsche, and V.E. Lyubovitskij, Phys. Rev. D **77**, 034007 (2008).
 - [2] W.M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
 - [3] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008). This edition of the PDG tables lists the same $\Gamma_{\sigma \rightarrow \gamma\gamma}$ data as the 2006 [2] edition, but including now also the 2006 result by Pennington [4].
 - [4] M.R. Pennington, Phys. Rev. Lett. **97**, 011601 (2006).
 - [5] M.R. Pennington, T. Mori, S. Uehara, and Y. Watanabe, Eur. Phys. J. C **56**, 1 (2008). In this very recent analysis, an estimate of the $\sigma \rightarrow \pi\pi$ coupling at the σ pole was taken from Ref. [15], resulting in a somewhat smaller value of $\Gamma_{\sigma \rightarrow \gamma\gamma}$ than in Ref. [4], viz. (3.1 ± 0.5) keV, for the favored solution.
 - [6] E. van Beveren, F. Kleefeld, G. Rupp, and M.D. Scadron, Mod. Phys. Lett. A **17**, 1673 (2002).
 - [7] F. Kleefeld, E. van Beveren, G. Rupp, and M.D. Scadron, Phys. Rev. D **66**, 034007 (2002).
 - [8] R. Delbourgo, D. s. Liu, and M.D. Scadron, Phys. Lett. B **446**, 332 (1999).
 - [9] M.D. Scadron, F. Kleefeld, and G. Rupp, arXiv:hep-ph/0601196.
 - [10] R. Delbourgo and M.D. Scadron, Mod. Phys. Lett. A **10**, 251 (1995); Int. J. Mod. Phys. A **13**, 657 (1998).
 - [11] A.S. Deakin, V. Elias, D.G.C. McKeon, M.D. Scadron, and A. Bramon, Mod. Phys. Lett. A **9**, 2381 (1994); L. Babukhadia, Ya.A. Berdnikov, A.N. Ivanov, and M.D. Scadron, Phys. Rev. D **62**, 037901 (2000); J.L. Lucio and M. Napsuciale, Phys. Lett. B **454**, 365 (1999).
 - [12] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
 - [13] M.D. Scadron, G. Rupp, F. Kleefeld, and E. van Beveren, Phys. Rev. D **69**, 014010 (2004); **69**, 059901 (2004).
 - [14] M.D. Scadron, AIP Conf. Proc. **1030**, 23 (2008).
 - [15] I. Caprini, G. Colangelo, and H. Leutwyler, Phys. Rev. Lett. **96**, 132001 (2006).
 - [16] D.V. Bugg, Phys. Lett. B **572**, 1 (2003); **595**, 556(E) (2004).
 - [17] E. van Beveren, <http://cft.fis.uc.pt/eef/sigkap.htm>.
 - [18] M.D. Scadron, Phys. Rev. D **57**, 5307 (1998).