

Solar system constraints on $f(\mathcal{G})$ gravity models

Antonio De Felice

*Theoretical and Mathematical Physics Group, Center for Particle Physics and Phenomenology, Louvain University,
2 Chemin du Cyclotron, 1348 Louvain-la-Neuve, Belgium*

Shinji Tsujikawa

*Department of Physics, Faculty of Science, Tokyo University of Science, 1-3, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan
(Received 10 July 2009; published 15 September 2009)*

We discuss solar system constraints on $f(\mathcal{G})$ gravity models, where f is a function of the Gauss-Bonnet term \mathcal{G} . We focus on cosmologically viable $f(\mathcal{G})$ models that can be responsible for late-time cosmic acceleration. These models generally give rise to corrections of the form $\epsilon(r/r_s)^p$ to the vacuum Schwarzschild solution, where $\epsilon = H_*^2 r_s^2 \ll 1$, r_s is the Schwarzschild radius of the Sun, and H_* is the Hubble parameter today. We generally estimate the strength of modifications to general relativity in order to confront models with a number of experiments, such as the deflection of light and the perihelion shift. We show that cosmologically viable $f(\mathcal{G})$ models can be consistent with solar system constraints for a wide range of model parameters.

DOI: [10.1103/PhysRevD.80.063516](https://doi.org/10.1103/PhysRevD.80.063516)

PACS numbers: 98.80.Cq, 04.50.Kd

I. INTRODUCTION

The modified gravity paradigm has been extensively studied over the last few years as a way to make gravity responsible for the observed acceleration of the Universe at large scales [1]. These models are attractive in the sense that cosmic acceleration can be realized without recourse to a dark energy matter component. Unlike the cosmological constant scenario, they generally give rise to a dynamical equation of state of dark energy that varies in time.

Among these modifications of gravity, the so-called $f(R)$ theory has become popular and has started to become a new branch of cosmology [2]. In this theory, the Lagrangian density f is a function of the Ricci scalar R . The $f(R)$ theory in the metric formalism can be proven to be classically equivalent to a class of Brans-Dicke theory with a Brans-Dicke parameter $\omega_{\text{BD}} = 0$ [3]. A scalar-field degree of freedom, called *scalaron* [4], can freely propagate to mediate a fifth force, unless the scalaron mass is heavy in the region of high density. For the compatibility with local gravity experiments, the $f(R)$ theory needs to approach the Lagrangian density $f(R) = R - 2\Lambda$ in general relativity (GR) for large values of R much larger than the present cosmological Ricci scalar R_0 [5]. Meanwhile, nontrivial deviation from the Λ CDM model can arise for R of the order of R_0 [6–11]. This leads to a number of interesting observational signatures, such as the modifications to the matter power spectrum [12] as well as to the weak lensing spectrum [13].

It is also possible to take into account a Gauss-Bonnet (GB) scalar \mathcal{G} that is a combination of the Ricci scalar R , the Ricci tensor $R_{\mu\nu}$, and the Riemann tensor $R_{\mu\nu\alpha\beta}$ [14,15]. This GB scalar, together with R , belongs to an infinite class of curvature invariants, the Lovelock scalars, which have the property that they do not introduce deriva-

tive terms higher than two into the equations of motion for the metric tensor. Among these scalars, R and \mathcal{G} are the only ones that do not identically vanish in four dimensions (apart from the cosmological constant). However, the term $\sqrt{-g}\mathcal{G}$ is a total derivative in four dimensions, where g is the determinant of the metric tensor. This means that the only way for the GB term to contribute to the equations of motion is to couple it to something else, e.g., a scalar field ϕ with the coupling of the form $F(\phi)\mathcal{G}$ [16]. This kind of coupling is present in the low energy effective action of string theory [17], due to the presence of dilaton-graviton mixing terms.

The dynamics of dark energy based on the dilatonic coupling $F(\phi) \propto e^{\mu\phi}$ with the exponential potential $V(\phi) \propto e^{-\lambda\phi}$ has been studied by a number of authors [18–22]. While it is possible to realize a scaling matter era [19,20] followed by a late-time cosmic acceleration, the region of viable parameter space to satisfy several observational constraints is restricted to be very small [19]. It was also shown in Refs. [20,23] that tensor perturbations tend to exhibit negative instabilities if the GB term is responsible for cosmic acceleration. Moreover, in such models, the energy fraction of the GB term needs to be strongly suppressed for the compatibility with local gravity experiments [24], which is at odds with the requirement of cosmic acceleration induced by the GB term.

There is another class of modified gravity models in which the Lagrangian density is described by $R + f(\mathcal{G})$ (so-called “ $f(\mathcal{G})$ gravity”), where f is the function in terms of the GB term \mathcal{G} [25]. Unlike $f(R)$ gravity, this theory does not have an action in the Einstein frame with a standard kinetic term of a scalar-field degree of freedom. The conditions for cosmological viabilities of $f(\mathcal{G})$ gravity have been studied in Refs. [26–28] (see also Refs. [29,30]). Li *et al.* [26] showed that the condition $0 < H^6 f_{,\mathcal{G}\mathcal{G}} \ll 1$

(where $f_{,\mathcal{G}\mathcal{G}} \equiv d^2f/d\mathcal{G}^2$) needs to be fulfilled in order to keep cosmological perturbations under control. In Ref. [27], the same condition has been derived to ensure the stability of a late-time de Sitter solution as well as the existence of standard radiation and matter-dominated epochs. In particular, the stability of the de Sitter point requires the condition $0 < H_1^6 f_{,\mathcal{G}\mathcal{G}} < 1/384$, where H_1 is the Hubble parameter at this point. In order to remove possible singularities in the cosmic expansion history, the second derivative $f_{,\mathcal{G}\mathcal{G}}$ should not change sign, i.e. $f_{,\mathcal{G}\mathcal{G}} > 0$ for all \mathcal{G} , together with the condition that $f_{,\mathcal{G}\mathcal{G}} \rightarrow 0$ as $|\mathcal{G}| \rightarrow \infty$. This removes the presence of unstable modes during the whole past evolution of the Universe. A number of cosmologically viable models satisfying these requirements have been proposed in Ref. [27].

In this paper, we will study the property of $f(\mathcal{G})$ gravity on solar system scales and investigate whether cosmologically viable $f(\mathcal{G})$ models can be consistent with solar system constraints. We first find approximate vacuum solutions for these models through an iterative method introduced in Ref. [31]. The solutions look like corrections to the Schwarzschild solution, where the corrections are typically in the form of positive powers in the ratio r/r_s , where r is the distance from the center of the compact object and r_s is the Schwarzschild radius. This behavior of positive powers is similar to the typical correction that the cosmological constant gives to the Schwarzschild solution [$\propto (r/r_s)^2$]. In the case of the Schwarzschild-de Sitter solution, the metric in the solar system is dominated by the term r_s/r , but one can put upper bounds on the value of the cosmological constant such that its contribution is allowed by experimental data. We follow a similar procedure in order to constrain the values of model parameters in $f(\mathcal{G})$ gravity. We will show that cosmologically viable $f(\mathcal{G})$ models can satisfy solar system constraints for a wide range of parameter space.

In Sec. II, we briefly review cosmologically viable $f(\mathcal{G})$ models. In Sec. III, we describe the method used to find approximate spherically symmetric solutions of the Einstein equations. In Sec. IV, we discuss solar system constraints in the presence of positive powers of the radius correction to the Schwarzschild solution. In Secs. V, VI, and VII, we apply the constraints to a number of $f(\mathcal{G})$ models. In Sec. VIII, we report our conclusions.

II. COSMOLOGICALLY VIABLE $f(\mathcal{G})$ MODELS

Let us first briefly review cosmologically viable $f(\mathcal{G})$ models proposed in Ref. [27]. The action describing this theory is given by

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} R + f(\mathcal{G}) \right] + S_m(g_{\mu\nu}, \Psi_m), \quad (1)$$

where R is the Ricci scalar, $\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta}$ is the GB term, and S_m is a matter action

that depends on a spacetime metric $g_{\mu\nu}$ and on matter fields Ψ_m . We use the unit $M_{\text{pl}} = 1/\sqrt{8\pi G_N} = 1$, but we restore the reduced Planck mass M_{pl} and the gravitational constant G_N if required.

The late-time cosmic acceleration can be realized by the presence of a de Sitter (dS) point satisfying the condition $3H_1^2 = \mathcal{G}_1 f_{,\mathcal{G}}(\mathcal{G}_1) - f(\mathcal{G}_1)$, where H_1 and \mathcal{G}_1 are the Hubble parameter and the GB term at the dS point, respectively. The stability of the dS point demands the condition $0 < H_1^6 f_{,\mathcal{G}\mathcal{G}}(H_1) < 1/384$ [27]. The GB term $\mathcal{G} = 24H^2(H^2 + \dot{H})$ changes sign from negative to positive during the transition from the matter era to the accelerated epoch. For the existence of standard radiation and matter eras, we require that $f_{,\mathcal{G}\mathcal{G}} > 0$ for $\mathcal{G} \leq \mathcal{G}_1$ and that $f_{,\mathcal{G}\mathcal{G}}$ approaches $+0$ in the limit $|\mathcal{G}| \rightarrow \infty$. We also need the regularities of $f(\mathcal{G})$ and its derivatives $f_{,\mathcal{G}}$, $f_{,\mathcal{G}\mathcal{G}}$. The following two models can satisfy these conditions [27]:

$$f(\mathcal{G}) = \lambda \frac{\mathcal{G}}{\sqrt{\mathcal{G}_*}} \arctan\left(\frac{\mathcal{G}}{\mathcal{G}_*}\right) - \frac{1}{2} \lambda \sqrt{\mathcal{G}_*} \ln\left(1 + \frac{\mathcal{G}^2}{\mathcal{G}_*^2}\right) - \alpha \lambda \sqrt{\mathcal{G}_*}, \quad (2)$$

$$f(\mathcal{G}) = \lambda \frac{\mathcal{G}}{\sqrt{\mathcal{G}_*}} \arctan\left(\frac{\mathcal{G}}{\mathcal{G}_*}\right) - \alpha \lambda \sqrt{\mathcal{G}_*}, \quad (3)$$

where α , λ , and \mathcal{G}_* are positive constants. Note that \mathcal{G}_* roughly corresponds to the scale H_*^4 for α and λ of the order of unity, where H_* is the Hubble parameter today. The second derivative of f with respect to \mathcal{G} is $f_{,\mathcal{G}\mathcal{G}} = \lambda/[\mathcal{G}_*^{3/2}(1 + \mathcal{G}^2/\mathcal{G}_*^2)]$ and $f_{,\mathcal{G}\mathcal{G}} = 2\lambda/[\mathcal{G}_*^{3/2}(1 + \mathcal{G}^2/\mathcal{G}_*^2)^2]$ for the models in Eqs. (2) and (3), respectively, so that $f_{,\mathcal{G}\mathcal{G}} > 0$ for $\lambda > 0$.

In the region of high density where local gravity experiments are carried out ($|\mathcal{G}| \gg \mathcal{G}_*$), the above models have the following asymptotic behavior:

$$f(\mathcal{G}) \simeq \frac{1}{2} \pi \lambda \frac{\mathcal{G}}{\sqrt{\mathcal{G}_*}} - (\alpha + 1) \lambda \sqrt{\mathcal{G}_*} - \frac{1}{2} \lambda \sqrt{\mathcal{G}_*} \ln\left(\frac{\mathcal{G}^2}{\mathcal{G}_*^2}\right) - \frac{\lambda \sqrt{\mathcal{G}_*}}{6} \frac{\mathcal{G}_*^2}{\mathcal{G}^2}, \quad (4)$$

$$f(\mathcal{G}) \simeq \frac{1}{2} \pi \lambda \frac{\mathcal{G}}{\sqrt{\mathcal{G}_*}} - (\alpha + 1) \lambda \sqrt{\mathcal{G}_*} + \frac{\lambda \sqrt{\mathcal{G}_*}}{3} \frac{\mathcal{G}_*^2}{\mathcal{G}^2}. \quad (5)$$

The first terms in Eqs. (4) and (5) are linear in \mathcal{G} , so they do not give rise to any contribution to the Einstein equation, whereas the second terms contribute to the field equation as a cosmological constant. The other terms in Eqs. (4) and (5) correspond to the corrections to the Λ CDM model. The difference between the models in Eqs. (4) and (5) is that the former has a logarithmic correction that mildly increases with the growth of $|\mathcal{G}|$. Note that the viable $f(R)$ models, such as (i) $f(R) = R - \lambda R_*(R/R_*)^{2n}/[(R/R_*)^{2n} + 1]$ and

(ii) $f(R) = R - \lambda R_* [1 - (1 + R^2/R_*^2)^{-n}]$ ($n > 0$), behave as $f(R) \simeq R - \lambda R_* + \lambda R_* (R_*/R)^{2n}$ in the region of high density ($R \gg R_*$). This asymptotic form is similar to the model in Eq. (5) given above by replacing R for \mathcal{G} .

In the following, we shall study solar system constraints on cosmologically viable $f(\mathcal{G})$ models. Before doing so, it is worth mentioning the difference between $f(R)$ and $f(\mathcal{G})$ theories. If we consider a spherically symmetric background, the Schwarzschild vacuum solution corresponds the vanishing Ricci scalar ($R = 0$). In the presence of nonrelativistic matter, R approximately equals the matter density ρ_m/M_{pl}^2 for viable $f(R)$ models [9–11]. Then the term $(R_*/R)^{2n}$ is roughly of the order of $(\rho_c/\rho_m)^{2n}$, where ρ_c is the cosmological density today. The ratio $(\rho_c/\rho_m)^{2n}$ becomes much smaller than 1 for $n > 0$ in the region of high density ($\rho_m \gg \rho_c$), so one has $(R_*/R)^{2n} \ll 1$. In the presence of nonrelativistic matter, the chameleon mechanism [32] can be at work for the $f(R)$ models that have the asymptotic form $f(R) \simeq R - \lambda R_* + \lambda R_* (R_*/R)^{2n}$ in the region $R \gg R_*$, which allows the possibility for consistency with local gravity tests. In fact, it was shown in Ref. [33] that these models can satisfy solar system and equivalence principle constraints for $n > 0.9$.

On the contrary, one has $\mathcal{G} = R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta} = 12r_s^2/r^6$ on the vacuum Schwarzschild solution, where $r_s = 2G_N M_\odot$ is the Schwarzschild radius and M_\odot is the mass of the star. Since \mathcal{G} does not vanish even in the vacuum, the term such as $(\mathcal{G}_*/\mathcal{G}^2)^n$ ($n > 0$) can be much smaller than 1 even in the absence of nonrelativistic matter. If matter is present, this will give rise to the contribution of the order of $R^2 \approx (8\pi G_N \rho_m)^2$ to the GB term. The ratio of the matter contribution to the vacuum GB value $\mathcal{G}^{(0)} = 12r_s^2/r^6 = 48(G_N M_\odot)^2/r^6$ can be estimated as

$$s \equiv \frac{R^2}{\mathcal{G}^{(0)}} \approx \frac{(8\pi)^2}{48} \frac{\rho_m^2 r^6}{M_\odot^2}. \quad (6)$$

As long as $s \ll 1$, we can neglect the matter contribution to the GB term.

At the surface of the Sun (radius $r_\odot = 6.96 \times 10^{10}$ cm = 3.53×10^{24} GeV $^{-1}$ and mass $M_\odot = 1.99 \times 10^{33}$ g = 1.12×10^{57} GeV), the density ρ_m drops down rapidly from the order $\rho_m \approx 10^{-2}$ g/cm 3 to the order $\rho_m \approx 10^{-16}$ g/cm 3 . If we take the value $\rho_m = 10^{-2}$ g/cm 3 , we have $s \approx 4 \times 10^{-5}$ (where we have used $1 \text{ g/cm}^3 = 4.31 \times 10^{-18} \text{ GeV}^4$). Taking the value $\rho_m = 10^{-16}$ g/cm 3 leads to a much smaller ratio: $s \approx 4 \times 10^{-33}$. The matter density approaches a constant value $\rho_m \approx 10^{-24}$ g/cm 3 around the distance $r = 10^3 r_\odot$ from the center of the Sun. Even at this distance we have $s \approx 4 \times 10^{-31}$, which means that the matter contribution to the GB term can be completely neglected in the solar system we are interested in. If we use the value $\rho_m \approx 10^{-24}$ g/cm 3 , s exceeds the order of 1 for the distance $r \geq 10^8 r_\odot$. However, this is out of the region where solar system experiments are concerned. Moreover, the

Schwarzschild solution has no meaning far away from the star where other contributions can arise, i.e. other close stars, the mean field of the galaxy, and so on. From the above discussion, we find that vacuum solutions can be used when we discuss solar system constraints on $f(\mathcal{G})$ gravity.

III. EXPANSION AROUND THE SPHERICALLY SYMMETRIC SPACETIME

The cosmologically viable $f(\mathcal{G})$ models (2) and (3) will consist of a numerical factor of order $\sqrt{\mathcal{G}_*}$ times a dimensionless function (because $f(\mathcal{G})$ has the dimension of [mass] 2). Meanwhile, the typical value of the GB term on the vacuum Schwarzschild solution is given by

$$\mathcal{G}_s \equiv 12/r_s^4. \quad (7)$$

When we discuss solar system constraints, it is convenient to define the following dimensionless ratio:

$$\varepsilon \equiv \sqrt{\frac{\mathcal{G}_*}{\mathcal{G}_s}}. \quad (8)$$

Since $\sqrt{\mathcal{G}_*}$ is of the order of the square of the present Hubble parameter H_* , the parameter ε is approximately given by $\varepsilon \approx (H_* r_s)^2 / (2\sqrt{3})$. In the subsequent sections, we shall discuss the case of the Sun with the Schwarzschild radius $r_s = 2.95 \times 10^3$ m. Using the value $H_* \approx 70 \text{ km sec}^{-1} \text{ Mpc}^{-1}$, the parameter ε for the Sun is approximately given by

$$\varepsilon \approx 10^{-46}. \quad (9)$$

The model (2) can be written in the form $f(\mathcal{G}) = \varepsilon[\lambda\sqrt{\mathcal{G}_s}g(x) - \alpha\lambda\sqrt{\mathcal{G}_s}]$, where $g(x) = x \arctan x - (1/2)\ln(1+x^2)$ and $x = \mathcal{G}/\mathcal{G}_* = \mathcal{G}/(\mathcal{G}_s \varepsilon^2)$. Hence, the function f can be replaced by the form $f = \varepsilon \tilde{f}$.

The equations of motion in the vacuum can be written as

$$G^\mu{}_\nu + \varepsilon \Sigma^\mu{}_\nu = 0, \quad (10)$$

where $G^\mu{}_\nu$ is the Einstein tensor, and

$$\begin{aligned} \Sigma_{\mu\nu} = & 8[R_{\mu\rho\nu\sigma} + R_{\rho\nu\sigma\mu} - R_{\rho\sigma\nu\mu} - R_{\mu\nu\sigma\rho} \\ & + R_{\mu\sigma\nu\rho} + R(g_{\mu\nu}g_{\sigma\rho} - g_{\mu\sigma}g_{\nu\rho})/2] \nabla^\rho \nabla^\sigma \tilde{f}_{,\mathcal{G}} \\ & + (\mathcal{G}\tilde{f}_{,\mathcal{G}} - \tilde{f})g_{\mu\nu}. \end{aligned} \quad (11)$$

In general, these theories will have vacuum solutions, which we wish to study here. Although exact solutions are not always found analytically, it is possible to obtain approximate solutions which reproduce the real ones very well at least on some scales. In fact, since $\varepsilon \ll 1$, we can try to use the iterative method introduced in Ref. [31].

We look for static spherical symmetric solutions of the kind

$$ds^2 = -A(r, \varepsilon)dt^2 + \frac{dr^2}{B(r, \varepsilon)} + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (12)$$

where the functions A and B are Taylor expanded in the form

$$A(r, \varepsilon) = A_0(r) + A_1(r)\varepsilon + A_2(r)\varepsilon^2 + \dots, \quad (13)$$

$$B(r, \varepsilon) = B_0(r) + B_1(r)\varepsilon + B_2(r)\varepsilon^2 + \dots \quad (14)$$

Using this expansion for A and B , both $G^\mu{}_\nu(r, \varepsilon)$ and $\Sigma^\mu{}_\nu(r, \varepsilon)$ can be expanded order by order in ε . At the lowest level, the equations of motion correspond to

$$G^\mu{}_\nu{}^{(0)} = 0, \quad (15)$$

which leads to the usual Schwarzschild solution $B_0 = A_0 = 1 - r_s/r$. At first order, one has

$$\varepsilon[G^\mu{}_\nu{}^{(1)}(A_1, B_1, A_0, B_0) + \Sigma^\mu{}_\nu{}^{(0)}(A_0, B_0)] = 0, \quad (16)$$

which reduces to ordinary differential equations for A_1, B_1 , where A_0, B_0 are the Schwarzschild terms found previously. This method can be reiterated to get the coefficients A_2, B_2 , as well as all other higher-order terms.

It should be noted that, for a general approach to the modification of gravity, including quintessence, there is no more analogue of the Birkhoff theorem regarding the unicity of the solution for a spherically symmetric vacuum solution. We can only impose to have a static metric. The bottom line is that the forms of A and B can be found at any order- ε approximation, by solving the Einstein equations iteratively with respect to the small parameter $0 < \varepsilon \ll 1$.

Suppose that we find such an iterative solution and write both A and B as power expansions of ε . Then the iterative solution should have in general a radius of convergence, inside which each correction is larger than the next-order ε term. In this case we expect that the dominant term corresponds to the Schwarzschild contribution. Therefore in the small ε limit we will have

$$A = 1 - 1/\rho + \varepsilon c_1 \rho^p + \varepsilon^2 c_3 \rho^m + \mathcal{O}(\varepsilon^3), \quad (17)$$

$$B = 1 - 1/\rho + \varepsilon c_2 \rho^q + \varepsilon^2 c_4 \rho^n + \mathcal{O}(\varepsilon^3), \quad (18)$$

where $\rho \equiv r/r_s$. In Secs. VI and VII we will show that the correction terms in Eqs. (17) and (18) in fact arise for the cosmologically viable $f(\mathcal{G})$ models given in (2) and (3). We will restrict ourselves to the region $\rho \gg 1$, which is generically satisfied in the solar system outside of the Sun. In other words, this corresponds to the weak limit of the theory. We have introduced $c_{1,2,3,4}$ as constants whose values need to be bounded experimentally, and also p, q, m, n as the powers of ρ . In the following, we will restrict our attention to the case $p = q > 0$ and $m = n > 0$. Later we will see that this assumption is quite general for cosmologically viable $f(\mathcal{G})$ models.

In order to have a meaningful ε expansion, one needs to verify that there exists a set of convergence, e.g.,

$$D = \{\rho | \rho \gg 1, \varepsilon \rho^p \ll \rho^{-1}\}. \quad (19)$$

This implies that the expansions (17) and (18) can be trusted only in this region. Outside of it, one must find the solutions of Einstein equations, both analytically or numerically, without using the ε expansion. This also implies that it does not make sense to study this solution in the limit $\rho \rightarrow \infty$. However, even if the full solution is known, it does not have physical meaning for values of r where the solar system begins to feel other stars' contributions. This situation is analogous to the Schwarzschild-de Sitter solution, where the same solution cannot be trusted any more at distances a few parsecs away from the Sun. In this region, the solar system cannot be treated as an entity isolated from the rest of the galaxy. Of course, for larger p , the set D becomes smaller. However, this is not enough: one should also verify that in the same set the ε^2 term is much smaller than the ε one. This implies that in D , one is also required to have

$$D_2 = \{\rho | \rho \gg 1, \varepsilon \rho^m \ll \rho^p\} \subseteq D. \quad (20)$$

If $0 \leq m \leq 2p + 1$, then this condition is verified. We will call a good ε expansion for the one where this last condition is valid. In general, we may have more complicated cases where there exists an order of expansion d such that $D_i \subseteq D_d$ for all $i \geq 1$ (and d might not be one), such that the expansion makes sense in D_d . If there is no such set, then the expansion does not make sense. If there is such a set, we can define $D = D_d$.

If $D_2 \subseteq D$, then, in D (or, more in general, in D_d), it is safe to approximate the perturbative solution as

$$A = 1 - 1/\rho + \varepsilon c_1 \rho^p, \quad B = 1 - 1/\rho + \varepsilon c_2 \rho^p. \quad (21)$$

In the next section we shall study a number of solar system bounds for the metric (21).

IV. SOLAR SYSTEM CONSTRAINTS

There are a number of solar system constraints on the deviation from general relativity, such as (a) deflection of light, (b) Cassini experiment, (c) Perihelion shift, (d) retardation of light, and (e) gravitational redshift. In the following, we discuss those constraints for a general metric in the form of (21).

A. Deflection of light

The first constraint we discuss is the deflection of light. The Lagrangian for a photon moving in the $\theta = \pi/2$ plane in the gravitational field of the metric (12) is given by

$$L = \frac{1}{2}A\dot{t}^2 - \frac{1}{2}B^{-1}\dot{r}^2 - \frac{1}{2}r^2\dot{\varphi}^2 = 0, \quad (22)$$

where a dot represents an affine parameter along the geo-

desics. There are two constants of motion, namely

$$E \equiv \dot{A}t \quad \text{and} \quad L \equiv r^2 \dot{\phi}. \quad (23)$$

Then we find

$$\dot{r}^2 = L^2 \left(\frac{E^2}{L^2} \frac{B}{A} - \frac{B}{r^2} \right). \quad (24)$$

The minimal distance r_0 can be defined such that $\dot{r}(r_0) = 0$, giving

$$\frac{L^2}{r_0^2 E^2} = \frac{\rho_0^2}{A(\rho_0)}, \quad (25)$$

where $\rho_0 \equiv r_0/r_s$.

Integrating $d\varphi/dr = \dot{\phi}/\dot{r}$ by using Eqs. (23)–(25), we obtain

$$\varphi(\rho) = \pm \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\rho_0}{\bar{\rho}} \sqrt{\frac{A(\bar{\rho})}{B(\bar{\rho})[A(\rho_0)\bar{\rho}^2 - A(\bar{\rho})\rho_0^2]}} \quad (26)$$

where $\bar{\rho}$ is used to avoid the confusion with the upper limit

$$F(\rho_0, \rho) = \left[\frac{\rho\rho_0}{\rho^3(\rho_0 - 1) - \rho\rho_0^3 + \rho_0^3} \right]^{3/2} \frac{\rho^p \{c_1 \rho^3(\rho_0 - 1) + c_2 [\rho^3(1 - \rho_0) + (\rho - 1)\rho_0^3]\} - c_1(\rho - 1)\rho^2 \rho_0^{p+1}}{2\rho(\rho - 1)}. \quad (28)$$

We can further simplify this expression by considering the limits $\rho, \rho_0 \gg 1$:

$$\begin{aligned} \varphi &= \pm \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\rho_0}{\bar{\rho} \sqrt{\bar{\rho}^2 - \rho_0^2}} \\ &\pm \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\bar{\rho}^2 + \rho_0 \bar{\rho} + \rho_0^2}{2\bar{\rho}^2(\bar{\rho} + \rho_0) \sqrt{\bar{\rho}^2 - \rho_0^2}} \\ &\pm \frac{1}{2} \varepsilon \rho_0 \int_{\rho_0}^{\rho} d\bar{\rho} \frac{c_1 \bar{\rho}^2(\bar{\rho}^p - \rho_0^p) - c_2 \bar{\rho}^p(\bar{\rho}^2 - \rho_0^2)}{\bar{\rho}(\bar{\rho}^2 - \rho_0^2)^{3/2}}. \end{aligned} \quad (29)$$

Taking the positive sign in front of each integral, the deviation angle in the region D is given by

$$\begin{aligned} \vartheta(\rho) &= 2 \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\rho_0}{\bar{\rho} \sqrt{\bar{\rho}^2 - \rho_0^2}} \\ &+ \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\bar{\rho}^2 + \rho_0 \bar{\rho} + \rho_0^2}{\bar{\rho}^2(\bar{\rho} + \rho_0) \sqrt{\bar{\rho}^2 - \rho_0^2}} \\ &+ \varepsilon \rho_0 \int_{\rho_0}^{\rho} d\bar{\rho} \frac{c_1 \bar{\rho}^2(\bar{\rho}^p - \rho_0^p) - c_2 \bar{\rho}^p(\bar{\rho}^2 - \rho_0^2)}{\bar{\rho}(\bar{\rho}^2 - \rho_0^2)^{3/2}} \\ &- \pi, \end{aligned} \quad (30)$$

in the limit that $\rho \gg \rho_0$. The first two contributions in Eq. (29), corresponding to the GR ones, give

ρ of the integral. The Schwarzschild-de Sitter solution corresponds to $c_1 = c_2$ and $p = 2$, in which case the integral (26) reduces to the standard GR contribution. Therefore, the cosmological constant does not give any modification to GR for light bending [34]. In standard GR, the integral is taken in the limit $\rho \rightarrow \infty$. This is a mathematical extrapolation, as the metric does not hold in the whole spacetime. Of course this property holds in our approach, where the approximate metric is sensible only in D , the domain of convergence. Up to first order in ε and in the domain D , it is possible to approximate the integral as

$$\begin{aligned} \varphi &= \pm \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\rho_0}{\bar{\rho}} \sqrt{\frac{\bar{\rho}\rho_0}{\bar{\rho}^3(\rho_0 - 1) - \rho_0^3(\bar{\rho} - 1)}} \\ &+ \varepsilon \int_{\rho_0}^{\rho} d\bar{\rho} F(\rho_0, \bar{\rho}), \end{aligned} \quad (27)$$

where

$$\vartheta_{\text{GR}} = \frac{2}{\rho_0} + \mathcal{O}(\rho_0/\rho). \quad (31)$$

Meanwhile, the ε contributions can be evaluated as

$$p = 1, \quad \vartheta_{\varepsilon} = -\varepsilon c_1 \rho_0 + \varepsilon \rho_0 (c_1 - c_2) \ln(2\rho/\rho_0) + \mathcal{O}(\rho_0^{-1}), \quad (32)$$

$$p = 2, \quad \vartheta_{\varepsilon} = (c_1 - c_2) \varepsilon \rho_0 \rho + \mathcal{O}(\rho_0^{-1}), \quad (33)$$

$$p = 3, \quad \vartheta_{\varepsilon} = \frac{1}{2}(c_1 - c_2) \varepsilon \rho_0 \rho^2 + \bar{c} \varepsilon \rho_0^3 [\ln(2\rho/\rho_0) - 1] + \mathcal{O}(\rho_0^{-1}), \quad (34)$$

$$p \geq 4,$$

$$\vartheta_{\varepsilon} = \frac{c_1 - c_2}{p - 1} \varepsilon \rho_0 \rho^{p-1} + \frac{\bar{c}}{p - 3} \varepsilon \rho_0^3 \rho^{p-3} + \mathcal{O}(\rho_0^{p-5}). \quad (35)$$

We have kept not only the dominant correction, but also the next one. Of course the latter becomes important only for $c_1 = c_2$, that is, when the first correction vanishes. Therefore, we have evaluated the second (smaller) contribution only when the first vanishes, that is, when $c_1 = c_2$. In other words, we define $\bar{c} = c_1$ if $c_1 = c_2$. The experimental bound on ϑ_{ε} relative to ϑ_{GR} is given by the very long baseline interferometry, which combines measurements taken by different radio telescopes on Earth [35]. The experimental value ϑ_{exp} relative to the theoretical

prediction has been constrained to be $\vartheta_{\text{exp}}/\vartheta_{\text{theor}} = 1.0001 \pm 0.0001$. Therefore, in order that ϑ_ε does not affect the GR result, we impose

$$\frac{\vartheta_\varepsilon}{\vartheta_{\text{GR}}} < 10^{-4}. \quad (36)$$

When $p \geq 2$, this condition translates into

$$\frac{c_1 - c_2}{2(p-1)} \varepsilon \rho_0^2 \rho^{p-1} < 10^{-4}. \quad (37)$$

Recall that in the domain D we have

$$\varepsilon \rho_0 \rho^{p-1} \ll \frac{1}{\rho_0} \frac{\rho_0^2}{\rho^2} \ll \frac{1}{\rho_0}, \quad (38)$$

in which case the condition (37) is satisfied for c_1, c_2, p of the order of unity. Hence, the deflection of light always remains a small correction in the domain of convergence.

B. Cassini experiment

Iess *et al.* [36] showed that the contribution to the fractional frequency shift of a stable and coherent two-way radio signal (Earth to spacecraft to Earth) y , due to the metric of a gravitational theory (which possesses a weak field limit), is proportional to the deviation angle ϑ of light, and it is given by the formula

$$y = 2 \frac{v_1 l_0 + v_0 l_1}{l_0 + l_1} \vartheta, \quad (39)$$

where v_0 and v_1 are transverse velocities of Earth and a spacecraft, and l_0 and l_1 are their distances from the Sun. Since $l_1 \gg l_0$, the ε contribution can be written as

$$\begin{aligned} y_\varepsilon &\approx 2v_{\text{Earth}}[\varphi(\rho_0, \rho_{\text{Earth}}) + \varphi(\rho_0, \rho_{\text{Cassini}})] \\ &\approx v_{\text{Earth}} \frac{c_1 - c_2}{p-1} \varepsilon \rho_0 (\rho_{\text{Cassini}}^{p-1} + \rho_{\text{Earth}}^{p-1}), \end{aligned} \quad (40)$$

where the approximate equality in the second line is valid for $p \geq 2$. Meanwhile, from Eq. (31), the GR contribution can be written as

$$y_{\text{GR}} \approx 4 \frac{v_{\text{Earth}}}{\rho_0}. \quad (41)$$

The ε contribution needs to be a negligible correction to GR. In order to have $y_\varepsilon \ll y_{\text{GR}}$, we require that

$$\varepsilon \rho_0^2 \rho_{\text{Cassini}}^{p-1} \ll 1, \quad (42)$$

which is satisfied, as we have seen before, in the region D , of course if $\rho_{\text{Cassini}} \in D$.

The signal due to GR detected by Cassini is $y \sim 10^{-10}$, within an experimental error of $\Delta y_{\text{exp}} \sim 10^{-14}$. Therefore, the contribution y_ε from the modifications of gravity needs to satisfy the condition $y_\varepsilon < \Delta y_{\text{exp}} \sim 10^{-14}$, or equivalently, $y_\varepsilon/y_{\text{GR}} < 10^{-4}$. If $p \geq 2$, this condition translates into

$$\frac{c_1 - c_2}{4(p-1)} \varepsilon \rho_0^2 \rho_{\text{Cassini}}^{p-1} < 10^{-4}. \quad (43)$$

As we have shown in Eq. (38), this relation is satisfied in the domain of convergence. The constraint (43) can be used to place experimental bounds on $f(\mathcal{G})$ models later.

C. Perihelion shift

Let us proceed to constraints coming from the perihelion shift of the inner planets, whose GR value is in extremely good agreement with experimental data. We will follow a similar procedure to the one discussed in Sec. IV A. The difference is that we now deal with the motion of a massive particle whose Lagrangian reads

$$L = \frac{1}{2} A \dot{t}^2 - \frac{1}{2} B^{-1} \dot{r}^2 - \frac{1}{2} r^2 \dot{\varphi}^2 = \frac{1}{2}. \quad (44)$$

As in the previous case, the constants of motion are

$$E = A \dot{t} \quad \text{and} \quad L = r^2 \dot{\varphi}. \quad (45)$$

Defining $u = r_s/r$, we can derive the following differential equation:

$$\begin{aligned} \frac{d^2 u}{d\varphi^2} + u - \frac{r_s^2}{2L^2} = \frac{r_s^2 E^2}{2L^2} \frac{B}{A} \left(\frac{1}{B} \frac{dB}{du} - \frac{1}{A} \frac{dA}{du} \right) - \frac{1}{2} \frac{dB}{du} u^2 \\ - (B-1)u - \frac{r_s^2}{2L^2} \left(\frac{dB}{du} + 1 \right). \end{aligned} \quad (46)$$

The flat-space solution can be obtained by setting the right-hand side of Eq. (46) to be zero, as

$$u^b = \frac{r_s^2}{2L^2} [1 + \delta \cos(\phi - \phi_0)], \quad (47)$$

where $0 < \delta < 1$ is the eccentricity of the closed orbit.

From Eq. (44), it follows that

$$\dot{r}^2 = B \left(\frac{E^2}{A} - \frac{L^2}{r^2} - 1 \right). \quad (48)$$

The minimum distance $\rho_0 = r_0/r_s$ satisfies

$$\begin{aligned} \frac{r_s^2 E^2}{2L^2} = A(\rho_0) \left(\frac{1}{2\rho_0^2} + \frac{r_s^2}{2L^2} \right) \\ = \left(1 - \frac{1}{\rho_0} + \varepsilon c_1 \rho_0^p \right) \left(\frac{1}{2\rho_0^2} + \frac{r_s^2}{2L^2} \right) \approx \frac{1}{2\rho_0^2} + \frac{r_s^2}{2L^2}. \end{aligned} \quad (49)$$

Since both E and L are constants, Eq. (49) is an algebraic condition for ρ_0 . Therefore, once we fix the orbit, that is r_s, E , and L , then we also fix r_0 . The same solution will be valid at all times; therefore, the orbit will always have the same condition for r_0 , i.e. the same perihelion and the same aphelion. In other words, the minimum (and the maximum as well) value for r will be unchanged at successive perihelia. Since $\dot{r} = 0$ also implies $dr/d\varphi = 0$, the same initial conditions for the differential equation at the perihelia are

identical at each perihelion and so the orbit repeats exactly, see Rindler's book [37].

Expanding Eq. (46) at linear order in ε , we find

$$\begin{aligned} \frac{d^2 u}{d\varphi^2} + u - \frac{r_s^2}{2L^2} &= \frac{3}{2}u^2 + \frac{\varepsilon u^{-p-1} r_s^2}{2L^2 \rho_0^3} \{c_1 p(\rho_0 - 1) \\ &\quad \times (L^2 r_s^{-2} + \rho_0^2) + c_2 \{L^2 r_s^{-2} [(p-2)u^2 \rho_0^3 \\ &\quad - p(\rho_0 - 1)] + p\rho_0^2\}\}. \end{aligned} \quad (50)$$

We evaluate the right-hand side of Eq. (50) on the Newtonian solution u^b . In this case, using Eq. (47), we have

$$\frac{L^2}{r_s^2} \approx \frac{1}{2} \rho_0 (1 + \delta). \quad (51)$$

By doing so, Eq. (50) reduces to

$$\begin{aligned} \frac{d^2 u}{d\varphi^2} + u - \frac{r_s^2}{2L^2} &= \frac{\varepsilon [\rho_0 (1 + \delta)]^p}{2(\delta + 1) \rho_0^2 (\delta \cos \varphi + 1)^{p+1}} \{c_1 p(\delta + 1) \\ &\quad \times (\rho_0 - 1)(\delta + 2\rho_0 + 1) \\ &\quad + c_2 (p-2) \delta^2 \rho_0 \cos^2 \varphi \\ &\quad + c_2 \{p[\delta^2 (1 - \rho_0) + 2\delta + 2\rho_0 + 1] \\ &\quad - 2\rho_0\} + 2c_2 (p-2) \delta \rho_0 \cos \varphi\}. \end{aligned} \quad (52)$$

In the limit $\rho_0 \gg 1$, the right-hand side can be simplified to

$$\begin{aligned} \frac{d^2 u}{d\varphi^2} + u - \frac{r_s^2}{2L^2} &= \frac{\varepsilon [\rho_0 (1 + \delta)]^p}{2(\delta + 1) \rho_0^2 (\delta \cos \varphi + 1)^{p+1}} \\ &\quad \times \{2c_1 p(1 + \delta) \rho_0^2 + \rho_0 [c_1 p(\delta^2 - 1) \\ &\quad + c_2 (p-2) \delta^2 \cos^2 \varphi - c_2 (p(\delta^2 - 2) + 2) \\ &\quad + 2c_2 (p-2) \delta \cos \varphi]\}. \end{aligned} \quad (53)$$

The second term on the right-hand side of Eq. (53) is subdominant unless $c_1 = 0$. In the following, we will focus on the case $c_1 \neq 0$.

The solutions can be written down and studied for each p . As it happens in GR, the homogeneous solution can be described by the Newtonian solution with some periodic corrections, i.e. only dependent on $\cos \varphi$ and higher harmonics ($\cos 2\varphi$ and so on). However, there will be terms which have a secular impact on the orbit, and here we are looking exactly for such terms. The solution for Eq. (53), valid for any p , is given by

$$\begin{aligned} u &= \frac{r_s^2}{2L^2} [1 + \delta \cos \varphi] + \frac{3\delta \varphi \sin \varphi}{2(1 + \delta)^2 \rho_0^2} - \delta \varepsilon p(p+1) \\ &\quad \times \frac{c_1 \rho_0^p P_1(\delta^2)}{\sqrt{1 - \delta^2} (1 - \delta)^p} \sin \varphi \arctan \left[\sqrt{\frac{1 - \delta}{1 + \delta}} \tan \frac{\varphi}{2} \right] \\ &\quad + P_2(\delta, \varphi), \end{aligned} \quad (54)$$

where $P_1(x)$ is a polynomial such that $P_1(0) = 1$ with

degree in x equal to the integer part of $p/2$, and P_2 is periodic in φ so that it does not have any secular contribution. The first contribution on the right-hand side of Eq. (54) is the Newtonian solution, the second one is the standard GR secular correction, whereas the third one is the secular contribution due to the ε modification of the metric. Let us examine orbits with small eccentricity. Then at lowest order in δ , one finds that Eq. (54) reduces to

$$\begin{aligned} u &\approx \frac{r_s^2}{2L^2} [1 + \delta \cos \varphi] + \frac{3\delta \varphi \sin \varphi}{2(1 + \delta)^2 \rho_0^2} - \frac{1}{2} \delta \varepsilon c_1 p(p+1) \\ &\quad \times \rho_0^p \varphi \sin \varphi, \end{aligned} \quad (55)$$

where we have neglected a possible constant contribution from the arctan which, multiplied by $\sin \varphi$, reduces to a periodic term.

The result (55) coincides with another simpler method at lowest order in δ . One can expand Eq. (53) in series of δ , giving

$$\begin{aligned} \frac{d^2 u}{d\varphi^2} + u - \frac{r_s^2}{2L^2} &= c_1 p \varepsilon \rho_0^p [1 - (p+1) \delta \cos \varphi + p \delta] \\ &\quad + \frac{6\delta (\cos \varphi - 1) + 3}{2\rho_0^2} + \mathcal{O}(\delta^2). \end{aligned} \quad (56)$$

The solution of this equation is

$$\begin{aligned} u &\approx \frac{r_s^2}{2L^2} [1 + \delta \cos \varphi] + \frac{3\delta \varphi \sin \varphi}{2\rho_0^2} - \frac{1}{2} \delta \varepsilon c_1 p(p+1) \\ &\quad \times \rho_0^p \varphi \sin \varphi + d_1 \cos(2\varphi) + d_2 \cos(3\varphi), \end{aligned} \quad (57)$$

where the coefficients $d_{1,2}$ do not need to be known for our purpose. The second term of Eq. (57) coincides with the second term of Eq. (55) at lowest order in δ . The corrections are due to higher orders of δ , which are not included in Eq. (57).

Using the relation (51), the approximate solution (55) can be rewritten as

$$u \approx \frac{r_s^2}{2L^2} [1 + \delta \cos(\varphi - \sigma \varphi)], \quad (58)$$

where $\sigma (\ll 1)$ is defined by

$$\sigma \equiv \frac{3}{2} \frac{1}{(1 + \delta) \rho_0} - \frac{1}{2} \varepsilon c_1 p(p+1) (1 + \delta) \rho_0^{p+1}. \quad (59)$$

From Eq. (58) we find that in one orbit the angle between two perihelia is larger than 2π approximately by $2\pi\sigma$, or

$$\begin{aligned} \Delta \varphi - 2\pi &= \frac{3\pi}{(1 + \delta) \rho_0} - \pi \varepsilon c_1 p(p+1) (1 + \delta) \rho_0^{p+1} \\ &\approx \frac{3\pi}{(1 + \delta) \rho_0} \left[1 - \frac{1}{3} \varepsilon c_1 p(p+1) (1 + 2\delta) \rho_0^{p+2} \right]. \end{aligned} \quad (60)$$

The experimental bound on the shift $\Delta \varphi - 2\pi$ for Earth, based on several thousands of optical observations of

planetary positions, is [38]

$$\Delta\varphi - 2\pi = 5 \pm 1 \text{ arcsec/century.} \quad (61)$$

For Mercury the bound is 43.1 ± 0.1 arcsec/century. Since the GR contribution is given by

$$\Delta\varphi_{\text{GR}} = \frac{3\pi}{(1 + \delta)\rho_0}, \quad (62)$$

one sets the modifications of gravity to contribute less than the experimental relative error, that is

$$\left| \frac{\Delta\varphi_\varepsilon}{\Delta\varphi_{\text{GR}}} \right| = \frac{1}{3} |c_1| \varepsilon p (p+1) (1+2\delta) \rho_0^{p+2} < \frac{1}{5}. \quad (63)$$

The correction term remains as such if

$$\varepsilon \rho_0^{p+2} \ll 1. \quad (64)$$

This condition is not necessarily satisfied in the whole domain of convergence D . Therefore, together with the bound (63), this can be used to constrain modified gravity models.

D. Retardation of light

Let us study the gravitational time-delay effect in light signals. For a light signal propagating from ρ_0 to ρ , the integration of Eq. (24) with respect to ρ gives

$$t = r_s \int_{\rho_0}^{\rho} d\bar{\rho} \left[AB \left(1 - \frac{A}{A_0} \frac{\rho_0^2}{\bar{\rho}^2} \right) \right]^{-1/2}. \quad (65)$$

Expanding the integrand in ε and assuming that both ρ_0 and ρ are much greater than unity, the integral (65) is approximately given by

$$\begin{aligned} t(\rho_0, \rho) \approx & \frac{r_s \sqrt{\rho_0 - 1}}{\sqrt{\rho_0}} \int_{\rho_0}^{\rho} d\bar{\rho} \frac{\bar{\rho}^2}{(\bar{\rho} - 1) \sqrt{\bar{\rho}^2 - \rho_0^2}} - \frac{\varepsilon r_s}{2} \int_{\rho_0}^{\rho} d\bar{\rho} \\ & \times \left\{ \rho \frac{\bar{\rho}^p [\bar{\rho}^2 (c_1 + c_2) - \rho_0^2 (2c_1 + c_2)] + c_1 \rho_0^{p+2}}{(\bar{\rho}^2 - \rho_0^2)^{3/2}} \right. \\ & \left. + \mathcal{O}(\bar{\rho}^{p-1}) \right\}, \quad (66) \end{aligned}$$

where ρ represents the position of the satellite. See Ref. [39] for the similar calculation about the gravitational time-delay effect.

The second term in Eq. (66) corresponds to the ε contribution t_ε . Under the condition $\rho \gg \rho_0$, the ε contribution can be evaluated as

$$\begin{aligned} p = 1, \\ r_s^{-1} t_\varepsilon(\rho_0, \rho) \approx & -\frac{1}{4} (c_1 + c_2) \varepsilon \rho^2 + \frac{1}{8} \varepsilon \rho_0^2 [(c_2 - 3c_1) \\ & + 2(c_1 - c_2) \ln(2\rho/\rho_0)], \quad (67) \end{aligned}$$

$$p = 2,$$

$$\begin{aligned} r_s^{-1} t_\varepsilon(\rho_0, \rho) \approx & -\frac{1}{6} (c_1 + c_2) \varepsilon \rho^3 + \frac{1}{4} (c_1 - c_2) \varepsilon \rho_0^2 \rho \\ & + \frac{1}{4} \bar{c} \varepsilon \rho_0^2, \quad (68) \end{aligned}$$

$$p = 3,$$

$$\begin{aligned} r_s^{-1} t_\varepsilon(\rho_0, \rho) \approx & -\frac{1}{8} (c_1 + c_2) \varepsilon \rho^4 + \frac{1}{8} (c_1 - c_2) \varepsilon \rho_0^2 \rho^2 \\ & - \frac{1}{32} \bar{c} \varepsilon \rho_0^4 [13 - 12 \ln(2\rho/\rho_0)], \quad (69) \end{aligned}$$

$$p \geq 4,$$

$$\begin{aligned} r_s^{-1} t_\varepsilon(\rho_0, \rho) \approx & -\frac{1}{2} \frac{c_1 + c_2}{p+1} \varepsilon \rho^{p+1} + \frac{1}{4} \frac{c_1 - c_2}{p-1} \varepsilon \rho_0^2 \rho^{p-1} \\ & - \frac{1}{4(p-3)} \bar{c} \varepsilon \rho_0^3 \rho^{p-3}, \quad (70) \end{aligned}$$

where we have introduced the constant \bar{c} defined as c_1 when $c_1 = c_2$.

If $p \geq 4$, then the time difference between two points ρ_1 and ρ_2 coming from the ε contribution is

$$\Delta t_\varepsilon \approx -\frac{r_s}{2} \frac{c_1 + c_2}{p+1} \varepsilon (\rho_1^{p+1} + \rho_2^{p+1}). \quad (71)$$

The two contributions add, because in the first integral one has propagation from the satellite to the Sun ($dt/d\rho < 0$), and in the other one from the Sun to Earth ($dt/d\rho > 0$). Since the standard GR contribution to Eq. (66) is $t_{\text{GR}} \approx r_s \ln[2\rho/\rho_0]$, the time difference between two points ρ_1 and ρ_2 can be estimated as

$$\Delta t_{\text{GR}} \approx r_s \ln \left[\frac{4\rho_1 \rho_2}{\rho_0^2} \right]. \quad (72)$$

The ratio among the two contributions is then given by

$$\frac{\Delta t_\varepsilon}{\Delta t_{\text{GR}}} = -\frac{1}{2} \varepsilon \frac{c_1 + c_2}{p+1} \frac{\rho_1^{p+1} + \rho_2^{p+1}}{\ln(4\rho_1 \rho_2 / \rho_0^2)}. \quad (73)$$

The bound regarding the ratio between the measured delay and the one predicted by GR comes from the Viking mission on Mars, which gives the result $\Delta t_{\text{exp}} / \Delta t_{\text{GR}} = 1.000 \pm 0.001$ [40]. Hence, this gives the bound

$$\left| \frac{\Delta t_\varepsilon}{\Delta t_{\text{GR}}} \right| < 10^{-3}. \quad (74)$$

Setting $\rho_1 \approx \rho_2 \approx \rho$, this condition translates into

$$\frac{\varepsilon |c_1 + c_2|}{2(p+1)} \frac{\rho^{p+1}}{\ln(2\rho/\rho_0)} < 10^{-3}. \quad (75)$$

This is generally satisfied in the domain of convergence, as the logarithmic term grows slowly with ρ .

E. Gravitational redshift and equivalence principle

Let us finally consider the gravitational redshift. In this case, for a light signal propagating at different heights r

and r_1 , the ratio of the frequencies ν and ν_1 for corresponding heights is given by

$$\frac{\nu}{\nu_1} = \sqrt{\frac{A(r)}{A(r_1)}} \approx 1 + \frac{1}{2}(\rho_1^{-1} - \rho^{-1}) + \frac{1}{2}c_1\varepsilon(\rho^p - \rho_1^p). \quad (76)$$

The ε -dependent term is much smaller than the standard GR one in the D domain. Defining $\Delta\nu = \nu - \nu_1$, it then follows that

$$\frac{\Delta\nu_\varepsilon/\nu}{\Delta\nu_{\text{GR}}/\nu} = \frac{c_1\varepsilon\rho\rho_1(\rho^p - \rho_1^p)}{\rho - \rho_1}. \quad (77)$$

The bound on $(\Delta\nu_\varepsilon/\nu)/(\Delta\nu_{\text{GR}}/\nu)$ comes from the experiment of a hydrogen-maser clock on a rocket launched to an altitude of about 10^7 m [41], which corresponds to $\Delta\nu_{\text{exp}}/\Delta\nu_{\text{GR}} = 1 \pm 0.0002$. This leads to the following bound:

$$\frac{\Delta\nu_\varepsilon/\nu}{\Delta\nu_{\text{GR}}/\nu} < 2 \times 10^{-4}. \quad (78)$$

In the nonrelativistic limit, the gravitational potential V can be identified as $g_{00} = 1 + 2V$, that is

$$V = -\frac{1}{2\rho} + \frac{1}{2}c_1\varepsilon\rho^p = -\frac{G_N M_\odot}{r} + \frac{1}{2}c_1\varepsilon\left(\frac{r}{2G_N M_\odot}\right)^p. \quad (79)$$

This only depends on the mass M_\odot of the Sun, not on the mass/properties of the test particle. Hence, all test particles with the same distance from the center will feel the same acceleration and the equivalence principle will not be violated. This appears evident as the theory of gravitation discussed here is still a metric theory.

V. POWER-LAW $f(\mathcal{G})$ MODEL

The approach we have used so far works only if the iterative parameter ε is much smaller than 1, and the method works better for smaller ε . As a result, this method cannot be evidently applied to all forms of $R + f(\mathcal{G})$. For example, let us consider the simple power-law case [42]

$$f(\mathcal{G}) = \lambda\sqrt{\mathcal{G}_*}\left(\frac{\mathcal{G}^2}{\mathcal{G}_*^2}\right)^k, \quad (80)$$

such that $f(\mathcal{G})$ is defined for all real values of k and \mathcal{G} . In fact, for the spherically symmetric spacetime, this Lagrangian will give rise to terms typically of order

$$\mathcal{G}f_{,\mathcal{G}} - f = \lambda(2k-1)\bar{\varepsilon}\sqrt{\mathcal{G}_s}\left[\left(\frac{\mathcal{G}}{\mathcal{G}_s}\right)^2\right]^k, \quad (81)$$

$$\mathcal{G}_s^2 f_{,\mathcal{G}\mathcal{G}} = 2\lambda k(2k-1)\bar{\varepsilon}\sqrt{\mathcal{G}_s}\left[\left(\frac{\mathcal{G}}{\mathcal{G}_s}\right)^2\right]^{k-1}, \quad (82)$$

where

$$\bar{\varepsilon} \equiv \varepsilon^{1-4k} = [\sqrt{\mathcal{G}_*/\mathcal{G}_s}]^{1-4k}. \quad (83)$$

The fact that $\bar{\varepsilon}$ can be larger than 1, implies that the corrections to the Schwarzschild metric may become large unless $1 - 4k > 0$, that is $k < 1/4$. The GR case corresponds to $k = 1/2$, in which the contribution of the GB term vanishes. If $k \neq 1/2$ but close to it, then we still require that $\lambda|2k - 1|/\varepsilon \ll 1$ in order to regard these terms as corrections to the equations of motion. In general, if $k \neq 1/2$ and $k > 1/4$, the iterative method cannot be used to find approximate solutions to the equations of motion. In such cases the solutions need to be obtained by numerical integrations.

If $k < 1/4$, then $f_{,\mathcal{G}\mathcal{G}}$ is proportional to $[(\mathcal{G}_s/\mathcal{G})^2]^{1-k}$, so that this term blows up as $\mathcal{G} \rightarrow 0$. In cosmological backgrounds, the GB term $\mathcal{G} = 24H^2(H^2 + \dot{H})$ changes sign from negative to positive during the transition from the matter era to the accelerated epoch [27]. This leads to the divergence of $f_{,\mathcal{G}\mathcal{G}}$ at $\mathcal{G} = 0$, which means that the pure power-law $f(\mathcal{G})$ model is not cosmologically viable. We also note that the GB term inside and outside of a spherically symmetric body (mass M_\odot and radius r_\odot) with homogeneous density is given by $\mathcal{G} = -48(G_N M_\odot)^2/r_\odot^6$ and $\mathcal{G} = 48(G_N M_\odot)^2/r^6$, respectively. As we move from the interior to the exterior of the star, the GB term also crosses 0 from negative to positive. Although it is possible to derive iterative spherically symmetric solutions for $k < 1/4$ by using the expansion in terms of $\bar{\varepsilon}$, the power-law $f(\mathcal{G})$ model is out of our interest because of the problems mentioned above.

VI. MODEL A

In this section we study the model given in Eq. (2), i.e.

$$f(\mathcal{G}) = \varepsilon[-\alpha\lambda\sqrt{\mathcal{G}_s} + \lambda\sqrt{\mathcal{G}_s}g(x)], \quad (84)$$

where

$$g(x) = x \arctan x - \frac{1}{2} \ln(1 + x^2) \quad \text{and} \quad x = \frac{\mathcal{G}}{\mathcal{G}_s \varepsilon^2}. \quad (85)$$

The Lagrangian is a function of $\mathcal{G}/\mathcal{G}_*$, but we choose to write it in this form, so the dependence on ε becomes explicit. Since $f_{,\mathcal{G}\mathcal{G}} > 0$ for positive λ and \mathcal{G}_* , there is no singularity of this quantity unlike the power-law $f(\mathcal{G})$ model.

Let us discuss then the different contributions to the equations of motion (11). For the model (84) we have

$$\begin{aligned} \mathcal{G}f_{,\mathcal{G}} - f &= \mathcal{G}\frac{\lambda}{\sqrt{\mathcal{G}_s}\varepsilon} \arctan x + \alpha\lambda\varepsilon\sqrt{\mathcal{G}_s} - \lambda\varepsilon\sqrt{\mathcal{G}_s}g(x) \\ &= \varepsilon\left[\alpha\lambda\sqrt{\mathcal{G}_s} + \frac{1}{2}\lambda\sqrt{\mathcal{G}_s}\ln\left(1 + \frac{\mathcal{G}^2}{\mathcal{G}_s^2\varepsilon^4}\right)\right], \end{aligned} \quad (86)$$

which is of the order of $\varepsilon\alpha\lambda\sqrt{\mathcal{G}_s}$ plus a logarithmic correction. The other terms that appear in the equations

of motion, i.e. $f_{,GG}$ and $f_{,GGG}$, can be written as follows:

$$f_{,GG} = \frac{\lambda\sqrt{G_s}}{G_s^2\varepsilon^3} \frac{d^2g}{dx^2} = \frac{\lambda\sqrt{G_s}\varepsilon}{G_s^2\varepsilon^4 + G^2}, \quad (87)$$

$$f_{,GGG} = \frac{\lambda\sqrt{G_s}}{G_s^3\varepsilon^5} \frac{d^3g}{dx^3} = -\frac{2\lambda\sqrt{G_s}G\varepsilon}{(G_s^2\varepsilon^4 + G^2)^2}, \quad (88)$$

which are both of the order of $\mathcal{O}(\varepsilon)$ (as typically $|G| \gg G_s$ in the solar system).

We write the metric in terms of the expansion parameter ε :

$$ds^2 = -\left[1 - \frac{r_s}{r} + \varepsilon\phi_1(r) + \varepsilon^2\phi_2(r)\right] dt^2 + \left[1 - \frac{r_s}{r} + \varepsilon\psi_1(r) + \varepsilon^2\psi_2(r)\right]^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2). \quad (89)$$

Although the second-order contribution is not used in order to obtain experimental bounds on the model, we will evaluate it to check whether the terms in the series become smaller for higher orders of ε and to verify that $D_2 \subseteq D$. Because of the Bianchi identities, we can use only two equations, i.e. the 0-0 and the 1-1 equations, as the others are automatically satisfied. The second-order equations follow after solving the first-order equations.

A. Spherically symmetric solutions and the domain of convergence

Linearizing the 0-0 component of the modified Einstein equations at first order in ε , we obtain the differential equation for ψ_1 in terms of $\rho = r/r_s$:

$$\rho \frac{d\psi_1}{d\rho} + \psi_1 = 32\sqrt{3}\lambda\rho^3 + 12\sqrt{3}\lambda\rho^2 \ln(\rho) + (4\ln\varepsilon - 2\alpha - 28)\sqrt{3}\lambda\rho^2, \quad (90)$$

whose particular solution is

$$\psi_1 = 8\sqrt{3}\lambda\rho^3 + 4\sqrt{3}\lambda\rho^2 \ln\rho + \frac{2}{3}\sqrt{3}(2\ln\varepsilon - \alpha - 16)\lambda\rho^2. \quad (91)$$

Here we have neglected the contribution coming from the homogeneous solution, as this would correspond to an order ε renormalization contribution to the mass of the system. Although $\varepsilon \ll 1$, the term in $\ln\varepsilon$ only contributes by a factor of order 10^2 . Hence, the first term on the right-hand side of Eq. (91) dominates over the last term.

The 1-1 component of the Einstein equations gives the equation used to determine ϕ as follows:

$$(\rho - \rho^2) \frac{d\phi_1}{d\rho} + \phi_1 = 8\sqrt{3}\lambda\rho^4 - 2\sqrt{3}(10 + 6\ln\rho + 2\ln\varepsilon - \alpha)\lambda\rho^3 - 2\sqrt{3}(\alpha - 6\ln\rho - 2\ln\varepsilon - 6)\lambda\rho^2 + \rho\psi_1. \quad (92)$$

Substituting the solution (91) into Eq. (92), we get the following solution for ϕ_1 :

$$\phi_1 = -\frac{16}{3}\sqrt{3}\lambda\rho^3 + \frac{2}{3}\sqrt{3}(4 - \alpha + 6\ln\rho + 2\ln\varepsilon)\lambda\rho^2. \quad (93)$$

Since $\rho \gg 1$, the largest contributions to ψ_1 and ϕ_1 correspond to the ones proportional to ρ^3 , which are different from the Schwarzschild-de Sitter contribution (which grows as ρ^2). Hence, the model (2) gives rise to the corrections larger than that in the cosmological constant case by a factor ρ .

In the Appendix, we present the equations for the second-order quantities ψ_2 and ϕ_2 . Discarding the homogeneous part of the solutions of Eqs. (A1) and (A2), we obtain

$$\psi_2 = -512\lambda^2\rho^7 - \frac{4}{21}\lambda^2(-3972 + 168\ln\varepsilon + 504\ln\rho - 84\alpha)\rho^6 - \frac{4}{21}\lambda^2(-322\ln\varepsilon + 161\alpha - 966\ln\rho + 1547)\rho^5, \quad (94)$$

$$\phi_2 = \frac{1216}{7}\lambda^2\rho^7 - \frac{4}{21}\lambda^2(-14\alpha + 28\ln\varepsilon + 1014 + 84\ln\rho)\rho^6 - \frac{4}{21}\lambda^2(294\ln\rho - 399 - 49\alpha + 98\ln\varepsilon)\rho^5, \quad (95)$$

where the dominant terms are the first terms in Eqs. (94) and (95).

So far we have found the solutions up to the order of ε^2 . This expansion is meaningful if each term of the expansion is smaller than the previous one. For the solar system experiments, we will consider the case $\rho \gg 1$. This automatically imposes that the Schwarzschild contribution r_s/r is smaller than the Minkowski value 1, which corresponds to the weak field approximation. The first-order correction in ε is smaller than the Schwarzschild contribution if

$$\varepsilon|\psi_1| \ll 1/\rho, \quad \text{and} \quad \varepsilon|\phi_1| \ll 1/\rho. \quad (96)$$

Since the dominant terms in ψ_1 and ϕ_1 have the dependence $\lambda\rho^3$, the conditions (96) translate into

$$\rho \ll |\lambda\varepsilon|^{-1/4}. \quad (97)$$

Therefore, at first order, the domain of convergence is

$$1 \ll \rho \ll 10^{11}|\lambda|^{-1/4}. \quad (98)$$

The consistency of this inequality requires that $|\lambda| \ll 10^{44}$. The second-order terms in ε can be neglected if the following conditions are satisfied in the same region:

$$\varepsilon^2\psi_2 \ll \varepsilon\psi_1 \quad \text{and} \quad \varepsilon^2\phi_2 \ll \varepsilon\phi_1. \quad (99)$$

Since the dominant contributions in ψ_2 and ϕ_2 have the dependence $\lambda^2 \rho^7$, the conditions (99) are equivalent to the requirement (97). Hence, our solutions using the expansion in ε can be justified in the domain of convergence D .

B. Solar system constraints

We have seen that in the domain D of convergence, the second-order term proportional to ε^2 can be neglected. Moreover, in this domain, we can approximate the solutions further, keeping only the highest power in $\rho = r/r_s$ at the order ε . Hence we have, for local gravity considerations, the metric (12) with

$$A = 1 - \frac{1}{\rho} - \frac{16}{3} \sqrt{3} \lambda \varepsilon \rho^3 [1 + \mathcal{O}(\varepsilon, \rho^{-1})], \quad (100)$$

$$B = 1 - \frac{1}{\rho} + 8\sqrt{3} \lambda \varepsilon \rho^3 [1 + \mathcal{O}(\varepsilon, \rho^{-1})]. \quad (101)$$

Therefore, in the domain of convergence, we will always regard the order ε quantity as a correction to the Schwarzschild contribution, and for the order ε we will always keep only the highest power in ρ . Since the solution given above is an approximate one, valid in the domain of convergence, it does not make sense to see whether or not this metric is asymptotically flat; this would correspond to the solution in a region outside of the domain of convergence. However, even having the real solution at hand (which is not the case), one should not trust it far away from the Sun, as other forces would provide large contributions.

In the following, let us place constraints on the model parameter λ by using a number of experimental bounds discussed in Sec. IV.

(i) Deflection of light

The constraint (37) coming from the deflection of light gives

$$\frac{|c_1 - c_2|}{2(p-1)} \varepsilon \rho_0^2 \rho_{\text{Earth}}^{p-1} < 10^{-4}. \quad (102)$$

For this model we have that

$$c_1 = -\frac{16}{3} \sqrt{3} \lambda, \quad c_2 = 8\sqrt{3} \lambda, \quad p = 3. \quad (103)$$

These numbers define the model and are the same for all remaining constraints. The radius of the Sun in units of the Schwarzschild radius is $\rho_0 = 2.35 \times 10^5$. The distance of Earth from the Sun in units of the Schwarzschild radius is $\rho_{\text{Earth}} = 5.08 \times 10^7$. This translates into the following bound:

$$\lambda < 1 \times 10^{15}, \quad (104)$$

where we have used the value (9) for ε .

(ii) Cassini experiment

As we have already seen, the Cassini experiment places the bound $y_\varepsilon < 10^{-14}$, i.e.

$$v_{\text{Earth}} \frac{|c_1 - c_2|}{p-1} \varepsilon \rho_0 (\rho_{\text{Cass}}^{p-1} + \rho_{\text{Earth}}^{p-1}) < 10^{-14}. \quad (105)$$

The speed of Earth in units of the speed of light is $v_{\text{Earth}} = 9.93 \times 10^{-5}$, whereas the distance of Saturn in units of the Schwarzschild radius is $\rho_{\text{Cass}} = 4.85 \times 10^8$. This gives

$$\lambda < 2 \times 10^{12}. \quad (106)$$

(iii) Perihelion shift

The bound (63) coming from the shift of the perihelion of Earth leads to

$$\frac{1}{3} |c_1| \varepsilon p (p+1) (1 + 2\delta_{\text{Earth}}) \rho_{\text{Earth}}^{p+2} < \frac{1}{5}. \quad (107)$$

For the eccentricity $\delta_{\text{Earth}} = 0.02$ and the perihelion of Earth in units of the Schwarzschild radius $\rho_{\text{Earth}} = 4.98 \times 10^7$, we obtain the constraint

$$\lambda < 2 \times 10^5. \quad (108)$$

For Mercury, $\delta_{\text{Hg}} = 0.2$ and $\rho_{\text{Hg}} = 1.56 \times 10^7$, the bound is slightly weaker, $\lambda < 5 \times 10^5$.

(iv) Retardation of light

The bound (74) coming from the retardation of light, together with Eq. (72), gives the following constraint:

$$\frac{1}{2} \varepsilon \frac{|c_1 + c_2|}{p+1} \frac{\rho_{\text{Mars}}^{p+1} + \rho_{\text{Earth}}^{p+1}}{\ln(4\rho_{\text{Mars}}\rho_{\text{Earth}}/\rho_0^2)} < 10^{-3}, \quad (109)$$

which gives, with $\rho_{\text{Mars}} = 7.71 \times 10^7$ in units of the Schwarzschild radius,

$$\lambda < 5 \times 10^{12}. \quad (110)$$

(v) Gravitational redshift

From the constraint (78) coming from the gravitational redshift, together with Eq. (77), it follows that

$$\frac{|c_1| \varepsilon_{\text{Earth}} \rho_2 \rho_1 (\rho_2^p - \rho_1^p)}{\rho_2 - \rho_1} < 2 \times 10^{-4}. \quad (111)$$

Here $\varepsilon_{\text{Earth}} \approx 10^{-57}$, ρ_1 is the radius of Earth in units of its Schwarzschild radius, i.e. $\rho_1 = 7.18 \times 10^8$, and $\rho_2 = 1.84 \times 10^9$ is the distance of the experimental apparatus (for a height of 10^4 km). We then obtain the following bound:

$$\lambda < 3 \times 10^{15}. \quad (112)$$

The tightest constraint on λ comes from the perihelion shift experiment. This bound is weak so that the $f(\mathcal{G})$ model (2) can be consistent with solar system constraints for a wide range of the model parameter.

VII. MODEL B

Let us next proceed to the constraints on the model (3). This model can be written as

$$f(\mathcal{G}) = \varepsilon[-\alpha\lambda\sqrt{\mathcal{G}_s} + \lambda\sqrt{\mathcal{G}_s}g(x)], \quad (113)$$

where

$$g(x) = x \arctan x, \quad \text{and} \quad x = \frac{\mathcal{G}}{\mathcal{G}_s \varepsilon^2}. \quad (114)$$

We shall derive vacuum solutions for the spherically symmetric metric (12) by using the same expansion parameter ε defined in (8). The term $\mathcal{G}f_{,\mathcal{G}} - f$ in the equations of motion can be estimated as

$$\mathcal{G}f_{,\mathcal{G}} - f = \varepsilon\lambda\sqrt{\mathcal{G}_s} \left[\alpha + \frac{\mathcal{G}^2}{\mathcal{G}_s^2 \varepsilon^4 + \mathcal{G}^2} \right]. \quad (115)$$

Since $\mathcal{G}^2 \gg \mathcal{G}_s^2 \varepsilon^4$, it follows that

$$\mathcal{G}f_{,\mathcal{G}} - f \approx \varepsilon\lambda\sqrt{\mathcal{G}_s}(\alpha + 1) + \mathcal{O}(\varepsilon^5), \quad (116)$$

which works as a cosmological constant at lowest order. Note that we have $f_{,\mathcal{G}\mathcal{G}} \sim \varepsilon^5 + \mathcal{O}(\varepsilon^9)$ and $f_{,\mathcal{G}\mathcal{G}\mathcal{G}} \sim \varepsilon^5 + \mathcal{O}(\varepsilon^9)$, so these terms are higher than the linear order in ε . These properties are different from those in the model (2).

For the model (3), the dominant contribution to the Schwarzschild metric comes from the linear term in ε and the next-order corrections correspond to terms in ε^5 . Hence, we look for a metric of the form

$$\begin{aligned} ds^2 = & - \left[1 - \frac{r_s}{r} + \varepsilon\phi_1(r) + \varepsilon^5\phi_2(r) \right] dt^2 \\ & + \left[1 - \frac{r_s}{r} + \varepsilon\psi_1(r) + \varepsilon^5\psi_2(r) \right]^{-1} dr^2 \\ & + r^2(d\theta^2 + \sin^2\theta d\varphi^2). \end{aligned} \quad (117)$$

Linearizing the 00 component of the Einstein Eq. (10) at first order in ε , we obtain the following differential equation:

$$\rho \frac{d\psi_1}{d\rho} - \psi_1 + 2\sqrt{3}\lambda(1 + \alpha)\rho^2 = 0, \quad (118)$$

which has the particular solution

$$\psi_1 = -\frac{2\sqrt{3}}{3}\lambda(1 + \alpha)\rho^2. \quad (119)$$

The 11 component of the linearized Einstein equation gives

$$\begin{aligned} (\rho^2 - \rho) \frac{d\phi_1}{d\rho} - \phi_1 + \frac{4\sqrt{3}}{3}\lambda(\alpha + 1)\rho^3 \\ - 2\sqrt{3}\lambda(\alpha + 1)\rho^2 = 0, \end{aligned} \quad (120)$$

whose particular solution is

$$\phi_1 = -\frac{2\sqrt{3}}{3}\lambda(1 + \alpha)\rho^2. \quad (121)$$

These expressions for the metric corrections represent the

contribution of an effective cosmological constant (as in the case of the Schwarzschild-de Sitter metric). This solves the equations of motion up to the order ε .

Let us derive next-order solutions $\psi_2(r)$ and $\phi_2(r)$. The differential equations for ψ_2 and ϕ_2 are

$$\rho \frac{d\psi_2}{d\rho} + \psi_2 - 2\sqrt{3}\lambda\rho^{14}(128\rho - 123) = 0, \quad (122)$$

$$(\rho^2 - \rho) \frac{d\phi_2}{d\rho} - \phi_2 + \frac{2\sqrt{3}}{5}\lambda\rho^{14}(80\rho^2 - 146\rho + 65) = 0, \quad (123)$$

which have the following particular solutions:

$$\psi_2 = \frac{2\sqrt{3}}{5}\lambda\rho^{14}(40\rho - 41), \quad (124)$$

$$\phi_2 = -\frac{2\sqrt{3}}{15}\lambda\rho^{14}(16\rho - 13). \quad (125)$$

In principle, one can repeat this method order by order in ε . It is then clear that the vacuum solution will not be strongly constrained by solar system experiments, as the first non-zero contribution will be that of a cosmological constant. To be more precise, the domain of convergence D is defined as

$$\varepsilon\lambda(1 + \alpha)\rho^2 \ll 1/\rho \ll 1, \quad (126)$$

which implies that

$$1 \ll \rho \ll \frac{\varepsilon^{-1/3}}{[\lambda(1 + \alpha)]^{1/3}} \approx \frac{10^{15}}{[\lambda(1 + \alpha)]^{1/3}}. \quad (127)$$

For the consistency of this inequality, we require that $\lambda(1 + \alpha) \ll 10^{45}$. If λ and α are of the order of unity, then the domain D corresponds to the distance 10^5 cm $\lesssim r \lesssim 10^{20}$ cm.

The contributions $\varepsilon^5\psi_2$ and $\varepsilon^5\phi_2$ can be negligible relative to the first-order contributions provided that $\lambda\varepsilon^5\rho^{15} \ll \lambda(1 + \alpha)\varepsilon\rho^2$, i.e.

$$\rho \ll (1 + \alpha)^{1/13}\varepsilon^{-4/13}. \quad (128)$$

If $\alpha = \mathcal{O}(1)$, then the domain D_2 of convergence corresponds to $r \lesssim 10^{19}$ cm. One may wonder if this trend continues at next order, that is, ε^6 . When the previous two contributions are coupled, this gives rise to terms of the order $\lambda^2(1 + \alpha)\varepsilon^6$. Introducing the corrections $\varepsilon^6\psi_3$ and $\varepsilon^6\phi_3$ to the metric and expanding the equations of motion at order ε^6 , we obtain the following differential equations:

$$\rho \frac{d\psi_3}{d\rho} + \psi_3 - \lambda^2(\alpha + 1)\rho^{17}(608\rho - 1136) = 0, \quad (129)$$

$$\begin{aligned} (\rho^2 - \rho) \frac{d\phi_3}{d\rho} - \phi_3 - \lambda^2(\alpha + 1)\rho^{17} \\ \times \left(32\rho^2 + \frac{1064}{45}\rho - \frac{304}{5} \right) = 0, \end{aligned} \quad (130)$$

whose solutions are given by

$$\psi_3 = \lambda^2(1 + \alpha)\rho^{17}(32\rho - \frac{568}{9}), \quad (131)$$

$$\phi_3 = \lambda^2(1 + \alpha)\rho^{17}(\frac{16}{9}\rho + \frac{152}{45}). \quad (132)$$

Therefore, the expansion is meaningful for $\lambda(1 + \alpha)\varepsilon^6\rho^{18} \ll \varepsilon^5\rho^{15}$, that is, $D_3 = D \subseteq D_2$. This shows that the domain of convergence at the order ε^6 coincides with D_2 .

Let us discuss solar system constraints on the model (3) by using the experimental bounds discussed in Sec. IV. We will show that the strongest bound comes from the shift of the perihelion of Earth.

(i) Deflection of light

For the model (3) we have that

$$c_1 = c_2 = -\frac{2\sqrt{3}}{3}\lambda(1 + \alpha), \quad p = 2. \quad (133)$$

Since $c_1 = c_2$, the constraint (37) coming from the deflection of light is trivially satisfied.

(ii) Cassini experiment

Since $c_1 = c_2$ at order ε , the bound (43) of the Cassini experiment is fulfilled.

(iii) Perihelion shift

The bound coming from the shift of the perihelion of Earth corresponds to (107) with $\delta_{\text{Earth}} = 0.02$ and $\rho_{\text{Earth}} = 4.98 \times 10^7$. This leads to

$$\lambda(1 + \alpha) < 1 \times 10^{14}. \quad (134)$$

From the bound of Mercury, we obtain a similar constraint.

(iv) Retardation of light

Using the bound (109) coming from the retardation of light with $\rho_{\text{Mars}} = 7.71 \times 10^7$, we obtain the following constraint:

$$\lambda(1 + \alpha) < 6 \times 10^{20}. \quad (135)$$

(v) Gravitational redshift

Using the constraint (111) of the gravitational redshift with $\varepsilon_{\text{Earth}} = 10^{-57}$, $\rho_1 = 7.18 \times 10^8$, and $\rho_2 = 1.84 \times 10^9$, it follows that

$$\lambda(1 + \alpha) < 3 \times 10^{25}. \quad (136)$$

The bottom line is that the model B, having $p = 2$, is less constrained than the model A.

VIII. CONCLUSIONS

In this paper, we have discussed solar system constraints on $f(\mathcal{G})$ gravity models that are cosmologically viable.

These models give rise to power-law corrections of the form $(r/r_s)^p$ to the Schwarzschild metric, which are characterized by an expansion parameter $\varepsilon = \sqrt{\mathcal{G}_*/\mathcal{G}_s} \approx 10^{-46}$ for the Sun. The smallness of this parameter allows us to find approximate vacuum solutions in a spherically symmetric spacetime.

In order to confront $f(\mathcal{G})$ models with a number of solar system experiments, we have carried out a general analysis for estimating their deviation from general relativity. These include the experiments such as deflection of light, Cassini tracking, perihelion shift of Earth, retardation of light, and gravitational redshift. The results we have derived can be generally applied to any modified gravity models which have power-law corrections to the Schwarzschild metric.

The $f(\mathcal{G})$ models given in Eqs. (2) and (3) are designed to give rise to a late-time cosmic acceleration preceded by a matter-dominated epoch. We find that these models can satisfy all of the solar system constraints discussed in literature for a wide range of model parameters. For the model (2), there exists a logarithmic correction $(\lambda/2) \times \sqrt{\mathcal{G}_*} \ln(\mathcal{G}^2/\mathcal{G}_*^2)$ to the Lagrangian density in the region of high density ($\mathcal{G}^2 \gg \mathcal{G}_*^2$). The tightest bound comes from the shift of perihelion of Earth, but the constraint on the parameter λ is weak: $\lambda < 2 \times 10^5$. In order to set stronger bounds on these theories, it is then necessary to have better measurement of the quadrupole moment of the Sun, as it affects the perihelion shift. For the model (3), the leading correction term to the Lagrangian density corresponds to $(\lambda/3)\sqrt{\mathcal{G}_*}\mathcal{G}_*^2/\mathcal{G}^2$, whose effect is very small even compared to the model (2). Hence, the model parameter for the model (3) is very weakly constrained: $\lambda(1 + \alpha) < 10^{14}$.

The main reason why the $f(\mathcal{G})$ models can satisfy solar system constraints fairly easily is that, even in the vacuum spherically symmetric background, the Gauss-Bonnet scalar takes a nonvanishing value $\mathcal{G} = 12r_s^2/r^6$, where r_s is the Schwarzschild radius. In the solar system, the GB term is much larger than the cosmological value $\mathcal{G}_* \sim H_*^4$. Hence, the inverse power-law terms such as $(\mathcal{G}_*^2/\mathcal{G}^2)^n$ ($n > 0$) are strongly suppressed even for the vacuum solution. This property is different from $f(R)$ gravity in which the Ricci scalar R vanishes in the vacuum spherically symmetric background. In this case, the presence of nonrelativistic matter with density ρ_m leads to a nonvanishing Ricci scalar R approximately proportional to ρ_m . Because of the existence of matter, the local gravity constraints can be satisfied for viable $f(R)$ models having the asymptotic behavior $f(R) = R - \lambda R_*[1 - (R_*/R^2)^n]$ in the region $R^2 \gg R_*^2$. In $f(\mathcal{G})$ gravity, we have shown that the contribution of matter density ρ_m to the GB term ($\sim \rho_m^2$) can be negligible relative to the vacuum contribution ($\sim r_s^2/r^6$) outside of the area of the Sun. Thus, our analysis based on the vacuum spherically symmetric solution is reliable to discuss the compatibility of $f(\mathcal{G})$ models with solar system experiments.

ACKNOWLEDGMENTS

The work of A. D. F. is supported by the Belgian Federal Office for Scientific, Technical, and Cultural Affairs through the Interuniversity Attraction Pole P6/11. S. T. thanks JSPS for financial support (Grant No. 30318802).

APPENDIX: SECOND-ORDER EQUATIONS FOR THE MODEL A

In this Appendix, we present second-order equations in terms of the expansion parameter ε for the model (2). The second-order quantities ψ_2 and ϕ_2 in the metric (89) obey the following equations of motion:

$$\begin{aligned} & (\rho^4 - \rho - 3\rho^3 + 3\rho^2) \frac{d\psi_2}{d\rho} + (3\rho - 3\rho^2 - 1 + \rho^3) \psi_2 + 4096\rho^{10} \lambda^2 + (672 \ln \rho + 224 \ln \varepsilon - 17488 - 112\alpha) \lambda^2 \rho^9 \\ & + (-1040 \ln \varepsilon + 520\alpha - 3120 \ln \rho + 29472) \lambda^2 \rho^8 + (-24448 + 1776 \ln \varepsilon - 888\alpha + 5328 \ln \rho) \lambda^2 \rho^7 \\ & + (664\alpha - 3984 \ln \rho + 9952 - 1328 \ln \varepsilon) \lambda^2 \rho^6 + (-1584 - 184\alpha + 368 \ln \varepsilon + 1104 \ln \rho) \lambda^2 \rho^5 = 0, \end{aligned} \quad (\text{A1})$$

and

$$\begin{aligned} & (-21\rho^2 + 21\rho^3 + 7\rho - 7\rho^4) \frac{d\phi_2}{d\rho} + (7 + 7\rho^2 - 14\rho) \phi_2 + 8512\rho^{10} \lambda^2 + (-34976 + 112\alpha - 224 \ln \varepsilon - 672 \ln \rho) \lambda^2 \rho^9 \\ & + (168 \ln \rho - 28\alpha + 56 \ln \varepsilon + 56260) \lambda^2 \rho^8 + (1344 \ln \varepsilon + 4032 \ln \rho - 44440 - 672\alpha) \lambda^2 \rho^7 \\ & + (-1960 \ln \varepsilon - 5880 \ln \rho + 980\alpha + 17444) \lambda^2 \rho^6 + (2352 \ln \rho - 392\alpha + 784 \ln \varepsilon - 2800) \lambda^2 \rho^5 = 0. \end{aligned} \quad (\text{A2})$$

The solutions to this equation, discarding the homogeneous parts, are given by Eqs. (94) and (95).

-
- [1] E. J. Copeland, M. Sami, and S. Tsujikawa, *Int. J. Mod. Phys. D* **15**, 1753 (2006); S. Nojiri and S. D. Odintsov, *Int. J. Geom. Methods Mod. Phys.* **4**, 115 (2007); T. P. Sotiriou and V. Faraoni, arXiv:0805.1726; R. Durrer and R. Maartens, arXiv:0811.4132.
- [2] S. Capozziello, *Int. J. Mod. Phys. D* **11**, 483 (2002); S. Capozziello, V. F. Cardone, S. Carloni, and A. Troisi, *Int. J. Mod. Phys. D* **12**, 1969 (2003); S. M. Carroll, V. Duvvuri, M. Trodden, and M. S. Turner, *Phys. Rev. D* **70**, 043528 (2004); S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **68**, 123512 (2003); L. Amendola, D. Polarski, and S. Tsujikawa, *Phys. Rev. Lett.* **98**, 131302 (2007); *Int. J. Mod. Phys. D* **16**, 1555 (2007); S. Capozziello and A. De Felice, *J. Cosmol. Astropart. Phys.* **08** (2008) 016.
- [3] T. Chiba, *Phys. Lett. B* **575**, 1 (2003).
- [4] A. A. Starobinsky, *Phys. Lett.* **91B**, 99 (1980).
- [5] G. J. Olmo, *Phys. Rev. D* **72**, 083505 (2005); *Phys. Rev. Lett.* **95**, 261102 (2005); A. L. Erickcek, T. L. Smith, and M. Kamionkowski, *Phys. Rev. D* **74**, 121501 (2006); V. Faraoni, *Phys. Rev. D* **74**, 023529 (2006); A. F. Zakharov *et al.*, *Phys. Rev. D* **74**, 107101 (2006); T. Faulkner *et al.*, *Phys. Rev. D* **76**, 063505 (2007); T. Chiba, T. L. Smith, and A. L. Erickcek, *Phys. Rev. D* **75**, 124014 (2007); I. Navarro and K. Van Acoleyen, *J. Cosmol. Astropart. Phys.* **02** (2007) 022; P. Brax, C. van de Bruck, A. C. Davis, and D. J. Shaw, *Phys. Rev. D* **78**, 104021 (2008).
- [6] L. Amendola, R. Gannouji, D. Polarski, and S. Tsujikawa, *Phys. Rev. D* **75**, 083504 (2007).
- [7] B. Li and J. D. Barrow, *Phys. Rev. D* **75**, 084010 (2007).
- [8] L. Amendola and S. Tsujikawa, *Phys. Lett. B* **660**, 125 (2008).
- [9] W. Hu and I. Sawicki, *Phys. Rev. D* **76**, 064004 (2007).
- [10] A. A. Starobinsky, *JETP Lett.* **86**, 157 (2007).
- [11] S. A. Appleby and R. A. Battye, *Phys. Lett. B* **654**, 7 (2007); *J. Cosmol. Astropart. Phys.* **05** (2008) 019; S. Tsujikawa, *Phys. Rev. D* **77**, 023507 (2008); S. Tsujikawa, K. Uddin, and R. Tavakol, *Phys. Rev. D* **77**, 043007 (2008); S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **657**, 238 (2007); G. Cognola *et al.*, *Phys. Rev. D* **77**, 046009 (2008); N. Deruelle, M. Sasaki, and Y. Sendouda, *Phys. Rev. D* **77**, 124024 (2008); A. Dev, D. Jain, S. Jhingan, S. Nojiri, M. Sami, and I. Thongkool, *Phys. Rev. D* **78**, 083515 (2008); Q. Exirifard, *Classical Quantum Gravity* **26**, 025001 (2009).
- [12] Y. S. Song, W. Hu, and I. Sawicki, *Phys. Rev. D* **75**, 044004 (2007); I. Sawicki and W. Hu, *Phys. Rev. D* **75**, 127502 (2007); S. M. Carroll, I. Sawicki, A. Silvestri, and M. Trodden, *New J. Phys.* **8**, 323 (2006); R. Bean *et al.*, *Phys. Rev. D* **75**, 064020 (2007); Y. S. Song, H. Peiris, and W. Hu, *Phys. Rev. D* **76**, 063517 (2007); L. Pogosian and A. Silvestri, *Phys. Rev. D* **77**, 023503 (2008); I. Laszlo and R. Bean, *Phys. Rev. D* **77**, 024048 (2008); H. Oyaizu, M. Lima, and W. Hu, *Phys. Rev. D* **78**, 123524 (2008); K. Koyama, A. Taruya, and T. Hiramatsu, *Phys. Rev. D* **79**, 123512 (2009).
- [13] S. Tsujikawa and T. Tatekawa, *Phys. Lett. B* **665**, 325 (2008); F. Schmidt, *Phys. Rev. D* **78**, 043002 (2008).
- [14] K. S. Stelle, *Gen. Relativ. Gravit.* **9**, 353 (1978); N. H. Barth and S. M. Christensen, *Phys. Rev. D* **28**, 1876 (1983); A. Hindawi, B. A. Ovrut, and D. Waldram, *Phys. Rev. D* **53**, 5583 (1996); N. Boulanger *et al.*, *Nucl. Phys. B* **597**, 127 (2001); S. M. Carroll, A. De Felice, V. Duvvuri,

- D. A. Easson, M. Trodden, and M. S. Turner, *Phys. Rev. D* **71**, 063513 (2005); M. Sami, A. Toporensky, P. V. Tretyakov, and S. Tsujikawa, *Phys. Lett. B* **619**, 193 (2005); G. Calcagni, S. Tsujikawa, and M. Sami, *Classical Quantum Gravity* **22**, 3977 (2005); A. De Felice, M. Hindmarsh, and M. Trodden, *J. Cosmol. Astropart. Phys.* 08 (2006) 005; G. Calcagni, B. de Carlos, and A. De Felice, *Nucl. Phys.* **B752**, 404 (2006); O. Mena, J. Santiago, and J. Weller, *Phys. Rev. Lett.* **96**, 041103 (2006); A. De Felice and M. Hindmarsh, *J. Cosmol. Astropart. Phys.* 06 (2007) 028; A. De Felice, P. Mukherjee, and Y. Wang, *Phys. Rev. D* **77**, 024017 (2008).
- [15] A. De Felice and T. Suyama, *J. Cosmol. Astropart. Phys.* 06 (2009) 034.
- [16] I. Antoniadis, J. Rizos, and K. Tamvakis, *Nucl. Phys.* **B415**, 497 (1994); M. Gasperini, M. Maggiore, and G. Veneziano, *Nucl. Phys.* **B494**, 315 (1997); R. Brustein and R. Madden, *Phys. Rev. D* **57**, 712 (1998); S. Kawai, M. a. Sakagami, and J. Soda, *Phys. Lett. B* **437**, 284 (1998); S. Kawai and J. Soda, *Phys. Lett. B* **460**, 41 (1999); C. Cartier, E. J. Copeland, and R. Madden, *J. High Energy Phys.* 01 (2000) 035; S. Tsujikawa, *Phys. Lett. B* **526**, 179 (2002); S. Tsujikawa, R. Brandenberger, and F. Finelli, *Phys. Rev. D* **66**, 083513 (2002); K. i. Maeda and N. Ohta, *Phys. Lett. B* **597**, 400 (2004); M. Sami *et al.*, *Phys. Lett. B* **619**, 193 (2005); R. Chingangbam *et al.*, *Phys. Lett. B* **661**, 162 (2008).
- [17] M. Gasperini and G. Veneziano, *Astropart. Phys.* **1**, 317 (1993); *Phys. Rep.* **373**, 1 (2003).
- [18] S. Nojiri, S. D. Odintsov, and M. Sasaki, *Phys. Rev. D* **71**, 123509 (2005).
- [19] T. Koivisto and D. F. Mota, *Phys. Lett. B* **644**, 104 (2007); *Phys. Rev. D* **75**, 023518 (2007).
- [20] S. Tsujikawa and M. Sami, *J. Cosmol. Astropart. Phys.* 01 (2007) 006.
- [21] A. K. Sanyal, *Phys. Lett. B* **645**, 1 (2007).
- [22] I. P. Neupane, *Classical Quantum Gravity* **23**, 7493 (2006); B. M. Leith and I. P. Neupane, *J. Cosmol. Astropart. Phys.* 05 (2007) 019.
- [23] Z. K. Guo, N. Ohta, and S. Tsujikawa, *Phys. Rev. D* **75**, 023520 (2007).
- [24] L. Amendola, C. Charmousis, and S. C. Davis, *J. Cosmol. Astropart. Phys.* 12 (2006) 020; 10 (2007) 004.
- [25] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **631**, 1 (2005).
- [26] B. Li, J. D. Barrow, and D. F. Mota, *Phys. Rev. D* **76**, 044027 (2007).
- [27] A. De Felice and S. Tsujikawa, *Phys. Lett. B* **675**, 1 (2009).
- [28] S. Y. Zhou, E. J. Copeland, and P. M. Saffin, *J. Cosmol. Astropart. Phys.* 07 (2009) 009.
- [29] G. Cognola, E. Elizalde, S. Nojiri, S. Odintsov, and S. Zerbini, *Phys. Rev. D* **75**, 086002 (2007); S. Nojiri, S. D. Odintsov, and P. V. Tretyakov, *Prog. Theor. Phys. Suppl.* **172**, 81 (2008).
- [30] K. Uddin, J. E. Lidsey, and R. Tavakol, arXiv:0903.0270.
- [31] I. Navarro and K. Van Acoleyen, *Phys. Lett. B* **622**, 1 (2005).
- [32] J. Khoury and A. Weltman, *Phys. Rev. Lett.* **93**, 171104 (2004); *Phys. Rev. D* **69**, 044026 (2004).
- [33] S. Capozziello and S. Tsujikawa, *Phys. Rev. D* **77**, 107501 (2008).
- [34] V. Kagramanova, J. Kunz, and C. Lammerzahl, *Phys. Lett. B* **634**, 465 (2006).
- [35] D. S. Robertson *et al.*, *Nature (London)* **349**, 768 (1991).
- [36] B. Bertotti, L. Iess, and P. Tortora, *Nature (London)* **425**, 374 (2003); L. Iess, G. Giampieri, J. D. Anderson, and B. Bertotti, *Classical Quantum Gravity* **16**, 1487 (1999).
- [37] W. Rindler, *Relativity* (Oxford University Press, Oxford, 2001).
- [38] J. D. Anderson, J. K. Campbell, M. A. Slade, R. F. Jurgens, E. L. Lau, X. X. Newhall, and E. M. Standish, in *Sixth Marcel Grossmann Meeting on General Relativity*, edited by H. Sato and T. Nakamura (World Scientific, Singapore, 1992).
- [39] H. Asada, *Phys. Lett. B* **661**, 78 (2008).
- [40] R. D. Reasenberg and I. I. Shapiro *et al.*, *Astrophys. J.* **234**, L219 (1979).
- [41] R. F. C. Vessot and M. W. Levine, *Gen. Relativ. Gravit.* **10**, 181 (1979).
- [42] S. C. Davis, arXiv:0709.4453.