

Signature of primordial non-Gaussianity of ϕ^3 type in the mass function and bias of dark matter haloes

Vincent Desjacques^{1,*} and Uroš Seljak^{1,2,3,†}

¹*Institute for Theoretical Physics, University of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

²*Physics and Astronomy Department, University of California, Berkeley, California 94720, USA
and Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

³*IEU, Ewha University, Seoul, S. Korea*

(Received 15 July 2009; published 15 January 2010)

We explore the effect of a cubic correction $g_{\text{NL}}\phi^3$ on the mass function and bias of dark matter haloes extracted from a series of large N-body simulations and compare it to theoretical predictions. Such cubic terms can be motivated in scenarios like the curvaton model, in which a large cubic correction can be produced while simultaneously keeping the quadratic $f_{\text{NL}}\phi^2$ correction small. The deviation from the Gaussian halo mass function is in reasonable agreement with the theoretical predictions. The scale-dependent bias correction $\Delta b_\kappa(k, g_{\text{NL}})$ measured from the auto- and cross-power spectrum of haloes, is similar to the correction in f_{NL} models, but the amplitude is lower than theoretical expectations. Using the compilation of LSS data in [A. Slosar *et al.*, J. Cosmol. Astropart. Phys. 08 (2008) 031], we obtain for the first time a limit on g_{NL} of $-3.5 \times 10^5 < g_{\text{NL}} < +8.2 \times 10^5$ (at 95% CL). This limit will improve with the future LSS data by 1–2 orders of magnitude, which should test many of the scenarios of this type.

DOI: 10.1103/PhysRevD.81.023006

PACS numbers: 98.65.–r, 95.36.+x, 98.70.Vc, 98.80.Cq

I. INTRODUCTION

In standard single field inflation, primordial curvature perturbations are produced by the inflaton field as it slowly rolls down its potential [1–4]. Most of these models predict a nearly scale-invariant spectrum of adiabatic curvature fluctuations in agreement with cosmological observations. In addition, very small deviations from Gaussianity are expected [5–7]. Therefore, any evidence for or against the detection of primordial non-Gaussianity would strongly constrain inflationary scenarios.

Non-Gaussianity can be produced by nonlinearities in the relation between the primordial curvature perturbation Φ (Here and henceforth, the usual Bardeen potential in matter-dominated era) and the inflaton field, interactions of scalar fields, a modified dispersion relation or a departure from the natural adiabatic vacuum state (see [8] for a review). Any non-Gaussianity that is generated outside the horizon induces a three-point function (or bispectrum) $B_\Phi(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$ that is peaked on squeezed triangles (i.e. $k_1 \ll k_2 \sim k_3$) for realistic values of the scalar spectral index. The resulting non-Gaussianity depends only on the local value $\Phi(\mathbf{x})$ of the Bardeen’s curvature potential and can thus be conveniently parametrized up to third order by

$$\Phi(\mathbf{x}) = \phi(\mathbf{x}) + f_{\text{NL}}[\phi^2(\mathbf{x}) - \langle \phi^2 \rangle] + g_{\text{NL}}\phi^3(\mathbf{x}), \quad (1)$$

where $\phi(\mathbf{x})$ is an isotropic Gaussian random field and f_{NL} , g_{NL} are dimensionless, phenomenological parameters. While the quadratic term generates the irreducible three-point function or bispectrum at leading order, the cubic

term does so for the irreducible four-point function or trispectrum. These statistics can be computed straightforwardly from a perturbative expansion of the homogeneous Robertson-Walker background [9,10]. Convolved with the appropriate transfer function [e.g. the radiation transfer function for the cosmic microwave background (CMB) temperature anisotropy], they can be used to constrain the value of the coupling parameters f_{NL} and g_{NL} . No significant detection of primordial non-Gaussianity has been reported from measurements of the three-point correlation function of the CMB anisotropies [11–15]. The tightest limits are $-4 < f_{\text{NL}} < 80$ at 95% confidence level [14].

If $\mathcal{O}(f_{\text{NL}}) \sim \mathcal{O}(g_{\text{NL}})$ then the cubic correction should always be negligibly small compared to the quadratic one since curvature perturbations are typically $\mathcal{O}(10^{-5})$. However, this condition is not satisfied by some multifield inflationary models such as the curvaton scenario, in which a large g_{NL} and a small f_{NL} can be simultaneously produced. In this model, curvature perturbations are generated by an additional scalar field, the curvaton, whose energy density is negligible during inflation [16–19]. Non-Gaussianity is generated by curvaton self-interactions which effectively contribute a nonquadratic term to the curvaton potential [20–24]. While the value and the sign of g_{NL} depend upon the exact form of the self-interaction term (which can dominate the mass term if the curvaton mass is small enough and the curvaton vacuum expectation value during inflation is large enough [25]), it is generically of magnitude $|g_{\text{NL}}| \sim 10^4$ – 10^5 for realistic models in which the ratio of the energy density of the curvaton to the total energy density at time of decay is small. There are other realizations where one can have large g_{NL} and small

*dvince@physik.uzh.ch

†seljak@physik.uzh.ch

f_{NL} [26,27]. In ekpyrotic and cyclic models, f_{NL} typically is of the order of a few tens while g_{NL} is of the order of a few thousand [28]. If f_{NL} were small, then the imprint of non-Gaussianity would be detected only in four-point statistics such as the CMB trispectrum [29–31]. Thus far, no observational limits have been set on g_{NL} by measuring the CMB trispectrum [32,33]. Nevertheless, since the current bound $|f_{\text{NL}}| \lesssim 100$ implies a relative contribution for the quadratic term of ~ 0.1 per cent, a third order coupling parameter $|g_{\text{NL}}| \sim 10^6$ should also be consistent with the data.

Large-scale structures offer another venue to test for the presence of primordial non-Gaussianity. Deviation from Gaussianity can significantly affect the high mass end of the mass function [34,35], the large-scale two-point correlation [36,37], the bispectrum [38–41] of dark matter haloes hosting the observed galaxies as well as void abundances [42,43] and topological measures of the cosmic web [44,45]. Recently, Refs. [46–48] showed that the local quadratic coupling $f_{\text{NL}}\phi^2$ induces a scale-dependent bias $\Delta b_\kappa(k, f_{\text{NL}})$ in the large-scale power spectrum of biased tracers,

$$\Delta b_\kappa(k, f_{\text{NL}}) = 3f_{\text{NL}}[b(M) - 1]\delta_c \frac{\Omega_m H_0^2}{k^2 T(k) D(z)}, \quad (2)$$

where $b(M)$ is the linear bias parameter, H_0 is the Hubble parameter, $T(k)$ is the matter transfer function normalized to unity as $k \rightarrow 0$, $D(z)$ is the growth factor normalized to $(1+z)^{-1}$ in the matter era and $\delta_c \sim 1.68$ is the present-day (linear) critical density threshold. Reference [48] applied Eq. (2) to constrain the value of f_{NL} using a compilation of large-scale structure data and found $-29 < f_{\text{NL}} < +69$ at 95% confidence. These limits are comparable with those from the CMB, demonstrating the competitiveness of the method. Forthcoming all sky surveys should achieve constraints of the order of $f_{\text{NL}} \sim 1$ [46,49–51] and should be sensitive to a possible scale-dependence of f_{NL} [52]. On the numerical side, however, while simulations of structure formation have confirmed the scaling $\Delta b_\kappa(k, f_{\text{NL}})$ with k [46,53–55], the exact amplitude of the non-Gaussian bias correction remains somewhat debatable.

All numerical studies to date have only implemented the quadratic term $f_{\text{NL}}\phi^2$. The purpose of this work is to quantify the impact of the cubic term $g_{\text{NL}}\phi^3$ on the mass function and bias of dark matter haloes extracted from cosmological simulations and assess the ability of forthcoming measurements of the large-scale bias of galaxies/quasars to constrain the size of a local cubic correction. This paper is organized as follows. We begin with a brief description of the N-body simulations and illustrate the extent to which the coupling $g_{\text{NL}}\phi^3$ affects the matter power spectrum and the halo mass function (Sec. II). We pursue with the non-Gaussian halo bias (Sec. III), to which we derive analytically the scale-dependent and scale-independent contribution, Δb_κ and Δb_1 , and demonstrates

the large suppression of the simulated Δb_κ relative to theory. We then place limits on the coupling parameter g_{NL} and forecast constraints from future large-scale surveys and CMB experiments (Sec. IV). We also show that our findings consistently apply to more general models with nonzero f_{NL} and g_{NL} (Sec. V). We conclude with a discussion of the results in Sec. VI.

II. THE NON-GAUSSIAN SIMULATIONS

A. Characteristics of the N-body runs

We utilize a series of large N-body simulations of the Λ CDM cosmology seeded with Gaussian and non-Gaussian initial conditions. The (dimensionless) power spectrum of the Gaussian part $\phi(\mathbf{x})$ of the Bardeen potential is the usual power-law $\Delta_\phi^2(k) \equiv k^3 P_\phi(k)/(2\pi^2) = A_\phi(k/k_0)^{n_s-1}$. The non-Gaussianity is of the ‘‘local’’ form $\Phi = \phi + g_{\text{NL}}\phi^3$. We adopt the standard (CMB) convention in which $\Phi(\mathbf{x})$ is primordial, and not extrapolated to present epoch. It is important to note that the local transformation is performed before multiplication by the matter transfer function. $T(k)$ is computed with CMBFAST [56] for the WMAP5 best-fitting parameters [13]: $h = 0.7$, $\Omega_m = 0.279$, $\Omega_b = 0.0462$, $n_s = 0.96$ and a normalization of the Gaussian curvature perturbations $A_\phi = 7.96 \times 10^{-10}$ at the pivot point $k_0 = 0.02 \text{ Mpc}^{-1}$. This yields a density fluctuations amplitude $\sigma_8 \approx 0.81$ when the initial conditions are Gaussian. Five sets of three 1024^3 simulations, each of which has $g_{\text{NL}} = 0, \pm 10^6$, were run with the N-body code GADGET2 [57]. We used the same Gaussian random seed field ϕ in each set of runs so as to minimize the sampling variance. We also explored scenarios with nonzero f_{NL} and g_{NL} and ran 2 realizations for each of the non-Gaussian models characterized by $(f_{\text{NL}}, g_{\text{NL}}) = (\pm 100, -3 \times 10^5)$. In all cases, the box size is $1600h^{-1} \text{ Mpc}$ with a force resolution of 0.04 times the mean interparticle distance. The particle mass of these simulations thus is $3.0 \times 10^{11} M_\odot/h$, enough to resolve haloes down to $10^{13} M_\odot/h$.

In the curvaton scenario, generic polynomial interaction terms of the form $\lambda m_\sigma^4 (\sigma/m_\sigma)^n$ (where λ is some coupling strength and m_σ is the curvaton mass) to the quadratic potential of the curvaton field σ yield $|g_{\text{NL}}| \gg 1$ even when the nonlinearity parameter f_{NL} is very small [21,22]. One typically finds $|g_{\text{NL}}| \sim \mathcal{O}(10^4) - \mathcal{O}(10^5)$ when f_{NL} varies in the range $-100 < f_{\text{NL}} < 100$. For practical reasons however, the values of g_{NL} adopted in our simulations are about an order of magnitude larger so as to produce an effect strong enough to be unambiguously measured despite the small simulated volume. Furthermore, we have also considered positive and negative values of g_{NL} so as to assess the sensitivity of the non-Gaussian bias to the sign of the coupling parameter. The simulations with $(f_{\text{NL}}, g_{\text{NL}}) = (-100, -3 \times 10^5)$ may be seen as a particular realisation of the curvaton model in

which the coupling constant λ is positive, and the non-quadratic term is very steep ($n \sim 5 - 10$) but contributes little to the total curvaton potential.

B. Properties of the initial density field

In order to ensure that the initial conditions are successfully generated, we measure at the redshift of our initial conditions, $z = 99$, the (normalized) skewness $S_3(R, z) = \langle \delta_{R,z}^3 \rangle / \sigma^4$ and kurtosis $S_4(R, z) = (\langle \delta_{R,z}^4 \rangle - 3\sigma^4) / \sigma^6$ of the density field $\delta_{R,z}$ smoothed with a (spherically symmetric) window function of characteristic radius R . We adopt a tophat filter throughout this paper. Note also that $\sigma(R, z)$ is the variance of smoothed density fluctuations at redshift z .

In the weakly nonlinear regime, the skewness and kurtosis of the density field may be written as the sum of a part generated by gravitational clustering and a part induced by primordial non-Gaussianity. For Zel'dovich initial conditions [58] and $\Omega_m(z) \approx 1$, the contribution generated by gravitational instabilities reads as [59–62]

$$\begin{aligned} S_3^{\text{Zel}}(R, z) &= 4 - (n_{\text{eff}} + 3) \\ S_4^{\text{Zel}}(R, z) &= \frac{272}{9} - \frac{50}{3}(n_{\text{eff}} + 3) + \frac{7}{3}(n_{\text{eff}} + 3)^2, \end{aligned} \quad (3)$$

where $n_{\text{eff}}(R)$ is the effective spectral index at the smoothing scale R ,

$$n_{\text{eff}}(R) \equiv -\frac{d \ln \sigma^2(R, z)}{d \ln R} - 3. \quad (4)$$

Note that the initial skewness and kurtosis given by the Zel'dovich approximation differs from the exact values predicted by perturbation theory, to which they asymptote in the limit $D(z) \rightarrow \infty$ [63]. In addition, the cubic coupling $g_{\text{NL}}\phi^3$ induces a nonzero kurtosis $S_4^{\text{Pri}}(R, z) \equiv g_{\text{NL}} S_4^{(1)}(R, z)$ at leading order which can be computed analytically from the relation

$$\begin{aligned} \sigma^6 S_4^{(1)}(R, z) &= 4! \left(\prod_{i=1}^3 \int \frac{d^3 k_i}{(2\pi)^3} \alpha_R(k_i, z) P_\phi(k_i) \right) \\ &\quad \times \alpha_R(|\mathbf{k}_1 + \dots + \mathbf{k}_{n-1}|, z), \end{aligned} \quad (5)$$

where

$$\alpha_R(k, z) = \frac{2}{3\Omega_m H_0^2} D(z) k^2 T(k) W_R(k) \quad (6)$$

is evaluated at redshift z and $W_R(k)$ is the Fourier transform of the tophat function. When $f_{\text{NL}} = 0$, primordial skewness is not generated at the first order and, therefore, may be neglected.

Figure 1 displays the initial skewness (top panel) and kurtosis (bottom panel) obtained upon distributing the dark matter particles onto a regular 512^3 mesh (i.e. of cell size $\approx 3h^{-1}$ Mpc) using the cloud-in-cell (CIC) interpolation scheme. Symbols represent the numerical results averaged over the realizations. Because S_4^{Zel} (which is the sole con-




FIG. 1 (color online). Skewness and kurtosis of the initial ($z = 99$) density field as a function of smoothing radius. While the top panel shows the sum of the contributions arising from the Zel'dovich dynamics and from primordial non-Gaussianity, $S_3 = S_3^{\text{Zel}} + S_3^{\text{Pri}}$, the bottom panel only shows the absolute value of the primordial kurtosis, $|S_4^{\text{Pri}}|$. Symbols represent the numerical results averaged over the realizations. They have been slightly shifted horizontally for clarity. Error bars show the scatter among the realizations for the models with $g_{\text{NL}} = 10^6$. Solid lines indicate the theoretical expectations.

tribution to the kurtosis in the Gaussian case) varies considerably between the realizations, we only show the absolute difference $|S_4^{\text{Pri}}|$ between the kurtosis in the non-Gaussian ($g_{\text{NL}} = \pm 10^6$) and the Gaussian ($g_{\text{NL}} = 0$) runs. As expected, the skewness is very similar among the Gaussian and non-Gaussian simulations. While S_3 in the simulations agrees well with the skewness induced by the Zel'dovich dynamics Eq. (3) (solid curve), it is gradually suppressed as the filtering radius R approaches the cell size, presumably because of the finite resolution which smooths also the fluctuations. Note that, in the simulations with nonzero g_{NL} and f_{NL} , there is a large primordial skewness which is of magnitude $|S_3^{\text{Pri}}| \sim$ a few on scale $R \lesssim 50h^{-1}$ Mpc, consistent with theory. As can also be seen, the absolute value of the primordial kurtosis increases sharply with R , in very good agreement with the theoretical prediction.

C. The matter power spectrum

Non-Gaussian corrections to the primordial curvature perturbation can renormalize the input power spectrum of fluctuations used to seed the simulations. Since our simulations implement its unrenormalized version $\Delta_\phi^2(k) = A_\phi(k/k_0)^{n_s-1}$, it is desirable to ascertain the effect of the local coupling term on the simulated density power spectrum before discussing the halo mass function and bias. For

f_{NL} models with $|f_{\text{NL}}| \lesssim 100$, renormalization effects are unlikely to be noticeable due to the limited dynamical range of current cosmological simulations [64]. As we will see shortly however, they can be significant in simulations of g_{NL} models with similar level of non-Gaussianity.

The cubic order term $g_{\text{NL}}\phi^3$ renormalizes the amplitude A_ϕ of the power spectrum of initial curvature perturbations to $A_\phi \rightarrow A_\phi + 6g_{\text{NL}}\langle\phi^2\rangle$, where

$$\langle\phi^2\rangle = \int \frac{d^3k}{(2\pi)^3} P_\phi(k). \quad (7)$$

For scale-invariant initial conditions, $\langle\phi^2\rangle$ has a logarithmic divergence at large and small scales (see [64] for a more detailed discussion of perturbative corrections in non-Gaussian cosmologies). In practice, a low- and high- k cutoff are naturally provided by the finite box size and the resolution of the simulations. Therefore, the effective amplitude of density fluctuations in non-Gaussian simulations with cubic coupling is $\sigma_8 + \delta\sigma_8$ with

$$\begin{aligned} \delta\sigma_8 &= 3g_{\text{NL}}\langle\phi^2\rangle \\ &= 3g_{\text{NL}}\left(\frac{k_0}{k_{\text{min}}}\right)^{1-n_s} \left[1 - \left(\frac{k_{\text{min}}}{k_{\text{max}}}\right)^{1-n_s}\right] \frac{A_\phi}{1-n_s}. \end{aligned} \quad (8)$$

Recall that $k_0 = 0.02h \text{ Mpc}^{-1}$ is our choice of normalization point, and k_{min} and k_{max} are the integration limits set by the fundamental mode and the Nyquist frequency of the periodic cubical box over which the initial conditions are generated. Equivalently,

$$\delta\sigma_8 = 3g_{\text{NL}}\left(\frac{Lk_0}{2\pi}\right)^{1-n_s} [1 - N^{n_s-1}] \frac{A_\phi}{1-n_s}, \quad (9)$$

where $N = 1024$ is the number of mesh points along one dimension. This result becomes $\delta\sigma_8 = 3g_{\text{NL}} \ln(N)A_\phi$ in the scale-invariant limit $n_s \rightarrow 1$. For the cosmological setup considered here, the absolute deviation is

$$\delta\sigma_8 \approx 0.015 \left(\frac{g_{\text{NL}}}{10^6}\right). \quad (10)$$

This correction is fairly large for the values of g_{NL} adopted here and, therefore, must be taken into account in the comparison between the theory and the simulations. As we will see below, this is especially important when studying the high mass tail of the halo mass function which is exponentially sensitive to the amplitude of density fluctuations.

The cubic coupling term $g_{\text{NL}}\phi^3$ can also induce a scale-dependent correction to the matter power spectrum which can be quantified by the fractional change $\beta_m(k, g_{\text{NL}}) = P_{\text{mm}}(k, g_{\text{NL}})/P_{\text{mm}}(k, g_{\text{NL}}=0) - 1$. In Fig. 2, symbols show the result of measuring $\beta_m(k, g_{\text{NL}})$ from the snapshots at $z = 0$ and 2 after correction of the normalization shift $|2\delta\sigma_8/\sigma_8| = 0.037$. There is some evidence for a scale-dependent correction at wave number $k \gtrsim$




FIG. 2 (color online). Non-Gaussian fractional correction $\beta_m(k, g_{\text{NL}}) = P_{\text{mm}}(k, g_{\text{NL}})/P_{\text{mm}}(k, 0) - 1$ to the matter power spectrum after subtracting a scale-independent normalization shift $6g_{\text{NL}}\langle\phi^2\rangle$ induced by the cubic coupling $g_{\text{NL}}\phi^3$.

$0.1h \text{ Mpc}^{-1}$ but the resulting deviation is broadly consistent with zero. We will thus neglect $\beta_m(k, g_{\text{NL}})$ henceforth.

D. The halo multiplicity function

Haloes were identified using the MPI parallelized version of the AHF halo finder [65] which is based on the spherical overdensity (SO) finder developed by [66]. The main reason for using a SO finder is that it is more closely connected to the predictions of the spherical collapse model, on which most of the analytic formulas presented in this paper are based. Namely, the virial mass M of a halo is defined by the radius at which the inner overdensity exceeds $\Delta_{\text{vir}}(z)$ times the background density $\bar{\rho}(z)$ [67,68]. The value of the overdensity threshold $\Delta_{\text{vir}}(z)$ is obtained from the collapse of a spherical tophat perturbation and has a dependence on redshift through the matter density $\Omega_m(z)$ [69,70]. We discard poorly resolved haloes and only study those containing at least 34 particles or, equivalently, with a mass larger than $M = 10^{13}M_\odot/h$.

Analytic arguments based on the Press-Schechter theory [71,72] predict that the halo mass function $n(M, z)$ is entirely specified by the distribution $\nu f(\nu)$ of first-crossings, or multiplicity function

$$\nu f(\nu) = M^2 \frac{n(M, z)}{\bar{\rho}} \frac{d \ln M}{d \ln \nu}. \quad (11)$$

The peak height $\nu(M, z) = \delta_c(z)/\sigma(M)$ is the typical amplitude of fluctuations that produce haloes of mass M by redshift z . Here and henceforth, $\sigma(M)$ denotes the variance of the density field δ_M smoothed on mass scale $M \propto R^3$ and linearly extrapolated to present epoch, whereas $\delta_c(z) \approx 1.68D(0)/D(z)$ is the critical linear overdensity for (spherical) collapse at redshift z .

Despite the lack of a compelling theoretical description of the multiplicity function for Gaussian initial conditions, the fractional deviation from Gaussianity can be modeled accurately using the Press-Schechter formalism. In this approach, the halo mass function $n(M, z)$ is related to the probability $P(>\delta_c, M)$ that a region of mass M exceeds the

critical density for collapse $\delta_c(z)$ through the relation $n(M, z) = -2(\bar{\rho}/M)dP/dM$. The non-Gaussian fractional correction to the multiplicity function then is $R(\nu, g_{\text{NL}}) \equiv f(\nu, g_{\text{NL}})/f(\nu, 0) = (dP/dM)(>\delta_c, M, g_{\text{NL}})/(dP/dM)(>\delta_c, M, 0)$. The level excursion probability $P(>\delta_c, M, g_{\text{NL}})$ can be computed once the probability distribution function (PDF) of the smoothed density field δ_M , $P(\delta_M)$, is known. Here, we will consider the simple extensions proposed by [73,74], in which $P(\delta_M)$ is generically expressed as the inverse transform of a cumulant generating function. Both extensions have been shown to give reasonable agreement with numerical simulations of non-Gaussian cosmologies [53,55,75].

In [73], the saddle-point technique is applied directly to $P(\delta_M)$. The resulting Edgeworth expansion is then used to obtain $P(>\delta_c, M, g_{\text{NL}})$. For f_{NL} non-Gaussianity, Ref. [53] found that the resulting non-Gaussian mass function agrees well with the simulations. For g_{NL} non-Gaussianity, neglecting cumulants other than the kurtosis $S_4^{\text{Pri}}(M)$ (Hereafter, we drop the superscript for conciseness) and truncating the series expansion at $S_4(M)$, the non-Gaussian fractional correction reads

$$\begin{aligned} R_{\text{LV}}(\nu, g_{\text{NL}}) &\approx \left\{ 1 + \frac{1}{4!} \sigma^2 S_4 (\nu^4 - 4\nu^2 - 3) - \frac{1}{4!} \sigma^2 \frac{dS_4}{d \ln \nu} \right. \\ &\quad \left. \times (\nu^2 - 3) \right\} \exp[\nu^2 \delta \sigma_8] \quad (12) \\ &= \left\{ 1 + \frac{1}{4!} \sigma^2 S_4 (\nu^4 - 6\nu^2 + 3) - \frac{1}{4!} \frac{d(\sigma^2 S_4)}{d \ln \nu} \right. \\ &\quad \left. \times (\nu^2 - 3) \right\} \exp[\nu^2 \delta \sigma_8] \quad (13) \end{aligned}$$

after integration over regions above $\delta_c(z)$. Note that we have omitted writing the redshift dependence explicitly. Strictly speaking however, $R(\nu, g_{\text{NL}})$ depends distinctly upon the variables M (or ν) and z due to the presence of $\sigma^2 S_4(M)$. Our notation is motivated by the fact that the measured non-Gaussian correction, as plotted in Fig. 4, appears to depend mostly on the peak height. The exponential factor in the right-hand side is the correction induced by the renormalization of the amplitude of linear density fluctuations, Eq. (10). For consistency, we have also used the Press-Schechter multiplicity function to derive this last term although a Sheth-Tormen mass function [76] may be more appropriate.

In [74], it is the level excursion probability $P(>\delta_c, M)$ that is calculated within the saddle-point approximation. Including only a cubic coupling $g_{\text{NL}} \phi^3$ and truncating the resulting expression at the kurtosis, we find

$$\begin{aligned} P(>\delta_c, M, g_{\text{NL}}) &\approx \frac{1}{\sqrt{2\pi}} \frac{\sigma}{\delta_c} \left(1 + 3g_{\text{NL}} \sigma^2 \langle \phi^2 \rangle - \frac{S_4}{12} \sigma^2 \delta_c^2 \right) \\ &\quad \times \exp \left\{ -\frac{\delta_c^2}{2\sigma^2} \left[1 - 6g_{\text{NL}} \langle \phi^2 \rangle - \frac{S_4}{12} \delta_c^2 \right] \right\} \quad (14) \end{aligned}$$

at first order in g_{NL} . Note that we have already included the renormalization of the fluctuation amplitude. For rare events, $\sigma \ll 1$ and the first parenthesis in the right-hand side can be neglected. To ensure that the resulting mass function is properly normalized, we follow [77] and use

$$\nu_* f(\nu_*) = M^2 \frac{n(M, z, g_{\text{NL}})}{\bar{\rho}} \frac{d \ln M}{d \ln \nu_*} \quad (15)$$

for the non-Gaussian mass function, where $\nu_* = \delta_*/\sigma$, $\delta_* = \delta_c \sqrt{1 - 2\delta\sigma_8 - S_4 \delta_c^2/12}$ ¹ and f is the same multiplicity function as in the Gaussian case. Taking the derivative of the level excursion probability then gives

$$\begin{aligned} \frac{(dP/dM)(>\delta_c, M, g_{\text{NL}})}{(dP/dM)(>\delta_c, M, 0)} &\approx \exp \left[\frac{S_4 \delta_c^4}{4! \sigma^2} + \nu^2 \delta \sigma_8 \right] \\ &\quad \times \left(\frac{\delta_*}{\delta_c} + \frac{1}{4!} \frac{\delta_c^3}{\delta_*} \frac{dS_4}{d \ln \sigma} \right). \quad (16) \end{aligned}$$

The fractional change in the multiplicity function eventually reads as

$$\begin{aligned} R_{\text{MVJ}}(\nu, g_{\text{NL}}) &\approx \exp \left[\frac{\nu^4}{4!} \sigma^2 S_4 + \nu^2 \delta \sigma_8 \right] \\ &\quad \times \left\{ 1 - \frac{\nu^2}{8} \sigma^2 S_4 - \frac{\nu^2}{4!} \frac{d(\sigma^2 S_4)}{d \ln \nu} \right\} \quad (17) \end{aligned}$$

after expanding δ_* at the first order and ignoring the shift in the normalization amplitude, i.e. $\delta_*^{\pm 1} \approx 1 \mp S_4 \delta_c^2/24$. In the limit $\sigma^2 S_4 \ll 1$ and $\nu \gg 1$, the two theoretical expectations reduce to $1 + \nu^4 \sigma^2 S_4/24$. However, they differ in the coefficient of the $\nu^2 \sigma^2 S_4$ term, which is $-1/4$ and $-1/8$ for the LV and MVJ formula, respectively. Therefore, we shall also consider the approximation

$$\begin{aligned} R(\nu, g_{\text{NL}}) &= \exp \left[\frac{\nu^4}{4!} \sigma^2 S_4 + \nu^2 \delta \sigma_8 \right] \\ &\quad \times \left\{ 1 - \frac{\nu^2}{4} \sigma^2 S_4 - \frac{\nu^2}{4!} \frac{d(\sigma^2 S_4)}{d \ln \nu} \right\}, \quad (18) \end{aligned}$$

which is designed to match better the Edgeworth expansion of [73] when the peak height is $\nu \sim 1$.

Calculating the fractional change in the mass function requires knowledge of the kurtosis $S_4(M) \equiv g_{\text{NL}} S_4^{(1)}(M)$ of

¹In local f_{NL} models, this formula only involves the skewness S_3 . As it is incorrectly quoted in some of the literature on non-Gaussian halo mass functions, let us write down its explicit expression:

$$R_{\text{MVJ}}(\nu, f_{\text{NL}}) = \exp \left[\frac{S_3 \delta_c^3}{6\sigma^2} \right] \left[\frac{\delta_c^2}{6\delta_*} \frac{dS_3}{d \ln \sigma} + \frac{\delta_*}{\delta_c} \right],$$

or, in terms of the peak height $\nu = \delta_c/\sigma$,

$$R_{\text{MVJ}}(\nu, f_{\text{NL}}) \approx \exp \left[\frac{\nu^3}{6} \sigma S_3 \right] \left[1 - \frac{\nu}{3} \sigma S_3 - \frac{\nu}{6} \frac{d(\sigma S_3)}{d \ln \nu} \right],$$

after expanding $\delta_* = \sqrt{1 - S_3 \delta_c/3}$ at the first order.

the smoothed density field δ_M , which we compute analytically using the general formula (valid for $n \geq 3$)

$$\sigma^{2n-2} S_n^{(1)}(M) = n! \left(\prod_{i=1}^{n-1} \int \frac{d^3 k_i}{(2\pi)^3} \alpha_M(k_i) P_\phi(k_i) \right) \times \alpha_M(|\mathbf{k}_1 + \dots + \mathbf{k}_{n-1}|), \quad (19)$$

where $\alpha_M(k) \equiv \alpha_R(k, z=0)$ in what follows. Over the mass range probed by our simulations, $10^{13} \leq M \leq 5 \times 10^{15} M_\odot/h$, the normalized kurtosis $\sigma^2 S_4^{(1)}(M)$ is a monotonic decreasing function of M that varies in the narrow range $\sim 4-6 \times 10^{-7}$ for the top-hat filter assumed here (see Fig. 3). Note also that the $\sigma^2 S_4$ term dominates the total contribution to the non-Gaussian correction Eqs. (13), (17), and (18) when the peak height is $\nu \gtrsim 2$ (One finds $|d(\sigma^2 S_4)/d \ln \nu| \lesssim 0.1 \sigma^2 S_4$).

The fractional correction is plotted in Fig. 4 for the haloes extracted from the simulations at redshift $z = 0.3, 0.5, 1$ and 2 . In the top panel, the data are compared to the theoretical predictions Eqs. (13), (17), and (18) evaluated at $z = 0$. As we can see, the level of non-Gaussianity in the halo multiplicity function is consistent with the theory. Our approximation (18) performs equally well regardless of the sign of g_{NL} . It agrees better with the measurements than the formulas of [74] which significantly overestimates the data for $g_{\text{NL}} = 10^6$, and than that of [73] which is not always positive definite for $g_{\text{NL}} = -10^6$. The bottom panel shows that the discrepancy somewhat worsens at higher redshift, especially in the case $g_{\text{NL}} = 10^6$. However, it is possible the agreement may be improved by adding higher order powers of $\sigma^2 S_4$ and higher order cumulants.

To conclude this section, one should keep in mind that all these extensions are based on Press-Schechter and, therefore, provide a bad fit to the Gaussian mass function of haloes. In this respect, excursion set approaches may be




FIG. 3 (color online). Variance σ (dotted line), skewness $\sigma S_3^{(1)}$ (dashed line) and kurtosis $\sigma^2 S_4^{(1)}$ (solid line) of the smoothed linear density field δ_M as a function of mass scale M .




FIG. 4 (color online). Top panel: Fractional correction to the Gaussian multiplicity function of dark matter haloes as a function of the peak height $\nu(M, z)$ for a coupling parameter $g_{\text{NL}} = \pm 10^6$. The dotted, dashed, and solid curves show the theoretical predictions Eqs. (13), (17), and (18), at $z = 0$, respectively. Error bars denote Poisson errors. For illustration, $M = 10^{15} M_\odot/h$ corresponds to $\nu = 3.2, 5.2, 7.7$ at redshift $z = 0, 1, 2$, respectively. Similarly, $M = 10^{14} M_\odot/h$ and $10^{13} M_\odot/h$ correspond to $\nu = 1.9, 3, 4.5$ and $1.2, 1.9, 2.9$, respectively. Bottom panel: a comparison with Eq. (18) evaluated at $z = 0$ and 2 .

more promising since they seem to reproduce both the Gaussian halo counts and the dependence on f_{NL} [78,79].

III. THE NON-GAUSSIAN BIAS SHIFT

A. Theoretical considerations

In order to calculate the scale-dependent bias correction induced by the g_{NL} coupling term to the correlation of haloes of mass M collapsing at redshift z , we follow [47] and consider the two-point correlation $\xi_{\text{hh}}(\mathbf{r})$ of regions of the smoothed density field δ_M above a threshold $\delta_c(z) = \nu(z)\sigma$. The two-point correlation function of this level excursion set, which was originally derived by [37], can be expressed in the high threshold approximation as

$$\xi_{\text{hh}}(\mathbf{r}) = -1 + \exp \left\{ \sum_{n=2}^{\infty} \sum_{j=1}^{n-1} \frac{\nu^n \sigma^{-n}}{j!(n-j)!} \xi^{(n)} \times \left(\begin{array}{cc} \mathbf{x}_1, \dots, \mathbf{x}_1, & \mathbf{x}_2, \dots, \mathbf{x}_2 \\ j \text{ times} & (n-j) \text{ times} \end{array} \right) \right\}, \quad (20)$$

where $\mathbf{r} = \mathbf{x}_1 - \mathbf{x}_2$. For the non-Gaussian model considered here, the leading-order correction induced by nonzero three-point and four-point correlations of the density field reads

$$\Delta\xi_{\text{hh}} = \frac{\nu^3}{\sigma^3} \xi^{(3)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) + \frac{\nu^4}{\sigma^4} \left[\frac{1}{3} \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) + \frac{1}{4} \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) \right]. \quad (21)$$

The non-Gaussian correction ΔP_{hh} to the power spectrum of biased tracers is obtained by Fourier transforming this expression.

In the case $f_{\text{NL}} = 0$ and $g_{\text{NL}} \neq 0$, only the four-point functions contribute at first order. It should also be noted that, at linear order, $\xi^{(2)}(\mathbf{x}_1, \mathbf{x}_2)$ amounts to a renormalization of the linear bias and, therefore, does not contribute to the scale-dependent correction. Details of the calculation can be found in Appendix A. In short, the non-Gaussian correction ΔP_{hh} in the limit of long wavelength $k \ll 1$ is given by the Fourier transform of $\nu^4/3\sigma^4 \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2)$,

$$\begin{aligned} \Delta P_{\text{hh}}(k) &= g_{\text{NL}} \nu^4 S_3^{(1)}(M) \alpha_M(k) P_\phi(k) \\ &= g_{\text{NL}} b_L^2(z) \delta_c^2(z) S_3^{(1)}(M) \alpha_M(k) P_\phi(k), \end{aligned} \quad (22)$$

where we have used $b_L(z) = \nu^2/\delta_c(z)$ as is appropriate for high density peaks. The smoothing window that appears in $\alpha_M(k)$ effectively makes little difference because we are considering the limit where k^{-1} is much larger than the smoothing radius, so we will omit it in the following. For small non-Gaussianity, we can also write $\Delta P_{\text{hh}} \approx 2b_L \Delta b_\kappa P_\delta(k)$ where $P_\delta(k) = \alpha_M^2 P_\phi(k)$ is the power spectrum of the smoothed density field. The scale-dependent bias correction $\Delta b_\kappa(k, g_{\text{NL}})$ can eventually be recast into the form

$$\begin{aligned} \Delta b_\kappa(k, g_{\text{NL}}) &= \frac{1}{2} g_{\text{NL}} b_L(z) \delta_c^2(z) S_3^{(1)}(M) \alpha_M^{-1}(k) \\ &= \frac{3}{4} g_{\text{NL}} b_L(z) \delta_c^2(0) \frac{D(0)}{D(z)^2} S_3^{(1)}(M) \frac{\Omega_m H_0^2}{k^2 T(k)} \\ &= \frac{1}{4} g_{\text{NL}} \delta_c(z) S_3^{(1)}(M) \Delta b_\kappa(k, f_{\text{NL}} = 1), \end{aligned} \quad (23)$$

where $\Delta b_\kappa(k, f_{\text{NL}})$ is the scale-independent bias induced by the quadratic coupling $f_{\text{NL}} \phi^2$, Eq. (2). We have also assumed the Eulerian bias prescription $b(M) = 1 + b_L(M)$.

The change in the mean number density of haloes also creates a scale-independent shift which we denote by $\Delta b_1(g_{\text{NL}})$. As shown in [53] for f_{NL} models, the inclusion of this correction noticeably improves the agreement with the simulations at wave number $k \lesssim 0.1 h \text{ Mpc}^{-1}$. Using a peak-background split and considering the limit of small non-Gaussianity, this contribution reads

$$\begin{aligned} \Delta b_1(g_{\text{NL}}) &= -\frac{1}{\sigma} \frac{\partial}{\partial \nu} \ln[R(\nu, g_{\text{NL}})] \\ &\approx -\frac{1}{\sigma} \left[\frac{1}{3!} (\nu^3 - 3\nu) \sigma^2 S_4 + 2\nu \delta\sigma_8 \right. \\ &\quad \left. + \frac{1}{4!} (\nu^3 - 8\nu) \frac{d(\sigma^2 S_4)}{d \ln \nu} - \frac{\nu}{4!} \frac{d^2(\sigma^2 S_4)}{d \ln \nu^2} \right] \end{aligned} \quad (24)$$

after truncating Δb_1 at first order in g_{NL} . This approximation should perform reasonably well for moderate values of the peak height, $\nu \lesssim 4$, for which the fractional change in the mass function, Eq. (18), matches well the numerical data. It is worth noticing that $\Delta b_1(g_{\text{NL}})$ has a sign opposite to that of g_{NL} because the bias decreases when the mass function goes up. $\Delta b_1(g_{\text{NL}})$ also includes a correction induced by the renormalization of σ_8 . In practice, we estimate $\Delta b_1(g_{\text{NL}})$ for a given halo sample by evaluating $\sigma^2 S_4$ and ν at the scale corresponding to the average halo mass \bar{M} of the sample, as it is unclear to which extent $R(\nu, g_{\text{NL}})$ agrees with the data in the limit $M \gg M_*$.

B. Comparison with the simulations

To assess the effect of primordial non-Gaussianity on the halo bias, we will consider the ratios

$$\begin{aligned} \frac{P_{\text{mh}}(k, g_{\text{NL}})}{P_{\text{mh}}(k, 0)} - 1 &= \frac{\Delta b(k, g_{\text{NL}})}{b(M)} + 2 \frac{\delta\sigma_8}{\sigma_8} \\ \frac{P_{\text{hh}}(k, g_{\text{NL}})}{P_{\text{hh}}(k, 0)} - 1 &= \left(1 + \frac{\Delta b(k, g_{\text{NL}})}{b(M)} \right)^2 + 2 \frac{\delta\sigma_8}{\sigma_8} - 1, \end{aligned} \quad (25)$$

where $\Delta b(k, g_{\text{NL}})$ is generally the sum of a scale-dependent and a scale-independent term. One should bear in mind that the scale-independent shift $2\delta\sigma_8/\sigma_8$ arises from the matter power spectrum and, therefore, is distinct from the term $-2\nu\delta\sigma_8/\sigma$ appearing in Δb_1 . Following [53], we shall also quantify the departure from the theoretical scaling as a function of wave-mode amplitude with the ratio $\Delta b^s/\Delta b^t$, where Δb^s is the non-Gaussian bias correction measured from the simulation and Δb^t is Eq. (27).

We interpolate the dark matter particles and halo centers onto a regular cubical mesh. The resulting dark matter and halo fluctuation fields are then Fourier transformed to yield the matter-matter, halo-matter, and halo-halo power spectra $P_{\text{mm}}(k)$, $P_{\text{mh}}(k)$, and $P_{\text{hh}}(k)$, respectively. These power spectra are measured for a range of halo masses and redshifts, covering the relevant range of statistical properties corresponding to the available data sets of galaxy or quasar populations with different luminosities and bias. Note that these quantities are computed on a 512^3 grid, whose Nyquist wave number is sufficiently large ($\approx 1 h \text{ Mpc}^{-1}$) to allow for an accurate measurement of the power in wave modes of amplitude $k \lesssim 0.1 h \text{ Mpc}^{-1}$. The halo power spectrum is corrected for the shot-noise due to the discrete nature of dark matter haloes, which we assume to be the

standard Poisson term $1/\bar{n}_h$. This discreteness correction is negligible for $P_{\text{mm}}(k)$ due to the large number of dark matter particles. Yet another important quantity is the linear halo bias $b(M)$, which must be measured accurately from the Gaussian simulations as it controls the magnitude of the scale-dependent shift. Here, we shall use the ratio $P_{\text{mh}}(k)/P_{\text{mm}}(k)$ as a proxy for the halo bias since it is less sensitive to shot-noise.

1. An effective non-Gaussian bias correction

Summarizing the analytical considerations of Sec. III A, non-Gaussianity of the g_{NL} type add a correction $\Delta b(k, g_{\text{NL}})$ to the bias $b(k)$ of dark matter haloes which is at leading order

$$\Delta b(k, g_{\text{NL}}) = \Delta b_{\kappa}(k, g_{\text{NL}}) + \Delta b_1(g_{\text{NL}}), \quad (26)$$

We found, however, that this theoretical expectation significantly overestimates the magnitude of the non-Gaussian bias shift measured from the simulations. This is exemplified in the top panel of Fig. 5, where $P_{\text{mh}}(k, g_{\text{NL}})/P_{\text{mh}}(k, 0) - 1$ is plotted for haloes of mass $M > 2 \times 10^{13} M_{\odot}/h$ identified at $z = 0.5$. Clearly, the predicted scale-dependent correction Δb_{κ} is much steeper than measured from the halo samples. In order to improve the agreement with the numerical data, we modify the




FIG. 5 (color online). Top panel: Non-Gaussian bias correction computed from the halo-matter power spectrum of haloes of mass $M > 2 \times 10^{13} M_{\odot}/h$ extracted from the snapshot at $z = 0.5$ (filled symbols). The solid curve represents $P_{\text{mh}}(k, g_{\text{NL}})/P_{\text{mh}}(k, 0) - 1$ with a non-Gaussian bias shift $\Delta b(k, g_{\text{NL}})$ given by Eq. (26). The dashed, dotted, and dotted-dashed curves show three separate contributions that arise at first order in g_{NL} . Bottom panel: $\Delta b(k, g_{\text{NL}})$ is replaced by the theoretical model Eq. (27). The shaded region indicates the data points used to fit the parameters ϵ_{κ} and ϵ_1 . Error bars indicate the scatter among 5 realizations.

above relation as follows:

$$\Delta b(k, g_{\text{NL}}) = \epsilon_{\kappa} \Delta b_{\kappa}(k, g_{\text{NL}}) + [\Delta b_1(g_{\text{NL}}) + \epsilon_1], \quad (27)$$

and treat ϵ_{κ} and ϵ_1 as free parameters that we fit to our measurements of the cross-power spectrum (weighted by the scatter among 5 realizations) in the range $0.005 \leq k \leq 0.03 h \text{ Mpc}^{-1}$ where the scale-dependent effect is largest. The bottom panel of Fig. 5 shows the resulting best-fit contributions $\epsilon_{\kappa} \Delta b_{\kappa}$ and $\Delta b_1 + \epsilon_1$ for the halo sample mentioned above. As seen in Fig. 6, ϵ_{κ} and ϵ_1 appear to depend mainly upon the linear halo bias $b(M)$ and the coupling parameter g_{NL} , although dependencies on redshift or other halo observables are not excluded (The data is too noisy for a reliable estimate of these). The most striking feature of Fig. 6 is the functional dependence of ϵ_{κ} on $b(M)$ and g_{NL} . First, ϵ_{κ} is a monotonically increasing function of the bias and never reaches unity, even for the most biased samples for which the high peak approximation should be valid. Second, ϵ_{κ} is noticeably larger for $g_{\text{NL}} = -10^6$, suggesting thereby that second (and higher) order contributions to the scale-dependent bias may be important. Furthermore, for $b \lesssim 1.5$ where the high peak approximation breaks down, there is some evidence that the effect reverses sign. The bottom panel of Fig. 6 shows that the scale-independent correction has sign opposite to that of g_{NL} , in agreement with theoretical expectations from the peak-background split (see Sec. III A). However, whereas for $b \lesssim 3$ the magnitude of the correction is comparable to that predicted by Eq. (24), it is




FIG. 6 (color online). Best-fitting ϵ_{κ} and $\Delta b_1 + \epsilon_1$ as a function of halo bias and $g_{\text{NL}} = \pm 10^6$, for two different mass cuts as indicated in the Figure. In the top panel, the dotted curve is our best fit to $\epsilon_{\kappa}(b, g_{\text{NL}})$. In the bottom panel, the solid and dashed lines show the scale-independent shift Δb_1 predicted by a peak-background split, Eq. (24), for the low and high mass samples, respectively.

considerably larger for $b \gtrsim 3$, reaching up to 5–10 per cent of the linear halo bias.

Assuming ϵ_κ is a function of $b(M)$ and g_{NL} only and asymptotes to a constant in the highly biased limit, we find that the following parametrized form

$$\epsilon_\kappa(b, g_{\text{NL}}) = c_1 - c_2 g_{\text{NL}} - \frac{c_3}{1 + \exp(c_4 b)} \quad (28)$$

captures reasonably well the increase of ϵ_κ with halo bias for $1.5 < b(M) < 7$. The best-fit values of the parameters are

$$\begin{aligned} c_1 &= 0.59 \pm 0.03, & c_2 &= (6.0 \pm 0.9) \times 10^{-8} \\ c_3 &= 2.1 \pm 0.5, & c_4 &= 0.88 \pm 0.13. \end{aligned} \quad (29)$$

We do not provide a fitting formula for ϵ_1 (or $\Delta b_1 + \epsilon_1$) since it is not directly measurable in real data.

Since both the kurtosis of the initial density field and the mass function of dark matter haloes are in fairly good agreement with theoretical expectations, the discrepancy in the scale-dependent bias $\Delta b_\kappa(k, g_{\text{NL}})$ indicates that it is the high peak approximation considered here which is inaccurate, even in the limit $b(M) \gg 1$ where it is supposed to work best. Our perturbative approach may be one




FIG. 7 (color online). Non-Gaussian bias correction measured in the simulation outputs at redshift $0 < z < 2$ for haloes of mass $M > 2 \times 10^{13} M_\odot/h$. In each panel, the upper plot shows the ratio $P_{\text{hh}}(k, g_{\text{NL}})/P_{\text{hh}}(k, 0) - 1$ (dashed curves, empty symbols) and $P_{\text{mh}}(k, g_{\text{NL}})/P_{\text{mh}}(k, 0) - 1$ (solid curves, filled symbols). The error bars represent the scatter among 5 realizations. The respective output redshift and linear halo bias are also quoted. The bottom of each panel displays the departure from the theoretical prediction, $\Delta b^s/\Delta b^t$. The shaded area indicates the domain where the deviation is less than 20 per cent. The parameters ϵ_κ and ϵ_1 are fitted individually to each sample. For illustration, ϵ_κ takes the best-fit values 0.06 and 0.60 for the lowest and highest biased samples, respectively.

of the reasons for this mismatch. Namely, we have derived the scale-dependent bias $\Delta b_\kappa(k, g_{\text{NL}})$ under the assumption that the non-Gaussian correction to the halo power spectrum is small whereas, for most of the bias range covered by our halo catalogues, the effect is in fact already very large at $k \lesssim 0.01 h \text{ Mpc}^{-1}$. Note, however, that for the quadratic coupling $f_{\text{NL}}\phi^2$, this perturbative treatment predicts an effect of the right magnitude [53]. Explaining these findings clearly requires a better theoretical understanding, which we leave for a future investigation. Notwithstanding this, the phenomenological prescription (27) with parameters ϵ_κ and ϵ_I fitted to the data provides, as we will see below, a good description of the large-scale halo power spectrum in simulations of g_{NL} models.

2. Non-Gaussian bias from auto- and cross-power spectra

We have measured auto- and cross-power spectra for a range of halo masses and redshifts spanning the range $0 < z < 2$. The ratios defined in Eq. (25) are shown in Figs. 7 and 8 as a function of wave number for the mass threshold $M > 2 \times 10^{13} M_\odot/h$ and the mass bin $10^{13} < M < 2 \times 10^{13} M_\odot/h$, respectively. The fractional deviation $\Delta b^s/\Delta b^t$ is also shown at the bottom of each panel. The shaded region indicates a departure less than 20 per cent. Error bars denote the scatter around the mean and, therefore, may underestimate the true errors as they are computed from a small number of realizations. Note that, in order to reduce the impact of sampling variance, we first

compute the ratios $P_{\text{mh}}(k, g_{\text{NL}})/P_{\text{mh}}(k, 0)$ and $P_{\text{hh}}(k, g_{\text{NL}})/P_{\text{hh}}(k, 0)$ for each realization before calculating the average.

As we can see, once ϵ_κ and ϵ_I are fitted to the ratio of cross-power spectra, the theoretical prediction Eq. (27) provides a reasonable description of the non-Gaussian bias in the halo power spectrum P_{hh} , indicating that non-Gaussianity does not generate much stochasticity and the predicted scaling $\Delta b_\kappa(k, g_{\text{NL}}) \propto k^{-2} T(k)^{-1}$ applies equally well for the auto- and cross-power spectrum. This was also found to be true in f_{NL} models [53]. The inclusion of a scale-independent correction $\Delta b_I + \epsilon_I$ significantly improve the agreement at $k \lesssim 0.03 h \text{ Mpc}^{-1}$. For the highly biased samples $b > 4$ however, this correction is so large that the non-Gaussian bias shift reverses sign at wave number $k \gtrsim 0.05 h \text{ Mpc}^{-1}$. Such an effect is not seen in the simulations, but we expect large deviations from the relation (27) in that range of wave number, where second- and higher-order corrections induced by the cubic coupling $g_{\text{NL}}\phi^3$ together with the nonlinear bias created by the gravitational evolution of matter density fluctuations may become important. Even though the data is noisier due to the low number density of haloes, it is worth noting that, for the highly biased samples at $k \lesssim 0.01 h \text{ Mpc}^{-1}$, the cross-power spectrum $P_{\text{mh}}(k, g_{\text{NL}} = -10^6)$ goes negative while $P_{\text{hh}}(k, g_{\text{NL}} = -10^6)$ remains positive and increases sharply, in agreement with the analytic prediction. Still, there is some evidence that the ratio $P_{\text{hh}}(k, g_{\text{NL}} = -10^6)/P_{\text{hh}}(k, 0) - 1$ saturates at a value noticeably larger




FIG. 8 (color online). Same as Fig. 7 but for haloes in the mass range $10^{13} < M < 2 \times 10^{13} M_\odot/h$.

than -1 before the sharp upturn, whereas our model predicts $2\delta\sigma_8/\sigma_8 - 1 \approx -0.96$ at the minimum.

Figure 8 further explore the effect in the low mass samples, for which the $z = 0$ haloes with $b(M) \approx 1.15$ constitute an almost unbiased sample of the density field. In this case, the sign of the scale-dependent contribution is reversed, namely, the large-scale halo power spectrum in simulations of $g_{\text{NL}} = -10^6$ is enhanced relative to that of the Gaussian ones. This is in rough agreement with the theory, which predicts a similar effect for $b(M) < 1$. Again, haloes with a similar bias also have a comparable scale-dependent and scale-independent bias regardless of mass or redshift. Finally, note that the sample at $z = 1.39$ shown in Fig. 8 corresponds closely to the quasar sample used by [48], for which $z = 1.8$ and $b = 2.7$.

IV. CONSTRAINTS ON THE COUPLING PARAMETER g_{NL}

A. Constraints on g_{NL} from current large-scale structure data

Reference [48] took advantage of the scale-dependence of the bias to constrain f_{NL} from a sample of highly biased luminous red galaxies (LRGs) and quasars (QSOs). It is straightforward to translate their $2\text{-}\sigma$ limit $-29 < f_{\text{NL}} < +69$ into a constraint on g_{NL} since the non-Gaussian scale-dependent bias $\Delta b_\kappa(k, g_{\text{NL}})$ has the same functional form as $\Delta b_\kappa(k, f_{\text{NL}})$.

Constraints will arise mostly from the QSO sample at median redshift $z = 1.8$, which covers a large comoving volume and is highly biased, $b = 2.7$. In light of our results (see Fig. 6), we expect the parameter $\epsilon_\kappa(b, g_{\text{NL}})$ to vary with g_{NL} . However, in order to simplify the analysis, we will assume that, at fixed b , $\epsilon_\kappa(b, g_{\text{NL}})$ is given by the mean of $\epsilon_\kappa(b, g_{\text{NL}} = \pm 10^6)$. For a sample with bias $b \sim 2.7$, this implies $\epsilon_\kappa \approx 0.4$. Furthermore, assuming the typical mass of QSO-hosting haloes is $\sim 10^{13} M_\odot/h$ yields $S_3^{(1)}(M) \approx 2.3 \times 10^{-4}$. Hence, the multiplicative factor $(1/4)\delta_c(z)\epsilon_\kappa S_3^{(1)}(M)$ is approximately $\approx 8.4 \times 10^{-5}$. Our limits on g_{NL} thus are

$$-3.5 \times 10^5 < g_{\text{NL}} < +8.2 \times 10^5 \quad (30)$$

at 95% confidence level. The scale-independent correction $\Delta b_1 + \epsilon_1$ is not directly measured as it adds up to the bias b which is fitted to the data. For the limits obtained here, $|\Delta b_1 + \epsilon_1|$ should be much smaller than b and can thus be ignored. Note also that, whereas the non-Gaussian bias scales as $D(z)^{-1}$ in f_{NL} models, we have $\Delta b(k, g_{\text{NL}}) \propto D(z)^{-2}$ for g_{NL} non-Gaussianity, so one can achieve relatively larger gains from measurements of high redshift tracers. In fact, the extent to which one can improve the observational bounds will strongly depend on our ability to minimize the impact of sampling variance caused by the random nature of the wave modes, and the shot-noise caused by the discrete nature of the tracers. By comparing

differently biased tracers of the same surveyed volume [51] and suitably weighting galaxies (e.g. by the mass of their host halo) [80,81], it should be possible to circumvent these problems and considerably improve the detection level.

B. Predictions for future LSS surveys

References [50,51,64,82] applied the Fisher matrix formalism to forecast constraints on f_{NL} from forthcoming galaxy redshift surveys. Here, we will simply try to estimate the detection limit for g_{NL} . Following [50,64], we consider a (nearly spherical) survey of volume V . Assuming the Fourier modes are still uncorrelated and Gaussian distributed, the total signal-to-noise squared reads

$$\left(\frac{S}{N}\right)^2 \approx \frac{V}{4\pi^2} \int_{k_{\min}}^{k_{\max}} dk k^2 \left[\left(1 + \frac{\Delta b_\kappa}{b}\right)^2 - 1 \right]^2 \quad (31)$$

in the limit where sampling variance dominates the errors. Here, $k_{\min} \sim \pi/V^{1/3}$ is the smallest wave mode accessible and k_{\max} is not necessarily finite since the integral does converge as one takes k_{\max} to infinity. Substituting the expression Eq. (23) for the scale-dependent bias $\Delta b_\kappa(k, g_{\text{NL}})$ and setting $T(k) \equiv 1$ over the wave number range across which the integral is performed, we arrive at

$$\left(\frac{S}{N}\right)^2 \approx \frac{V}{\pi^2} \left(k_\star^2\right)^2 \left(\frac{1}{k_{\min}} - \frac{1}{k_{\max}}\right), \quad (32)$$

where

$$k_\star^2 \approx 5.0 \times 10^{-12} g_{\text{NL}} \epsilon_\kappa \frac{(1 - 1/b)}{D^2(z)} \left(\frac{S_3^{(1)}}{10^{-4}}\right) h^2 \text{ Mpc}^{-2}. \quad (33)$$

We have also assumed $|g_{\text{NL}}| \lesssim 10^5$, such that $|k_\star^2|$ is at most of the order of k_{\min}^2 and the term linear in $\Delta b_\kappa/b$ dominates the signal. When $k_{\min} \ll k_{\max}$, we can further simplify $(S/N)^2$ to

$$\begin{aligned} \left(\frac{S}{N}\right)^2 &\approx 8.1 \times 10^{-13} g_{\text{NL}}^2 \epsilon_\kappa^2 \left(1 - \frac{1}{b}\right)^2 D(z)^{-4} \left(\frac{S_3^{(1)}}{10^{-4}}\right)^2 \\ &\times \left(\frac{V}{h^{-3} \text{ Gpc}^3}\right)^{4/3}. \end{aligned} \quad (34)$$

Note the strong sensitivity of the signal-to-noise squared to the growth factor $D(z)$ (For f_{NL} non-Gaussianity, this dependence is only $D(z)^{-2}$).

To highlight the improvement one could achieve with future galaxy surveys, it is useful to first calculate the detection limit for the SDSS LRG sample centered at $z \sim 0.3$ and covering a volume $V \approx 2h^{-3} \text{ Gpc}^3$. Assuming a linear bias $b = 2$ and a skewness parameter $S_3^{(1)} \sim 2 \times 10^{-4}$ appropriate for haloes of mass $M \sim 10^{12} - 10^{13} M_\odot/h$, the minimum g_{NL} detectable at the $1\text{-}\sigma$ level is $\approx 10^6$ for a correction factor $\epsilon_\kappa = 0.3$ which we

read off from Fig. 6. For a survey configuration analogous to SDSS-III/BOSS,² with central redshift $z = 0.5$ and a comoving volume $V = 6h^{-3} \text{ Gpc}^3$, the minimum g_{NL} would be $\sim 4 \times 10^5$ for galaxies tracing haloes of similar mass and bias. Finally, for a configuration like EUCLID³ with a $V = 100h^{-3} \text{ Gpc}^3$ survey centered at $z = 1.4$, the detection limit would be $\sim 2.1 \times 10^4$. Clearly, these limits are only indicative: they may be significantly improved by selecting highly biased, high redshift (single- or multi-) tracers. Nevertheless, this shows that future galaxy surveys should furnish interesting constraints on the size of the cubic coupling $g_{\text{NL}} \phi^3$.

C. Predictions for CMB temperature anisotropies

The CMB trispectrum provides an alternative probe of local, nonquadratic correction to the Gaussian curvature perturbations, so it is interesting to assess the sensitivity of this statistics to the nonlinear parameter g_{NL} .

The temperature anisotropy field is conveniently decomposed into spherical harmonics, $\Delta T(\hat{\mathbf{n}})/T = \sum_{lm} a_l^m Y_l^m(\hat{\mathbf{n}})$. As shown in [29,30], statistical isotropy and invariance under parity transformation $\hat{\mathbf{n}} \rightarrow -\hat{\mathbf{n}}$ implies that the 4-point correlation of the spherical harmonic coefficients a_l^m takes the form

$$\langle a_{l_1}^{m_1} a_{l_2}^{m_2} a_{l_3}^{m_3} a_{l_4}^{m_4} \rangle = \sum_{LM} (-1)^M \begin{pmatrix} l_1 & l_2 & L \\ m_1 & m_2 & -M \end{pmatrix} \times \begin{pmatrix} l_3 & l_4 & L \\ m_3 & m_4 & M \end{pmatrix} Q_{l_3 l_4}^{l_1 l_2}(L). \quad (35)$$

Here, $Q_{l_3 l_4}^{l_1 l_2}(L)$ is the angular average trispectrum and brackets are Wigner-3j symbols. Statistical homogeneity also implies that $Q_{l_3 l_4}^{l_1 l_2}(L)$ is independent of position. The connected part of the trispectrum, $T_{l_3 l_4}^{l_1 l_2}(L)$, encodes information about non-Gaussianity and is obtained by subtracting a Gaussian piece constructed from the power spectra C_l . Equation (35) can be inverted with the aid of the orthogonality of the Wigner-3j symbols to form an estimator for the CMB trispectrum.

The signal-to-noise for the CMB trispectrum $T_{l_3 l_4}^{l_1 l_2}(L)$ summed up to a certain l_{max} is [30]

$$\left(\frac{S}{N}\right)^2 (<l_{\text{max}}) \approx \sum_{l_1 > l_2 > l_3 > l_4}^{l_{\text{max}}} \sum_L \frac{|T_{l_3 l_4}^{l_1 l_2}(L)|^2}{(2L+1)C_{l_1}C_{l_2}C_{l_3}C_{l_4}} \quad (36)$$

when cosmic variance dominates the errors. Otherwise, one shall include a contribution from the power spectrum of the detector noise to the C_l . Galactic foreground subtraction on a fraction $1 - f_{\text{sky}}$ would further reduce $(S/N)^2$ by a factor of f_{sky} .

Neglecting the ISW effect, the Sachs-Wolfe provides a useful order-of-magnitude estimate of the signal-to-noise as long as l_{max} does not exceed $\lesssim 100$ [5,30,31,83]. The calculation is performed in Appendix B. We find that the signal-to-noise can be recast into the compact form

$$\left(\frac{S}{N}\right)^2 (<l_{\text{max}}) = \frac{9}{2} g_{\text{NL}}^2 A_\phi^2 \left\{ \frac{1}{6} \int_{-1}^{+1} dx s_{l_{\text{max}}}^3(x) t_{l_{\text{max}}}(x) + \frac{1}{2} \int_{-1}^{+1} dx r_{l_{\text{max}}}^2(x) s_{l_{\text{max}}}^2(x) \right\} \quad (37)$$

where the auxiliary functions $r_l(x)$, $s_l(x)$ and $t_l(x)$ are defined as

$$r_l(x) = \sum_{k=2}^l (2k+1) P_k(x) \quad (38)$$

$$s_l(x) = \sum_{k=2}^l \frac{(2k+1)}{k(k+1)} P_k(x) \quad (39)$$

$$t_l(x) = \sum_{k=2}^l (2k+1)k(k+1) P_k(x). \quad (40)$$

Here, $P_l(x)$ are Legendre polynomials. Note that we have excluded the monopole and dipole from the summation since these modes are unobservable. We have also assumed a nearly scale-invariant spectrum $n_s \approx 1$.

Figure 9 shows the signal-to-noise ratio squared in the Sachs-Wolfe approximation for the CMB trispectrum assuming $g_{\text{NL}} = 1$. Although this approximation breaks down for $l_{\text{max}} \gtrsim 100$, we have extended the calculation up to $l_{\text{max}} = 200$ so as to extrapolate more robustly the l_{max} -dependence to small angular resolution. A power-law fit to $(S/N)^2$ in the range $50 \leq l_{\text{max}} \leq 200$ gives

$$\left(\frac{S}{N}\right)^2 (<l_{\text{max}}) \approx 2.43 \times 10^{-17} g_{\text{NL}}^2 \left(\frac{A_\phi}{10^{-9}}\right)^2 l_{\text{max}}^{2.6}. \quad (41)$$

Our results appear consistent with the findings of [30] shown in their Fig. 2. However, our constant of proportionality is about 20–30 times larger, presumably because they adopted a lower fluctuation amplitude (compare also their prediction for the f_{NL} model with that of [31]). Adding the information encoded in temperature-polarization trispectra may enhance $(S/N)^2$ by a factor of a few [29].

Assuming the scaling Eq. (41) persists well beyond the range over which the Sachs-Wolfe effect dominates, the minimum g_{NL} detectable at 1- σ level is $g_{\text{NL}} \approx 20, 7.9, 3.2, 1.9,$ and 1.3×10^4 for $l_{\text{max}} = 250, 500, 1000, 1500,$ and 2000 . A more realistic calculation should include the full radiation transfer function, detector noise etc. In this respect, detailed calculations have shown that, for the quadratic coupling $f_{\text{NL}} \phi^2$, $(S/N)^2$ of the CMB bispectrum and trispectrum closely follows the behavior obtained in the Sachs-Wolfe approximation [30,31]. It seems reasonable, then, to expect that this is also true for $g_{\text{NL}} \phi^3$.

²www.sdss3.org

³http://sci.esa.int/science-e/www/object/index.cfm?fobjectid=42266




FIG. 9 (color online). Signal-to-noise ratio squared for the CMB trispectrum as a function of the maximum multipole l_{\max} . We have assumed $g_{\text{NL}} = 1$ and $f_{\text{sky}} = 1$.

While our predictions are qualitative they show that, for the WMAP CMB temperature measurement⁴ (which we approximate as a noise-free experiment with $l_{\max} \sim 250$), no detection of a significant trispectrum implies $|g_{\text{NL}}| \leq 2 \times 10^5$ at the $1\text{-}\sigma$ level. This is of the same order as the limit we derived from the QSO sample analyzed by [48]. For a PLANCK-like experiment⁵ ($l_{\max} \sim 1500$), no evidence for a trispectrum would imply $|g_{\text{NL}}| \leq 1.3 \times 10^4$ at the $1\text{-}\sigma$ level. This is comparable to the detection limit that could be achieved with an all-sky survey such as EUCLID.

V. EFFECT OF NON-GAUSSIANITY WITH NONZERO f_{NL} AND g_{NL}

In this section, we examine the halo multiplicity function and large-scale bias in numerical simulations of structure formation with nonzero coupling parameters $(f_{\text{NL}}, g_{\text{NL}}) = (\pm 100, -3 \times 10^5)$. We show that the results are consistent with those obtained from the simulations with nonvanishing g_{NL} solely.

A. Mass function

It is straightforward to calculate the fractional deviation from the Gaussian mass function, Eq. (18), to nonzero f_{NL} and g_{NL} . Again, we start with the MVJ formula and neglect second order corrections such as $(\sigma S_3)^2$, etc. Adjusting the coefficient of the terms $\nu \sigma S_3$ and $\nu^2 \sigma^2 S_4$ to that of the small ν expansion obtained by [73], we arrive at

$$R(\nu, \text{NL}) = \exp \left[\frac{\nu^3}{3!} \sigma S_3 + \frac{\nu^4}{4!} \sigma^2 S_4 + \nu^2 \delta \sigma_8 \right] \left\{ 1 - \frac{\nu}{2} \sigma S_3 - \frac{\nu}{6} \frac{d(\sigma S_3)}{d \ln \nu} - \frac{\nu^2}{4} \sigma^2 S_4 - \frac{\nu^2}{4!} \frac{d(\sigma^2 S_4)}{d \ln \nu} \right\}, \quad (42)$$

⁴<http://map.gsfc.nasa.gov/>

⁵<http://sci.esa.int/science-e/www/area/index.cfm?fareaid=17>

where the shorthand notation NL designates the combination $(f_{\text{NL}}, g_{\text{NL}})$. In Fig. 10, this theoretical prediction is compared $R(\nu, \text{NL})$ measured in non-Gaussian simulations of $(f_{\text{NL}}, g_{\text{NL}}) = (\pm 100, -3 \times 10^5)$. We account for the fact that the amplitude of density fluctuations is renormalized by $\delta \sigma_8 \approx 0.0045$. Figure 10 demonstrates that our approximation is in good agreement with the data, although it slightly overestimates the effect at $z = 2$ when $f_{\text{NL}} = -100$. For $f_{\text{NL}} = 100$, the positive and negative contributions from the quadratic and cubic coupling, respectively, almost cancel each other and flatten the fractional deviation over most of the mass range probed by the simulations.

The scale-independent bias shift which arises from the change in the mean number density of haloes can again be estimated using the peak-background split. We find

$$\begin{aligned} \Delta b_1(\text{NL}) \approx & -\frac{1}{\sigma} \left[\frac{1}{3!} (\nu^3 - 3\nu) \sigma^2 S_4 + \frac{1}{2} (\nu^2 - 1) \sigma S_3 \right. \\ & + \frac{1}{4!} (\nu^3 - 8\nu) \frac{d(\sigma^2 S_4)}{d \ln \nu} - \frac{\nu}{4!} \frac{d^2(\sigma^2 S_4)}{d \ln \nu^2} + \frac{1}{6} \\ & \left. \times (\nu^2 - 4) \frac{d(\sigma S_3)}{d \ln \nu} - \frac{1}{6} \frac{d^2(\sigma S_3)}{d \ln \nu^2} + 2\nu \delta \sigma_8 \right] \end{aligned} \quad (43)$$

at the first order in the nonlinear parameters f_{NL} and g_{NL} .

B. Bias

Having checked that the amount of non-Gaussianity in the mass function is also consistent with our simple theoretical expectation when both f_{NL} and g_{NL} are nonzero, we now turn to the clustering of dark matter haloes. As shown in Appendix A, the non-Gaussian correction to the halo




FIG. 10 (color online). Fractional correction to the Gaussian multiplicity function of dark matter haloes as a function of the peak height $\nu(M, z)$ for $f_{\text{NL}} = \pm 100$ and a cubic coupling parameter $g_{\text{NL}} = -3 \times 10^5$. The solid and dashed curves show the theoretical prediction Eq. (42) at $z = 0$ and 2 , respectively. Error bars denote Poisson errors.

power spectrum can eventually be written down as

$$\begin{aligned} \Delta P_{\text{hh}}(k) &= 4f_{\text{NL}}b_L^2\delta_c(z)\alpha_M(k)P_\phi(k) \\ &+ 4f_{\text{NL}}^2b_L^2\delta_c^2(z)P_\phi(k) \\ &+ \left(g_{\text{NL}} + \frac{4}{3}f_{\text{NL}}^2\right)b_L^2\delta_c^2(z)S_3^{(1)}(M)\alpha_M(k)P_\phi(k). \end{aligned} \quad (44)$$

If we set $g_{\text{NL}} = 0$ and keep only the first two terms in the right-hand side, then the non-Gaussian (Eulerian) halo power spectrum can be cast into the form

$$P_{\text{hh}}(k) = [b(M) + f_{\text{NL}}b_\phi(k)]^2 P_\delta(k) \quad (45)$$

where the scale-dependent bias parameter $b_\phi(k)$ is

$$b_\phi(k) = 2[b(M) - 1]\delta_c(z)\alpha_M^{-1}(k). \quad (46)$$

Note that Ref. [64] obtained this relation by considering the halo power spectrum implied by a bias relation that is a local mapping of the density field. In practice, the term proportional to $P_\phi(k)$ is negligible as it contributes only at very small wave number $k \lesssim 0.001h^{-1}$ Mpc. The third term in the right-hand side of Eq. (44) is derived in this paper for the first time. In the case $g_{\text{NL}} = 0$, its magnitude relative to the term linear in f_{NL} is $(1/3)f_{\text{NL}}\delta_c(z)S_3^{(1)}(M)$, which is approximately ~ 0.03 at redshift $z = 1.8$ and for a mass scale $M = 10^{13}M_\odot/h$. Although its contribution be-

comes increasingly important at higher redshift, it is fairly small for the values of f_{NL} considered here. Consequently, we shall employ the approximation

$$\begin{aligned} \Delta b(k, \text{NL}) &= \epsilon_\kappa \Delta b_\kappa(k, g_{\text{NL}}) + \Delta b_\kappa(k, f_{\text{NL}}) \\ &+ [\Delta b_1(\text{NL}) + \epsilon_1]. \end{aligned} \quad (47)$$

to describe the non-Gaussian bias of dark matter haloes.

The quadratic coupling $f_{\text{NL}}\phi^2$ also affects the matter power spectrum at leading order [38,84], positive values of f_{NL} increasing the small scale power. However, the relative size of this k -dependent correction, $\beta_m(k, f_{\text{NL}})$, is at a per cent level only in the weakly nonlinear regime $k \lesssim 0.1h$ Mpc $^{-1}$ [53,85] and fades rapidly as one goes to larger scales. We will thus neglect it in what follows.

In Fig. 11, the result of measuring ratios of auto- and cross-power spectra in the simulations with $(f_{\text{NL}}, g_{\text{NL}}) = (\pm 100, -3 \times 10^5)$ is shown at $0 < z < 1.5$ for the haloes with $M > 2 \times 10^{13}M_\odot/h$. We do not quote error bars since the data points are obtained by averaging over two realizations only. The solid and dashed curves show the theoretical prediction Eq. (47). The value of the multiplicative factor $\epsilon_\kappa(b, g_{\text{NL}})$ was obtained from the four-parameter formula Eq. (28), while $\Delta b_1 + \epsilon_1$ was individually fitted for each halo sample over the wave number range $0.005 < k < 0.03h$ Mpc $^{-1}$. As can be seen, the theoretical expectation Eq. (47) agrees reasonably well with the numerical data. This demonstrates that the range of validity of the




FIG. 11 (color online). Non-Gaussian bias correction measured in the simulations with $(f_{\text{NL}}, g_{\text{NL}}) = (\pm 100, -3 \times 10^5)$ for haloes of mass $M > 2 \times 10^{13}M_\odot/h$. Error bars are not shown as the data points are averaged over two realizations solely.

non-Gaussian bias formula Eq. (27) extends to smaller values of g_{NL} as well as models with nonvanishing f_{NL} and g_{NL} .

The lowest order, k -dependent corrections to the Gaussian bias induced by the quadratic and the cubic coupling are fully degenerated in the halo power spectrum as they both scale as $\alpha_M^{-1}(k) \propto k^{-2}T(k)^{-1}$. For the values of f_{NL} and g_{NL} and the halo mass range considered here, the ratio $\Delta b_\kappa(k, g_{\text{NL}})/\Delta b_\kappa(k, f_{\text{NL}})$ increases approximately from 0.25 to 0.5 when the redshift increases from $z = 0$ to 2. It is unclear whether higher-order corrections could help breaking such a degeneracy. A more promising alternative may be to measure the bispectrum of dark matter haloes, which carries much more information about the shape of the primordial three-point function than the power spectrum of bias tracers [40,41]. However, this is beyond the scope of this paper.

VI. DISCUSSION

In this paper we explored the effect of a local cubic coupling $g_{\text{NL}}\phi^3$ on the mass function and bias of dark matter haloes. We derived analytical expressions for these statistics and tested them against the outcome of numerical simulations.

We showed that current theoretical predictions of the non-Gaussian correction to the mass function reasonably agree with the simulations. The LV formula [73] appears to provide a better fit to the data than the MVJ formula [74], in agreement with some of the literature on the subject [46,86]. The two approximations can be combined to provide an accurate description if one adjusts the low- ν expansion of the latter so as to match that of the former.

We found that the magnitude of the non-Gaussian scale-dependent bias $\Delta b_\kappa(k, g_{\text{NL}})$ is suppressed relative to a theoretical prediction based on the statistics of highly overdense regions, even on linear scales $k \lesssim 0.01h \text{ Mpc}^{-1}$. This suppression is stronger for the low biased samples $b \lesssim 3$ and, at fixed value of b , for positive values of g_{NL} . We were able to fit the measured halo bias at the expense of introducing two free parameters, ϵ_κ and ϵ_Γ , that depend mostly on the halo bias $b(M)$ and the coupling parameter g_{NL} . These parameters quantify the departure from the theoretical scale-dependent and scale-independent non-Gaussian bias correction, respectively. We provide a simple fitting formula for $\epsilon_\kappa(b, g_{\text{NL}})$, Eq. (28), which should be used when analyzing observational data. In non-Gaussian simulations of the f_{NL} type, the data also hint at a (possibly f_{NL} -dependent) suppression of the non-Gaussian scale-dependent bias relative to theory for wave modes $k \lesssim 0.03h \text{ Mpc}^{-1}$ [53–55], but the effect is much weaker than seen in our simulations of g_{NL} models. Clearly, these results require a better theoretical modelling of the non-Gaussian halo bias.

Reference [55] argued that both the MVJ and LV analytic formula can be reconciled with measurements of the

non-Gaussian fractional correction to the mass function once nonspherical collapse is included. In practice, the critical density for collapse is replaced by $\delta_c \rightarrow \sqrt{q}\delta_c$, where the value $q = 0.75$ is obtained from a fit to the mass function measured in simulations [87]. Reference [88] claimed that such a relation is a consequence of the diffusive nature of the critical threshold for collapse. Their model predicts $q \simeq 0.8$, in good agreement with the findings of [55]. However, we found that substituting $\delta_c \rightarrow \sqrt{q}\delta_c$ in Eq. (18) only modestly improve the agreement with the data. Regarding the non-Gaussian bias, it is not obvious how one could justify the replacement $\delta_c \rightarrow \sqrt{q}\delta_c$ given that the linear bias of our (Gaussian) halo samples converges towards the spherical collapse prediction ν^2/δ_c for large peak height.

A important ingredient is the choice of the halo identification algorithm. While we used a spherical overdensity (SO) finder, Ref. [88] considered a Friends-of-Friends (FoF) finder with a linking length $b = 0.2$. The question of how the spherical overdensity masses can be mapped onto friends-of-friends masses remains a matter of debate [89]. Clearly however, since the peak height depends on the halo mass M through the variance $\sigma(M)$, any systematic difference will be reflected in the value of ν associated to a specific halo sample. This will in turn affects the size of the fractional deviation from the Gaussian mass function at some specified peak height. The sensitivity of the non-Gaussian mass function and bias to the halo finder will be presented elsewhere.

The observational bound on f_{NL} inferred by [48] from the clustering of a high redshift sample of quasars can be straightforwardly translated into a limit on g_{NL} since $\Delta b_\kappa(k, g_{\text{NL}})$ also scales as $k^{-2}T(k)^{-1}$ at low wave number. We have obtained

$$-3.5 \times 10^5 < g_{\text{NL}} < +8.2 \times 10^5 (95\%). \quad (48)$$

These are the first limits derived on g_{NL} . While they are too weak to provide interesting constraints on inflationary scenarios such as the curvaton model, future all-sky redshift surveys should improve them by a factor of ~ 100 . Future CMB observations, including PLANCK, should also improve the limits derived here by an order of magnitude. With these improvements we expect that, in the next decade, realistic models of cubic type non-Gaussianity will be tested with real observations.

ACKNOWLEDGMENTS

We thank Paolo Creminelli and Leonardo Senatore for useful discussions. We acknowledge support from the Swiss National Foundation under Contract No. 200021-116696/1 and WCU Grant No. R32-2008-000-10130-0.

APPENDIX A: NON-GAUSSIAN BIAS IN THE HIGH PEAK LIMIT

In this appendix, we detail the derivation of the scale-dependent bias correction induced by the g_{NL} coupling to the two-point correlation of dark matter haloes. We follow [47] and approximate the latter by the two-point correlation $\xi_{\text{hh}}(\mathbf{r})$ of regions of the smoothed density field δ_M with a peak height $\nu \gg 1$.

1. Case $g_{\text{NL}} \neq 0$ only

In this case, only the four-point correlations of the density field contribute at the first order. The leading-order correction to the correlation of thresholded regions thus reads as

$$\Delta \xi_{\text{hh}} = \frac{\nu^4}{\sigma^4} \left[\frac{1}{3} \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) + \frac{1}{4} \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) \right]. \quad (\text{A1})$$

The four-point correlations $\xi^{(4)}$ are Fourier transform of the trispectrum of the density field, $T_\delta(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4)$, with conservation of momentum enforced. Linearly extrapolating the density field to present epoch, the latter can be expressed as

$$T_\delta(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) = \left(\prod_{i=1}^4 \alpha_M(k_i) T_\Phi(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) \right), \quad (\text{A2})$$

where the expression for the trispectrum of primordial curvature perturbation

$$T_\Phi(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) = 6g_{\text{NL}} [P_\phi(k_1)P_\phi(k_2)P_\phi(k_3) + (\text{cyclic})] \quad (\text{A3})$$

follows straightforwardly from the Fourier mode relation

$$\Phi(\mathbf{k}) = \phi(\mathbf{k}) + g_{\text{NL}} \int \frac{d^3 k_1}{(2\pi)^3} \int \frac{d^3 k_2}{(2\pi)^3} \phi(\mathbf{k}_1)\phi(\mathbf{k}_2) \times \phi(\mathbf{k} - \mathbf{k}_1 - \mathbf{k}_2). \quad (\text{A4})$$

Density and curvature perturbations are related through the Poisson equation, whose scale-dependence is reflected in the function $\alpha_M(k) = \alpha_R(k, z=0)$ defined in Eq. (6). Combining these relations gives

$$\xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) = 6g_{\text{NL}} \left(\prod_{i=1}^3 \int \frac{d^3 k_i}{(2\pi)^3} \alpha_M(k_i) P_\phi(k_i) \right) \times \left[1 + 3 \frac{P_\phi(k_{123})}{P_\phi(k_3)} \right] \alpha_M(k_{123}) e^{i\mathbf{k}_{123} \cdot \mathbf{r}} \quad (\text{A5})$$

$$\xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) = 6g_{\text{NL}} \left(\prod_{i=1}^3 \int \frac{d^3 k_i}{(2\pi)^3} \alpha_M(k_i) P_\phi(k_i) \right) \times \left[1 + 2 \frac{P_\phi(k_{123})}{P_\phi(k_1)} + \frac{P_\phi(k_{123})}{P_\phi(k_3)} \right] \times \alpha_M(k_{123}) e^{i\mathbf{k}_{12} \cdot \mathbf{r}}, \quad (\text{A6})$$

where we have defined $\mathbf{k}_{ij\dots l} = \mathbf{k}_i + \mathbf{k}_j + \dots + \mathbf{k}_l$ for shorthand convenience. Since we will examine the effect of non-Gaussianity on Fourier space statistics only, we take the Fourier transform of the four-point functions. After some simplification, we arrive at

$$\int d^3 r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) e^{-i\mathbf{k} \cdot \mathbf{r}} = 6g_{\text{NL}} \alpha_M(k) P_\phi(k) \int \frac{d^3 k_1}{(2\pi)^3} \alpha_M(k_1) P_\phi(k_1) \times \int \frac{d^3 k_2}{(2\pi)^3} \alpha_M(k_2) P_\phi(k_2) \alpha_M(|\mathbf{k} + \mathbf{k}_{12}|) \times \left[3 + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_{12}|)}{P_\phi(k)} \right] \quad (\text{A7})$$

$$\int d^3 r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) e^{-i\mathbf{k} \cdot \mathbf{r}} = 6g_{\text{NL}} \int \frac{d^3 k_1}{(2\pi)^3} \alpha_M(k_1) \alpha_M(|\mathbf{k} + \mathbf{k}_1|) P_\phi(k_1) \times P_\phi(|\mathbf{k} + \mathbf{k}_1|) \int \frac{d^3 k_2}{(2\pi)^3} \alpha_M(k_2) \alpha_M(|\mathbf{k} + \mathbf{k}_2|) \times P_\phi(k_2) \left[1 + 2 \frac{P_\phi(|\mathbf{k} + \mathbf{k}_2|)}{P_\phi(|\mathbf{k} + \mathbf{k}_1|)} + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_2|)}{P_\phi(k_2)} \right]. \quad (\text{A8})$$

For realistic values of the spectral index ($n_s \sim 1$), the products $\alpha_M(|\mathbf{k} + \mathbf{k}_i|) P_\phi(|\mathbf{k} + \mathbf{k}_i|)$ appearing in the right-hand side of the above equalities formally diverges whenever $\mathbf{k} + \mathbf{k}_1 = 0$ due to the ultraviolet divergence of $P_\phi(k)$. To cure this problem, one can set $P_\phi(k) = 0$ for sufficiently small wave numbers or excise a thin shell centered at wave number k_i from the integral. In the large-scale limit $k \ll k_i$, the ratio $P_\phi(|\mathbf{k} + \mathbf{k}_i|)/P_\phi(k)$ vanishes whereas $P_\phi(|\mathbf{k} + \mathbf{k}_i|)/P_\phi(k_i)$ tends towards unity. In this case, the above expressions reduce to

$$\int d^3 r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) e^{-i\mathbf{k} \cdot \mathbf{r}} \approx 3g_{\text{NL}} \sigma^4 S_3^{(1)}(M) \alpha_M(k) P_\phi(k) + 6g_{\text{NL}} \alpha_M(k) \times \int \frac{d^3 k_1}{(2\pi)^3} \alpha_M(k_1) P_\phi(k_1) \times \int \frac{d^3 k_2}{(2\pi)^3} \alpha_M(k_2) P_\phi(k_2) \alpha_M(k_{12}) P_\phi(k_{12}), \quad (\text{A9})$$

and

$$\int d^3r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) e^{-i\mathbf{k}\cdot\mathbf{r}} \approx 24g_{\text{NL}}\sigma^2 \int \frac{d^3k_1}{(2\pi)^3} \alpha_M^2(k_1) P_\phi^2(k_1). \quad (\text{A10})$$

Only the first term in the right-hand side of Eq. (A9) is not well behaved in the limit $k \rightarrow 0$ where it becomes proportional to k^{n_s-2} . The second scales as k^2 , while the Fourier transform of $\xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2)$ asymptotes to a constant. A similar decomposition also arises in f_{NL} models. For this type of non-Gaussianity, the first order correction is furnished by the three-point function $\xi^{(3)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2)$, whose Fourier transform can be split into the familiar term $2f_{\text{NL}}\sigma^2\alpha_M(k)P_\phi(k)$, and a second piece given by

$$\frac{1}{2}f_{\text{NL}}\alpha_M(k) \int \frac{d^3k_1}{(2\pi)^3} \alpha_M^2(k_1) P_\phi^2(k_1). \quad (\text{A11})$$

In both quadratic and cubic local non-Gaussianity, the term proportional to $\alpha_M(k)$ can be neglected since, at the pivot point $k = k_0$, its magnitude relative to the term involving $\alpha_M(k)P_\phi(k)$ is only $\mathcal{O}(0.01)$ and $\mathcal{O}(10^{-6})$, respectively. Moreover, it decreases as one goes to larger scales. By contrast, it is not so obvious how to handle the term (A10). In the non-Gaussian halo power spectrum, this term would appear multiplied by $\nu^4/(4\sigma^4)$,

$$6g_{\text{NL}}b_L^2 \frac{\delta_c^2(z)}{\sigma^2} \int \frac{d^3k_1}{(2\pi)^3} \alpha_M^2(k_1) P_\phi^2(k_1) \approx 6 \times 10^{-4} g_{\text{NL}} b_L^2 \frac{D^2(0)}{D^2(z)} \left(\frac{M}{10^{13} M_\odot/h} \right)^{0.375}. \quad (\text{A12})$$

The approximation (second line) holds for $10^{13} \leq M \leq 10^{14} M_\odot/h$. For $g_{\text{NL}} = 10^6$ and $b_L \geq 3$, this can be much larger than the typical shot-noise correction applied to the halo power spectra we measure in the simulations (see below). For $g_{\text{NL}} = -10^6$, this will certainly produce a halo power spectrum which is negative at sufficiently low wave number. It is plausible that higher-order counterterms in the expansion Eq. (21) renormalizes its value. However, such a calculation is beyond the scope of this paper, so the simplest choice is to ignore this term in the following of the analysis. Hence, we can approximate the non-Gaussian correction ΔP_{hh} in the limit of long wavelength $k \ll 1$ by the Fourier transform of $\nu^4 \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2)/3\sigma^4$, Eq. (23).

2. Case g_{NL} and f_{NL} nonzero

In addition to the first order trispectrum induced by $g_{\text{NL}}\phi^3$, Eq. (A3), the quadratic coupling $f_{\text{NL}}\phi^2$ generates the bispectrum at leading order

$$B_\Phi(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) = 2f_{\text{NL}}[P(k_1)P(k_2) + (\text{cyclic})], \quad (\text{A13})$$

and an additional, albeit second order, contribution to the

trispectrum [90–92],

$$T_\Phi(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4) = 4f_{\text{NL}}^2 [P_\phi(k_{13})P_\phi(k_3)P_\phi(k_4) + 11 \text{ permutations}]. \quad (\text{A14})$$

The bispectrum (A13) induces a three-point contribution $(\nu^3/\sigma^3)\xi^{(3)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2)$ to the power spectrum of biased tracers which is calculated in [47]. Upon Fourier transformation, it reads as

$$\Delta P_{\text{hh}}(k) = 4f_{\text{NL}}b_L^2 \delta_c(z) \alpha_M(k) P_\phi(k). \quad (\text{A15})$$

We follow the steps outlined above to calculate the contribution from the second order trispectrum (A14). After some algebra, the Fourier transform of the four-point correlations of the density field can be written down as

$$\begin{aligned} & \int d^3r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) e^{-i\mathbf{k}\cdot\mathbf{r}} \\ &= 8f_{\text{NL}}^2 \alpha_M(k) P_\phi(k) \int \frac{d^3k_1}{(2\pi)^3} \alpha_M(k_1) P_\phi(k_1) \\ & \quad \times \int \frac{d^3k_2}{(2\pi)^3} \alpha_M(k_2) P_\phi(k_2) \alpha_M(|\mathbf{k} + \mathbf{k}_{12}|) \\ & \quad \times \left\{ \frac{P_\phi(k_{12})}{P_\phi(k_2)} \left[1 + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_{12}|)}{P_\phi(k)} \right] + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)}{P_\phi(k)} \right. \\ & \quad + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)}{P_\phi(k_1)} + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)P_\phi(|\mathbf{k} + \mathbf{k}_{12}|)}{P_\phi(k_1)P_\phi(k_2)} \\ & \quad \left. + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)P_\phi(|\mathbf{k} + \mathbf{k}_{12}|)}{P_\phi(k)P_\phi(k_2)} \right\}, \quad (\text{A16}) \end{aligned}$$

and

$$\begin{aligned} & \int d^3r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) e^{-i\mathbf{k}\cdot\mathbf{r}} \\ &= 4f_{\text{NL}}^2 \int \frac{d^3k_1}{(2\pi)^3} \alpha_M(k_1) \alpha_M(|\mathbf{k} + \mathbf{k}_1|) P_\phi(k_1) \\ & \quad \times \int \frac{d^3k_2}{(2\pi)^3} \alpha_M(k_2) \alpha_M(|\mathbf{k} + \mathbf{k}_2|) P_\phi(k_2) \left\{ P_\phi(|\mathbf{k} + \mathbf{k}_{12}|) \right. \\ & \quad \times \left[1 + 2 \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)}{P_\phi(k_2)} + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)P_\phi(|\mathbf{k} + \mathbf{k}_2|)}{P_\phi(k_1)P_\phi(k_2)} \right] \\ & \quad + P_\phi(k) \left[1 + 2 \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)}{P_\phi(k_1)} \right. \\ & \quad \left. + \frac{P_\phi(|\mathbf{k} + \mathbf{k}_1|)P_\phi(|\mathbf{k} + \mathbf{k}_2|)}{P_\phi(k_1)P_\phi(k_2)} \right] + 2P_\phi(|\mathbf{k} + \mathbf{k}_1|) \\ & \quad \left. \times \left[\frac{P_\phi(k_{12})}{P_\phi(k_1)} + \frac{P_\phi(k_{12})}{P_\phi(k_2)} \right] \right\}. \quad (\text{A17}) \end{aligned}$$

In the large-scale limit $k \rightarrow 0$, these expressions asymptote to

$$\begin{aligned}
& \int d^3 r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2) e^{-i\mathbf{k}\cdot\mathbf{r}} \\
& \approx 4f_{\text{NL}}^2 \sigma^4 S_3^{(1)}(M) \alpha_M(k) P_\phi(k) + 8f_{\text{NL}}^2 \alpha_M(k) \\
& \quad \times \int \frac{d^3 k_1}{(2\pi)^3} \alpha_M^2(k_1) P_\phi(k_1) \int \frac{d^3 k_2}{(2\pi)^3} \alpha_M^2(k_2) \alpha_M(k_{12}) \\
& \quad \times [P_\phi(k_1) P_\phi(k_2) + P_\phi(k_1) P_\phi(k_{12}) + P_\phi^2(k_{12})]
\end{aligned} \tag{A18}$$

$$\begin{aligned}
& \int d^3 r \xi^{(4)}(\mathbf{x}_1, \mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_2) e^{-i\mathbf{k}\cdot\mathbf{r}} \\
& \approx 16f_{\text{NL}}^2 \sigma^4 P_\phi(k) + 16f_{\text{NL}}^2 \int \frac{d^3 k_1}{(2\pi)^3} \alpha_M^2(k_1) \\
& \quad \times \int \frac{d^3 k_2}{(2\pi)^3} \alpha_M^2(k_2) P_\phi(k_2) P_\phi(k_{12}) [P_\phi(k_1) + P_\phi(k_2)].
\end{aligned} \tag{A19}$$

Ignoring the second piece in the right-hand side of Eqs. (A18) and (A19), the non-Gaussian correction to the halo power spectrum is then easily recast as Eq. (44).

APPENDIX B: SIGNAL-TO-NOISE FOR THE CMB TRISPECTRUM

The formalism for the CMB trispectrum has been established in [29,30]. The invariance of the 4-point harmonic function of the CMB temperature anisotropy field under the 4! permutations of the coefficients $a_{l_i}^{m_i}$ imposes constraints on the CMB trispectrum $T_{l_3 l_4}^{l_1 l_2}(L)$ which can be enforced by defining

$$\begin{aligned}
T_{l_3 l_4}^{l_1 l_2}(L) &= P_{l_3 l_4}^{l_1 l_2}(L) + (2L+1) \sum_{L'} \left[(-1)^{l_2+l_3} \right. \\
& \quad \times \left\{ \begin{matrix} l_1 & l_2 & L \\ l_4 & l_3 & L' \end{matrix} \right\} P_{l_1 l_3}^{l_2 l_4}(L') + (-1)^{L+L'} \\
& \quad \times \left. \left\{ \begin{matrix} l_1 & l_2 & L \\ l_3 & l_4 & L' \end{matrix} \right\} P_{l_1 l_4}^{l_3 l_2}(L') \right],
\end{aligned} \tag{B1}$$

where curly brackets are Wigner-6j symbols,

$$\begin{aligned}
P_{l_3 l_4}^{l_1 l_2}(L) &= t_{l_3 l_4}^{l_1 l_2}(L) + (-1)^{\sum_i l_i} t_{l_3 l_4}^{l_1 l_2}(L) + (-1)^{L+l_3+l_4} t_{l_4 l_3}^{l_1 l_2}(L) \\
& \quad + (-1)^{L+l_1+l_2} t_{l_3 l_4}^{l_2 l_1}(L),
\end{aligned} \tag{B2}$$

and the reduced trispectrum $t_{l_3 l_4}^{l_1 l_2}(L)$ is symmetric under the exchange of its upper and lower indices and fully characterizes the model.

The expansion coefficients a_l^m are related to the primordial curvature perturbation $\Phi(\mathbf{k})$ through

$$a_l^m = 4\pi(-i)^l \int \frac{d^3 k}{(2\pi)^3} \Phi(\mathbf{k}) g_{Tl}(k) Y_l^{m*}(\hat{\mathbf{k}}), \tag{B3}$$

where $g_{Tl}(k)$ is the radiation transfer function. The reduced

trispectrum can be calculated from this relation once the four-point function $T_\Phi(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4)$ is specified. For a local cubic coupling $g_{\text{NL}} \phi^3$,

$$\begin{aligned}
t_{l_3 l_4}^{l_1 l_2}(L) &= \int_0^\infty dr r^2 \beta_{l_2}(r) \beta_{l_4}(r) h_{l_1 L l_2} h_{l_3 L l_4} [\mu_{l_1}(r) \beta_{l_3}(r) \\
& \quad + \beta_{l_1}(r) \mu_{l_3}(r)]
\end{aligned} \tag{B4}$$

with

$$\beta_l(r) = \frac{2}{\pi} \int_0^\infty dk k^2 P_\phi(k) g_{Tl}(k) j_l(kr) \tag{B5}$$

$$\mu_l(r) = \frac{2}{\pi} \int_0^\infty dk k^2 g_{\text{NL}} g_{Tl}(k) j_l(kr), \tag{B6}$$

and

$$h_{l_1 L l_2} = \frac{1}{\sqrt{4\pi}} \pi_{l_1 L l_2} \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix}. \tag{B7}$$

We also use the notation

$$\pi_{l_1 \dots l_j} = \sqrt{(2l_1+1) \times \dots \times (2l_j+1)}. \tag{B8}$$

Note that most of the contribution to $t_{l_3 l_4}^{l_1 l_2}(L)$ comes from a small volume centered at the comoving distance r_* to the surface of last scattering.

The Sachs-Wolfe approximation $g_{Tl}(k) \approx -j_l(kr_*)/3$ valid at low multipoles $l \ll 100$ provides a useful order-of-magnitude estimate [5,30,31,83]. In this limit, we can approximate $\mu_l(r)$ as $-g_{\text{NL}} \delta_D(r-r_*)/(3r_*^2)$ since we assume g_{NL} independent of wave number. Hence, the reduced trispectrum simplifies to

$$t_{l_3 l_4}^{l_1 l_2}(L) \approx 9g_{\text{NL}} C_{l_2}^{\text{SW}} C_{l_4}^{\text{SW}} (C_{l_1}^{\text{SW}} + C_{l_3}^{\text{SW}}) h_{l_1 L l_2} h_{l_3 L l_4}, \tag{B9}$$

Inserting this expression successively into Eqs. (B1) and (B2), the CMB trispectrum eventually reads as

$$\begin{aligned}
T_{l_3 l_4}^{l_1 l_2}(L) &= \frac{27}{2\pi} g_{\text{NL}} (2L+1) \pi_{l_1 l_2 l_3 l_4}^2 \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix} \\
& \quad \times \begin{pmatrix} l_3 & l_4 & L \\ 0 & 0 & 0 \end{pmatrix} [C_{l_1}^{\text{SW}} C_{l_2}^{\text{SW}} C_{l_3}^{\text{SW}} + (\text{cyclic})].
\end{aligned} \tag{B10}$$

where

$$C_l^{\text{SW}} = \frac{2}{9\pi} \int_0^\infty dk k^2 P_\phi(k) j_l^2(kr_*) \approx \frac{2\pi A_\phi}{9l(l+1)}. \tag{B11}$$

The last equality assumes a nearly scale-invariant spectrum $n_s \approx 1$. The following relation between the Wigner-3j and 6j symbols (e.g., Appendix A of [29]),

$$\begin{aligned}
& \sum_{l'_3} (2l'_3 + 1) (-1)^{\Sigma + l'_3 - l_3 - m_1 - m'_1} \begin{Bmatrix} l_1 & l_2 & l_3 \\ l'_1 & l'_2 & l'_3 \end{Bmatrix} \\
& \times \begin{pmatrix} l_2 & l'_1 & l'_3 \\ m_2 & m'_1 & -m'_3 \end{pmatrix} \begin{pmatrix} l_1 & l'_3 & l'_2 \\ m_1 & m_3 & -m'_2 \end{pmatrix} \\
& = \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & -m \end{pmatrix} \begin{pmatrix} l_3 & l'_1 & l'_2 \\ m & m'_1 & -m'_2 \end{pmatrix} \quad (\text{B12})
\end{aligned}$$

where $\Sigma = l_1 + l_2 + l'_1 + l'_2$ and the value of m is set by the triangle condition, can be useful to derive Eq. (B10). The signal-to-noise summed up to multipole l_{\max} Eq. (36) then becomes

$$\begin{aligned}
\left(\frac{S}{N}\right)^2 (< l_{\max}) &= \left(\frac{27}{2\pi}\right)^2 g_{\text{NL}}^2 \sum_{l_1 > l_2 > l_3 > l_4}^{l_{\max}} \pi_{l_1 l_2 l_3 l_4}^2 \\
& \times \frac{[C_{l_1}^{\text{SW}} C_{l_2}^{\text{SW}} C_{l_3}^{\text{SW}} + (\text{cyclic})]^2}{C_{l_1}^{\text{SW}} C_{l_2}^{\text{SW}} C_{l_3}^{\text{SW}} C_{l_4}^{\text{SW}}} \sum_{L=0}^{2l_{\max}} (2L+1) \\
& \times \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix}^2 \begin{pmatrix} l_3 & l_4 & L \\ 0 & 0 & 0 \end{pmatrix}^2. \quad (\text{B13})
\end{aligned}$$

We can recast the sum over the diagonal modes L into a manifestly symmetric form with the aid of the Gaunt integral

$$\frac{1}{2} \int_{-1}^{+1} dx P_{l_1}(x) P_{l_2}(x) P_{l_3}(x) = \begin{pmatrix} l_2 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix}^2 \quad (\text{B14})$$

and the orthogonality relation

$$\sum_{k=0}^{\infty} (2k+1) P_k(x) P_k(y) = 2\delta_D(x-y), \quad (\text{B15})$$

where δ_D is the Dirac delta. We find

$$\begin{aligned}
& \sum_{L=0}^{2l_{\max}} (2L+1) \begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix}^2 \begin{pmatrix} l_3 & l_4 & L \\ 0 & 0 & 0 \end{pmatrix}^2 \\
& = \frac{1}{2} \int_{-1}^{+1} dx P_{l_1}(x) P_{l_2}(x) P_{l_3}(x) P_{l_4}(x). \quad (\text{B16})
\end{aligned}$$

There is a strict equality because the Wigner-3j symbols vanish for $L > l_1 + l_2$. Equation (37) for the signal-to-noise then follows by replacing the above equality into Eq. (B13) and summing over all the $4!$ permutations of the quadruplet (l_1, l_2, l_3, l_4) . Although Eq. (37) becomes computationally expensive when $l_{\max} \gg 100$ (because we are summing over redundant configurations), we found that it is quite efficient for $l_{\max} \lesssim 200$.

Following [31], we can roughly estimate the dependence of the signal-to-noise squared on l_{\max} by considering only the contribution of the $L=1$ mode in Eq. (B13). Consequently, the product of the Wigner-3j symbols squared reduces to $\sim l_1 l_3 \delta_{l_1-1, l_2} \delta_{l_3-1, l_4}$ and yields $(S/N)^2 \propto l_{\max}^2$. However, including all L modes as in Eq. (37) gives a steeper dependence, $(S/N)^2 \propto l_{\max}^{2.6}$ (see Fig. 9), due to the fact that the Wigner-3j symbols decay slowly with increasing L . This is quite apparent in the classical limit $l_1, l_2, L \gg 1$, where

$$\begin{aligned}
\begin{pmatrix} l_1 & l_2 & L \\ 0 & 0 & 0 \end{pmatrix}^2 &\approx \frac{1}{\pi} [(l_1 + l_2)^2 - L^2]^{-1/2} \\
& \times [L^2 - (l_1 - l_2)^2]^{-1/2}. \quad (\text{B17})
\end{aligned}$$

Clearly, the terms in the summation Eq. (B13) decay only as $1/L$ for $L \gg l_1, l_2$. By contrast, the second order contribution to the CMB trispectrum induced by the quadratic coupling $f_{\text{NL}} \phi^2$ adds an additional multiplicative factor of $(C_L^{\text{SW}})^2$ in the summation over the L modes which increases the relative contribution of the low- L modes (since these now decay as L^{-5}). This is the reason why considering only $L \leq 10$ modes as done in [31] still provides a good approximation to the signal-to-noise of the CMB trispectrum for the f_{NL} model.

-
- [1] V.F. Mukhanov and G.V. Chibisov, JETP Lett. **33**, 532 (1981).
[2] A.A. Starobinsky, Phys. Lett. **117B**, 175 (1982).
[3] S.W. Hawking, Phys. Lett. **115B**, 295 (1982).
[4] A.H. Guth and S. Pi, Phys. Rev. Lett. **49**, 1110 (1982).
[5] A. Gangui, F. Lucchin, S. Matarrese, and S. Mollerach, Astrophys. J. **430**, 447 (1994).
[6] T.J. Allen, B. Grinstein, and M.B. Wise, Phys. Lett. B **197**, 66 (1987).
[7] T. Falk, R. Rangarajan, and M. Srednicki, Phys. Rev. D **46**, 4232 (1992).
[8] N. Bartolo, E. Komatsu, S. Matarrese, and A. Riotto, Phys. Rep. **402**, 103 (2004).
[9] V. Acquaviva, N. Bartolo, S. Matarrese, and A. Riotto, Nucl. Phys. **B667**, 119 (2003).
[10] J. Maldacena, J. High Energy Phys. 5 (2003) 013.
[11] E. Komatsu, A. Kogut, M.R. Nolta, C.L. Bennett, M. Halpern, G. Hinshaw, N. Jarosik, M. Limon, S. S. Meyer, L. Page, D.N. Spergel, G. S. Tucker, L. Verde, E. Wollack, and E.L. Wright, Astrophys. J. Suppl. Ser. **148**, 119 (2003).
[12] P. Creminelli, L. Senatore, M. Zaldarriaga, and M. Tegmark, J. Cosmol. Astropart. Phys. 03 (2007) 005.
[13] E. Komatsu, J. Dunkley, M.R. Nolta, C.L. Bennett, B.

- Gold, G. Hinshaw, N. Jarosik, D. Larson, M. Limon, L. Page, D. N. Spergel, M. Halpern, R. S. Hill, A. Kogut, S. S. Meyer, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright, *Astrophys. J. Suppl. Ser.* **180**, 330 (2009).
- [14] K. M. Smith, L. Senatore, and M. Zaldarriaga (2009).
- [15] A. Curto, E. Martínez-González, P. Mukherjee, R. B. Barreiro, F. K. Hansen, M. Liguori, and S. Matarrese, *Mon. Not. R. Astron. Soc.* **393**, 615 (2009).
- [16] D. H. Lyth, C. Ungarelli, and D. Wands, *Phys. Rev. D* **67**, 023503 (2003).
- [17] N. Bartolo, S. Matarrese, and A. Riotto, *Phys. Rev. D* **69**, 043503 (2004).
- [18] K. Enqvist and S. Nurmi, *J. Cosmol. Astropart. Phys.* **10** (2005) 013.
- [19] K. A. Malik and D. H. Lyth, *J. Cosmol. Astropart. Phys.* **09** (2006) 008.
- [20] M. Sasaki, J. Väliviita, and D. Wands, *Phys. Rev. D* **74**, 103003 (2006).
- [21] K. Enqvist and T. Takahashi, *J. Cosmol. Astropart. Phys.* **09** (2008) 012.
- [22] Q.-G. Huang and Y. Wang, *J. Cosmol. Astropart. Phys.* **09** (2008) 025.
- [23] P. Chingangbam and Q.-G. Huang, *J. Cosmol. Astropart. Phys.* **04** (2009) 031.
- [24] K. Ichikawa, T. Suyama, T. Takahashi, and M. Yamaguchi, *Phys. Rev. D* **78**, 023513 (2008).
- [25] Q.-G. Huang, *J. Cosmol. Astropart. Phys.* **11** (2008) 005.
- [26] Q.-G. Huang, *J. Cosmol. Astropart. Phys.* **06** (2009) 035.
- [27] C. T. Byrnes and G. Tasinato, *J. Cosmol. Astropart. Phys.* **08** (2009) 016.
- [28] J.-L. Lehners and S. Renaux-Petel, *Phys. Rev. D* **80**, 063503 (2009).
- [29] W. Hu, *Phys. Rev. D* **64**, 083005 (2001).
- [30] T. Okamoto and W. Hu, *Phys. Rev. D* **66**, 063008 (2002).
- [31] N. Kogo and E. Komatsu, *Phys. Rev. D* **73**, 083007 (2006).
- [32] E. Komatsu (2002).
- [33] M. Kunz, A. J. Banday, P. G. Castro, P. G. Ferreira, and K. M. Górski, *Astrophys. J. Lett.* **563**, L99 (2001).
- [34] F. Lucchin and S. Matarrese, *Astrophys. J.* **330**, 535 (1988).
- [35] S. Colafrancesco, F. Lucchin, and S. Matarrese, *Astrophys. J.* **345**, 3 (1989).
- [36] B. Grinstein and M. B. Wise, *Astrophys. J.* **310**, 19 (1986).
- [37] S. Matarrese, F. Lucchin, and S. A. Bonometto, *Astrophys. J. Lett.* **310**, L21 (1986).
- [38] R. Scoccimarro, E. Sefusatti, and M. Zaldarriaga, *Phys. Rev. D* **69**, 103513 (2004).
- [39] E. Sefusatti and E. Komatsu, *Phys. Rev. D* **76**, 083004 (2007).
- [40] E. Sefusatti (2009).
- [41] D. Jeong and E. Komatsu (2009).
- [42] M. Kamionkowski, L. Verde, and R. Jimenez, *J. Cosmol. Astropart. Phys.* **01** (2009) 010.
- [43] T. Y. Lam and R. K. Sheth, *Mon. Not. R. Astron. Soc.* **395**, 1743 (2009).
- [44] C. Hikage, E. Komatsu, and T. Matsubara, *Astrophys. J.* **653**, 11 (2006).
- [45] C. Hikage, P. Coles, M. Grossi, L. Moscardini, K. Dolag, E. Branchini, and S. Matarrese, *Mon. Not. R. Astron. Soc.* **385**, 1613 (2008).
- [46] N. Dalal, O. Doré, D. Huterer, and A. Shirokov, *Phys. Rev. D* **77**, 123514 (2008).
- [47] S. Matarrese and L. Verde, *Astrophys. J. Lett.* **677**, L77 (2008).
- [48] A. Slosar, C. Hirata, U. Seljak, S. Ho, and N. Padmanabhan, *J. Cosmol. Astropart. Phys.* **08** (2008) 031.
- [49] N. Afshordi and A. J. Tolley, *Phys. Rev. D* **78**, 123507 (2008).
- [50] C. Carbone, L. Verde, and S. Matarrese, *Astrophys. J. Lett.* **684**, L1 (2008).
- [51] U. Seljak, *Phys. Rev. Lett.* **102**, 021302 (2009).
- [52] E. Sefusatti, M. Liguori, A. P. S. Yadav, M. G. Jackson, and E. Pajer (2009).
- [53] V. Desjacques, U. Seljak, and I. T. Iliev, *Mon. Not. R. Astron. Soc.* **396**, 85 (2009).
- [54] A. Pillepich, C. Porciani, and O. Hahn (2008).
- [55] M. Grossi, L. Verde, C. Carbone, K. Dolag, E. Branchini, F. Iannuzzi, S. Matarrese, and L. Moscardini (2009).
- [56] U. Seljak and M. Zaldarriaga, *Astrophys. J.* **469**, 437 (1996).
- [57] V. Springel, *Mon. Not. R. Astron. Soc.* **364**, 1105 (2005).
- [58] Y. B. Zel'Dovich, *Astron. Astrophys.* **5**, 84 (1970).
- [59] J. N. Fry, *Astrophys. J.* **279**, 499 (1984).
- [60] M. H. Goroff, B. Grinstein, S.-J. Rey, and M. B. Wise, *Astrophys. J.* **311**, 6 (1986).
- [61] F. R. Bouchet, R. Juszkiewicz, S. Colombi, and R. Pellat, *Astrophys. J. Lett.* **394**, L5 (1992).
- [62] F. Bernardeau, *Astrophys. J.* **433**, 1 (1994).
- [63] R. Scoccimarro, *Mon. Not. R. Astron. Soc.* **299**, 1097 (1998).
- [64] P. McDonald, *Phys. Rev. D* **78**, 123519 (2008).
- [65] S. R. Knollmann and A. Knebe (2009).
- [66] S. P. D. Gill, A. Knebe, and B. K. Gibson, *Mon. Not. R. Astron. Soc.* **351**, 399 (2004).
- [67] M. S. Warren, P. J. Quinn, J. K. Salmon, and W. H. Zurek, *Astrophys. J.* **399**, 405 (1992).
- [68] C. Lacey and S. Cole, *Mon. Not. R. Astron. Soc.* **271**, 676 (1994).
- [69] V. R. Eke, S. Cole, and C. S. Frenk, *Mon. Not. R. Astron. Soc.* **282**, 263 (1996).
- [70] G. L. Bryan and M. L. Norman, *Astrophys. J.* **495**, 80 (1998).
- [71] W. H. Press and P. Schechter, *Astrophys. J.* **187**, 425 (1974).
- [72] J. R. Bond, S. Cole, G. Efstathiou, and N. Kaiser, *Astrophys. J.* **379**, 440 (1991).
- [73] M. Lo Verde, A. Miller, S. Shandera, and L. Verde, *J. Cosmol. Astropart. Phys.* **04** (2008) 014.
- [74] S. Matarrese, L. Verde, and R. Jimenez, *Astrophys. J.* **541**, 10 (2000).
- [75] M. Grossi, K. Dolag, E. Branchini, S. Matarrese, and L. Moscardini, *Mon. Not. R. Astron. Soc.* **382**, 1261 (2007).
- [76] R. K. Sheth and G. Tormen, *Mon. Not. R. Astron. Soc.* **308**, 119 (1999).
- [77] P. Valageas (2009).
- [78] T. Y. Lam and R. K. Sheth (2009).
- [79] T. Y. Lam, R. K. Sheth, and V. Desjacques, *Mon. Not. R. Astron. Soc.* **399**, 1482 (2009).
- [80] A. Slosar, *J. Cosmol. Astropart. Phys.* **03** (2009) 004.
- [81] U. Seljak, N. Hamaus, and V. Desjacques, *Phys. Rev. Lett.* **103**, 091303 (2009).
- [82] P. McDonald and U. Seljak (2008).

- [83] E. Komatsu and D.N. Spergel, *Phys. Rev. D* **63**, 063002 (2001).
- [84] M. Grossi, E. Branchini, K. Dolag, S. Matarrese, and L. Moscardini, *Mon. Not. R. Astron. Soc.* **390**, 438 (2008).
- [85] A. Taruya, K. Koyama, and T. Matsubara, *Phys. Rev. D* **78**, 123534 (2008).
- [86] X. Kang, P. Norberg, and J. Silk, *Mon. Not. R. Astron. Soc.* **376**, 343 (2007).
- [87] R.K. Sheth and G. Tormen, *Mon. Not. R. Astron. Soc.* **329**, 61 (2002).
- [88] M. Maggiore and A. Riotto (2009).
- [89] Z. Lukić, D. Reed, S. Habib, and K. Heitmann, *Astrophys. J.* **692**, 217 (2009).
- [90] C.T. Byrnes, M. Sasaki, and D. Wands, *Phys. Rev. D* **74**, 123519 (2006).
- [91] L. Boubekeur and D.H. Lyth, *Phys. Rev. D* **73**, 021301 (2006).
- [92] M.-X. Huang and G. Shiu, *Phys. Rev. D* **74**, 121301 (2006).