

Cosmological constant and local gravity

José Bernabéu

Departamento de Física Teórica and IFIC, Universidad de Valencia-CSIC, E-46100, Burjassot (Valencia), Spain, and CERN, Theory Division, CH-1211 Geneva 23, Switzerland

Catalina Espinoza

Departamento de Física Teórica and IFIC, Universidad de Valencia-CSIC, E-46100, Burjassot (Valencia), Spain

Nick E. Mavromatos

King's College London, University of London, Department of Physics, Strand WC2R 2LS, London, United Kingdom
(Received 15 November 2009; published 1 April 2010)

We discuss the linearization of Einstein equations in the presence of a cosmological constant, by expanding the solution for the metric around a flat Minkowski space-time. We demonstrate that one can find consistent solutions to the linearized set of equations for the metric perturbations, in the Lorentz gauge, which are not spherically symmetric, but they rather exhibit a cylindrical symmetry. We find that the components of the gravitational field satisfying the appropriate Poisson equations have the property of ensuring that a scalar potential can be constructed, in which both contributions, from ordinary matter and $\Lambda > 0$, are attractive. In addition, there is a novel tensor potential, induced by the pressure density, in which the effect of the cosmological constant is repulsive. We also linearize the Schwarzschild–de Sitter exact solution of Einstein's equations (due to a generalization of Birkhoff's theorem) in the domain between the two horizons. We manage to transform it first to a gauge in which the 3-space metric is conformally flat and, then, make an additional coordinate transformation leading to the Lorentz gauge conditions. We compare our non-spherically symmetric solution with the linearized Schwarzschild–de Sitter metric, when the latter is transformed to the Lorentz gauge, and we find agreement. The resulting metric, however, does not acquire a proper Newtonian form in terms of the unique scalar potential that solves the corresponding Poisson equation. Nevertheless, our solution is stable, in the sense that the physical energy density is positive.

DOI: [10.1103/PhysRevD.81.084002](https://doi.org/10.1103/PhysRevD.81.084002)

PACS numbers: 04.20.Jb, 04.25.Nx, 04.40.Nr

I. INTRODUCTION

De Sitter space has become physically relevant in the last decade, after the extraordinary discovery, based on astrophysical and cosmological observations, that our Universe is accelerating at late eras. A diverse range of results [1], from cosmic microwave background temperature fluctuation measurements to high-redshift supernova measurements, baryon acoustic oscillation measurements and weak lensing techniques (such as cosmic shear and red-shift space distortion), has indicated that a global best fit for cosmology is provided by a simple Friedman-Robertson-Walker Universe with a positive cosmological constant Λ [2], whose dominance over matter at late eras is held responsible for the Universe acceleration. In fact, according to this simple and successful scenario, the best fitting of the data says that over 70% of the current-epoch Universe energy-density budget is dominated by this mysterious form of dark energy, which is compatible with a cosmological constant Λ .

Once Λ is included in the Einstein equations for general relativity, its implications are not limited to cosmological problems. The observable effects of the cosmological constant for orbits in the solar system and double pulsars exist [3], but are too small to be detected. Its effect becomes

more important for extended galaxy clusters through a correction to the Virial relation [4]. In this paper we address the problem of understanding the consistency of the theory, in the presence of Λ , when applied to local gravity and whether the steps of linearizing gravity and going to the Lorentz gauge lead to a sound description of the gravitational field within special relativity, and to a Newtonian form for the metric, in terms of a unique scalar potential. The prevalent view [5] is that the effects of the cosmological constant are equivalent locally, i.e., within the distances of galaxies or galaxy clusters, to those corresponding to a repulsive tidal force, of a conservative nature, being derived from a unique scalar potential Φ_Λ of the form

$$\Phi_\Lambda/c^2 = -\frac{1}{6}\Lambda r^2. \quad (1)$$

The correspondence principle demands that the theory should contain the Newtonian limit for weak classical gravity, with a relative motion of the source much smaller than c and with the material stresses much smaller than the mass-energy density. In absence of Λ , general relativity satisfies this limit. The requirement of the Newtonian limit is made explicit by linearizing the theory for weak gravity, expanding around the flat Minkowski space-time

$\eta_{\mu\nu} = (1, -1, -1, -1)$, writing in Cartesian components

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (2)$$

and assuming that both $h_{\mu\nu}$ and the derivatives are small with respect to unity, so only first-order terms are kept. With the transformation to the Lorentz gauge and the dominance of the mass-energy density for the gravity source, one derives the “*Newtonian metric*” for the gravitational field

$$ds^2 = (1 + 2\Phi/c^2)c^2 dt^2 - (1 - 2\Phi/c^2)d\vec{x}^2, \quad (3)$$

where Φ is the unique scalar potential, obtained as a solution of the Poisson equation with the appropriately linearized source. One should notice that the 3-space metric in (3) is *conformally flat*. If these conditions are met, the constant κ of general relativity in the Einstein equations, $R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -\kappa T_{\mu\nu}$, describing the coupling of the source to the gravitational field, is identified with Newton constant G as

$$\kappa(\Lambda = 0) = 8\pi G/c^4. \quad (4)$$

In using (3) and (4) for the geodesics of a test particle in nonrelativistic motion, the result is Newton equation of motion with a force $\vec{F} = -\vec{\nabla}\Phi$. In that nonrelativistic limit, only the h_{00} component of the gravitational field is operative. Equation (3), however, is more general and it applies, for example, to the problem of light bending within special relativity.

How is this picture modified, if any, by the presence of the (positive) cosmological constant $\Lambda \neq 0$? It is quite frequent to see in the literature that the above results are preserved and, then, one adds the Λ -term *a posteriori*. We emphasize, however, that the conditions to arrive to (3) are *not guaranteed* in the presence of Λ and, in particular, that the theory contains two fundamental constants: κ and Λ . Without the Newtonian limit, the identification (4) finds no justification.

The language of a gravitational field necessitates the introduction of a space-time background metric. For the purpose of this work, the background will still be Minkowski’s, so that weak gravity means first order in both κ and Λ . In this view, Λ is sitting in the right-hand side of Einstein equations, with the meaning of an unavoidable gravity source as “dark energy.” We first find in Sec. II the new solutions for linearized gravity assuming that the conditions for the choice of coordinates in the Lorentz gauge are satisfied. We shall see that we have to depart from strict spherical symmetry, finding some components of the field with cylindrical symmetry. In Sec. III we linearize the Schwarzschild–de Sitter solution and proceed, first, to transform it to a gauge in which the 3-space metric is conformally flat and, then, make an additional coordinate transformation leading to the Lorentz gauge conditions. We compare the solution found in Sec. II with that for the linearized Schwarzschild–de Sitter metric

[6] when the latter is appropriately transformed to the Lorentz gauge. We investigate whether the resulting metric is Newtonian, i.e., with the structure shown in (3) in terms of a unique scalar potential. In the physically meaningful coordinate frame where the metric of 3-space is conformally flat, we examine effects to first order in both Λ and m , where m is the mass of a massive celestial object, on the photon orbit, thus demonstrating the existence of nontrivial effects of the cosmological constant on it. These findings are relevant for the ongoing debate in the literature about the rôle of Λ on the deflection of light. Finally, Sec. IV presents a discussion of the results and some open issues.

II. LINEARIZED GRAVITY IN THE PRESENCE OF A COSMOLOGICAL CONSTANT

The Newtonian gravitational field is described by a single scalar potential $\Phi(\vec{x})$ which is a solution of the Poisson equation, with a matter source given by the mass density ρ_m

$$\nabla^2\Phi(\vec{x}) = 4\pi G\rho_m. \quad (5)$$

The coupling of the source to the gravitational field is determined by the Newton constant G .

Einstein equations, derived from the Einstein-Hilbert four-dimensional action, read

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = -\kappa T_{\mu\nu}, \quad (6)$$

where $\Lambda > 0$ is the cosmological constant, which has the dimensions of a curvature, namely $[\text{length}]^{-2}$, $T_{\mu\nu}$ is the stress-energy tensor of matter in the gravitational field generated by the metric tensor $g_{\mu\nu}$ and κ is a dimensionful constant defining the coupling of gravity to matter.

Clearly Eqs. (6) do *not* permit a flat spacetime in the absence of matter sources, when they reduce to

$$R_{\mu\nu} = \Lambda g_{\mu\nu} \quad (7)$$

which implies curvature. In spite of this, we still insist on the possibility of connecting (6) with a linearized theory around Minkowski flat space-time, within special relativity, using the expansion (2): $h_{\mu\nu}$ would be the metric perturbation, a tensor under global Lorentz coordinate transformations, This approach necessarily implies that the cosmological constant Λ is *not* a mere geometric effect but has to be put on the right-hand side of (6) and interpreted as a form of *dark energy*, i.e., as an additional source of gravity beyond matter. The consistency of the linearized theory requires the limit of *weak gravity*, with the source being of first order, as is the field $h_{\mu\nu}$ and its derivatives. As a consequence, we have to stay in first order in both κ and Λ , without any back-reaction of the gravitational field on $T_{\mu\nu}$ and $g_{\mu\nu}$ in the right-hand side of Eq. (6).

Upon defining the trace-reversed version of the h ’s as

$$\tilde{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h, \quad (8)$$

it follows that:

$$h \equiv h^\alpha{}_\alpha = -\tilde{h}^\mu{}_\mu \equiv -\tilde{h} \quad h_{\mu\nu} = \tilde{h}_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\tilde{h}, \quad (9)$$

where it is understood the indices are raised and lowered with the Minkowski metric. One observes that the Ricci tensor $R_{\mu\nu}$ can be written as

$$R_{\mu\nu} = \frac{1}{2}(\square h_{\mu\nu} - h_{,\mu\nu} - h^\lambda{}_{,\nu\lambda} - h^\lambda{}_{,\mu\lambda}) \quad (10)$$

to leading order in the field and its derivatives, with $\square \equiv \eta^{\lambda\rho}\partial_\lambda\partial_\rho$ denoting the D'Alembertian operator.

A suitable coordinate transformation can get rid of the last two terms of Eq. (10) if we choose the following condition

$$\tilde{h}^\mu{}_{,\nu,\mu} = 0. \quad (11)$$

This is called the *Lorentz gauge*, in analogy with electromagnetism.

In the Lorentz gauge, the Einstein tensor $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$ becomes

$$G_{\mu\nu} = \frac{1}{2}\square\tilde{h}_{\mu\nu}, \quad (12)$$

so that the full field equations (6) now reduce to

$$\tilde{h}_{\mu\nu} = \begin{pmatrix} -\frac{4m}{r} + \frac{1}{3}\Lambda r^2 & 0 & 0 & 0 \\ 0 & -\frac{1}{2}\Lambda(y^2 + z^2) & 0 & 0 \\ 0 & 0 & -\frac{1}{2}\Lambda(z^2 + x^2) & 0 \\ 0 & 0 & 0 & -\frac{1}{2}\Lambda(x^2 + y^2) \end{pmatrix} \quad (15)$$

We observe that these components of the field, each of them solution of Poisson's equations, satisfy the following:

- (1) The scalar potential $\Phi = \frac{1}{4}c^2\tilde{h}_{00}$ gets modified by an *attractive effect* induced by a *positive energy density* due to Λ , besides that generated by (14) due to ordinary matter;
- (2) there is a *novel tensor potential* $\tau_{ij} = \frac{1}{4}c^2\tilde{h}_{ij}$ with a *repulsive effect* induced by the *negative pressure density* due to Λ ;
- (3) whereas the scalar potential presents a rotational symmetry, the components of the tensor potential have a *cylindrical symmetry* around the corresponding principal axis. The cylindrical symmetry is, in fact, the same for the three principal axes, a remnant

$$h_{\mu\nu} = \begin{pmatrix} 2\Phi - 2\tau & 0 & 0 & 0 \\ 0 & 2\Phi + 2\tau + 4\tau_{11} & 0 & 0 \\ 0 & 0 & 2\Phi + 2\tau + 4\tau_{22} & 0 \\ 0 & 0 & 0 & 2\Phi + 2\tau + 4\tau_{33} \end{pmatrix} \quad (17)$$

with the property $h = -\tilde{h}$ for the complete trace, as required. Equation (17) gives the solution for the metric within special relativity. It is neither conformally flat nor of the form we have for $\Lambda = 0$, i.e. $g_{00} = 1 + 2\Phi/c^2$, $g_{ii} = -1 + 2\Phi/c^2$ (no sum over i), $i = 1, 2, 3$. The con-

$$\square\tilde{h}_{\mu\nu} = -2\kappa T_{\mu\nu} - 2\Lambda\eta_{\mu\nu} \quad (13)$$

Equations (13), plus the gauge condition (11), constitute the basic ingredients for our study. These linear field equations are also decoupled when they are written in terms of $\tilde{h}_{\mu\nu}$. As a consequence, the trace-reversed $\tilde{h}_{\mu\nu}$ has the meaning of the gravitational field and Eq. (13) is the relativistic generalization of the Newtonian gravity equation (5).

We are interested in the solution for the linearized metric, in the Lorentz gauge, for a weak static field generated by a point mass M located at $r = 0$: all components of $T_{\mu\nu}$ vanish except

$$T_{00} = Mc^2\delta(\vec{x}) \quad (14)$$

In terms of $m \equiv \kappa Mc^2/8\pi$, we find the static solutions to (13) and (11) in diagonal form in the presence of Λ . Each diagonal component $\tilde{h}_{\mu\mu}$ (no sum over μ) is independent of the specific coordinate x^μ . This condition implies that the space components of the gravitational field \tilde{h}_{ii} (no sum over i), if they exist, have to depart from spherical symmetry. The solution is

of the rotational symmetry. This *breaking* of the rotational symmetry is an *artifact* of the gauge fixing imposed by the Lorentz condition.

From Eq. (15) we find the complete trace of the field and the 3-space trace of the tensor potential as

$$\tilde{h}^\alpha{}_\alpha \equiv \tilde{h} = -\frac{4m}{r} + \frac{4}{3}\Lambda r^2 \quad \tau^i{}_i \equiv \tau = +\frac{1}{4}c^2\Lambda r^2. \quad (16)$$

From these results it becomes already apparent that the metric is *not Newtonian* when $\Lambda \neq 0$. In going from the trace-reversed gravitational field $\tilde{h}_{\mu\nu}$ to the metric perturbation in the Lorentz gauge, we find

tribution of the modified scalar potential Φ to the metric (17) is still as in the Newtonian form (3), but all the diagonal components get additional contributions from the tensor potential. We realize, in particular, that the h_{00} component acquires an *effective Λ -term*, which originates

from both the attractive scalar potential and the trace of the repulsive tensor potential: its net effect is repulsive and given by (1). One should notice that, whereas

$$h_{00} = -2\frac{m}{r} - \frac{1}{3}\Lambda r^2, \quad (18)$$

is the only component of the metric entering the geodesic equation for the motion of a nonrelativistic body, given by $\frac{d^2x^i}{dt^2} \simeq -c^2\Gamma_{00}^i$, where $\Gamma_{\mu\nu}^\rho$ is the Christoffel symbol, the 3-space components of the metric

$$h_{ii} = -2\frac{m}{r} + \frac{1}{6}\Lambda(r^2 + 3x_i^2), \quad (19)$$

$i = 1, 2, 3$ (no sum over i)

are also intervening for a test-body in relativistic motion. For the case of light bending, we shall discuss the effects of the cosmological constant $\Lambda > 0$ on the photon orbit in Sec. III. We note at this point that within the Schwarzschild–de Sitter metric, the magnitude of the effects of the cosmological constant $\Lambda > 0$ on the bending of light from distant galaxies is still an open issue. At present, there are several approaches to the subject, giving different answers, in which the order of magnitude of the effects ranges from zero [7] or unobservably small [8] to appreciable one [9,10].

Our considerations in this paper pertain to local effects of the cosmological constant, which are discussed within the special relativity framework. As we show in the next Sec. III, our linearized solution (17) can be obtained from the Schwarzschild–de Sitter solution upon appropriate coordinate transformations.

III. THE LINEARIZED SCHWARZSCHILD–DE SITTER SOLUTION

At this point one should investigate the connection of the linearized gravity solution found in Sec. II with the Schwarzschild–de Sitter metric. We do know that, under the conditions we have imposed for the source of coupling with gravity, there is a theorem analogous to *Birkhoff's* for the Schwarzschild metric, stating [6] that there is a unique static solution with spherical symmetry of the form¹:

¹According to the analysis in [6], if one relaxes the assumption of staticity, the spherical symmetry implies two types of solutions in the case of positive $\Lambda > 0$: (i) a *static* solution (21), expressing the field around a spherically symmetric mass, where, as in the Schwarzschild case, the mass is an integration constant of the Einstein equations; (ii) a *nonstatic*, time-dependent solution, distinct from that due to the field around a spherical distribution of masses, which consists of successive identical spheres, that is, a *cylindrical* Universe of Bertotti-Kasner type:

$$ds^2 = dt^2 - e^{2\sqrt{\Lambda}t} dr^2 - \frac{1}{\Lambda}(d\theta^2 + \sin^2\theta d\phi^2), \quad (20)$$

In view of the last term, the reader should understand now why this solution exists only in the case of positive Λ .

$$ds^2 = \left(1 - 2\frac{m}{\bar{r}} - \Lambda\bar{r}^2/3\right)c^2 dt^2 - \left(1 - 2\frac{m}{\bar{r}} - \Lambda\bar{r}^2/3\right)^{-1} \times d\bar{r}^2 - \bar{r}^2(d\bar{\theta}^2 + \sin^2\bar{\theta}d\bar{\phi}^2). \quad (21)$$

This is the Schwarzschild–de Sitter metric, in which the Schwarzschild space coordinates define \bar{r} as the “*area distance*,” i.e., the distance for which the surface is given by the Euclidean measure $4\pi\bar{r}^2$. The presence of horizons at the approximate values $\bar{r} = 2m$ and $\bar{r} = \sqrt{3/\Lambda}$ should be noted along with the fact that the observer is actually required to live in the space between them, for our *weak-gravity* analysis to be valid. Indeed, in the domain

$$2m \ll \bar{r} \ll \sqrt{3/\Lambda} \quad (22)$$

we can linearize the components of the metric (21) around Minkowski background space-time. Even for the case $\Lambda = 0$ and *a fortiori* for $\Lambda \neq 0$, the result for the metric (21) is neither conformally flat nor of the Newtonian form (3) with a single scalar potential. We notice, in particular, that the lack of a conformally flat metric in these Schwarzschild coordinates implies that the metric is not diagonal when written in Cartesian components, even if it is so in spherical components. One can check that the corresponding linearized metric does *not* satisfy the *Lorentz gauge* condition.

At this point it is natural to ask whether it is possible to reach the Lorentz gauge by a coordinate transformation. To this end, we proceed in two steps:

(i) First, we move from the Schwarzschild coordinates to fully symmetric spherical coordinates leading to a conformally flat metric in 3-space. This transformation exists because, with $\Lambda \neq 0$, we are in a case of *constant curvature*. With the transformation

$$\bar{r} \rightarrow r' = \bar{r} \left(1 - \frac{m}{\bar{r}} + \frac{1}{12}\Lambda\bar{r}^2\right) \quad (23)$$

the resulting metric becomes

$$ds^2 = \left(1 - 2\frac{m}{r'} - \frac{1}{3}\Lambda r'^2\right)c^2 dt^2 - \left(1 + 2\frac{m}{r'} - \frac{1}{6}\Lambda r'^2\right)d\vec{x}'^2 \quad (24)$$

In Eq. (24), both Cartesian and spherical components of the 3-space metric are diagonal. In spite of this property, the metric is not of the form of Eq. (3) in terms of the r' coordinate, i.e. $(1 + 2\Phi'(r'))$ vs $(1 - 2\Phi'(r'))$, if $\Lambda \neq 0$. One can check, in fact, that the *Lorentz gauge* condition is *not* satisfied for the metric (24) in the presence of a nonzero cosmological constant, $\Lambda \neq 0$, which is to be contrasted with the $\Lambda = 0$ case, for which the two gauges coincide. The trace of the linearized metric perturbation is given by

$$h' = 4\frac{m}{r'} - \frac{5}{6}\Lambda r'^2 \quad (25)$$

so that the trace-reversed gravitational field (8) is given, in these coordinates, by

$$\tilde{h}'_{\mu\nu} = \begin{pmatrix} -4\frac{m}{r'} + \frac{1}{12}\Lambda r'^2 & 0 & 0 & 0 \\ 0 & -\frac{1}{4}\Lambda r'^2 & 0 & 0 \\ 0 & 0 & -\frac{1}{4}\Lambda r'^2 & 0 \\ 0 & 0 & 0 & -\frac{1}{4}\Lambda r'^2 \end{pmatrix} \quad (26)$$

Notice that, for $\Lambda = 0$, this field would be described by a unique scalar potential solution of the Laplace equation for $r' > 0$. However, for $\Lambda \neq 0$, the field components do not satisfy the Lorentz gauge condition for the coordinates (x', y', z') .

In this fully symmetric gauge, the scalar Φ' and tensor $\tau'_{ij} = \tau' \eta_{ij}/3$ potentials, with τ' the corresponding 3-space trace, are given by

$$\Phi'/c^2 = -\frac{m}{r'} + \frac{1}{48}\Lambda r'^2 \quad \tau'/c^2 = +\frac{3}{16}\Lambda r'^2. \quad (27)$$

They satisfy the following coupled Equations in terms of the sources

$$\nabla'^2(\Phi' + \frac{1}{3}\tau') = \frac{1}{2}\kappa c^2 T_{00} + \frac{1}{2}\Lambda c^2, \quad \nabla'^2 \tau' = \frac{9}{8}\Lambda c^2. \quad (28)$$

The corresponding 3-space conformally flat metric (24) can then be written in the simple form

$$ds^2 = \left(1 + \frac{2(\Phi' - \tau')}{c^2}\right) c^2 dt^2 - \left(1 - \frac{2\Phi'}{c^2} - \frac{2\tau'}{3c^2}\right) d\vec{x}'^2. \quad (29)$$

This coordinate frame is the appropriate one for measurements using standard clocks and rods, due to the fully isotropic 3-space metric components. In the case of photons, the null geodesics, derived from (24), can be easily constructed, leading to a direction-independent speed of light, a feature that is not valid on other coordinate frames.. The geodesics encompass, as usual, the two first integrals of motion related to the photon energy and angular momentum. The orbit $r'(\phi, \theta = \frac{\pi}{2})$ can be expressed as an equation for the function $u(\phi) \equiv \frac{R}{r'}$, which to first order in m and Λ reads:

$$\left(\frac{du}{d\phi}\right)^2 + u^2 - 1 = \frac{4m}{R}(u - 1) + \frac{1}{6}\Lambda R^2\left(\frac{1}{u^2} - 1\right), \quad (30)$$

where R is the radius of the spherical mass distribution and we consider a path of the photon that grazes its surface at $\phi = \frac{\pi}{2}$. A symmetric solution about the axis $\phi = \pi/2$, as required by the geometry, can be found analytically around $\phi \simeq 0$ and $\phi \simeq \pi$, to leading order in $\sin\phi \ll 1$ for the first-order terms proportional to m and Λ :

$$u(\phi) \simeq \sin\phi + \frac{2m}{R} - \frac{1}{12}\frac{\Lambda R^2}{\sin\phi}. \quad (31)$$

For the case $\Lambda = 0$, there is an asymptotic $u \rightarrow 0$ solution, leading to the one-sided bending angle $\phi_\infty = -\frac{2m}{R}$, thus reproducing the standard Einstein's result in general relativity.

However, in the presence of the Λ -term, there is no asymptotic limit, due to the de-Sitter horizon. As we observe from Eq. (31), there are nontrivial effects of order Λ to the photon orbit in our 3-space conformally flat coordinate system, which come with opposite sign to the $\Lambda = 0$ General Relativity contribution. Indeed, the one-sided bending angle $\beta_\Lambda(r')$ obtained from (31), with $\Lambda > 0$, is:

$$\beta_\Lambda(r') \simeq -\frac{2m}{R} + \frac{1}{12}\Lambda R r', \quad r' \gg R. \quad (32)$$

Given that a conformal transformation of the 3-space coordinates preserves the angles, the above result would also be valid in flat space-time, which is the background space-time in our linearized gravity approximation. On the other hand, as shown in Ref. [11], if one writes the orbit of the photon in the original (nonconformal) Schwarzschild coordinates, the orbit appears to be independent of Λ .

Our conclusion that Λ is contributing through Eq. (32) to the deflection of light by a spherical mass distribution is in line with the claims [8–10] that the effect exists, in spite of the Λ -independent equation in Schwarzschild coordinates. A more detailed analysis of this and other observational effects of a gravitating Λ will be presented elsewhere.

(ii) As a second step, we make the coordinate transformation

$$\begin{aligned} x' \rightarrow x &= x' + \frac{1}{12}\Lambda x'^3 & y' \rightarrow y &= y' + \frac{1}{12}\Lambda y'^3 \\ z' \rightarrow z &= z' + \frac{1}{12}\Lambda z'^3, \end{aligned} \quad (33)$$

which definitively leads to the Cartesian components of the field (15) in the coordinates (ct, x, y, z) . These ones are the coordinates associated with the Lorentz gauge and the discussion after Eq. (15) follows. We conclude that the solution found in Section II is precisely the linearized Schwarzschild–de Sitter metric, written in a new set of appropriate coordinates that correspond to the Lorentz gauge.

IV. CONCLUSIONS

In this work we have discussed a first-order solution to the Einstein equations in the presence of a small positive cosmological constant $\Lambda > 0$, in the case where the equations are linearized about flat Minkowski space-time. Usually, in the literature, this problem is associated straightforwardly with the presence of a repulsive force in the ‘‘Newtonian limit.’’ As we emphasized in this note, however, since there is no proper Newtonian metric in the case $\Lambda \neq 0$, the linearization procedure has to be applied with care, and the identification of what plays the rôle of a repulsive ‘‘potential’’ presents subtleties. In particular, we have found that in this case there are scalar and tensor

potentials, which *both* describe the gravitational field in special relativity in the presence of a nontrivial (no matter how small) cosmological constant. As we have discussed, with $\Lambda \neq 0$ there is *no Newtonian form* for the metric in any gauge. The Lorentz gauge, for $\Lambda \neq 0$, does not lead to a 3-space conformally flat metric.

If one insists on the requirement of a Newtonian form for the metric, in agreement with the correspondence principle, then one has to conclude that Λ cannot be a *classical geometric effect*, but rather a *relativistic quantum effect*, which should vanish (formally) for $\hbar \rightarrow 0$. From dimensional arguments, by writing $\Lambda = \kappa \rho_\Lambda$, the dark energy density ρ_Λ is given by

$$\rho_\Lambda \sim \hbar c / \ell^4 \quad (34)$$

where ℓ should be a characteristic length, whose microscopic origin, and hence its order of magnitude, is a mystery. In this case, the relation (4) of κ with Newton constant G would still be valid for $\Lambda \neq 0$.

Our discussion in this note has been concentrated solely on local effects of the cosmological constant. At this point we feel that we should contrast our findings with the traditional point of view adopted in the literature, in which a “Newtonian scalar potential” is constructed from the h_{00} component of the metric perturbation by looking only at the nonrelativistic geodesics, i.e. from Eq. (18). Using the metric for the Schwarzschild–de Sitter solution (21) [5,6], in the limit of $m \rightarrow 0$, or equivalently in the case where the Λ term dominates over the mass term, it is often stated that the resulting “energy density” appears *negative*, thereby indicating an instability. Indeed, this line of thinking prompted the elevation of the Schwarzschild–de Sitter solution from a local to a global one, relevant for an expanding Universe, where this instability is remedied, in the sense that the cosmological de Sitter solution has still positive energy, but negative pressure.

However, as our analysis in this article has shown [cf. discussion following Eq. (15)], this “instability” arises from a misinterpretation of what energy density means in the linearized solution. In our case, precisely due to the absence of a proper Newtonian form of the metric, the correct identification of the energy density can only be made via the Poisson equation for the gravitational field $\tilde{h}_{\mu\nu}$, Eq. (13). From this point of view, our solution appears *stable*. The corresponding dark energy density is indeed proportional to $\Lambda > 0$, and originates from an attractive contribution to the scalar potential defined via \tilde{h}_{00} in (15), even in the case $m \ll \Lambda$ [i.e. when $|\kappa T_{\mu\nu}| \ll |\Lambda \eta_{\mu\nu}|$ in (13)], and hence is positive.

A final comment before closure. The cosmological de Sitter solution, in the context of a Friedman–Robertson–Walker universe, is an entirely different problem. The precise connection from the global solution to the local one, by means of space-dependent cosmological perturbations, is a complicated issue, which is still unsolved. One might hope that, by perturbing the Λ -de Sitter–FRW appropriately, it would be possible to make a connection with the local gravity case and thus compare the results with the analysis presented here.

ACKNOWLEDGMENTS

We wish to thank D. Espriu, V. A. Mitsou, M. Quirós, R. Rebolo, and M. Sereno for interesting discussions. We also acknowledge helpful suggestions and comments by the anonymous referee. The research work of J.B. and C.E. is supported in part by the Grants of the Spanish Ministry of Science and Innovation FPA 2008-02878 and of the Valencia Generalitat PROMETEO 2008/004, while that of N.E.M. is partly supported by the European Union, through the FP6 Marie Curie Research and Training Network *UniverseNet* (MRTN-CT-2006-035863).

-
- [1] A. G. Riess *et al.* (Supernova Search Team Collaboration), *Astron. J.* **116**, 1009 (1998); S. Perlmutter *et al.* (Supernova Cosmology Project Collaboration), *Astrophys. J.* **517**, 565 (1999); A. G. Riess, *Publ. Astron. Soc. Pac.* **112**, 1284 (2000); E. Linder and S. Perlmutter, *Phys. World* **20N12**, 24 (2007); M. Kowalski *et al.* (Supernova Cosmology Project Collaboration), *Astrophys. J.* **686**, 749 (2008); W. M. Wood-Vasey *et al.* (ESSENCE Collaboration), *Astrophys. J.* **666**, 694 (2007); A. G. Riess and M. Livio, *Astrophys. J.* **648**, 884 (2006); A. G. Riess *et al.* (Supernova Search Team Collaboration), *Astrophys. J.* **607**, 665 (2004); D. N. Spergel *et al.* (WMAP Collaboration), *Astrophys. J. Suppl. Ser.* **170**, 377 (2007); D. J. Eisenstein *et al.* (SDSS Collaboration), *Astrophys. J.* **633**, 560 (2005); H. J. Seo and D. J. Eisenstein, *Astrophys. J.* **598**, 720 (2003), and references therein; L. Fu *et al.*, *Astron. Astrophys.* **479**, 9 (2008); L. Guzzo *et al.*, *Nature (London)* **451**, 541 (2008).
- [2] See, for instance: S. Weinberg, *Cosmology* (Oxford University Press, Oxford, England, 2008).
- [3] M. Sereno and P. Jetzer, *Phys. Rev. D* **73**, 063004 (2006); A. Balaguera-Antolinez, C. G. Boehmer, and M. Nowakowski, *Classical Quantum Gravity* **23**, 485 (2006); L. Iorio, *Adv. Astron. Astrophys.*, doi: 10.1155/2008/268647 (2008); *New Astron. Rev.* **14**, 196 (2009).
- [4] Y. Suto, *Prog. Theor. Phys.* **90**, 1173 (1993); P. J. Peebles, *Astrophys. Space Sci.* **45**, 3 (1976); *The Large Scale Structure of The Universe* (Princeton University Press, New Jersey, 1980); G. S. Adkins, J. McDonnell, and R. N. Fell, *Phys. Rev. D* **75**, 064011 (2007).

- [5] See, for instance: W. Rindler, *Relativity: Special, General, and Cosmological* (Oxford University Press, New York, 2006), 2nd ed., ISBN-13: 978-0-19-856731-8.
- [6] W. Rindler, *Phys. Lett. A* **245**, 363 (1998); the spherically-symmetric Schwarzschild–de Sitter solution was first discussed in F. Kottler, *Ann. Phys. (Leipzig)* **361**, 401 (1918).
- [7] I. B. Khriplovich and A. A. Pomeransky, *Int. J. Mod. Phys. D* **17**, 2255 (2008); M. Park, *Phys. Rev. D* **78**, 023014 (2008); F. Simpson, J. A. Peacock, and A. F. Heavens, arXiv:0809.1819.
- [8] M. Sereno, *Phys. Rev. Lett.* **102**, 021301 (2009); *Phys. Rev. D* **77**, 043004 (2008); See also: R. Kantowski, B. Chen, and X. Dai, arXiv:0909.3308.
- [9] W. Rindler and M. Ishak, *Phys. Rev. D* **76**, 043006 (2007); M. Ishak, W. Rindler, and J. Dossett, arXiv:0810.4956; G. W. Gibbons, C. M. Warnick, and M. C. Werner, *Classical Quantum Gravity* **25**, 245009 (2008); T. Schucker, arXiv:0903.2940.
- [10] M. Ishak, W. Rindler, J. Dossett, J. Moldenhauer, and C. Allison, *Mon. Not. R. Astron. Soc.* **388**, 1279 (2008).
- [11] N. J. Islam, *Phys. Lett. A* **97A**, 239 (1983).