

Motion of pole-dipole and quadrupole particles in nonminimally coupled theories of gravity

Morteza Mohseni*

Physics Department, Payame Noor University, 19395-4697 Tehran, Iran

(Received 12 March 2010; published 22 June 2010)

We study theories of gravity with nonminimal coupling between polarized media with pole-dipole and quadrupole moments and an arbitrary function of the space-time curvature scalar R and the squares of the Ricci and Riemann curvature tensors. We obtain the general form of the equation of motion and show that an induced quadrupole moment emerges as a result of the curvature tensor dependence of the function coupled to the matter. We derive the explicit forms of the equations of motion in the particular cases of coupling to a function of the curvature scalar alone, coupling to an arbitrary function of the square of the Riemann curvature tensor, and coupling to an arbitrary function of the Gauss-Bonnet invariant. We show that in these cases the extra force resulting from the nonminimal coupling can be expressed in terms of the induced moments.

DOI: 10.1103/PhysRevD.81.124039

PACS numbers: 04.20.Cv, 04.20.Fy, 04.40.-b

I. INTRODUCTION

Various extensions of the general theory of relativity have been proposed in recent years to explain astrophysical observations like the accelerated expansion of the Universe and rotation curves of spiral galaxies. A widely known class of such proposals comprises variants of the so-called $f(R)$ theories of gravity; see [1–3] for reviews. Even these theories have been themselves subject to different modifications in the hope of obtaining a more satisfactory picture, namely, by considering nonminimal couplings with matter [4–7]. In the models investigated in these references, some powers of the curvature scalar R have been coupled to the matter Lagrangian which have resulted in modified equations of motion. A more general case was considered in Ref. [8] in which an arbitrary function of the curvature scalar nonminimally coupled with the matter Lagrangian. Taking perfect fluids and scalar fields as matter fields, several implications of such a coupling were subsequently studied in Refs. [9–17]. The effects of such a coupling on the dynamics of test particles have been studied in Ref. [18], where it has been shown that an extra force emerges from the nonminimal coupling. The explicit form of the force depends on the form of the matter Lagrangian, and it has been shown in Refs. [19,20] that, for the case of the perfect fluid for which there are several equivalent Lagrangians, one may end up with vanishing or nonvanishing extra forces depending on the chosen form of the Lagrangian. In Ref. [21], it has been argued that the Lagrangians which are equivalent for a noncoupled perfect fluid are not in fact equivalent when the perfect fluid couples nonminimally to curvature. The author of Ref. [22] has argued that the form of the coupling fixes the matter Lagrangian.

The extra force affects the world lines of test particles. In fact within the framework of models with nonminimal

coupling, test particles move along nongeodesic trajectories in general [18]. Taking the effect of extra force into account, the motion of test pole-dipole particles, i.e. test bodies with certain microstructures, has been studied in Ref. [23] by using the method of multipoles [24,25]. In this work we consider the motion of pole-dipole and quadrupole particles in extended gravity models with more general coupling to matter. To this end we consider a class of matter fields with a general dependence on a tetrad field, its covariant derivative, a velocity field, curvature tensors, and a generic set of matter fields, coupled to $f(R)$ gravity action. In particular, we show that when the Lagrangian depends on curvature tensor through nonminimal coupling to geometry, induced quadrupole moments emerge. We obtain the equations of motion for media with pole-dipole and quadrupole moments nonminimally coupled to an arbitrary function of the curvature scalar, and as a special case we consider a Weyssenhoff-type spinning fluid from which we extract the equations of motion of pole-dipole particles. The method can then be easily generalized to particles with electric charges. We also investigate the motion of test bodies nonminimally coupled to an arbitrary function of the square of the Riemann curvature tensor or to an arbitrary function of the Gauss-Bonnet invariant and show that the extra forces can be expressed in terms of the induced quadrupole moments. The problem of induced quadrupole moments in media due to an applied gravitational field has been studied in Refs. [26–30] in the context of general relativity. The study of motion of quadrupole particles is well motivated by their role in gravitational radiation. Several examples of motion of quadrupole particles in curved space-times of general relativity may be found in Refs. [31–35]. The effect of multipole moments of planets on bending of light rays has been discussed in Ref. [36], and it has been suggested in Refs. [37,38] that such effects may be detected by using microarcsecond astronomical interferometers.

*m-mohseni@pnu.ac.ir

In the subsequent sections, we first give a brief review of the polarized media as formulated in Refs. [39,40] and then deploy it to obtain the equations of motion of polarized media with pole-dipole and quadrupole moments nonminimally coupled to an arbitrary function constructed out of the curvature scalar and the squares of the Ricci and Riemann curvature tensors. We subsequently consider three particular cases: the case of nonminimal $f(R)$ gravity, the case of coupling with an arbitrary function of the Riemann curvature tensor squared, and the case of coupling to an arbitrary function of the Gauss-Bonnet invariant. In the first case, our investigation results in equations of motion of particles with pole-dipole and quadrupole moments in nonminimal $f(R)$ gravity. In the second and third cases in which for simplicity no intrinsic moments (i.e. those originating from the structure of the body) are assumed, we show that the extra force of the nonminimal coupling can be expressed in terms of the induced moments. In the last section we discuss the results. Throughout the work we use the following (anti)symmetrization conventions: $A^{\cdots(\mu\nu)\cdots} \equiv A^{\cdots\mu\nu\cdots} + A^{\cdots\nu\mu\cdots}$, $A^{\cdots(\mu\cdots B^{\cdots\nu)\cdots} \equiv A^{\cdots\mu\cdots B^{\cdots\nu\cdots} + A^{\cdots\nu\cdots B^{\cdots\mu\cdots}$, and $A^{\cdots[\mu\cdots B^{\cdots\nu]\cdots} \equiv A^{\cdots\mu\cdots B^{\cdots\nu\cdots} - A^{\cdots\nu\cdots B^{\cdots\mu\cdots}$.

II. PARTICLES WITH POLE-DIPOLE AND QUADRUPOLE MOMENTS IN GENERAL RELATIVITY

It has been shown in Refs. [39,40] that, by starting from a Lagrangian with the general form $\mathcal{L}(u^\mu, e_a^\mu, \dot{e}_a^\mu, R^\mu_{\alpha\beta\gamma})$ in which u^μ is a velocity field, e_a^μ is a tetrad field satisfying $e_a^\mu e_b^\nu \eta_{ab} = g_{\mu\nu}$, with $\eta_{ab} = \text{diag}(-1, 1, 1, 1)$, $\dot{e}_a^\mu = u^\alpha \nabla_\alpha e_a^\mu$, and $R^\mu_{\alpha\beta\gamma} = \partial_\alpha \Gamma_{\beta\nu}^\mu - \partial_\beta \Gamma_{\alpha\nu}^\mu + \Gamma_{\alpha\delta}^\mu \Gamma_{\beta\nu}^\delta - \Gamma_{\beta\delta}^\mu \Gamma_{\alpha\nu}^\delta$ is the Riemann curvature tensor, one can arrive at the following energy-momentum tensor:

$$T_{\mu\nu} = p_\mu u_\nu + \mathcal{P}h_{\mu\nu} - B_{\mu\nu} + A_{\mu\nu} \quad (1)$$

describing a polarized fluid with pole-dipole and quadrupole moments. Here u^μ is the four-velocity of a fluid element, p^μ its four-momentum, \mathcal{P} is the pressure,

$$h_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu, \quad (2)$$

$$B_{\mu\nu} = \frac{1}{2} \nabla_\alpha (s^\alpha_{(\mu} u_{\nu)}) + s_{\mu\nu} u^\alpha, \quad (3)$$

$$A_{\mu\nu} = q_{\mu\alpha\beta\gamma} R_\nu^{\alpha\beta\gamma} - 3q_{\nu\alpha\beta\gamma} R_\mu^{\alpha\beta\gamma} - 2\nabla_\beta \nabla_\alpha q^{\alpha(\mu\nu)\beta}, \quad (4)$$

where $s^{\mu\nu}$ is the fluid spin density and

$$q^\mu_{\alpha\beta\gamma} = \frac{\partial \mathcal{L}}{\partial R^\mu_{\alpha\beta\gamma}} \quad (5)$$

is the quadrupole moment tensor, which is assumed to share the symmetries of the Riemann curvature tensor.

The above energy-momentum tensor is symmetric provided the spin evolution equation

$$\nabla_\alpha (u^\alpha s^{\mu\nu}) = p^\mu u^\nu - p^\nu u^\mu - 4R^{[\mu\alpha\beta\gamma} q^{\nu]}_{\alpha\beta\gamma} \quad (6)$$

holds. The translational equation of motion

$$u^\nu \nabla_\nu p^\mu + p^\mu \nabla_\nu u^\nu + h^{\mu\nu} \nabla_\nu \mathcal{P} + \mathcal{P} \nabla_\nu (u^\mu u^\nu) = -\frac{1}{2} R^\mu_{\nu\alpha\beta} u^\nu s^{\alpha\beta} + q_{\alpha\beta\gamma\delta} \nabla^\mu R^{\alpha\beta\gamma\delta} \quad (7)$$

can be obtained from $\nabla_\nu T^{\mu\nu} = 0$ or by variation of world lines [40]. The so-called Frenkel condition

$$u_\mu s^{\mu\nu} = 0 \quad (8)$$

ensures that spin remains spacelike. Imposing the energy conditions may put further restrictions on the spin tensor. By neglecting the quadrupole moments, the equations of motion of a spinning fluid (see e.g. [41] and references therein) are recovered. By turning the spin off and setting the quadrupole moments equal to zero, the standard perfect fluid energy-momentum tensor is recovered from Eq. (1).

As has been shown in Refs. [39,40], the above procedure can be extended to Lagrangians of more general dependence of the form

$$\mathcal{L} \equiv \mathcal{L}(u^\mu, e_a^\mu, \dot{e}_a^\mu, \phi^A) \quad (9)$$

in which

$$\phi^A = \{R^\mu_{\alpha\beta\gamma}, \nabla_{\mu_1} \cdots \nabla_{\mu_m} R^\mu_{\alpha\beta\gamma}, \varphi^A, \nabla_\alpha \varphi^A\}, \quad (10)$$

with φ^A (A is a collective index) being a generic set of matter fields whose free field Lagrangians are also included in the total Lagrangian. In particular, by taking φ^A to be the electromagnetic field, the equations of motion for charged particles would be obtained. Thus in the special case where particles have only pole-dipole moments, the Mathisson-Papapetrou-Dixon [42] and Dixon-Souriau equations [43] for chargeless and charged particles are recovered, respectively. Multipole moments higher than the quadrupole moment are incorporated via dependence on covariant derivatives of the Riemann curvature tensor.

III. MULTIPOLE PARTICLES IN NONMINIMAL EXTENDED GRAVITY

The action for $f(R)$ gravity with nonminimal coupling to matter is given by [18]

$$S = \int \left(\frac{1}{2} f_1(R) + \{1 + \lambda f_2(R)\} \mathcal{L}_m \right) \sqrt{-g} d^4x, \quad (11)$$

where \mathcal{L}_m represents the matter Lagrangian, λ is a coupling constant, and $f_1(R)$ and $f_2(R)$ are arbitrary functions of the curvature scalar $R = g^{\alpha\gamma} g^{\beta\delta} R_{\alpha\beta\gamma\delta}$. For $\lambda = 0$ this reduces to the usual $f(R)$ gravity. As has been shown in Ref. [18], one can obtain the following relation from the equations of motion associated with the above action:

$$\nabla_\nu T^{\mu\nu} = \frac{\lambda f'_2(R)}{1 + \lambda f_2(R)} (g^{\mu\nu} \mathcal{L}_m - T^{\mu\nu}) \nabla_\nu R, \quad (12)$$

where $f'_2(R) = \frac{df_2(R)}{dR}$ and $T^{\mu\nu}$ is the matter energy-momentum tensor

$$T^{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta(\mathcal{L}_m \sqrt{-g})}{\delta g_{\mu\nu}}. \quad (13)$$

Here we consider a more general extension of the above action with the following form:

$$S = \int \sqrt{-g} \left[\frac{1}{2} R + \frac{1}{2} f + (1 + \lambda F) \mathcal{L} \right] d^4x, \quad (14)$$

where $f \equiv f(R, P, Q)$, $F \equiv F(R, P, Q)$, and $P = R_{\mu\nu} R^{\mu\nu}$, $R^{\mu\nu}$ being the Ricci tensor, $Q = R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta}$ is the Kretschmann invariant, and \mathcal{L} is a matter Lagrangian (see [44] for a cosmological model with this type of action but with $\lambda = 0$). In fact, one may think of $(1 + \lambda F)\mathcal{L}$ as a Lagrangian of the type described by Eq. (9). Various models including minimal or nonminimal $f(R)$ gravity, modified Gauss-Bonnet gravity, the so-called $f(R, \mathcal{G})$ models [45,46], where $\mathcal{G} = R^2 - 4P + Q$ is the Gauss-Bonnet invariant, and the model introduced in Ref. [47] can be recovered as special cases by choosing appropriate forms for f and F . Functions with more general dependence on curvature invariants (e.g. those introduced in Ref. [48]) may also be considered as toy models.

The equations of motion can be obtained by varying the above action with respect to metric components. Defining

$$\begin{aligned} H^{\mu\nu} = & (R^{\mu\nu} - \nabla^\mu \nabla^\nu + g^{\mu\nu} \square) f_R + 2R^{\mu\alpha} R_{\alpha}^{\nu} f_P \\ & - \nabla_\alpha \nabla^\nu (f_P R^{\alpha\mu}) + g^{\mu\nu} \nabla_\alpha \nabla_\beta (f_P R^{\alpha\beta}) \\ & + \square (f_P R^{\mu\nu}) + 2R^{\mu\alpha\beta\gamma} R_{\alpha\beta\gamma}^{\nu} f_Q \\ & - 2\nabla_\alpha \nabla_\beta (f_Q R^{\alpha(\mu\nu)\beta}), \end{aligned}$$

we have

$$G^{\mu\nu} + H^{\mu\nu} - \frac{1}{2} f g^{\mu\nu} = (1 + \lambda F) T^{\mu\nu} - 2\lambda \mathcal{H}^{\mu\nu}, \quad (15)$$

where $\square = \nabla_\mu \nabla^\mu$, $\frac{df}{dR}$, $\frac{df}{dP}$, $\frac{df}{dQ}$, $\frac{dF}{dR}$, $\frac{dF}{dP}$, and $\frac{dF}{dQ}$ stand for f_R , f_P , f_Q , F_R , F_P , and F_Q , respectively, and $\mathcal{H}^{\mu\nu}$ is obtained from $H^{\mu\nu}$ by substituting

$$f_R \rightarrow \mathcal{L} F_R, \quad f_P \rightarrow \mathcal{L} F_P, \quad f_Q \rightarrow \mathcal{L} F_Q.$$

The matter equation of motion may be obtained by taking the divergence of both sides of Eq. (15), which results in

$$\nabla_\nu T^{\mu\nu} = \frac{-\lambda}{1 + \lambda F} (\nabla_\nu F T^{\mu\nu} - 2\nabla_\nu \mathcal{H}^{\mu\nu}). \quad (16)$$

Now if one chooses a Lagrangian of the form given by Eq. (9), the contributions from intrinsic pole-dipole and quadrupole moments can be tracked through $T^{\mu\nu}$ terms in the above equation. However, even if we choose a Lagrangian involving no pole-dipole or quadrupole mo-

ments, quadrupole moment contributions would still appear, through F_R , F_P , and F_Q terms, in the dynamics. Thus the nonminimal coupling induces quadrupole moments in the medium. The appearance of these induced moments is an essential feature of the models with nonminimal coupling between geometry and matter. It is a novel phenomenon not observed in other modified theories of gravity. To proceed further, we consider some rather interesting special cases of this model.

IV. POLE-DIPOLE AND QUADRUPOLE PARTICLES IN NONMINIMAL $f(R)$ GRAVITY

In this section we restrict ourselves to a particular case of the model described above where $f = f(R)$, $F = F(R)$, and $\mathcal{L} = \mathcal{L}(u^\mu, e_a^\mu, \dot{e}_a^\mu, R^\mu_{\alpha\beta\gamma})$; i.e. we consider particles with pole-dipole and quadrupole moments in nonminimal $f(R)$ gravity.

Inserting Eq. (1) into Eq. (16) we obtain

$$\begin{aligned} u^\nu \nabla_\nu p^\mu + p^\mu \nabla_\nu u^\nu + h^{\mu\nu} \nabla_\nu \mathcal{P} + \mathcal{P} \nabla_\nu (u^\mu u^\nu) \\ + \frac{1}{2} R^\mu_{\nu\alpha\beta} u^\nu s^{\alpha\beta} - q_{\alpha\beta\gamma\delta} \nabla^\mu R^{\alpha\beta\gamma\delta} = f^\mu, \end{aligned} \quad (17)$$

with

$$f^\mu = \frac{\lambda F_R \nabla^\nu R}{1 + \lambda F} (\delta_\nu^\mu \mathcal{L}_m - p^\mu u_\nu - \mathcal{P} h_\nu^\mu + B^\mu_\nu - A^\mu_\nu) \quad (18)$$

being the extra force resulting from the nonminimal coupling. There are some quadrupole contributions to this extra force coming from dependence on curvature. These induced moment contributions can be computed with the aid of the following relation:

$$q^{\alpha\beta\gamma\delta} = (1 + \lambda F) \frac{\partial \mathcal{L}}{\partial R_{\alpha\beta\gamma\delta}} + \frac{1}{2} g^{[\alpha\gamma} g^{\beta]\delta} \lambda F_R \mathcal{L}, \quad (19)$$

which is obtained from Eq. (5) by replacing \mathcal{L} with $(1 + \lambda F)\mathcal{L}$. In this equation the first term in the right-hand side is proportional to the medium intrinsic quadrupole moment, while the second one represents the quadrupole moment induced as a result of the nonminimal coupling. One may use the above type of equations to obtain a rough estimate of the magnitude of these induced moments. In the case of the model considered in this section, the magnitude of the induced quadrupole moment relative to the intrinsic one is of the same order as of $\frac{\lambda F_R}{1 + \lambda F}$.

To obtain the equations of motion of particles with pole-dipole and quadrupole moments, in Eq. (17) we set

$$\mathcal{P} = 0, \quad p^\mu = nP^\mu, \quad s^{\mu\nu} = nS^{\mu\nu}, \quad q^\mu_{\alpha\beta\gamma} = nQ^\mu_{\alpha\beta\gamma}$$

and use $\nabla_\alpha (nu^\alpha) = 0$, where n is the particle number density and P^μ , $S^{\mu\nu}$, and $Q^\mu_{\alpha\beta\gamma}$ are the single-particle momentum, spin, and quadrupole moment, respectively. This results in

$$u^\nu \nabla_\nu P^\mu = f^\mu - \frac{1}{2} R^\mu{}_{\nu\alpha\beta} u^\nu S^{\alpha\beta} + Q_{\alpha\beta\gamma\delta} \nabla^\mu R^{\alpha\beta\gamma\delta}. \quad (20)$$

If we neglect the quadrupole moments, this equation is the nonminimal $f(R)$ gravity counterpart of the Mathisson-Papapetrou-Dixon equation and agrees with the results of Ref. [23]. By turning the spin off, the nonminimal coupling of $f(R)$ -perfect fluid considered in Ref. [18] is achieved. For pole-dipole particles with electric charges one can add the Maxwell energy-momentum tensor to the right-hand side of Eq. (1) and proceed as above. This would result in the nonminimal $f(R)$ gravity counterpart of the Dixon-Souriau equations.

A particular case of the model described above is the case where $F(R) = R$. If we also assume, for simplicity, that $\mathcal{P} = 0$ and the medium has no intrinsic pole-dipole or quadrupole moments, Eq. (17) together with Eqs. (18) and (19) gives

$$\nabla_\nu (u^\nu p^\mu) = \frac{\lambda \nabla^\nu R}{1 + \lambda R} \left(\frac{1}{6} \delta_\nu^\mu q^{\alpha\beta}{}_{\alpha\beta} - p^\mu u_\nu \right) \quad (21)$$

in which the following term:

$$\frac{1}{6} q^{\alpha\beta}{}_{\alpha\beta} \nabla^\mu \ln(1 + \lambda R) \approx \frac{\lambda}{6} q^{\alpha\beta}{}_{\alpha\beta} \nabla^\mu R$$

gives the contribution from the induced moment. One may omit the term $p^\mu u_\nu$ in the right-hand side of Eq. (21) by projecting normal to u^μ . Note that according to Ref. [12] this particular toy model is not cosmologically viable.

We further remark that, as has been shown in Ref. [40], in deriving the energy-momentum tensors of the type given in Eq. (1), it is not necessary to have the explicit form of the Lagrangian. Thus it is basically possible to start with different Lagrangians and end up with the same energy-momentum tensor. In this sense, the case of spinning fluids is even more degenerate than the perfect fluid. On the other hand, for the explicit form of the Lagrangians proposed in the literature, e.g. in Ref. [49], the extra force does not vanish.

V. PARTICLES IN NONMINIMAL $f(Q)$ GRAVITY

Now we take $f = f(R)$, $F = F(Q)$, and for simplicity take \mathcal{L} the same as that of a perfect dust; i.e. we consider particles with no intrinsic pole-dipole or quadrupole moments. Inserting these data back into Eq. (16) and projecting the resulting equation normal to u^μ , we get

$$\rho u^\nu \nabla_\nu u^\mu = \frac{4\lambda}{1 + \lambda F} h_\kappa^\mu \nabla_\nu \{ R^{\kappa\alpha\beta\gamma} R^\nu{}_{\alpha\beta\gamma} F_Q \mathcal{L} - \nabla_\alpha \nabla_\beta (F_Q \mathcal{L} (R^{\alpha(\kappa\nu)\beta})) \}. \quad (22)$$

Projection parallel to u^μ would result in a continuity equation. In the above equation, one can express F_Q in terms of the quadrupole moments $q^\mu{}_{\alpha\beta\gamma}$. In fact, by using Eq. (5) with \mathcal{L} being replaced by $(1 + \lambda F(Q)) \mathcal{L}$, we have

$$q^\mu{}_{\alpha\beta\gamma} = 2\lambda \mathcal{L} F_Q R^\mu{}_{\alpha\beta\gamma} \quad (23)$$

from which we obtain

$$\lambda \mathcal{L} F_Q = \frac{1}{2Q} q_{\mu\alpha\beta\gamma} R^{\mu\alpha\beta\gamma}. \quad (24)$$

Inserting this back into Eq. (22), we will arrive at an equation which shows that the particles do not follow geodesics of the space-time due an extra force resulting from the nonminimal coupling. The extra force, which is proportional to the right-hand side of Eq. (22), is normal to the particle trajectories and involves quadrupole moments. Thus, as in the previous case, the nonminimal coupling induces quadrupole moments in the medium. The magnitude of these induced moments depends on the strength of the coupling (controlled by λ) and the function F .

For the particular case of $F(Q) = Q$, we have

$$\rho u^\nu \nabla_\nu u^\mu = \frac{2\lambda}{1 + \lambda Q} h_\kappa^\mu \nabla_\nu \left\{ R^{\kappa\alpha\beta\gamma} R^\nu{}_{\alpha\beta\gamma} \frac{q_{\rho\sigma\tau\omega} R^{\rho\sigma\tau\omega}}{Q} - \nabla_\alpha \nabla_\beta \left(\frac{q_{\rho\sigma\tau\omega} R^{\rho\sigma\tau\omega}}{Q} R^{\alpha(\kappa\nu)\beta} \right) \right\}, \quad (25)$$

which shows explicitly dependence on the induced moments.

VI. PARTICLES IN NONMINIMAL GAUSS-BONNET GRAVITY

Another interesting model sharing the above-mentioned property is the one in which a function of the Gauss-Bonnet invariant is coupled to matter. Such models are of special interest partly due to the fact that $F(R, \mathcal{G})$ are the only subclass of $F(R, P, Q)$ models which are free of ghosts [50,51].

By taking $f = f(R)$, $F = F(\mathcal{G})$ and performing calculations similar to those in the previous section, we obtain

$$\rho u^\nu \nabla_\nu u^\mu = \frac{4\lambda}{1 + \lambda F} h_\kappa^\mu \nabla_\nu \mathcal{K}^{\kappa\nu}, \quad (26)$$

where

$$\begin{aligned} \mathcal{K}^{\kappa\nu} = & (R^{\kappa\nu} - \nabla^\kappa \nabla^\nu + g^{\kappa\nu} \square) \mathcal{L} R F' - 4R^{\kappa\alpha} R^\nu{}_{\alpha} \mathcal{L} F' \\ & + 2\nabla_\alpha \nabla^\nu (\mathcal{L} F' R^{\alpha\kappa}) - 2g^{\kappa\nu} \nabla_\alpha \nabla_\beta (\mathcal{L} F' R^{\alpha\beta}) \\ & - 2\square (\mathcal{L} F' R^{\kappa\nu}) + R^{\kappa\alpha\beta\gamma} R^\nu{}_{\alpha\beta\gamma} \mathcal{L} F' \\ & - \nabla_\alpha \nabla_\beta (\mathcal{L} F' R^{\alpha(\kappa\nu)\beta}) \end{aligned}$$

and F' stands for $\frac{dF(\mathcal{G})}{d\mathcal{G}}$. Also from Eq. (5) with \mathcal{L} replaced by $(1 + \lambda F) \mathcal{L}$ we have

$$\lambda \mathcal{L} F' = \frac{1}{2\mathcal{G}} q_{\mu\alpha\beta\gamma} R^{\mu\alpha\beta\gamma}, \quad (27)$$

which can be inserted back into the expression for $\mathcal{K}^{\kappa\nu}$ to express the right-hand side of Eq. (26) in terms of the induced quadrupole moments.

VII. CONCLUSIONS

We have obtained equations of motion of polarized media with nonminimal coupling to an arbitrary function of the curvature scalar and the squares of the Ricci and Riemann curvature tensors. As a special case, we considered the nonminimal $f(R)$ coupling and obtained the equations of motion of particles with pole-dipole and quadrupole moments which agree with the results of Ref. [23] when the quadrupole moments are absent. Such equations might be used to study the motion of astrophysical objects with pole-dipole or quadrupole moments in the context of modified gravity with nonminimal coupling.

We have shown that, for coupling to a function of rather general dependence as described above, the nonminimal coupling induces quadrupole moments. The magnitude of these induced moments depends on the strength of the coupling and the function coupled to the matter field. The extra force resulting from the coupling can be expressed in terms of the induced moments. We have shown this explicitly for the example case where the coupling function depends only on the Kretschmann invariant and also for the case where the matter Lagrangian couples to an

arbitrary function of the Gauss-Bonnet invariant. By taking dependence on covariant derivatives of the Riemann curvature tensor, higher order intrinsic and induced moments can also be accounted for.

The emergence of induced moments is an essential property differentiating nonminimal modified gravity theories from other modified gravity theories. These induced moments are model-dependent, and possible detection of them via solar system experiments such as the bending of light rays using high precession measurements like those provided by microarcsecond astronomical interferometers might be a test for these models. Keeping the interrelations between gravity and geometry in mind, the quadrupole moments induced by nonminimal coupling to geometry may be compared with the induced quadrupole moments from gravitational fields applied to the media.

ACKNOWLEDGMENTS

I thank Donato Bini for helpful comments on an early draft of this manuscript. I am also indebted to two anonymous referees of Physical Review D for valuable comments.

-
- [1] S. Capozziello and M. Francaviglia, *Gen. Relativ. Gravit.* **40**, 357 (2008).
 - [2] T.P. Sotiriou and V. Faraoni, *Rev. Mod. Phys.* **82**, 451 (2010).
 - [3] A. De Felice and S. Tsujikawa, [arXiv:1002.4928](https://arxiv.org/abs/1002.4928).
 - [4] H.F.M. Goenner, *Found. Phys.* **14**, 865 (1984).
 - [5] S. Nojiri and S.D. Odintsov, *Phys. Lett. B* **599**, 137 (2004).
 - [6] S. Nojiri, S.D. Odintsov, and P.V. Tretyakov, *Prog. Theor. Phys. Suppl.* **172**, 81 (2008).
 - [7] C. Corda, *Mod. Phys. Lett. A* **23**, 109 (2008).
 - [8] G. Allemandi, A. Borowiec, M. Francaviglia, and S.D. Odintsov, *Phys. Rev. D* **72**, 063505 (2005).
 - [9] V. Faraoni, *Phys. Rev. D* **76**, 127501 (2007).
 - [10] O. Bertolami and J. Páramos, *Phys. Rev. D* **77**, 084018 (2008).
 - [11] O. Bertolami and J. Páramos, *Classical Quantum Gravity* **25**, 245017 (2008).
 - [12] T.P. Sotiriou, *Phys. Lett. B* **664**, 225 (2008).
 - [13] T. Harko, *Phys. Lett. B* **669**, 376 (2008).
 - [14] O. Bertolami, C. Carvalho, and J.N. Laia, *Nucl. Phys. B* **807**, 56 (2009).
 - [15] O. Bertolami and M. C. Sequeira, *Phys. Rev. D* **79**, 104010 (2009).
 - [16] S. Nesseris, *Phys. Rev. D* **79**, 044015 (2009).
 - [17] T. Harko, *Phys. Rev. D* **81**, 084050 (2010).
 - [18] O. Bertolami, C. G. Böhmer, T. Harko, and F. S. N. Lobo, *Phys. Rev. D* **75**, 104016 (2007).
 - [19] O. Bertolami, F. S. N. Lobo, and J. Páramos, *Phys. Rev. D* **78**, 064036 (2008).
 - [20] T.P. Sotiriou and V. Faraoni, *Classical Quantum Gravity* **25**, 205002 (2008).
 - [21] V. Faraoni, *Phys. Rev. D* **80**, 124040 (2009).
 - [22] T. Harko, *Phys. Rev. D* **81**, 044021 (2010).
 - [23] D. Puetzfeld and Y.N. Obukhov, *Phys. Rev. D* **78**, 121501 (R) (2008).
 - [24] D. Puetzfeld and Y.N. Obukhov, *Phys. Rev. D* **76**, 084025 (2007).
 - [25] D. Puetzfeld and Y.N. Obukhov, *Phys. Lett. A* **372**, 6711 (2008).
 - [26] P. Szekeres, *Ann. Phys. (N.Y.)* **64**, 599 (1971).
 - [27] G. Montani, R. Ruffini, and R. Zalaletdinov, *Classical Quantum Gravity* **20**, 4195 (2003).
 - [28] W.-M. Suen, *Phys. Rev. D* **34**, 3617 (1986).
 - [29] W.-M. Suen, *Phys. Rev. D* **34**, 3633 (1986).
 - [30] H. Fang and G. Lovelace, *Phys. Rev. D* **72**, 124016 (2005).
 - [31] R. Tammelo, *Gen. Relativ. Gravit.* **8**, 313 (1977).
 - [32] D. Bini, P. Fortini, A. Geralico, and A. Ortolan, *Classical Quantum Gravity* **25**, 035005 (2008).
 - [33] D. Bini, P. Fortini, A. Geralico, and A. Ortolan, *Classical Quantum Gravity* **25**, 125007 (2008).
 - [34] D. Bini, A. Geralico, O. Luongo, and H. Quevedo, *Classical Quantum Gravity* **26**, 225006 (2009).
 - [35] D. Bini, C. Cherubini, S. Filippi, and A. Geralico, *Gen. Relativ. Gravit.* **41**, 2781 (2009).
 - [36] S.M. Kopeikin, *J. Math. Phys. (N.Y.)* **38**, 2587 (1997).
 - [37] M. T. Crosta and F. Mignard, *Classical Quantum Gravity* **23**, 4853 (2006).
 - [38] S.M. Kopeikin and V.V. Makarov, *Phys. Rev. D* **75**, 062002 (2007).

- [39] I. Bailey and W. Israel, *Commun. Math. Phys.* **42**, 65 (1975).
- [40] I. Bailey, *Ann. Phys. (N.Y.)* **119**, 76 (1979).
- [41] M. Mohseni, *Phys. Lett. B* **663**, 165 (2008).
- [42] W.G. Dixon, *Proc. R. Soc. A* **314**, 499 (1970).
- [43] J.M. Souriau, *Ann. Inst. Henri Poincaré, A* **20**, 315 (1974).
- [44] S.M. Carroll, A. De Felice, V. Duvvuri, D.A. Easson, M. Trodden, and M.S. Turner, *Phys. Rev. D* **71**, 063513 (2005).
- [45] G. Cognola, E. Elizalde, S. Nojiri, S.D. Odintsov, and S. Zerbini, *Phys. Rev. D* **73**, 084007 (2006).
- [46] A. De Felice and T. Suyama, *J. Cosmol. Astropart. Phys.* **06** (2009) 034.
- [47] M. Mohseni, *Phys. Lett. B* **682**, 89 (2009).
- [48] G. Cognola and S. Zerbini, *Int. J. Theor. Phys.* **47**, 3186 (2008).
- [49] Y.N. Obukhov and O.B. Piskareva, *Classical Quantum Gravity* **6**, L15 (1989).
- [50] D. Comelli, *Phys. Rev. D* **72**, 064018 (2005).
- [51] I. Navarro and K. Van Acoleyen, *J. Cosmol. Astropart. Phys.* **03** (2006) 008.