

Anti-de Sitter-space/conformal-field-theory correspondence and large- N volume independenceErich Poppitz¹ and Mithat Ünsal²¹*Department of Physics, University of Toronto, Toronto, Ontario M4Y 3B6, Canada*²*SLAC and Physics Department, Stanford University, Stanford, California 94025/94305, USA*

(Received 16 June 2010; published 2 September 2010)

We study the Eguchi-Kawai reduction in the strong-coupling domain of gauge theories via the gravity dual of $\mathcal{N} = 4$ super-Yang-Mills on $\mathbb{R}^3 \times \mathbb{S}^1$. We show that D -branes geometrize volume independence in the center-symmetric vacuum and give supergravity predictions for the range of validity of reduced large- N models at strong coupling.

DOI: 10.1103/PhysRevD.82.066002

PACS numbers: 11.25.Tq, 11.15.Ha, 11.15.Pg, 11.15.Tk

I. INTRODUCTION

Gauge-gravity duality is a powerful method to study strongly coupled gauge dynamics. It relates a weakly coupled theory of gravity to a lower dimensional large- N gauge theory at strong coupling [1–3]. The best-understood example is the AdS₅/CFT₄ correspondence between $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) theory on \mathbb{R}^4 and weakly coupled type-IIB supergravity on AdS₅ × S₅, where all available evidence suggests that the correspondence is exact, at least to leading order in $1/N$.

Another method to study gauge dynamics in lattice and continuum formulations is the large- N volume independence [4–9]. While much less appreciated than the AdS/CFT correspondence, large- N volume independence is one of the few exact results in gauge theories. The statement of the volume-independence theorem is that large- N non-Abelian quantum gauge theories toroidally compactified on four-manifolds, $M_4 = \mathbb{R}^{4-k} \times (\mathbb{S}^1)^k$, have properties that are independent of the $(\mathbb{S}^1)^k$ compactification radii. More precisely, expectation values and connected correlators of single-trace operators are the same in the reduced and infinite-volume theories, to leading order in $1/N$ —if the operators are neutral under the $(\mathbb{Z}_N)^k$ center symmetry and carry momenta in the compact directions quantized in units of the inverse compactification radii. Volume independence holds provided two basic quantum mechanical conditions are satisfied: (i) translation symmetry is not spontaneously broken, and (ii) $(\mathbb{Z}_N)^k$ center symmetry is not spontaneously broken.

In lattice-regularized gauge theories, where the lattice is reduced to a single site, this equivalence is known as “large- N reduction” or “Eguchi-Kawai (EK) reduction” [4]. The necessary and sufficient conditions for the validity of volume independence have been known since the early 1980s. However, the first examples of gauge theories which satisfy them to arbitrarily small volumes were found only recently [8,9]. Because of this, there has been a recent resurgence of interest in this subject, particularly in the lattice community—not only because small volume large- N simulations are more cost effective, but also for other reasons, such as lattice supersymmetry [10–15] (for

volume independence of torus compactifications of non-commutative theories, see [16]). Furthermore, any gauge theory which satisfies volume independence admits a complementary volume-dependent domain, obtained by first fixing N and taking the radii small, where subtle non-perturbative aspects, such as the existence of a mass gap, can be analyzed by semiclassical methods; see, e.g., [9,17,18]. The existence of a semiclassical domain is the main advantage of studying the compactified theory, instead of the theory on \mathbb{R}^4 . For some center-symmetric theories, there is evidence suggesting that the small radius domain is the analytic continuation of the large or infinite radius [9] (also see [19,20]), and the size of the circle times N may be used as an analytic expansion parameter.

Volume independence holds for arbitrary values of the coupling, including the strong-coupling limit of the gauge-gravity correspondence. It is thus interesting to examine the consistency of the two correspondences; at the very least, this provides a consistency check on their exactness. In this paper, we exhibit the simplest setup where volume independence and AdS/CFT should hold simultaneously (see the concluding section for comments on related earlier work [21]). We consider the gravity dual of strongly coupled $\mathcal{N} = 4$ SYM compactified on $\mathbb{R}^3 \times \mathbb{S}^1$ and study how volume independence arises. We show that in the center-symmetric vacuum D -branes “geometrize” volume independence, ensuring that the expectation value of, e.g., a Wilson loop in the uncompactified (\mathbb{R}^3) directions is independent of the \mathbb{S}^1 compactification radius, for arbitrary interquark separation and in accordance with the volume-independence theorem.

II. CENTER-SYMMETRY BROKEN VACUUM: VOLUME DEPENDENCE

The type-IIB background dual to $\mathcal{N} = 4$ SYM compactified on $\mathbb{R}^3 \times \mathbb{S}^1$ of radius R_0 is

$$ds^2 = \frac{u^2}{R_3^2} \left(-dt^2 + \sum_{i=1}^2 dx_i^2 + R_0^2 d\theta^2 \right) + \frac{R_3^2}{u^2} du^2 + R_3^2 d\Omega_5^2, \quad (1)$$

This is compactified AdS⁵ × S⁵ of radius $R_3 \sim \lambda^{1/4}$, in

local Poincaré coordinates, expressed in terms of the energy variable $u \equiv r/l_s^2$. We use string units $l_s = 1$ and denote the 't Hooft coupling of the dual SYM theory by $\lambda \equiv g_{\text{YM}}^2 N$. Compactification of a worldvolume direction of AdS_5 leads to a conical singularity: As seen from (1), the proper radius of \mathbb{S}^1 , equal to uR_0/R_3 , becomes of the order of the string scale at $uR_0 \sim \lambda^{1/4}$. The masses of Kaluza-Klein excitations and string winding modes become comparable, invalidating the supergravity approximation. Thus, for energy scales $uR_0 < \lambda^{1/4}$ the nonsingular gravity description is given by the T-dual (along the $x_3 = R_0\theta$ direction) type-IIA background of N $D2$ -branes located on a dual circle of size $1/R_0$.

The positions of the $D2$ -branes on the dual circle correspond to the eigenvalues of the Wilson loop $\Omega \equiv \exp[i \oint_{\mathbb{S}^1} A]$ of the gauge field around the compact direction. Thus, a vacuum where all N $D2$ -branes are located at the same point on the dual circle breaks the center symmetry, as $\text{tr}\Omega = N$. The type-IIA gravity background corresponding to the center-broken vacuum is easy to determine by the method of images and knowledge of the background of N $D2$ -branes in $\mathbb{R}^{1,9}$:

$$ds^2 = H_2(\bar{r})^{-(1/2)} \left(-dt^2 + \sum_{i=1}^2 dx_i^2 \right) + H_2(\bar{r})^{1/2} (d\bar{r}^2 + \bar{r}^2 d\Omega_6^2). \quad (2)$$

Here $H_2(\bar{r}) = 6\pi^2 g_s N / \bar{r}^5$ is a harmonic function in the seven dimensions transverse to the stack of $D2$ -branes, and g_s is the type-IIA string coupling. Instead of presenting detailed formulas (given in, e.g., [22]), for our purposes it suffices to only picture the brane arrangement. Taking x_3 as the compact direction, the metric of the center-broken BPS-brane configuration is determined by a harmonic function equal to the sum of the harmonic functions due to each stack of N $D2$ -branes separated a distance $1/R_0$ along x_3 , as shown in Fig. 1(a). Each stack of branes creates an $1/\bar{r}^5$ “potential,” where $\bar{r}^2 = x_3^2 + r^2$ and $r (= u)$ denotes the radial direction transverse to both the $D2$ -branes and the compact direction. It is clear from the picture (and intuition from electrostatics) that when $u \gg 1/R_0$ the x_3 -translational invariance of the background is recovered and that at $u = r \gg 1/R_0$ the harmonic function becomes $\sim 1/r^4$, identical to that of the corresponding stack of $D3$ -branes (recall that the type-IIA coupling is $g_s = g_{\text{YM}}^2/R_0$). Thus the type-IIA metric in the center-broken vacuum reads, for $uR_0 \gg 1$:

$$ds^2 = \frac{u^2}{R_3^2} \left(-dt^2 + \sum_{i=1}^2 dx_i^2 \right) + \frac{R_3^2}{R_0^2 u^2} d\theta^2 + \frac{R_3^2}{u^2} du^2 + \hat{R}_3^2 d\Omega_5^2, \quad (3)$$

up to exponentially small corrections. The metric (3) is the T-dual metric of (1) (in the sense of [23]) as evidenced by the fact that only the $d\theta^2$ terms are different (we do not




FIG. 1 (color online). Center-symmetric and center-broken $D2$ -branes on the dual \mathbb{S}^1 of radius $1/R_0$.

show the relation between the type-IIA and type-IIB dilatons, which is trivial to obtain). It is also clear that in the center-broken vacuum the type-IIA metric will differ from (3) once uR_0 becomes of order unity or smaller.

We conclude that in the center-broken vacuum the backgrounds (3) and (1) are equivalent for $uR_0 \gg 1$, where the x_3 isometry is restored. Thus, for example, a calculation of the expectation value of a Wilson loop of size $R \times T$, positioned in the $x_1 - t$ plane of the noncompact \mathbb{R}^3 can be made via (3) so long as $R \ll R_0$ —so that the string world sheet only probes the bulk geometry in the $uR_0 \gg 1$ region, close to the “UV-brane” (recall the “energy-distance” relation $u_* R \sim 1$ for the minimum value u_* of u probed by a Wilson loop of size R [24]). Thus, Wilson loops of interquark separation $R \ll R_0$ are unaffected by the compactification, as one would naively expect. However, the world sheet relevant for Wilson loops with $R \gg R_0$ probes the bulk geometry further away from the UV, as now $u_* R_0 \ll 1$, a region where (3) receives corrections due to the compactification.

Hence, in the center-symmetry breaking vacuum, the Wilson loop (and other correlators) exhibit volume dependence. This is consistent with expectations from compactified field theory that the Wilson loop with interquark separation $R \gg R_0$ should be sensitive to the $\mathbb{R}^3 \times \mathbb{S}^1$ compactification. In fact, a dual gravity analysis [25] of the Wilson loop in the compactified $\mathcal{N} = 4$ SYM theory shows that the behavior of the quark-antiquark potential changes from $1/R$, at short distances $R \ll R_0$, to $1/R^{2/3}$ in an intermediate $D2$ -brane region, and back to $1/R$ in the far-infrared $M2$ -brane region (the latter describes the three-dimensional 16 supercharge CFT that $\mathcal{N} = 4$ SYM flows to upon a center-symmetry breaking compactification [26]).

III. CENTER-SYMMETRIC VACUUM: VOLUME INDEPENDENCE

Consider now the center-symmetric vacuum of the $\mathcal{N} = 4$ SYM theory on $\mathbb{R}^3 \times \mathbb{S}^1$. According to EK reduction, appropriate observables should now exhibit \mathbb{S}^1 -size independence.

In the type-IIA picture, a center-symmetric vacuum corresponds to a configuration of N $D2$ -branes distributed equidistantly on the dual circle—since their positions on the dual \mathbb{S}^1 correspond to the eigenvalues of Ω , now clearly $\text{tr}\Omega^k = 0$, for all $k \neq 0(\text{mod}N)$. The metric dual to the center-symmetric $D2$ -brane configuration can similarly be computed using the method of images. The difference is that now single $D2$ -branes are spaced a distance $1/(NR_0)$ apart along the compact x_3 direction, as shown in Fig. 1(b). The harmonic function determining the background is, again, the sum of the $1/\bar{r}^5$ “potentials” of the individual $D2$ -branes (with $\bar{r}^2 = x_3^2 + r^2$, as before), resulting in the $D2$ -brane harmonic function which determines the metric, as in (2):

$$H_2^{\text{sym}}(r, x_3) = \sum_{n=-\infty}^{\infty} \sum_{k=1}^N \frac{6\pi^2 g_s}{[r^2 + (x_3 - \frac{2\pi}{R_0 N} k - \frac{2\pi n}{R_0})^2]^{5/2}}. \quad (4)$$

By Poisson resummation, (4) takes the form:

$$\frac{R_3^4}{u^4} \left[1 + \sum_{m=1}^{\infty} (muNR_0)^2 K_2(muNR_0) \cos(mx_3NR_0) \right], \quad (5)$$

where K_2 is the modified Bessel function.

The crucial difference with respect to the center-broken vacuum discussed in [22] is the appearance of a factor of N in the correction term in (5). Hence, the x_3 isometry is now recovered for much smaller values of r ($= u$, the energy scale). It is clear [from (5) or from electrostatics] that now the condition for isometry restoration is $uNR_0 \gg 1$, instead of $uR_0 \gg 1$ in the center-broken vacuum. Thus the background dual to the center-symmetric vacuum is also given by (3) but is now valid for $uNR_0 \gg 1$. We note that while near each individual $D2$ -brane the supergravity approximation is not to be trusted (because large curvatures occur and physics is described by an IR free Abelian theory), no large curvatures appear in the background (3) and (5) at $u \gg 1/(NR_0)$, i.e., at any finite distance away from the $D2$ -branes (similar backgrounds are also considered in [27]).

The fact that the center-symmetric vacuum is described by (3) for any $u \gg 1/(NR_0)$ immediately implies that a Wilson loop of any size (strictly speaking of size $R \ll NR_0$) will be insensitive to the compactification. Thus, the potential between two static quarks exhibits the behavior characteristic of the four-dimensional $\mathcal{N} = 4$ CFT, $V(R) \sim \lambda^{1/2}/R$, at all scales, despite the fact that one dimension is compactified and conformal symmetry of the background is explicitly broken.


In the language of nonperturbative orbifold equivalences [8], the neutral-sector observables in the compactified “daughter” theory enjoy the conformal symmetry of its “parent” theory on \mathbb{R}^4 , at leading order in N . However, it should also be noted that the daughter theory also possesses a non-neutral sector aware of the compactification radius. The main point is that for neutral-sector observables, the space may be viewed as having an effective size $R_{\text{eff}} = R_0 N$ and thus $N = \infty$ is a decompactification limit.

It is clear that other quantities will also exhibit volume independence—for example, correlation functions of single-trace operators that only carry momentum in the noncompact directions will also be insensitive to the compactification, as required by EK reduction. EK reduction also requires that expectation values of Wilson loops extending also in \mathbb{S}^1 , but not winding around the compact direction (i.e., center-symmetry neutral ones), exhibit volume independence; however, explicitly verifying their volume independence in the gravity dual appears more challenging to us than for the observables we consider.

Finally, we briefly note a slight refinement of the condition for volume independence inferred from the gravity dual. In our discussion above, we did not consider the behavior of the type-IIA dilaton. In fact, examining its behavior shows that the effective string coupling becomes large when $uNR_0 \sim \lambda^{5/4}$ —thus, the region of validity of volume independence that can be inferred from the type-IIA dual of the center-symmetric vacuum would be $uNR_0 \gg \lambda^{5/4}$, instead of simply $uNR_0 \gg 1$. However, the type-IIA description can be uplifted to 11-dimensional supergravity (M theory) [25]. The size of the 11th direction, parameterized by x_{11} , is related to the type-II couplings as $R_{11} = g_s = \lambda/(NR_0)$ and the center-symmetric $D2$ brane configuration is replaced by a similar configuration of $M2$ -branes located a distance $1/(NR_0)$ apart along x_3 , as shown in Fig. 2. For $u \gg R_{11} = \frac{\lambda}{NR_0} \gg 1/(NR_0)$ the $M2$ -brane background can be written in a form:

$$ds_{11} = e^{-2\Phi/3} ds_{10}^2 + e^{4\Phi/3} dx_{11}^2, \quad (6)$$

where ds_{10}^2 is the type-IIA metric (3) and Φ is the type-IIA dilaton. As this background is dual to (3), it follows that the regime of volume independence is $uNR_0 \gg \lambda$, improving on the type-IIA bound $uNR_0 \gg \lambda^{5/4}$. A bound on the validity of volume independence of the form $uNR_0 \gg \max(1, \lambda)$ can also be inferred from field-theory considerations and is consistent with the strong-coupling bound from supergravity obtained here (see [28] for a field-theory analysis of $\mathcal{N} = 4$ SYM in the volume-independence context). At asymptotically low energies $uNR_0 \ll 1$, 11-dimensional supergravity breaks down for center-symmetric $M2$ -branes, consistent with the free Abelian long-distance dynamics, unlike the coincident $M2$ -branes case where the IR physics is non-Abelian and superconformal.

FIG. 2 (color online). Center-symmetric $M2$ -branes.

IV. CONCLUSIONS

We considered the simplest case where EK reduction of a four-dimensional gauge theory is valid simultaneously with gauge-gravity duality. Our considerations indicate that $\mathcal{N} = 4$ SYM compactified on $\mathbb{R}^3 \times S^1$ indeed exhibits volume independence in the center-symmetry preserving vacuum. The gravity dual of four-dimensional $\mathcal{N} = 4$ SYM gives the first explicitly solvable realization of volume independence above two dimensions (where EK reduction is manifest in the large- N limit of the exactly solvable pure YM lattice theory [29]), at least in the present context—previously, large- N volume (temperature) independence of the confined phase has been exhibited in studies of gravity duals beginning with [30]. Our findings can also be viewed as providing a check on the weakest form of the AdS/CFT correspondence.

It would be interesting to consider how EK reduction works when more than one dimension is compactified, especially with regard of how center-symmetry preservation is reflected in the brane and gravity setups. The $\mathbb{R}^3 \times S^1$ case is special in this respect, as one is free to choose a

classical center-symmetric vacuum state, not washed away by quantum fluctuations which become strong as more dimensions are compactified. We note that Ref. [21] previously considered large- N reductions in the holographic picture with all dimensions compactified, but the matching of observables and the question of fluctuations raised above were not studied.

It may also be interesting to study volume independence for confining gauge theories with known gravity duals, as well as by exploiting the analogy between the $1/N$ and genus expansion in gauge and string theories. For example, [31] showed that free energy of YM theory receives contributions only from Riemann surfaces of genus ≥ 1 in the confined phase [$O(N^0)$], but it receives a contribution from genus zero in the deconfined phase [$O(N^2)$]. This is nothing but the temperature independence of confined phase and temperature dependence of the deconfined phase, to leading order in N . In this context, for example, the double-trace deformation stabilizing the theory to the confined phase [9,18] may have a stringy interpretation in terms of analytic continuation of the winding-number unbroken phase of [32] to small radii.

ACKNOWLEDGMENTS

We are indebted to J. Maldacena, O. Aharony, and L. Yaffe for their crucial help. We also thank B. Burrington, M. Hanada, A. Karch, H. Neuberger, A. Peet, A. Rajaraman, S. Shenker, and E. Witten for useful discussions. M. Ü. thanks the Weizmann Institute of Science, where portions of this work was done, for hospitality. This work was supported by the U.S. Department of Energy Grant No. DE-AC02-76SF00515 and by the National Science and Engineering Council of Canada (NSERC).

-
- [1] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998); *Int. J. Theor. Phys.* **38**, 1113 (1999).
 - [2] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Phys. Lett. B* **428**, 105 (1998).
 - [3] E. Witten, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
 - [4] T. Eguchi and H. Kawai, *Phys. Rev. Lett.* **48**, 1063 (1982).
 - [5] L. G. Yaffe, *Rev. Mod. Phys.* **54**, 407 (1982).
 - [6] G. Bhanot, U. M. Heller, and H. Neuberger, *Phys. Lett.* **113B**, 47 (1982).
 - [7] A. Gonzalez-Arroyo and M. Okawa, *Phys. Rev. D* **27**, 2397 (1983).
 - [8] P. Kovtun, M. Ünsal, and L. G. Yaffe, *J. High Energy Phys.* **06** (2007) 019.
 - [9] M. Ünsal and L. G. Yaffe, *Phys. Rev. D* **78**, 065035 (2008).
 - [10] B. Bringoltz and S. R. Sharpe, *Phys. Rev. D* **80**, 065031 (2009).
 - [11] A. Hietanen and R. Narayanan, [arXiv:0911.2449](https://arxiv.org/abs/0911.2449).
 - [12] M. Hanada, L. Mannelli, and Y. Matsuo, *Phys. Rev. D* **80**, 125001 (2009).
 - [13] T. Ishii, G. Ishiki, S. Shimasaki, and A. Tsuchiya, *Phys. Rev. D* **78**, 106001 (2008).
 - [14] G. Ishiki, S. W. Kim, J. Nishimura, and A. Tsuchiya, *J. High Energy Phys.* **09** (2009) 029.
 - [15] M. Hanada, S. Matsuura, and F. Sugino, [arXiv:1004.5513](https://arxiv.org/abs/1004.5513).
 - [16] A. Connes, M. R. Douglas, and A. S. Schwarz, *J. High Energy Phys.* **02** (1998) 003.
 - [17] E. Poppitz and M. Ünsal, *J. High Energy Phys.* **09** (2009) 050.
 - [18] P. N. Meisinger and M. C. Ogilvie, *Phys. Rev. D* **81**, 025012 (2010).
 - [19] J. C. Myers and M. C. Ogilvie, *Phys. Rev. D* **77**, 125030 (2008).
 - [20] R. D. Pisarski, *Phys. Rev. D* **74**, 121703 (2006).
 - [21] K. Furuuchi, *Phys. Rev. D* **74**, 045027 (2006).

- [22] E. Pontón and E. Poppitz, *J. High Energy Phys.* **02** (2001) 042.
- [23] T.H. Buscher, *Phys. Lett. B* **201**, 466 (1988).
- [24] J.M. Maldacena, *Phys. Rev. Lett.* **80**, 4859 (1998).
- [25] N. Izhaki *et al.*, *Phys. Rev. D* **58**, 046004 (1998).
- [26] N. Seiberg, *Nucl. Phys. B, Proc. Suppl.* **67**, 158 (1998).
- [27] H. Lin and J.M. Maldacena, *Phys. Rev. D* **74**, 084014 (2006).
- [28] M. Ünsal and L.G. Yaffe, *arXiv:1006.2101*.
- [29] D.J. Gross and E. Witten, *Phys. Rev. D* **21**, 446 (1980).
- [30] E. Witten, *Adv. Theor. Math. Phys.* **2**, 505 (1998).
- [31] J.J. Atick and E. Witten, *Nucl. Phys.* **B310**, 291 (1988).
- [32] J. Polchinski, *Phys. Rev. Lett.* **68**, 1267 (1992).