

Quasilocal Hamiltonians in general relativity

Michael T. Anderson*

Department of Mathematics, Stony Brook University, Stony Brook, N.Y. 11794-3651, USA

(Received 1 September 2010; published 27 October 2010)

We analyze the definition of quasilocal energy in general relativity based on a Hamiltonian analysis of the Einstein-Hilbert action initiated by Brown-York. The role of the constraint equations, in particular, the Hamiltonian constraint on the timelike boundary, neglected in previous studies, is emphasized here. We argue that a consistent definition of quasilocal energy in general relativity requires, at a minimum, a framework based on the (currently unknown) geometric well-posedness of the initial boundary value problem for the Einstein equations.

DOI: 10.1103/PhysRevD.82.084044

PACS numbers: 04.20.Fy

The analysis of the gravitational field by Arnowitt-Deser-Misner (ADM) [1] has led to a clear and well-defined construction of the Hamiltonian, and resulting definitions of energy, linear, and angular momentum in the context of asymptotically flat spacetimes. These concepts are obviously of basic importance in understanding the physics of such (infinite) isolated gravitating systems. Nevertheless, infinite systems are idealizations of more realistic physical situations, and it is desirable to have available a similar analysis in the case of physical systems of finite extent.

However, the understanding of this issue for domains of finite extent is much less satisfactory. Despite numerous proposals, from a number of different viewpoints, a consensus has not yet been reached on a suitable definition of the Hamiltonian or energy of a finite system, i.e., a quasilocal Hamiltonian; cf. [2] for an excellent detailed survey of the current state of the art.

In this paper, we first examine and comment on the approach to the definition of energy of a finite region of spacetime based on the Hamiltonian formulation of general relativity (GR). This is essentially based on a localization of the approach taken by ADM [1] and Regge-Teitelboim [3], keeping careful track of the boundary terms that arise in a Hamiltonian or Hamilton-Jacobi analysis. This approach was initiated and pioneered by Brown-York (BY) [4]. To keep the discussion focused on the central issue, we only consider the gravitational field, (so other matter fields are set to zero); in addition, we consider only the energy and not related concepts such as linear and angular momentum, although this could be done without undue difficulty. Finally, most of the discussion below applies also to more recent modifications of the BY approach by several authors, cf. [5–8], however, again for clarity and simplicity we focus on the Brown-York Hamiltonian and leave it to the reader to extend the analysis to the more recent alternatives.

We first recall the setup. Let M be a spacetime region, topologically of the form $I \times \Sigma$, with $I = [0, 1]$ parametrizing time and Σ a compact 3-manifold with boundary S ; typically $S = \partial\Sigma$ is a 2-sphere and Σ a 3-ball. The boundary ∂M of M is a union of two spatial hypersurfaces $\Sigma_0 \cup \Sigma_1$ and the timelike boundary $\mathcal{T} = I \times \partial S = \cup S_t$. These boundaries meet at the seams or corners S_0 and S_1 . The Einstein-Hilbert action is then given by (setting $8\pi G = 1$),

$$I_{\text{EH}}(\mathfrak{g}) = \int_M R_{\mathfrak{g}} dV_{\mathfrak{g}}, \quad (1)$$

where \mathfrak{g} is a smooth Lorentz metric on M .

The Hamiltonian in GR plays two important but *a priori* distinct roles. In classical field theories without dynamical gravity and based on a fixed background (Minkowski) spacetime, these two roles coincide. It is not at all clear, at least in the case of finite domains, whether they can be made to coincide in GR.

A Hamiltonian \mathcal{H} for the action (1) depends on a choice of time function t and associated vector field ∂_t , giving a foliation Σ_t of the spacetime, i.e., a $3 + 1$ decomposition. Given the spacetime metric \mathfrak{g} , this is equivalent to specifying a lapse u and shift X , so that $\partial_t = uT + X$, where T is the unit timelike normal to the foliation. Thus, $\mathcal{H} = \mathcal{H}_{(u,X)}$.

A full-fledged Hamiltonian analysis requires a well-defined phase space T^*Q , the cotangent bundle of the configuration space Q , with variables (g, π) where π is the momentum conjugate to g ; g is a Riemannian metric on Σ . In the case of finite boundaries, boundary conditions for the variables (g, π) must be specified in such a way that the Hamiltonian $\mathcal{H}_{(u,X)}: T^*Q \rightarrow \mathbb{R}$ is first a smooth function on T^*Q , and second the Hamiltonian vector field on T^*Q (generated from \mathcal{H} by the symplectic structure on T^*Q) generates exactly the equations of motion; integral curves of the Hamiltonian vector field give vacuum solutions of the Einstein equations satisfying the boundary conditions. For this to be consistent, boundary conditions for the lapse shift (u, X) must be determined, and these must also be preserved under the equations of motion. If the above can

*anderson@math.sunysb.edu
http://www.math.sunysb.edu/~anderson

be accomplished, one obtains a Hamiltonian depending on the gauge choice (u, X) .

On the other hand, in other classical field theories, the energy is understood as the Noether charge associated to time-translation symmetries; for Minkowski backgrounds, the energy is thus the time component of an invariantly defined energy-momentum 4-vector. In GR, one has typically no symmetries, i.e., Killing fields, and so no conserved charges. Nevertheless, one can attempt to define preferred or distinguished “quasisymmetries” or “quasi-Killing fields.” This gives a preferred choice of the lapse shift (u, X) , and the Hamiltonian is then taken with respect to such a choice, cf. [2,9] for further discussion.

The approach of Brown-York is to choose Dirichlet boundary conditions for the metric $\gamma = \gamma_{\mathcal{T}}$ induced on \mathcal{T} . This choice naturally conforms to the modification of the Einstein-Hilbert (EH) action by the addition of a boundary term—the well-known Gibbons-Hawking-York or $\text{tr}K$ boundary term [10,11]. Thus, consider the modified Lagrangian

$$I(\mathfrak{g}) = \int_M R_{\mathfrak{g}} dV_{\mathfrak{g}} + 2 \int_{\mathcal{T}} k dV_{\gamma}, \quad (2)$$

where $k = \text{tr}K$ is the mean curvature of the boundary \mathcal{T} in (M, \mathfrak{g}) with respect to the outward unit normal and γ is the metric on \mathcal{T} induced by \mathfrak{g} . A straightforward calculation shows that the variation of I at \mathfrak{g} in the direction h is given by

$$\delta_{\mathfrak{g}} I(h) = - \int_M \langle E, h \rangle dV_{\mathfrak{g}} - \int_{\mathcal{T}} \langle \pi_{\mathcal{T}}, h^T \rangle dV_{\gamma}, \quad (3)$$

where $E(\mathfrak{g}) = \text{Ric}_{\mathfrak{g}} - \frac{R}{2} \mathfrak{g}$ is the Einstein tensor, $\pi_{\mathcal{T}} = K - k\gamma$ is momentum conjugate to γ on \mathcal{T} , and h^T is the induced variation of γ on \mathcal{T} ; $\langle \cdot, \cdot \rangle$ is the pairing induced by the metric \mathfrak{g} . Thus, the variation vanishes on-shell, when the metric γ is held fixed at \mathcal{T} , ($h^T = 0$). The action is functionally differentiable (in fact C^∞ smooth) with respect to variations of the metric fixed on the boundary. In other words, one has a well-defined variational principle for Dirichlet boundary data. [The variational formula (3) should also include other terms at Σ_0 and Σ_1 and the corners, but these will be ignored here since they play no role in the analysis.]

On-shell, i.e., on the space of solutions of the vacuum Einstein equations, the BY quasilocal Hamiltonian (or quasilocal energy) is formed by taking the on-shell variation of the action (2) in the direction of a timelike unit normal vector field T on the boundary $S = \partial\Sigma$; one assumes here that T is tangent to \mathcal{T} . Since $\pi(T, T) = H$, where H is the mean curvature of $S \subset \Sigma$, the on-shell variation in the (T, T) direction is given by

$$- \int_S H dV_{\gamma_S}, \quad (4)$$

where γ_S is the metric induced on S . More generally, let (u, X) be the lapse-shift of the foliation Σ_t . A standard Hamiltonian analysis gives

$$\mathcal{H}_{(u,X)} = - \int_S [uH - \pi(X, \nu)] dV_{\gamma_S}, \quad (5)$$

on-shell, where ν is the outward unit normal of $S \subset \Sigma$. The case (4) is recovered by setting $(u, X) = (1, 0)$. There remains a freedom in specifying the zero-point energy; the prescription of Brown-York is to define

$$\mathcal{H}_{\text{BY}}(S) = \int_S (H_0 - H) dV_{\gamma_S}, \quad (6)$$

where H_0 is the mean curvature of an isometric embedding of (S, γ_S) into Euclidean 3-space \mathbb{R}^3 . Hence $\mathcal{H}_{\text{BY}}(S)$ is well defined only if there exists a unique isometric embedding into \mathbb{R}^3 ; this is the case if for instance the Gauss curvature K_{γ_S} is positive, by the Weyl embedding theorem.

The BY quasilocal Hamiltonian has a number of important and interesting properties, both physically and mathematically. The expression (4) is local on S and easily computable, although the subtraction term in (6) is more complicated since it depends on the global structure of (S, γ_S) . Particularly noteworthy is the result of Shi-Tam [12] that for time-symmetric data ($\pi_{\Sigma} = 0$), if the Gauss curvature $K_{\gamma_S} > 0$ then $\mathcal{H}_{\text{BY}}(S) \geq 0$ with equality if and only if Σ is flat. However, for general surfaces S , $\mathcal{H}_{\text{BY}}(S)$ may be negative.

Observe that $\mathcal{H}_{\text{BY}}(S)$ depends only on the Cauchy data on the initial surface Σ ; in fact it depends only on the metric g on Σ near S . It depends on the choice of Σ , (within the domain of dependence) and one obtains different energies for different Cauchy surfaces, i.e. the Brown-York energy is gauge dependent. However, it does not in fact depend on the actual unit normal T at S ; this is due to a cancellation of boundary terms. Note that the Cauchy data (g, π) of Σ do not determine the unit normal T at S ; equivalently the lapse shift (u, X) at S is undetermined by Cauchy data.

It follows that given an initial data surface Σ , the BY Hamiltonian is independent of the structure of the space-time outside the domain of dependence $D(\Sigma)$ of Σ ; it is the same no matter what the metric is outside $D(\Sigma)$. It is not *a priori* clear why the energy of the gravitational field should be independent of its structure at \mathcal{T} .

Although certainly natural, there is a basic problem with a definition of the Hamiltonian depending on the choice of Dirichlet boundary data as in and following (2). As stated explicitly in [4], the action evaluated on a classical solution (i.e., the on-shell action) is understood to be a function of the boundary metric γ on \mathcal{T} . The space \mathcal{E} of solutions of the Einstein equations is thus to be smoothly parametrized by the space of metrics $\text{Met}(\mathcal{T})$ on the boundary \mathcal{T} , together with the space \mathcal{C} of Cauchy data on Σ (satisfying the constraint equations):

$$\mathcal{E} \simeq \mathcal{C} \times \text{Met}(\mathcal{T}).$$

However, this is not the case. In fact for generic choices of boundary metric γ , there will be *no* solution of the

equations of motion inducing γ on \mathcal{T} . This is due the constraint equations, and, in particular, to the Hamiltonian or scalar constraint, on \mathcal{T} . This issue, that there may be no solutions to the equations of motion on the phase space without the correct boundary conditions, is exactly the underlying issue and theme in the Regge-Teitelboim analysis of the ADM Hamiltonian.

To illustrate this point clearly, we discuss it in three different but related situations.

(I) Consider first the simpler case of the pure Cauchy problem for the Einstein equations. Here the Cauchy data (g, π) on Σ parametrize the space of solutions, and one may ask if one may specify the metric g (Dirichlet data) arbitrarily on Σ to generate a solution. However, this is not the case since the data must satisfy the momentum constraint $\delta\pi = 0$ and, more importantly here, the Hamiltonian or Gauss constraint $|\pi|^2 - \frac{1}{2}(\text{tr } \pi)^2 - R_g = 0$. The initial data (g, π) are usually assumed to lie in Sobolev spaces $H^s \times H^{s-1}$; the arguments below are equally valid in simpler spaces such as $C^k \times C^{k-1}$. The Hamiltonian constraint then implies $R_g \in H^{s-1}$. For generic $g \in H^s$, $R_g \in H^{s-2}$, not in H^{s-1} . Hence, $g \in H^s$ cannot be freely prescribed. Thus, as in the Lichnerowicz approach to solving the constraint equations, only, for example, the conformal class $[g]$ can be prescribed.

(II) Consider next the Euclidean situation, i.e., the Euclidean Einstein equations on a compact domain M with boundary ∂M . Here the Einstein equations $E(g) = 0$ with Dirichlet boundary conditions $\gamma = g|_{\partial M}$ given, do *not* form a well-posed elliptic boundary value problem (for any choice of gauge) essentially for the same reasons described in (I), namely, the Gauss or Hamiltonian constraint. This is in spite of the fact that the variational problem for the action (2) is well defined for Dirichlet boundary data, exactly as in the Lorentzian case.

There are many choices of boundary data that can be used to obtain a well-posed elliptic boundary value problem; geometrically perhaps the most natural is that given by

$$([\gamma], k), \quad (7)$$

i.e., prescribing the conformal class $[\gamma]$ of the metric on ∂M and the mean curvature k of ∂M in M ; cf. [13] for proof of these results and further discussion.

(III) The same argument applies for a timelike boundary \mathcal{T} . The ‘‘Hamiltonian constraint’’ $|K|^2 - k^2 + R_\gamma = 0$ along \mathcal{T} again constrains the freedom of the metric γ on \mathcal{T} . For example, given a 3 + 1 decomposition of the spacetime, let (g_t, π_t) be the curve of Cauchy data on the surfaces Σ_t . Working again in Sobolev spaces H^s , the trace of the metric

(or any function) on the boundary, loses half a derivative, so on \mathcal{T} , $\gamma_S \in H^{s-1/2}$. The second fundamental form K of \mathcal{T} involves a (spatial) derivative of g , so $K \in H^{s-3/2}$. The constraint equation then gives $R_\gamma \in H^{s-3/2}$. But for generic $\gamma \in H^{s-1/2}$, one will not have $R_\gamma \in H^{s-3/2}$ but instead $R_\gamma \in H^{s-5/2}$ —the same behavior as in (I) or (II).

As noted above, the quasilocal energy $\mathcal{H}_{\text{BY}}(S)$ is the value of the on-shell Hamiltonian that generates unit time translation orthogonal to S at the boundary Σ and is given by the variation of the $\text{Tr}K$ action in a unit timelike normal direction along the boundary. From the point of view of the initial boundary value problem (IBVP) for the vacuum Einstein equations, this presupposes that there exists a solution of the Einstein equations in the bulk M , whose boundary metric is of the form

$$\gamma_{\mathcal{T}} = -dt^2 + \gamma_t, \quad (8)$$

on \mathcal{T} , at least to 1st order in t , so that one is prescribing the form of the metric, i.e., Dirichlet boundary data, on \mathcal{T} . The Cauchy data (g, π) at $t = 0$ determine the derivative $\partial_t \gamma_{\mathcal{T}}|_{t=0}$. However, the boundary value problem (8) cannot be solved in general; the IBVP is not well posed (even at the linearized level) for Dirichlet boundary data. The Hamiltonian constraint serves as an obstruction to solvability in general.

In sum, the constraint equations on the Cauchy surface Σ and boundary \mathcal{T} ‘‘generate’’ the diffeomorphisms and impose constraints on the allowed Cauchy and boundary data. The Hamiltonian constraint on \mathcal{T} generates diffeomorphisms normal to \mathcal{T} and so is related to the location of \mathcal{T} in the spacetime which thus cannot be fully prescribed *a priori*. The presence of the constraints on \mathcal{T} and the difficulties they present in obtaining a well-behaved quasilocal Hamiltonian has not been addressed previously in the literature.

For the flow of the Hamiltonian vector field to be well-defined, it is necessary that the IBVP for the Einstein equations is well-posed. It is this criterion one should choose to seek suitable boundary conditions on \mathcal{T} .

Before discussing the problem in general, we consider some constructions of possible alternatives to the BY Hamiltonian. First, the linearization of the scalar curvature R_g in the direction of a variation h of g is given by $L(h) = R'_h = -\square \text{tr}h + \delta\delta h - \langle \text{Ric}, h \rangle$, where δh is the divergence of h . Hence,

$$\begin{aligned} \delta \mathcal{I}_{\text{EH}}(h) &= \int_M -\square \text{tr}h + \delta\delta h - \langle \text{Ric}, h \rangle + \frac{1}{2}R \text{tr}h \\ &= -\int_M \langle E, h \rangle + \int_{\partial M} -\nu(\text{tr}h) - (\delta h)(\nu), \end{aligned} \quad (9)$$

where ν is the unit outward normal to ∂M in M . One has $-\nu(\text{tr}h) - (\delta h)(\nu) = -2k'_h - \langle K, h \rangle + \delta(h(\nu))^T$. The last

term is a divergence term, which integrates to 0 on the boundary, so

$$\delta I_{\text{EH}}(h) = - \int_M \langle E, h \rangle - \int_{\partial M} 2k'_h + \langle K, h \rangle.$$

Since $(k dV_g)'_h = k'_h dV_g + \frac{1}{2} k \text{tr} h dV_g$, this gives the variational formula for I in (3).

Consider next for instance the variational problem for a fixed conformal class and mean curvature, as in (7) above. One has $\langle K, h \rangle = \langle K_0, h_0 \rangle + \frac{1}{3} k \text{tr} h$ where K_0 and h_0 denote the trace-free parts. On other hand, $\frac{2}{3} (k dV)'_h = \frac{2}{3} k'_h + \frac{1}{3} k \text{tr} h$, and so for the action

$$I_{Ck}(g) = \int_M R + \frac{2}{3} \int_{\mathcal{T}} k,$$

one has

$$\delta I_{Ck}(h) = - \int_M \langle E, h \rangle - \int_{\mathcal{T}} \left[\langle \pi_0, h_0 \rangle + \frac{4}{3} k'_h \right]. \quad (10)$$

This gives a well-defined variational problem with prescribed conformal class $[\gamma]$ and mean curvature k on \mathcal{T} , i.e., this action is a smooth function on the configuration space of 4d metrics with boundary data (7) fixed.

In the case of Euclidean signature, the boundary data $([\gamma], k)$ form a well-posed elliptic boundary value problem, so that one may expect to find a unique solution, (at least under mild conditions). Whether this holds for the Lorentzian signature is unknown, (but unlikely). Suppose nevertheless for the sake of argument that the Lorentzian problem is well posed. If one takes a timelike vector field ∂_t to form a Hamiltonian then linearized boundary data (h_0, k'_h) on \mathcal{T} determine a unique bulk h solution along Σ (or Σ_t); from this one may then read off Dirichlet boundary data h^T on \mathcal{T} . To evaluate the Hamiltonian, one then chooses h (if possible) so that $h^T = T \cdot T$ at S . The corresponding data (h_0, k'_h) are then paired with the coefficients $(\pi_0, \frac{4}{3})$ above, allowing one to determine the corresponding quasilocal Hamiltonian. The determination of the Hamiltonian is thus "global" on the solution and is rather complicated.

The simplest solutions of the Einstein equations are the time-independent static solutions, of the form

$$g = -u^2 dt^2 + g,$$

where ∂_t is a (hypersurface orthogonal) Killing field. Consider then a Hamiltonian analysis for static metrics. In place of the EH action (1) or the BY action (2), consider here

$$I_{\text{St}}(g) = \int_M R_g dV_g + 2 \int_{\partial M} \nu(u) dV_\gamma,$$

where ν is the outward unit normal. The term R_g may be computed in terms of R_g and u , and via an integration by parts it is easily verified that

$$I_{\text{St}}(g) = \int_\Sigma u R_g dV_g, \quad (11)$$

so that the boundary term disappears on passage to the Cauchy surface. A straightforward computation along the lines of (9) and (10) above gives

$$\begin{aligned} \delta_g I_{\text{St}}(h, u') &= \int_\Sigma \left[\left\langle L^* u + \frac{1}{2} u R \gamma, h \right\rangle + R u' \right] \\ &+ \int_S \langle u K - \nu(u) \gamma_S, h^T \rangle + 2u k'_h, \end{aligned} \quad (12)$$

cf. [14] for a proof. Here $L^* u = D^2 u - \Delta u \cdot \gamma - u \text{Ric}$ is the adjoint of the linearization L of the scalar curvature. The vanishing of the bulk term in (12) gives exactly the static vacuum Einstein equations, as expected. The boundary term vanishes when $h^T = 0$ and $k'_h = 0$. The mean curvature k of \mathcal{T} in M is the same as the mean curvature H of S in Σ .

This gives a well-defined variational problem for the boundary data

$$(\gamma_S, H) \quad \text{at } S = \partial \Sigma, \quad (13)$$

so that the Lagrangian is a smooth function on the configuration space with these boundary data. Note that one may (trivially) Wick rotate static spacetimes to Euclidean signature. In contrast to Dirichlet boundary data (γ_S, u) , the boundary data (13) are well posed, i.e., elliptic, for the static Einstein equations, (cf. [15]).

The time function t and vector field ∂_t give the natural lapse shift $(u, 0)$ for static metrics. With respect to this, the static Hamiltonian \mathcal{H}_{St} , given by the variation of the action in the direction of the unit normal, is just given by (11). In more detail, for $h = T \cdot T$ on \mathcal{T} , one has $h^T = 0$ and $k'_h = 0$, and so, on-shell, one has

$$\mathcal{H}_{\text{St}} = 0.$$

Thus, in this situation, the quasilocal energy is zero. Physically, the simplest and most natural energy of a static vacuum system where one has a preferred timelike Killing field is given by the Komar energy or mass. It is clear that the Komar mass vanishes for finite boundaries $S = \partial \Sigma$ of compact static vacuum domains as above; (vacuum implies there are no interior source terms). Thus, the energy \mathcal{H}_{St} agrees with the Komar energy. However, by the result of Shi-Tam [12], the BY energy of a static vacuum solution is strictly positive (when $K_{\gamma_S} > 0$) unless the solution is flat.

Turning now to the general problem one would like to find a boundary term $B(g, K)$ such that the action

$$\int_M R + \int_{\mathcal{T}} B(\gamma, K) \quad (14)$$

gives a well-defined variational problem, for a choice of boundary data on \mathcal{T} . Next, one would like to find an associated Hamiltonian

$$\mathcal{H}_{(u,X)} = \int_{\Sigma} uC + X^{\mu}C_{\mu} + \int_S B_{(u,X)}(g, \pi), \quad (15)$$

where C and C_{μ} are the Hamiltonian and momentum constraints, respectively. Then, boundary conditions are specified for the variables (g, π) on the phase space T^*Q such that the Hamiltonian is functionally differentiable (smooth as a function on T^*Q) and the Hamiltonian vector field generates the equations of motion $E = 0$, so that one has a well-posed IBVP (in some gauge). Finally, if possible, one would like to select a preferred choice of time-like vector field ∂_t (i.e. a preferred lapse-shift) giving a choice of quasi-Killing field.

Given a choice of lapse-shift (u, X) the EH action (1) decomposes into a time-space integral when the scalar curvature R is expressed in terms of the data (g, π) on the phase space T^*Q ; the spatial integral is then just the integrated constraint operator given by

$$C(u, X) = \int_{\Sigma} u \left(R - \pi^2 + \frac{1}{2}(\text{tr}\pi)^2 \right) - \langle \delta\pi, X \rangle.$$

Integrating the last term by parts gives

$$C(u, X) = \int_{\Sigma} \left[u \left(R - \pi^2 + \frac{1}{2}(\text{tr}\pi)^2 \right) - \langle \pi, \delta^*X \rangle \right] - \int_S \langle \pi(X), \nu \rangle.$$

Now consider the variation of C on the phase space, so with respect to (g, π) . Calculating the variation of R and performing an integration by parts gives rise to the usual Einstein evolution equations in Hamiltonian form in the bulk, together with a boundary term equal to

$$\int_S u \langle \nabla_{\nu} h, g \rangle + u \langle \nu, \delta h \rangle + \langle h, du \cdot \nu \rangle + \langle h, 2X \cdot \pi(\nu) \rangle - \langle h, \pi \rangle \langle \nu, X \rangle + 2 \langle \pi', X \cdot \nu \rangle, \quad (16)$$

cf. [9] for example. Observe that the first three terms involve only the lapse u while the last three terms involve only the shift X . A well-defined variational problem then holds for (14) or (15) with $B = 0$ provided the boundary term (16) vanishes. In general, sum of the variation of B in (15) and (16) must vanish.

The first three terms in (16) can be expressed in terms of the induced metric γ_S on S and the mean curvature H of $S \subset \Sigma$. In fact, (up to signs) the first three terms can be rewritten as

$$\int_S \langle uK - \nu(u)\gamma_S, h^T \rangle + 2uk'_h,$$

exactly as the boundary term in (12). Consider then as configuration space the space of metrics with shift $X = 0$, i.e., the space of metrics of the form

$$g = -u^2 dt^2 + g_{\nu},$$

with $u = u(t, x)$. Imposing the boundary conditions (13) then gives a well-defined variational problem for (14) with $B = 0$, again with zero Hamiltonian on-shell.

This example, as well as the examples discussed previously, give quasilocal Hamiltonians coming from a well-defined variational principle with corresponding boundary conditions. However, in each case the Hamiltonian vector field is not integrable, i.e., the associated flow equation is not generally solvable.

If the Hamiltonian vector field $(\frac{\partial H}{\partial \pi}, -\frac{\partial H}{\partial \gamma})$ is to have a well-defined flow on the phase space T^*Q , then there must exist a gauge choice (and, in particular, a choice of lapse-shift (u, X)) such that the IBVP for the Hamiltonian evolution equations are well posed, i.e., one has existence and uniqueness of solutions with given Cauchy and boundary data, and smooth dependence of the solutions on such data. Further, the IBVP must be geometric in the sense that solutions are isometric if and only if the Cauchy and boundary data differ by the action of diffeomorphisms.

However, as clearly stressed by Friedrich [16], it is currently a basic open problem if in fact there exists a choice of gauge and boundary data such that the IBVP is geometrically well posed in this sense. There is a well-posed formulation of the IBVP first discovered by Friedrich-Nagy [17] and a more recent formulation due to Kreiss-Winicour [18,19]. The results in [18,19], in particular, are naturally formulated in harmonic gauge for the spacetime metric $g_{\alpha\beta}$; however, in both [17] and [18,19] the boundary data imposed are not geometric, but incorporate or assume an extraneous choice of timelike unit vector T along the boundary \mathcal{T} .

To obtain a well-posed geometric IBVP, one expects that it is necessary to choose maximally dissipative boundary conditions on \mathcal{T} , (cf. also [5] for discussion of various boundary conditions). The exact form of these will depend on the choice of gauge (and is currently unknown) but typically such boundary conditions have the schematic form

$$\partial_t g_{ij} + \partial_{\nu} g_{ij} = F_{ij}, \quad (17)$$

where ν is the outward unit normal to $S \subset \Sigma$ and F_{ij} is given. Such boundary conditions are not close to Dirichlet (or Neumann) boundary conditions and so are far from the BY prescription. In fact, no component of the metric γ itself appears in (17). As discussed following (12), the timelike unit normal T to the Cauchy surface Σ is determined only globally, by solving the IBVP for given data (17), and then reading off the value of T at S . Only at that point can the Hamiltonian be actually computed as the unit-time variation of the action. [There is also the issue of finding a boundary term B as in (15) so that boundary conditions analogous to (17) give a well-defined variational problem.]

The notion of quasilocal energy is difficult to make precise since energy is to be defined for an “isolated system” and it not *a priori* clear how to isolate a given region from its surroundings. (This issue bears some resemblance to certain versions of Mach’s principle.) Typically one would impose conditions such as no incoming radiation or absorbing boundary conditions, cf. [20] for example. However, due partly to the general covariance of GR, such boundary conditions are notoriously difficult to identify and implement in practice. Although simple and natural, it is unclear in what manner Dirichlet boundary conditions effectively model isolated physical systems.

There are now several very interesting and useful geometric notions associated to local spacetimes, such as the Hawking mass, the Brown-York energy, the Bel-Robinson energies and many others. These concepts are clearly very useful tools in understanding the physics, geometry and analysis of such spacetimes. However, as discussed above, they do not provide a fully satisfactory notion of quasilocal energy or Hamiltonian.

In closing, two brief remarks. First, the definition of the BY Hamiltonian, as well as its more recent modifications [5–8], requires a choice of subtraction term to normalize the zero-point of the energy. These subtraction terms are

typically determined by choices of isometric embedding of (S, γ_S) into either Euclidean space \mathbb{R}^3 or Minkowski space $\mathbb{R}^{1,3}$. On the other hand, one might hope that a correct choice of gauge and boundary conditions would obviate the need for such subtraction terms (which are somewhat artificial and *ad hoc* given the intrinsic nature of GR).

Finally, the approach of Brown-York is used in the determination of the energy, mass, and other charges for asymptotically anti-de Sitter (AdS) spacetimes in the AdS/CFT correspondence, cf. [21,22], for instance. These concepts, which are of basic importance in the aspects of the correspondence related to thermodynamics of black holes, are global, and the charges are given by suitably renormalized integrals at conformal infinity. The difficulties discussed above in the quasilocal case do not apply in this context where limits at infinity are taken (as in the AF case). For example, in contrast to the finite case, the Dirichlet boundary value problem is well-posed at conformal infinity in the AdS context (for Euclidean metrics); the constraint equations “disappear” in the limit at infinity as restrictions on the form of the conformal metric at infinity.

This work is partially supported by NSF Grant No. DMS 0905159).

-
- [1] R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation, An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962), p. 227–265.
 - [2] L. B. Szabados, *Living Rev. Relativity* **12**, 4 (2004), <http://relativity.livingreviews.org>.
 - [3] T. Regge and C. Teitelboim, *Ann. Phys. (N.Y.)* **88**, 286 (1974).
 - [4] J. D. Brown and J. W. York, Jr., *Phys. Rev. D* **47**, 1407 (1993).
 - [5] J. Kijowski, *Gen. Relativ. Gravit.* **29**, 307 (1997).
 - [6] R. J. Epp, *Phys. Rev. D* **62**, 124018 (2000).
 - [7] C.-C.M. Liu and S.-T. Yau, *Phys. Rev. Lett.* **90**, 231102 (2003).
 - [8] M.-T. Wang and S.-T. Yau, *Phys. Rev. Lett.* **102**, 021101 (2009).
 - [9] L. B. Szabados, *Classical Quantum Gravity* **26**, 125013 (2009).
 - [10] G. W. Gibbons and S. W. Hawking, *Phys. Rev. D* **15**, 2752 (1977).
 - [11] J. W. York Jr., *Found. Phys.* **16**, 249 (1986).
 - [12] Y. Shi and L.-F. Tam, *J. Diff. Geom.* **62**, 79 (2002).
 - [13] M. T. Anderson, *Geom. Topol.* **12**, 2009 (2008).
 - [14] M. T. Anderson and M. A. Khuri, [arXiv:1003.2803](https://arxiv.org/abs/1003.2803).
 - [15] M. T. Anderson and M. A. Khuri, [arXiv:0909.4550](https://arxiv.org/abs/0909.4550).
 - [16] H. Friedrich, *Gen. Relativ. Gravit.* **41**, 1947 (2009).
 - [17] H. Friedrich and G. Nagy, *Commun. Math. Phys.* **201**, 619 (1999).
 - [18] H.-O. Kreiss and J. Winicour, *Classical Quantum Gravity* **23**, S405 (2006).
 - [19] H.-O. Kreiss, O. Reula, O. Sarbach, and J. Winicour, *Commun. Math. Phys.* **289**, 1099 (2009).
 - [20] O. Sarbach, *J. Phys. Conf. Ser.* **91**, 012005 (2007).
 - [21] S. de Haro, K. Skenderis, and S. Solodukhin, *Commun. Math. Phys.* **217**, 595 (2001).
 - [22] V. Balasubramanian and P. Kraus, *Commun. Math. Phys.* **208**, 413 (1999).