Static $\bar{Q}Q$ potentials and the magnetic component of QCD plasma near T_c

Jinfeng Liao^{1,2,*} and Edward Shuryak^{3,†}

¹Physics Department, Brookhaven National Laboratory, Upton, New York 11973, USA

²Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

³Department of Physics and Astronomy, State University of New York, Stony Brook, New York 11794, USA (Received 9 August 2010; published 10 November 2010)

Static quark-antiquark potential encodes important information on the chromodynamical interaction between color charges, and recent lattice results show its very nontrivial behavior near the deconfinement temperature T_c . In this paper we study such potential in the framework of the "magnetic scenario" for the near T_c QCD plasma, and particularly focus on the linear part (as quantified by its slope, the tension) in the potential as well as the strong splitting between the free energy and internal energy. By using an analytic "ellipsoidal bag" model, we will quantitatively relate the free energy tension to the magnetic condensate density and relate the internal energy tension to the thermal monopole density. By converting the lattice results for static potential into density for thermal monopoles we find the density to be very large around T_c and indicate at quantum coherence, in good agreement with direct lattice calculation of such density. A few important consequences for heavy ion collisions phenomenology will also be discussed.

DOI: 10.1103/PhysRevD.82.094007

PACS numbers: 12.38.Mh, 12.38.Gc, 25.75.Nq

I. INTRODUCTION

A traditional observable for studying the QCD confinement is the interaction potential between static quark and antiquark. It was originally inferred from heavy meson spectrum and Regge trajectories, and has then been extensively studied in lattice gauge theories. (For reviews see e.g. [1,2].) Its vacuum (T = 0) form is well-known, usually represented as a sum of a Coulomb part $V \sim 1/r$, dominant at small separation between $Q\bar{Q}$, and a linear part $V = \sigma r$ dominant at large separation (see the black solid curve in Fig. 1). The latter implies the confinement of quarks and has been interpreted in terms of chromo-electric flux tube (or "string") formation between well-separated QQ pair. The so-called string tension σ in the vacuum (T = 0) has been consistently determined by different methods to be

$$\sigma_{\rm vac} \approx (426 \text{ MeV})^2 \approx 0.92 \text{ GeV/fm.}$$
 (1)

With current Relativistic Heavy Ion Collider (RHIC) and future LHC experimental programs exploring excited hadronic matter and quark-gluon plasma (QGP) at increasing temperature T, it is very important to know the finite Tform of the static QQ potential, which has recently been calculated by means of the lattice QCD. (See e.g. [3–5].) At finite temperature, there are actually two potentials associated with a QQ pair separated by distance r: one is the free energy F(T, r) and the other is the internal energy V(T, r), with the difference related to the entropy generated in the medium by the $\bar{Q}Q$ pair, i.e.

$$V(T,r) = F(T,r) - T(\partial F/\partial T) = F(T,r) + TS(T,r).$$
(2)

What is directly evaluated on the lattice is the free energy F(T, r) from which the corresponding V(T, r) and S(T, r)can be inferred [3]. While at T = 0 there is no entropy and the free and internal energies are identical, splitting between the two shall be expected at T > 0 and may carry key information about the medium and deconfinement transition near T_c .

The lattice results indeed show remarkably different potentials F(T, r) and V(T, r) near T_c (see e.g. Fig. 1–4 in [6] and also here Fig. 1 adapted from [3]). In particular let us emphasize two important points. (i) The tensions (slopes of the potentials at r about 0.3-1 fm) have very different temperature dependences: while the tension of the



FIG. 1 (color online). The static $\bar{Q}Q$ potential at $T \approx T_c$ (adapted from [3]). The boxes (blue) are for the internal energy V(r) while the diamonds (red) are for free energy F(r), with the dashed line (green) indicating the strong linear rise in V(r) for $r \in (0.5, 1)$ fm and the solid curve (black) showing the vacuum $\bar{Q}Q$ potential.

^{*}jliao@bnl.gov

shuryak@tonic.physics.sunysb.edu

free energy σ_F decreases with *T*, to near zero at T_c (an expected signal of deconfinement), the tension of the internal energy σ_V remains nonzero till about $T = 1.3T_c$, with a peak value at T_c about 5 times (!) the vacuum tension σ_{vac} (see Fig. 2). (ii) This drastically different behavior persists to very large distances, where linear behavior changes to saturated values. Near T_c the internal energy flattens to a huge asymptotic value at large $r \rightarrow \infty$, e.g. $V(T, \infty) \sim 4$ GeV at T_c with the corresponding entropy $S(T_c, \infty) \approx 20$ implying huge number of states involved, $\sim \exp(20)$. These features indicate a strikingly strong interaction between the static color charges and the medium near T_c , which persists into the deconfined phase.

Such static $\bar{Q}Q$ potentials at finite T are closely connected with a number of phenomenological issues. For example, the consequence of these features for the survival of quarkonium in deconfined plasma is much debated, e.g. on what/which potential should be used [7–9]. If, as suggested in [10,11], the internal energy is used, J/ψ state would exist even in the deconfined plasma in $1-2T_c$. Persistence of some baryonic states above T_c is also indicated by other observable like the baryonic susceptibilities [12,13]. These potentials also imply significant interaction energy in the quark-gluon plasma and in the many body context this may lead to a large classical plasma parameter Γ (defined as the ratio of average interaction energy to average kinetic energy): indeed the Γ value in strongly coupled QGP has been estimated to be above one (about three) and thus in a typical liquid regime (see for example [14-16]). If so, OGP would be a strongly coupled Coulombic liquid, in agreement with the strong collective flow observed at RHIC (see more in reviews [17,18]). Apart from QGP phenomenology, it is important to understand the microscopic origin of the potentials, especially the strong splitting between two potentials and the large energy/entropy associated with the static $\bar{Q}Q$ pair near T_c . Earlier attempts can be found in e.g. [19–21].



FIG. 2 (color online). Effective string tensions in the free energy $\sigma_F(T)$ (from [4]) and the internal energy $\sigma_V(T)$ (extracted from [3]).

In this paper we will specifically focus on the "tensions" σ_F and σ_V (as shown in Fig. 2) related to the linear part of the potentials (while leaving the discussion of "screening" behavior at very large distances to further studies). We will provide an explanation in the framework of the *magnetic* scenario of QCD plasma near T_c [15,22–25]. In such a scenario, the near T_c QCD plasma is strongly influenced by the magnetic component, made of relatively light and abundant chromo-magnetic monopoles. Those are quasiparticles above T_c which undergo the Bose-Einstein condensation (BEC) below T_c , enforcing color confinement (for reviews see e.g. [26]). Two key points of the present model for the potentials are: First, we identify the $\bar{Q}Q$ free energy as been probed by an adiabatically "slow separation" process while the internal energy by a "fast separation" process. Second, we further relate the linear part of potentials with the flux tube formation, enabled by condensed monopoles below T_c while thermal monopoles above T_c , between the $Q\bar{Q}$ pair during the separation process [27], and relate the free/internal energy tensions with the condensed/thermal monopoles, respectively. These ideas will be elaborated more in Sec. II and III.

The rest of the paper is structured as follows. In Sec. IV we will develop an analytic "elliptic flux bag" model for a static charge-anti-charge pair by solving the Laplace equation for electric field inside it. This allows to get the potentials correctly interpolating between a Coulomb at short distance and linear behavior at larger distance. The model will then be used in Sec. V to determine the free and potential energies and relate the extracted $\sigma_F(T)$ and $\sigma_V(T)$ with the monopole condensate and the thermal monopole density, respectively. Finally we summarize the results in Sec. VI.

II. FREE VS INTERNAL ENERGY AND SLOW VS FAST SEPARATION

Let us start by examining the difference between the free energy and the internal energy. We already introduced the effective string tensions $\sigma_F(T)$ and $\sigma_V(T)$ as the slopes of linear parts in F(T, r) and V(T, r) respectively, and emphasized their quite different T dependencies shown in Fig. 2. While σ_F vanishes at $T > T_c$, σ_V survives to at least $1.3T_c$. While σ_F monotonously decreases with T, σ_V peaks at T_c to a maximal value of 5 times the vacuum string tension σ_{vac} . What is the difference in the meaning of F and V, and why do they have such different T dependence? As has been emphasized in [10], the free and internal energies actually correspond to slow and fast (relative) motion of the charges, respectively. Let us explain this idea in more details.

Consider the system made of the medium as a thermal bath and the static $\overline{Q}Q$ pair separated by a distance L. Because of interaction between the medium particles and the static pair, the medium's various eigentstates have their energies depending on the distance L. When the medium is in equilibrium these states should be populated according to thermal distributions. This means at different L, the thermal equilibrium requires different population of medium's eigenstates. Now suppose the pair of static charges (held by external "hands") are moved apart in thermal medium at certain speed $v = \dot{L}$. Clearly when L is changed, the medium has to rearrange the population of various eigenstates in order to reach again the thermal equilibrium and that can happen only on the time scale of the medium relaxation time. If the motion of static charges is adiabatically slow, then all the processes for rearrangement of population will happen; in thermodynamical context this leads to maintained equilibrium and maximal entropy/heat generation. If however the pair is separated very fast, then the time is just not enough for the medium to "adjust" the eigenstates' distribution and the medium can no longer stay in equilibrium; in thermodynamical context this means the medium does not yet maximize its total entropy for the given new L, and during the course of such fast separation the amount of entropy generated is therefore less than in the adiabatic case. In the extreme case of the pair separation on a time scale much much shorter than the medium relaxation time scale, one may expect negligible entropy generation.

In summary, the essence of the above arguments is about the separation of time scales between the probing time scale versus the medium relaxation time scale. In the slow case the medium is allowed ample time to repopulate various states according to equilibrium distribution and achieve maximal entropy, while in the fast case with the needed processes for repopulating the medium *not* occurring the medium no longer maintains thermal equilibration and generates negligible entropy. It is plausible, therefore, to identify the adiabatic limit as probing the free energy F(T, L) measured on the lattice with the presence of static $\overline{Q}Q$ pair. The "internal energy" V(T, L), on the other hand, is different from F(T, L) by subtracting the entropy term and thus can be probed in the extremely fast limit.

III. STABLE AND META-STABLE FLUX TUBES

We now turn to the possible microscopic origin of the linear rise in both potentials. Let us start with the "dual superconductor" model for QCD confinement in the vacuum, introduced by t'Hooft-Mandelstam [28] and well supported by extensive studies in lattice QCD. In this model, certain "magnetically charged" condensate (i.e. a magnetic superconductor) occupies the vacuum and expels the electric flux between QQ into a stable flux tube by forming magnetic supercurrent on tube surface, known as (dual) Meissner effect. Such flux tube naturally gives rise to a linear potential and the vacuum string tension is thus identified with energy per unit length of such flux tube, mathematically described by the well-known Abrikosov-Nielsen-Olesen (ANO) solution [29] (for reviews and further references see e.g. [1,2,30,31]). What

happens at finite T then? With increasing T, the free energy tension decreases and eventually the linear part in $\overline{Q}Q$ free energy disappears at T_c , signaling the deconfinement transition. Since the flux tube and free energy tension is a direct consequence of the magnetic condensate, the decrease of σ_F toward T_c is naturally interpreted as the gradual "melting" of the magnetic condensate due to thermal excitations; similar phenomena is known for the usual superconductor in condensed matter systems.

Now, where does the linear part in internal energy (and the associated tension σ_V) come from? In particular, why does it persist even above T_c ? The answer first proposed in [15] relates it to the "normal" monopoles, as opposed to the "supercomponent" which form a Bose-Einstein Condensation (BEC) and exist only below T_c . Such thermal monopoles can also expel the electric flux into the (meta-stable) flux tube by forming a magnetic current (which may suffer from dissipation) on the tube surface; its dual phenomenon, i.e. magnetic flux tube formation in thermal electron plasma is well-known in classical (e.g. solar) plasma physics. The specific condition for the persistence of the electric flux tube in a magnetic plasma was further developed in [27], for infinitely long flux tubes. There it has been found that "normal" monopoles are much less effective for this task as compared with "super" monopoles, but nevertheless are able to mechanically stabilize the flux tube provided high enough density of these thermal monopoles. What has not been previously considered is the mechanism for the dynamical formation of the flux tube between a $\bar{Q}Q$ pair with finite separation.

Here we provide a dynamical explanation of why large energy, growing approximately linearly with length, appears in a magnetic plasma when a pair of two electric charges are separated with certain speed v. (See sketch of the setting in Fig. 3.) The answer lies in the Maxwell equations (with the presence of magnetic sources, see e.g. [32]), in particularly the *dual* Faraday's law which relates the circulation of the magnetic field $\int \vec{B} \, dl$ over a closed contour with the change of electric flux penetrating the enclosed area. As an electric charge moves through the loop, rotating magnetic field in the magnetic medium leads to solenoidal magnetic current (a "magnetic coil"). In the confined phase $T < T_c$ this current, after relaxation, becomes the persistent supercurrent, remaining forever without loss. Thus, the free energy F has a linear term for $T < T_c$. In a deconfined plasma phase $T > T_c$ this is impossible, thus $\sigma_F = 0$. The solenoidal "magnetic coil" created in the fast process has only normal magnetic current, which is a meta-stable flux tube and eventually disappears due to dissipation. Yet it is still generated; thus $\sigma_V - \sigma_F$ is nonzero and there is splitting between free and internal energy.

Let us emphasize again the different roles of the *super* and *normal* magnetic components. The former responds quantum mechanically as a whole and does not generate



FIG. 3 (color online). Schematic demonstration of magnetic solenoidal by Dural Faraday's law, see text.

any entropy nor contribute to the splitting. The latter, however, has finite relaxation time and nonzero dissipation, "feels" the different time scales involved in the slow/ fast processes, and therefore is responsible for entropy generation and the splitting between free and internal energy. In short, the σ_F tells us about the super component only, while the difference $in\sigma_V - \sigma_F$ tells us about the normal component. One arrives at the following picture for an evolving magnetic medium: with increasing T the monopole ensemble starts as a monopole condensate and continuously evaporates into a mixture of both condensed and thermal monopoles; at $T > T_c$ the condensate melts entirely into a normal component of thermally excited monopoles. If so, the thermal monopoles are expected to be most important in the temperature range $0.8-1.3T_{c}$ where the splitting is most significant.

We end this Sec. with discussions on a few important phenomenological implications of the "dual Faraday effect" and the meta-stable flux tube. First, it means that magnetic monopoles may induce new mechanism of (electric) jet energy loss particularly near T_c . In the jet quenching process, a very *fast* electric parton (quark or gluon) penetrates the bulk medium through various phases and thus may create behind it the above discussed "magnetic coil" in the near T_c region where there are abundant thermal monopoles to be accelerated solenoidally by *B* field due to the fast moving electric jet and thus take enormous amount of energy away from the jet. Such near T_c enhancement of jet quenching has been first suggested by us in [33] and found to be strongly favored by the azimuthal anisotropy data of jet quenching. Second, it also implies specific patterns of multiparticle correlations if such flux tubes can be created and protected by monopoles in heavy ion collisions, as elaborated first in [34]. One example is related to what happens to the flux tube created by a fast jet: clearly the monopoles forming the "coil" will subsequently collide with the bulk thermal matter, with their energy being converted and distributed into the bigger volume. This may possibly be the beginning of "conical flow" process suggested in [35]. The other example concerns the narrow-azimuthal-angle long-rangerapidity correlations known as "ridge" which seems to originate from certain local initial fluctuation seeds, but its narrowness in angle may possibly be preserved till the end of long bulk evolution only if certain mechanism like the flux tube by the thermal monopoles protects the initial seed from acoustic expansion (see detailed discussions in [34]). The existence of meta-stable flux tubes in the near T_c plasma (and their associated large entropy) may also bear relevance to the observed cluster correlations [36]. While the existence and dynamical formation of such flux tubes are studied in [27] and here, another very important question (particularly for phenomenology) is its lifetime, i.e. the flux tube decay and the end products. This problem has recently been partially addressed in [37] where a relatively short lifetime is found in the classical treatment. On one hand, from hydrodynamic modeling we know the near T_c plasma has very small shear viscosity which indicates short mean free path and frequent scattering, while on the other hand for the magnetic currents to last long and hold the flux tube they better not scatter too often. Such a dilemma might be resolved if the thermal monopoles become really coherent over a large distance at T close to T_c , and the lattice study supports such coherence [38]. These questions will be studied further elsewhere.

IV. ELECTRIC FIELD SOLUTION IN THE ELLIPSOIDAL BAG

In this Sec. we will solve the Maxwell equation for the electric field induced by a static charge-anti-charge pair separated by a distance L = 2a along \hat{z} axis ($\pm Q_e$ sitting at $\mp a\hat{z}$), with a special "tangent boundary condition" (T.B.C.) on the boundary surface Σ_B , i.e.

$$\vec{\nabla}^2 \Phi(\mathbf{r}) = Q_e [\delta^3 (\mathbf{r} - a\hat{\mathbf{z}}) - \delta^3 (\mathbf{r} + a\hat{\mathbf{z}})]$$
$$\vec{\nabla} \Phi \cdot \hat{\mathbf{n}}_{\Sigma_B}|_{\Sigma_B} = 0.$$
(3)

The model itself is a version of an old idea known as the Bag Model used for light hadrons [39] at T = 0, now generalized to give an approximate description of the

STATIC $\bar{Q}Q$ POTENTIALS AND THE MAGNETIC ...

electric field configuration between static $\overline{Q}Q$ in the chromo-magnetic medium at finite temperature.

A simplification we use is that the boundary Σ_B is approximated by a rotational ellipsoid with the two charges at its focal points. This boundary shape can be specified by a single parameter ξ_B , the ellipticity. Such boundary Σ_B is very conveniently parameterized in terms of the parabolic coordinates system (ξ, η, ϕ) , which we use (see the Appendix for necessary formulas related to it). In Fig. 4 we show a few ellipsoidal shapes with parameters (from inside to outside) (L, ξ_B) to be (0.1, 6.62), (1, 1.68), (2, 1.29), (3, 1.16), respectively. The dashed lines indicate constant η curves (for L = 3 case) with (from top to bottom) $\eta = 0.8$, 0.5, 0.2, -0.2, -0.5, -0.8, the solid/ empty circles indicate the positions of positive/negative charges, and the arrows indicate the tangent electric fields on the boundary.

We follow the standard method in classical electrostatics (see for example [32]). First, we rewrite (3) in (ξ, η, ϕ) coordinates for solutions with axial symmetry, i.e. assuming $\Phi = \Phi(\xi, \eta)$ independent of angle ϕ :

$$\frac{\partial}{\partial \xi} \left[(\xi^2 - 1) \frac{\partial \Phi}{\partial \xi} \right] + \frac{\partial}{\partial \eta} \left[(1 - \eta^2) \frac{\partial \Phi}{\partial \eta} \right]$$

$$= \frac{Q_e \delta(\xi - 1)}{\pi L} \left[\delta(\eta - 1) - \delta(\eta + 1) \right]$$

$$= \sum_{\nu = 1,3,5,\dots} \frac{Q_e \delta(\xi - 1)}{\pi L} (2\nu + 1) P_{\nu}[\eta]. \quad (4)$$

The last line in the above is an expansion of the η dependence in terms of Legendre functions $P_{\nu}[\eta]$, which in the interval $\eta \in [-1, 1]$ form a set of orthogonal and complete basis functions. Similarly, we do the expansion for the η dependence of Φ :



$$\Phi_{\xi,\eta} = \sum_{\nu=1,3,5,\dots} \frac{Q_e f_\nu [\xi]}{\pi L} (2\nu + 1) P_\nu[\eta].$$
(5)

Then by simply comparing the coefficients of $P_{\nu}[\eta]$ on both sides of Eq. (4) we obtain the equations for the functions $f_{\nu}[\xi]$ defined in $\xi \in (1, \infty)$:

$$\frac{d}{d\xi} \left[(1 - \xi^2) \frac{df_{\nu}}{d\xi} \right] + \nu(\nu + 1) f_{\nu} = -\delta(\xi - 1), \quad (6)$$

while the boundary condition in Eq. (3) now becomes

$$f'[\xi = \xi_B] = 0 \tag{7}$$

with the parameter ξ_B specifying the boundary surface Σ_B . The solutions are given in terms of the Legendre functions of the first and second kinds:

$$f_{\nu}[\xi] = -k_{\nu}^{B} P_{\nu}[\xi] - Q_{\nu}[\xi]$$

$$k_{\nu}^{B} = -\frac{Q_{\nu}'[\xi_{B}]}{P_{\nu}'[\xi_{B}]} = -\frac{\xi_{B} Q_{\nu}[\xi_{B}] - Q_{\nu-1}[\xi_{B}]}{\xi_{B} P_{\nu}[\xi_{B}] - P_{\nu-1}[\xi_{B}]}.$$
 (8)

The full electrostatic potential is then given by

$$\Phi(\vec{r}|L,\xi_B) = -\frac{Q_e}{4\pi L} \sum_{\nu=1,3,5,\dots} \times (8\nu+4)P_{\nu}[\eta](k_{\nu}^{B}P_{\nu}[\xi] + Q_{\nu}[\xi])$$
$$= \frac{Q_e}{4\pi L} \frac{2}{\xi+\eta} + \frac{(-Q_e)}{4\pi L} \frac{2}{\xi-\eta} - \frac{Q_e}{4\pi L}$$
$$\times \sum_{\nu=1,3,5,\dots} (8\nu+4)k_{\nu}^{B}P_{\nu}[\xi]P_{\nu}[\eta].$$
(9)

We have used the Neumann expansion of Legendre functions (see e.g. [40]) to write down the second equality; in there the first two terms are nothing but the usual Coulomb potentials by the $\pm Q_e$ charges, while the last summation term reflects the nontrivial boundary contribution. At very large ν the summand terms go asymptotically like $\nu \xi^{\nu} / \xi_B^{2\nu+2}$, so with ξ satisfying $1 < \xi \leq \xi_B$, the summation is guaranteed to converge. The electric field $\vec{E} = -\vec{\nabla}\Phi$ has been calculated using (A4) and the expression is quite lengthy, which we skip showing here.

The volume occupied by the electric field (i.e. the ellipsoid bulk within ξ_B) is given by

$$V_E(L, \xi_B) = \int_1^{\xi_B} d\xi \int_{-1}^1 d\eta \int_0^{2\pi} d\phi H_{\xi} H_{\eta} H_{\phi}$$
$$= \frac{\pi L^3}{6} \xi_B(\xi_B^2 - 1).$$
(10)

FIG. 4 (color online). The ellipsoidal shapes we use for solving the electric field equations, see text for detailed explanations.

And the total electric field energy in this volume is given by

$$\mathcal{E}_{\text{total}}(L, \xi_B) = \int_1^{\xi_B} d\xi \int_{-1}^1 d\eta \int_0^{2\pi} d\phi H_{\xi} H_{\eta} H_{\phi}$$

$$\times \frac{\rho_e \times \Phi(\xi, \eta)}{2} = \mathcal{E}_{\text{self}} + \mathcal{E}_E$$

$$\mathcal{E}_{\text{self}} = \frac{Q_e^2}{4\pi L} \frac{1}{(\xi + \eta) \to 0} + \frac{Q_e^2}{4\pi L} \frac{1}{(\xi - \eta) \to 0}$$

$$\mathcal{E}_E = -\frac{Q_e^2}{4\pi L} + \frac{Q_e^2}{4\pi L} \sum_{\nu=1,3,5,\dots} (8\nu + 4) k_{\nu}^B$$

$$\equiv \frac{Q_e^2}{4\pi L} \bar{\mathcal{E}}_E(\xi_B). \tag{11}$$

The \mathcal{E}_{self} is the familiar self-interaction of the two charges which we discard. The "real" interactional energy \mathcal{E}_E consists (again) of a Coulomb piece and a boundary modification.

We conclude this Sec. by one remark: so far the two key variables L and ξ_B remain free parameters. They will be related in the next Sec..

V. THE FREE AND INTERNAL ENERGY OF THE CHARGE PAIR

With the solutions of the electric field in the ellipsoidal bag (characterized by two parameters L and ξ_B) from the preceding Sec., we now examine the dynamic formation of such bag when separating a pair of QQ from zero to a finite distance L. The key point is that for a given L, the bag boundary ξ_B shall be optimized so that the "cost" for creating such a configuration is minimized. Furthermore, we study two settings: slow and fast separation of the QQto a finite distance L, with the outcome being, respectively, the free and internal energy associated with the pair. With slow separation, the free energy increase associated with the pair shall be minimized, and the dominant contributions to the free energy include both the electric field energy stored inside the bag and the energy needed to exclude the monopole condensate out of the bag volume (noting that for both there is no entropy associated and for thermal monopoles their contribution to free energy in the slow separation process largely cancels out between energy and entropy). With fast separation, the energy increase shall be minimized, and the dominant contributions to the energy include both the electric field energy stored inside the bag and the energy deposited to the thermal monopoles via the dual Faraday effect. We will calculate both processes in the rest of this Sec. and make connections with the lattice data.

A. Free energy from slow separation

As aforementioned, when the $Q\bar{Q}$ pair is separated in an adiabatically slow way, the *super* component of the magnetic medium, i.e. the monopole condensate, will be expelled entirely (in an idealized picture) out of the volume V_E occupied by electric field. Suppose the condensate has a

negative energy density $-\epsilon_C$ (thus a positive "bag pressure"), then the overall change in free energy brought about by separating the pair will be

$$\Delta F = \mathcal{E}_E(L, \xi_B) + \epsilon_C(T) \cdot V_E(L, \xi_B).$$
(12)

Now for given separation distance *L* and bulk temperature *T*, we determine the physical boundary of flux bag ξ_B^{phy} by minimizing the above ΔF , i.e. the physical boundary $\xi_B^{\text{phy}}(L, T)$, satisfies

$$\frac{\partial \Delta F}{\partial \xi_B} \bigg|_{\xi_B = \xi_B^{\text{phy}}} = 0.$$
(13)

Combining the above with Eq. (10) and (11) we then obtain

$$\left[\frac{1}{3\xi_B^2 - 1}\frac{d\bar{\mathcal{E}}_E}{d\xi_B}\right]\Big|_{\xi_B = \xi_B^{\rm phy}} = -\left(\frac{L}{l_C}\right)^4,\tag{14}$$

where we have introduced a length scale

$$l_C \equiv (6\alpha_E/\pi\epsilon_C)^{1/4} \tag{15}$$

with $\alpha_E \equiv Q_e^2/4\pi$. This equation could be solved easily by numerics. For each L with the above determined ξ_{R}^{phy} , we obtain via (12) the free energy associated with the pair as a function of separation L, shown in Fig. 5(a). It turns out to be a Coulomb at short distance (see the dashed curve (magenta)) plus linear at large distance (see the dashed line (blue)). The occurrence of a linear part is due to the physical effect that for large L the medium pressure (with which the electric field has to balance) limits the transverse size of flux bag (where the field gets weak as L increases) to saturate rather than grow forever. Thus, the bag shape approaches a cylinder. Mathematically, as $L \rightarrow \infty$ one finds $\xi_B^{\text{phy}} \to 1$ but $L \cdot \sqrt{(\xi_B^{\text{phy}})^2 - 1} \to \text{finite. In Fig. 4}$ the four bag shapes are at growing L with ξ_B determined as in the above, which clearly shows the shape becomes more and more cylindrical at large L.

By fitting the dimensionless slop of the linear part in Fig. 5(a) we obtain the free energy string tension σ_F :

$$\sqrt{\sigma_F} = 2.32 \times \alpha_E^{1/4} \times \epsilon_C^{1/4}.$$
 (16)

(See [41] for a note about the numerical coefficient). Inversely, since we know $\sigma_F(T)$ from lattice as shown in Fig. 2, from the above formula we can infer the *T* dependence of the monopole condensate energy density ϵ_C (see Fig. 5(b)). The two curves are for α_E being 0.5 (upper) and 1 (lower), respectively. In both cases, ϵ_C decreases with *T* and drops abruptly close to T_c . The interpretation is natural: toward T_c the monopole condensate becomes less and less due to increasing thermal excitations and eventually dies out around T_c .

A connection can be made between our result (16) and the dual superconductor model (also known as Abelian Higgs model) of vacuum confinement [30]. In that model,



FIG. 5 (color online). (a) (Left) free energy F (in unit of α_E/l_C) versus separation L/l_C ; (b) (Right) monopole condensate energy density (ϵ_C)^{1/4} in unit of $\sqrt{\sigma_{\text{vac}}}$ with the two curves for α_E being 0.5 (upper, red) and 1 (lower, blue) in Eq. (16), respectively.

a quadratic Higgs potential leads to a Higgs condensate (the prototype of postulated monopole condensate) ϕ_0 (with dimension of mass). By solving ANO flux tube a string tension is obtained in the form $\sqrt{\sigma} = c_1 \phi_0$ with the coefficient determined by gauge and Higgs coupling constants λ and g. On the other hand, the Higgs potential implies that the condensate has a negative energy density $-\epsilon_C = -\lambda \phi_0^4/2$. Thus, one arrives at a similar relation between string tension and condensate energy density: $\sqrt{\sigma} = c_2 \epsilon_C^{1/4}$ in that model with the coefficient to be determined numerically for given coupling parameters (see e.g. [42]). While that model works primarily at T = 0, our model for σ_F extends to finite T.

B. Internal energy from fast separation

Now we study the case of separating the two charges to a finite distance L within a time much smaller than the relaxation time of the surrounding thermal bath. In particular we focus on the region about $0.8-1.3T_c$, in which the *normal* component of thermal monopoles becomes substantial and dominant while the *super* component becomes less and less.

During such fast process, each monopole originally in the volume to be occupied by the electric filed (i.e. the ellipsoidal bag) will get a "kick" due to the dual Faraday effect (see Fig. 3) but have no time to release this energy into the surrounding medium. Suppose the positive charge is moved along \hat{z} axis from z = C to $z = C + \delta z$ in δt (and correspondingly the negative one from z = -C to z = $-(C + \delta z)$), then the electric flux penetrating the plane z = C changes from 0 to Q_e , thus generating a magnetic dynamical voltage $Q_e/\delta t$. For a monopole at a transverse distance ρ from \hat{z} axis, the force is $Q_m(Q_e/\delta t)/(2\pi\rho)$, thus it gets the "kick" and obtains a momentum $\delta p =$ $Q_m Q_e / (2\pi\rho)$, forming strong nonthermal and nonsuper magnetic currents. For a bag (L, ξ_B) formed after separation, the total kinetic energy passed to the monopoles in the flux bag is obtained by integration over the bag volume (with $D \equiv Q_m Q_e / 4\pi = 1$):

$$\Delta K_{M} = \int_{1}^{\xi_{B}} d\xi \int_{-1}^{1} d\eta \int_{0}^{2\pi} d\phi H_{\xi} H_{\eta} H_{\phi}$$
$$\times \frac{4n_{M}D}{L\sqrt{(\xi^{2}-1)(1-\eta^{2})}}$$
$$= \frac{\pi^{2}Dn_{M}}{2}L^{2}\xi_{B}(\xi_{B}^{2}-1)^{1/2}.$$
(17)

We emphasize in the above only the monopole density $n_M(T)$ enters as a property of the medium depending on T, while other equilibrium properties of the medium shall not be "felt" in such fast process. In obtaining the above, we have estimated the acquired kinetic energy by simply using the momentum kick which is a reasonable approximation for sufficiently light monopoles, expected to be the case in the near T_c region [15,38].

Now the total energy change during such process includes the electric field energy in the bag volume, the energy for expelling the monopole condensate (if there is any) out of the volume, and the kinetic energy delivered to the normal monopoles, which are summed to be $\Delta E = \mathcal{E}_E(L, \xi_B) + \epsilon_C V_E + \Delta K_M(L, \xi_B)$. The last new term due to the thermal monopoles will then partly convert into entropy after they interact with the medium particles at large for a time longer than the relaxation, ultimately causing the splitting between free/internal energy. Since the condensate term $\epsilon_C V_E$ becomes very small close to T_c (as we showed in the previous subsection) and vanishes above T_c , we neglect it here for simplicity, i.e. $\Delta E \approx \mathcal{E}_E(L, \xi_B) + \Delta K_M(L, \xi_B)$. To obtain the physical value ξ_B^{phy} , we need to minimize ΔE according to ξ_B , which leads to

$$\frac{\partial \Delta E}{\partial \xi_B} \bigg|_{\xi_B = \xi_B^{\text{phy}}} = 0.$$
(18)

This can be written as

$$\left[\frac{\sqrt{\xi_B^2 - 1}}{2\xi_B^2 - 1} \frac{d\bar{\mathcal{E}}_E}{d\xi_B}\right] \bigg|_{\xi_B = \xi_B^{\text{phy}}} = -\left(\frac{L}{l_M}\right)^3.$$
(19)



FIG. 6 (color online). (a) (Left) internal energy V (in unit of α_E/l_M) versus separation L/l_M ; (b) (Right) thermal monopole density n_M/T^3 with the two curves connecting box symbols for α_E being 0.5 (upper, red) and 1 (lower, blue) in Eq. (20), respectively, while the two dashed curves (green) connecting diamond symbols showing SU(2) (lower) lattice data and their SU(3) extrapolation (upper) for $T > 1.3T_c$ from [25] (see text for more details).

Here we introduced a different length scale $l_M \equiv (2\alpha_E/\pi^2 Dn_M)^{1/3}$. Not surprisingly we find the internal energy, shown in Fig. 6(a), to be a Coulomb at short distance (see the dashed curve (magenta)) plus linear at large distance (see the dashed line (blue)).

Now the string tension in the internal energy is given by the following formula:

$$\sqrt{\sigma_V} = 3.88 \times \alpha_E^{1/6} \times n_M^{1/3}.$$
 (20)

(See [41] for a note about the numerical coefficient). Since we know σ_V from lattice data in the 0.8–1.3 T_c region, by the above formula we can convert σV into the thermal monopole density $n_M(T)$ in the same region; see the two curves (connecting box symbols) for α_E being 0.5 (upper, red) and 1 (lower, blue), respectively, in Fig. 6(b). We also show an independent information on the thermal monopole density above $1.3T_c$ from lattice study in [25] in Fig. 6(b) as green curves (connecting diamond symbols): the lower one is the original data for SU(2) in [25], while the upper one is an extrapolation to SU(3) by the simple $N_c - 1$ scaling for monopole species (i.e. twice more monopoles in SU(3) than in SU(2), with both curves extended toward T_c according to the fitting formula $n_m/T^3 = 0.48/\log(2.48 \cdot T/T_c)^{1.89}$ (and twice in the upper one for SU(3) in [25]. The comparison shows reasonably good agreement between our estimates for the thermal monopole density from string tension and the measured density by directly identifying the thermal monopoles on the lattice.

A few comments are in order: (i) for $0.8-1T_c$ the density quickly grows toward T_c while at the same time results from previous subsection show rapid dropping of condensate density in the same region, which strongly indicates the scenario that close to T_c monopole condensate is continuously and substantially getting excited into thermal monopoles; (ii) around $1.3T_c$ we see our results connect well to the higher T lattice data with reasonable values of coupling α_E ; (iii) cooling down to T_c we find the monopole density n_M/T^3 quickly rising almost by an order of magnitude; (iv) the strongly increasing density also suggests rapid increase of magnetic screening toward T_c , which is in agreement with lattice results [43].

A particularly interesting feature is that the (normalized) density n_M/T^3 increases roughly by 1 order of magnitude from $1.3T_c$ down to T_c , with the number much larger than even a Stefan-Boltzman gas. This indicates that near T_c the monopoles should be very light, and furthermore their interactions should make it beneficial in energy to have a large number of monopole-anti-monopole pairs. The monopoles are so dense and light that they become the dominant component in the near T_c plasma and presumably become quantum coherent, and eventually reach the condensation point at T_c (see most recent lattice results in [38] showing evidences for such a scenario). It has been suggested that these thermal monopoles near T_c seem to form a densely packed liquid [22,25].

VI. SUMMARY

In this paper, we have argued that the free energy F(r, T)and internal energy V(r, T) can be probed by slow and fast separation of the $\bar{Q}Q$ pair, respectively. Furthermore we have identified the linear part in both potentials with flux tube formation between the pair: for free energy as probed by slow separation, there is stable flux tube protected by magnetic supercurrent due to condensed monopoles which has no dissipation and exists below T_c ; for internal energy as probed by fast separation, there is meta-stable flux tube protected by magnetic normal current due to thermal monopoles which are very dense in the region $0.8-1.3T_{c}$ and generate large entropy (the splitting between free and internal energy) via dissipation on a longer time scale. Based on these ideas we have solved analytically the elliptic bags and provided expressions for the potentials at all separations, which happen to describe the data very well.

STATIC $\overline{Q}Q$ POTENTIALS AND THE MAGNETIC ...

The main outcome from our study of the static $\bar{Q}Q$ potentials is the particular relations we suggest between the free energy F(r, T) and internal energy V(r, T) measured on the lattice and the densities of the condensed and "normal" monopoles (see Eqs. (16) and (20)). Since those densities can be directly obtained from the lattice configurations, one may check if these relations are correct or not. Such further tests of the "magnetic scenario"[15,23] for the near T_c QCD plasma are rather straightforward and should be performed.

ACKNOWLEDGMENTS

The work of J.L. was partially supported by the US Department of Energy under DOE Contract No. DE-AC02-98CH10886 and No. DE-AC02-05CH11231. The work of E.S. was supported in parts by the US-DOE grant DE-FG-88ER40388.

APPENDIX

In this Appendix we briefly list the parabolic coordinates formulae needed for the calculation in Sec. II.

The coordinates we use are (ξ, η, ϕ) with two focal points at $\pm a\hat{\mathbf{z}}$, which are related to cylindrical coordinates (ρ, ϕ, z) by

$$\rho = a\sqrt{(\xi^2 - 1)(1 - \eta^2)}, \quad \phi = \phi, \quad z = a\xi\eta.$$
(A1)

The variables are defined in the following domains: $\xi \in (1, \infty)$, $\eta \in [-1, 1]$, $\phi \in [0, 2\pi)$. Writing $ds^2 = H_{\xi}^2 d\xi^2 + H_{\eta}^2 d\eta^2 + H_{\phi}^2 d\phi^2$, we have

$$H_{\xi} = a \frac{\sqrt{\xi^2 - \eta^2}}{\sqrt{\xi^2 - 1}},$$

$$H_{\eta} = a \frac{\sqrt{\xi^2 - \eta^2}}{\sqrt{1 - \eta^2}},$$

$$H_{\phi} = a \sqrt{(\xi^2 - 1)(1 - \eta^2)}.$$
(A2)

The Laplacian is given by

$$\vec{\nabla}^2 = \frac{1}{a^2(\xi^2 - \eta^2)} \left\{ \frac{\partial}{\partial \xi} \left[(\xi^2 - 1) \frac{\partial}{\partial \xi} \right] + \frac{\partial}{\partial \eta} \\ \times \left[(1 - \eta^2) \frac{\partial}{\partial \eta} \right] + \left[\frac{1}{\xi^2 - 1} + \frac{1}{1 - \eta^2} \right] \frac{\partial^2}{\partial \phi^2} \right\}.$$
(A3)

Finally the gradient is given by

$$\vec{\nabla} = \hat{\xi} \frac{\partial}{H_{\xi} \partial \xi} + \hat{\eta} \frac{\partial}{H_{\eta} \partial \eta} + \hat{\phi} \frac{\partial}{H_{\phi} \partial \phi}.$$
 (A4)

For more details one could consult books such as [40].

- [1] G.S. Bali, Phys. Rep. 343, 1 (2001).
- [2] J. Greensite, Prog. Part. Nucl. Phys. 51, 1 (2003); R. Alkofer and J. Greensite, J. Phys. G 34, S3 (2007).
- [3] O. Kaczmarek and F. Zantow, Phys. Rev. D 71, 114510 (2005); arXiv:hep-lat/0506019.
- [4] O. Kaczmarek, F. Karsch, E. Laermann, and M. Lutgemeier, Phys. Rev. D 62, 034021 (2000); O. Kaczmarek, F. Karsch, P. Petreczky, and F. Zantow, Phys. Lett. B 543, 41 (2002); Nucl. Phys. B, Proc. Suppl. 129, 560 (2004).
- [5] P. Petreczky and K. Petrov, Phys. Rev. D **70**, 054503 (2004).
- [6] O. Kaczmarek and F. Zantow, Proc. Sci., LAT2005 (2006) 192.
- [7] C. Young and E. Shuryak, Phys. Rev. C 79, 034907 (2009); 81, 034905 (2010).
- [8] X. Zhao and R. Rapp, Phys. Lett. B 664, 253 (2008).
- [9] A. Mocsy and P. Petreczky, Phys. Rev. D 77, 014501 (2008).
- [10] E. V. Shuryak and I. Zahed, Phys. Rev. C 70, 021901 (2004); Phys. Rev. D 70, 054507 (2004).
- [11] J. Liao and E. V. Shuryak, Nucl. Phys. A775, 224 (2006).
- [12] C. R. Allton *et al.*, Phys. Rev. D **71**, 054508 (2005); R. V. Gavai and S. Gupta, Phys. Rev. D **72**, 054006 (2005).

- [13] J. Liao and E. V. Shuryak, Phys. Rev. D 73, 014509 (2006).
- [14] B. A. Gelman, E. V. Shuryak, and I. Zahed, Phys. Rev. C 74, 044908 (2006); 74, 044909 (2006).
- [15] J. Liao and E. Shuryak, Phys. Rev. C 75, 054907 (2007).
- [16] C. Ratti and E. Shuryak, Phys. Rev. D 80, 034004 (2009).
- [17] M. Gyulassy and L. McLerran, Nucl. Phys. A750, 30 (2005); E. V. Shuryak, Prog. Part. Nucl. Phys. 53, 273 (2004); Nucl. Phys. A750, 64 (2005).
- [18] E. V. Shuryak, arXiv:hep-ph/0608177; Prog. Theor. Phys. Suppl. 168, 320 (2007).
- [19] D. Antonov, S. Domdey, and H.J. Pirner, Nucl. Phys. A789, 357 (2007).
- [20] E. Megias, E. Ruiz Arriola, and L. L. Salcedo, Phys. Rev. D 75, 105019 (2007).
- [21] A. Beraudo, J. P. Blaizot, and C. Ratti, Nucl. Phys. A806, 312 (2008).
- [22] J. Liao and E. Shuryak, Phys. Rev. Lett. 101, 162302 (2008).
- [23] M. N. Chernodub and V. I. Zakharov, Phys. Rev. Lett. 98, 082002 (2007); arXiv:hep-ph/0702245.
- [24] M. N. Chernodub, K. Ishiguro, A. Nakamura, T. Sekido, T. Suzuki, and V. I. Zakharov, Proc. Sci., LAT2007 (2007) 174.

- [25] A. D'Alessandro and M. D'Elia, Nucl. Phys. B799, 241 (2008).
- [26] E. Shuryak, J. Phys. G 35, 104044 (2008); Prog. Part. Nucl. Phys. 62, 48 (2009); Nucl. Phys. B, Proc. Suppl. 195, 111 (2009); J. Liao and E. Shuryak, J. Phys. G 35, 104058 (2008); arXiv:0809.2419.
- [27] J. Liao and E. Shuryak, Phys. Rev. C 77, 064905 (2008).
- [28] S. Mandelstam, Phys. Rep. 23, 245 (1976); G. 't Hooft, Nucl. Phys. B190, 455 (1981).
- [29] A. A. Abrikosov, Sov. Phys. JETP 5, 1174 (1957);H.B. Nielsen and P. Olesen, Nucl. Phys. B61, 45 (1973).
- [30] G. Ripka, *Lecture notes in Physics* (Springer, Berlin/ Heidelberg, 2004), Vol. 639; arXiv:hep-ph/0310102.
- [31] A. Di Giacomo, Prog. Theor. Phys. Suppl. 131, 161 (1998); arXiv:hep-lat/0310023.
- [32] J. D. Jackson, *Classical Electrodynamics* (John Wiley & Sons, Inc., New York, 1999), 3rd ed..
- [33] J. Liao and E. Shuryak, Phys. Rev. Lett. 102, 202302 (2009).
- [34] E. Shuryak, Phys. Rev. C 80, 054908 (2009); 80, 069902
 (E) (2009).
- [35] J. Casalderrey-Solana, E. V. Shuryak, and D. Teaney, J. Phys. Conf. Ser. 27, 22 (2005); Nucl. Phys. A774, 577 (2006).

- [36] B. Alver *et al.* (PHOBOS Collaboration), Phys. Rev. C 75, 054913 (2007); 81, 024904 (2010).
- [37] C. Faroughy and E. Shuryak, Phys. Rev. C 82, 014902 (2010).
- [38] A. D'Alessandro, M. D'Elia, and E. V. Shuryak, Phys. Rev. D 81, 094501 (2010).
- [39] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn, and V. F. Weisskopf, Phys. Rev. D 9, 3471 (1974).
- [40] Z. X. Wang and D. R. Guo, *Special Functions* (World Scientific, Singapore, 1989).
- [41] It has been noticed that the numerical coefficients here in Eq. (16) and later in Eq. (20) are different from those quoted in a review paper [E. Shuryak, J. Phys. G 35, 104044 (2008)].11/8 The numbers in the review were from a very early version of the current study when it was not finalized. The corrections had been made before the first arXiv version of this paper appeared. The present paper and all its arXiv versions consistently give the right coefficients.
- [42] M. Baker, N. Brambilla, H. G. Dosch, and A. Vairo, Phys. Rev. D 58, 034010 (1998).
- [43] A. Nakamura, T. Saito, and S. Sakai, Phys. Rev. D 69, 014506 (2004).