

Equivalence between Euclidean and in-in formalisms in de Sitter QFTAtsushi Higuchi,^{1,*} Donald Marolf,^{2,†} and Ian A. Morrison^{2,‡}¹*Department of Mathematics, University of York Heslington, York, YO10 5DD, United Kingdom*²*Physics Department, UCSB, Santa Barbara, California 93106, USA*

(Received 16 January 2011; published 15 April 2011)

We study the relation between two sets of correlators in interacting quantum field theory on de Sitter space. The first are correlators computed using in-in perturbation theory in the expanding cosmological patch of de Sitter space (also known as the conformal patch, or the Poincaré patch), and for which the free propagators are taken to be those of the free Euclidean vacuum. The second are correlators obtained by analytic continuation from Euclidean de Sitter; i.e., they are correlators in the fully interacting Hartle-Hawking state. We give an analytic argument that these correlators coincide for interacting massive scalar fields with any $m^2 > 0$. We also verify this result via direct calculation in simple examples. The correspondence holds diagram by diagram, and at any finite value of an appropriate Pauli-Villars regulator mass M . Along the way, we note interesting connections between various prescriptions for perturbation theory in general static spacetimes with bifurcate Killing horizons.

DOI: 10.1103/PhysRevD.83.084029

PACS numbers: 04.62.+v

I. INTRODUCTION

While free quantum fields in de Sitter space (dS_D) have been well understood for some time (see [1] for scalar fields), interacting de Sitter quantum field theory continues to be a topic of much discussion. In particular, there has been significant interest in the possibility of large infrared (IR) effects in interacting de Sitter quantum field theories [2–17], both with and without dynamical gravity. Most of these discussions have been in Lorentzian signature, using some form of in-in perturbation theory. (See, e.g., [18–21] for early use of in-in perturbation theory in QFT in curved space.) A popular choice is to choose the initial surface to be a cosmological horizon, so that the perturbation theory involves integrals over the region to the future of this horizon (see Fig. 1). This region of de Sitter space is also known as the expanding cosmological patch, the conformal patch, or the Poincaré patch. We will therefore refer to the associated perturbation scheme as the Poincaré in-in formalism, especially when the initial state is chosen to be the free Bunch-Davies (i.e., Euclidean) vacuum.

On the other hand, IR effects are often easier to control and analyze in Euclidean-signature de Sitter space, which is just the D sphere S^D . Analytic continuation of such correlators to Lorentz signature defines the so-called Hartle-Hawking vacuum of the theory [22]. The fact that S^D is compact means that no IR divergences can arise in perturbation theory unless they are already present at order zero. With appropriate techniques one can often analytically continue the resulting IR-finite Euclidean correlators to Lorentzian signature while maintaining control over the

IR behavior. This was done in [23–25] for massive scalar fields using standard perturbation theory. For massless scalars, [26] used the Euclidean setting to introduce a new form of perturbation theory which again yields IR-finite Euclidean correlators whose continuation to Lorentz signature can be controlled.

One would therefore like to understand precisely how correlators analytically continued from Euclidean signature are related to those computed using an intrinsically Lorentz-signature technique. On general grounds, the analytically continued correlators will satisfy the Lorentz-signature Schwinger-Dyson equations. So long as they satisfy appropriate positivity requirements to define a positive-definite Hilbert space, this means that the analytically continued (Hartle-Hawking) correlators define a valid state of the theory. Recall that positivity will generally follow from the de Sitter analogue [27] of reflection positivity and the Osterwalder-Schröder construction, and that reflection positivity holds formally when the Euclidean action is bounded below.¹ In such cases, it remains only to ask how the Hartle-Hawking state relates to other states of interest, such as the state defined by in-in perturbation theory in the Poincaré patch.

A hint was given by [12] which studied a free scalar field but treated the mass term as a perturbation about the conformally coupled value. The Euclidean and Poincaré in-in formalisms were found to agree, and in fact to both give the exact result once all orders in perturbation theory had been included. (There are no UV divergences due to the fact that the theory has only quadratic terms and thus only tree diagrams.) This may at first seem surprising. Indeed, for in-in perturbation theory defined using a

*ah28@york.ac.uk

†marolf@physics.ucsb.edu

‡ian_morrison@physics.ucsb.edu

¹This has been rigorously shown in $D = 2$ dimensions for standard kinetic terms and polynomial potentials; see e.g., [28].

Cauchy surface at finite time as the initial surface, a result of this form would be impossible. Since the past light cone of any external point of a Feynman diagram is cut off by the initial surface, all integrals are over regions of finite spacetime volume. Furthermore, the volume of any such region would shrink to zero when the external point approaches the initial surface. As a result, the in-in correlators would necessarily approach the correlators of the zeroth-order theory as all arguments approach the initial slice. On the other hand, analytic continuation of Euclidean correlators gives a de Sitter-invariant interacting state that cannot approach the zeroth-order state on any surface, so the two formalisms could not agree.

In contrast, in the Poincaré in-in formalism the initial surface is a null cosmological horizon. In particular, it has the important property that there is an infinite volume of spacetime that lies both to the future of this surface and to the past of any given point in the interior of the Poincaré patch.² This means that the integrals which compute perturbative corrections to the zeroth-order correlators need not become small as the arguments of correlators approach the initial surface and no contradiction with the Euclidean formalism arises.

Indeed, symmetry arguments suggest that this correspondence holds more generally. Since both the free propagators and the Poincaré patch is invariant under translations, rotations, and dilations, the results of Poincaré in-in perturbation theory will be similarly invariant so long as all integrals converge. But for free fields on dS_D the only Hadamard state which is invariant under these symmetries is the Euclidean vacuum. One therefore expects a similar result to hold in perturbation theory, suggesting that the Poincaré in-in approach generally computes correlators in the interacting Euclidean vacuum.

An independent motivation comes from the work of Gibbons and Perry [29], who pointed out that interacting Euclidean field theory on S^D describes thermal field theory inside the cosmological horizon of de Sitter space (i.e., in the static patch) with Gibbons-Hawking temperature [30]. While the Euclidean formalism is commonly used to study thermal field theory, there is a Lorentzian version called the Schwinger-Keldysh formalism [31,32]. This formalism agrees with what is usually called the in-in formalism in relativistic field theory if the property called *factorization* is satisfied (see, e.g., [33]). The physical content of this property is that generic states thermalize if given sufficient time, so that one need not take particular care to prepare a thermal state so long as the initial state is taken to be sufficiently far in the past. Since it is known that correlators in a wide class of states approach those of the Euclidean vacuum at late times [23–25], it is reasonable to conjecture

²This follows immediately from the fact that the Poincaré patch is a homogeneous space in and of itself. Any spacetime point in the patch can be mapped to any other using only the symmetries of the patch.

that the Euclidean and in-in formalisms agree at least in the static patch of de Sitter space.

We argue below that the Euclidean and Poincaré in-in approaches in fact agree for general interacting scalar field theories with $m^2 > 0$. The argument can be sketched in three steps. Step 1 is to relate the analytic continuation of Euclidean correlators to in-in perturbation theory in the static patch of de Sitter. This amounts to checking that conditions are right for the usual relation between Euclidean field theory and Lorentz-signature thermal field theory, i.e., factorization, to hold. Step 2 is to note that, for position-space correlators with all arguments in the static patch, in-in perturbation theory is the same whether one thinks of it as perturbation theory in the static patch or as perturbation theory in the Poincaré patch. This follows from the well-known fact that in-in perturbation theory can be expressed in terms of integrals over the region that is i) to the past of all external points of a Feynman diagram and ii) to the future of the initial surface; see Fig. 1. As a result, analytic continuation from the Euclidean reproduces Poincaré in-in calculations at least when the arguments are restricted to a single static patch. Finally, step 3 is to show that both sets of correlators are appropriately analytic, so that their extension to the full spacetime is uniquely determined by their values in the static patch. We consider Pauli-Villars regulated correlators and show agreement at each value of the Pauli-Villars regulator masses. It follows that the fully renormalized correlators must agree as well.

The majority of this paper is devoted to the details of this argument and to providing some simple checks of the results. Section II quickly reviews the relevant features of de Sitter geometry. Section III then verifies that analytic continuation of Euclidean correlators does indeed give in-in correlators in the static patch for massive scalar fields, while Sec. IV argues that the correlators are sufficiently analytic so as to be determined by their restriction to the static patch. Since the arguments are somewhat involved, we explicitly compute some simple in-in loop diagrams in Sec. V and demonstrate agreement with the Euclidean




FIG. 1. Standard Carter-Penrose diagram of de Sitter space. Region I is the static patch, and the Poincaré patch consists of regions I and II. The causal pasts of points X_1 and X_2 are the shaded regions. See Sec. II A for details.

results computed in [23]. We close with a discussion in Sec. VI. In the Appendix we describe a more direct way for the analytic continuation of Euclidean correlators, which gives a slightly different method for demonstrating their equivalence to Poincaré in-in correlators.

II. PRELIMINARIES

This section serves to briefly review various features of both Lorentzian and Euclidean de Sitter space, and to introduce notation and conventions. After discussing the geometry and relevant coordinate systems in Sec. II A we review aspects of de Sitter propagators in Sec. II B.

A. de Sitter geometry and coordinates

Let us begin with Euclidean de Sitter space. As is well known, this is just the sphere S^D . Throughout this work, we set the de Sitter length ℓ to 1 and work on the unit sphere. We may thus describe S^D using the metric

$$ds_{S^D}^2 = d\Omega_D^2 = d\vartheta^2 + \sin^2\vartheta d\Omega_{D-1}^2, \quad \vartheta \in [0, \pi], \quad (2.1)$$

where $d\Omega_d^2$ is the line element of the unit S^d .

It is useful to consider the complexified manifold \mathbb{S}^D , which may be thought of as the surface $X \cdot X = 1$ in \mathbb{C}^{D+1} . Wick rotations of various coordinates correspond to passing from one real section of \mathbb{S}^D to another, e.g., from S^D to dS_D . One useful Wick rotation is given by defining

$$\Theta = i\left(\vartheta - \frac{\pi}{2}\right) \quad (2.2)$$

and taking Θ real; i.e., by Wick rotating the polar angle. This yields

$$ds_{\text{global } dS_D}^2 = -d\Theta^2 - \cosh^2\Theta^2 d\Omega_{D-1}^2, \quad \Theta \in \mathbb{R}, \quad (2.3)$$

which is the metric of dS_D in the so-called global coordinates. Indeed, these coordinates are regular on all of dS_D . Making a further coordinate transformation

$$\tan T = \sinh\Theta \quad (2.4)$$

and writing $d\Omega_{D-1}^2 = d\chi^2 + \sin^2\chi d\Omega_{D-2}^2$, we have

$$ds_{\text{global } dS_D}^2 = \sec^2 T (-dT^2 + d\chi^2 + \sin^2\chi d\Omega_{D-2}^2), \quad T \in (-\pi/2, \pi/2), \quad (2.5)$$

where the factor inside the parentheses is the metric on a piece of the Einstein static universe. Note that this piece extends only for a finite amount of Einstein static universe time. Figure 1 is the corresponding Carter-Penrose diagram.

However, one may also arrive at the same real section dS_D by defining

$$t = i\phi, \quad \text{for } \tan\phi = \frac{X^1}{X^2}, \quad (2.6)$$

where $X = (X^1, X^2, \dots, X^{D+1})$, and taking t real; i.e., by Wick rotating the azimuthal angle. This yields

$$ds_{\text{static } dS_D}^2 = -\cos^2\theta dt^2 + d\theta^2 + \sin^2\theta d\Omega_{D-2}^2, \quad t \in \mathbb{R}, \quad \theta \in [0, \pi/2), \quad (2.7)$$

with

$$\tan\theta = \sqrt{\frac{(X^3)^2 + \dots + (X^{D+1})^2}{(X^1)^2 + (X^2)^2}}, \quad (2.8)$$

which is the metric of dS_D in the so-called static coordinates. The coordinate range $t \in \mathbb{R}$, $\theta \in [0, \pi/2)$ describes the static patch of de Sitter. The coordinates t and θ can be expressed in terms of T and χ as

$$\tanh t = \sin T \sec\chi, \quad (2.9)$$

$$\sin\theta = \sec T \sin\chi. \quad (2.10)$$

The boundary at $\theta = \pi/2$ is a coordinate singularity that coincides with the past and future cosmological horizons, $T = \pm(\chi - \frac{\pi}{2})$, defined by the observer at $\theta = 0$; see Fig. 2.

We will also make use of the so-called Poincaré (also known as conformally flat) coordinates on dS_D in which the metric takes the form

$$ds^2 = \frac{1}{\lambda^2} (-d\lambda^2 + d\mathbf{x}^2), \quad (2.11)$$

where $\mathbf{x} = (x^1, \dots, x^{D-1})$. These coordinates are related to the global ones via

$$\lambda = \frac{\cos T}{\sin T + \cos\chi}, \quad (2.12)$$

$$x^i = \frac{\sin\chi}{\sin T + \cos\chi} \hat{X}^i, \quad (2.13)$$

where $\hat{X}^i = X^{i+2}/\sqrt{(X^3)^2 + \dots + (X^{D+1})^2}$. The expanding cosmological patch is the region $0 < \lambda < \infty$ with $\mathbf{x} \in \mathbb{R}^{D-1}$, which we also call the conformal or Poincaré patch. Here $\lambda = \infty$ is the (past) cosmological horizon defined by the observer at $\mathbf{x} = 0$, which we take to coincide with the geodesic $\theta = 0$. With this convention, the Poincaré patch contains the static patch as shown in Fig. 1.




FIG. 2. Carter-Penrose diagram of de Sitter space with $\theta = \text{const}$ surfaces (schematically) indicated by solid lines and $t = \text{const}$ surfaces by dashed lines.

We also take $\lambda = 0$ to coincide with both $t = +\infty$ and $\Theta = +\infty$ on this geodesic. (Thus, the variable λ runs backwards in time. It is more common to use the variable $\eta = -\lambda$ in the cosmology community.)

The remaining relation between Poincaré coordinates and those discussed before is best summarized by using the concept of embedding coordinates. Recall that dS_D can be defined as the locus of points $X \cdot X = 1$ in $D + 1$ -dimensional Minkowski space. Given two such points, X and Y , one may treat them as vectors and compute the invariant Minkowski scalar product $Z = X \cdot Y$, which gives a de Sitter-invariant measure of the separation between X and Y . In the above coordinate systems one finds

$$Z = -\sinh\Theta_x \sinh\Theta_y + \cosh\Theta_x \cosh\Theta_y \cos\gamma^{D-1}, \tag{2.14}$$

(global)

$$= \cos\theta_x \cos\theta_y \cosh(t_x - t_y) + \sin\theta_x \sin\theta_y \cos\gamma^{D-2}, \tag{2.15}$$

(static)

$$= 1 - \frac{\|\mathbf{x} - \mathbf{y}\|^2 - (\lambda_y - \lambda_x)^2}{2\lambda_x \lambda_y}, \tag{2.16}$$

(Poincaré)

where γ^d is the angle between the X and Y on the relevant S^d . It is useful to note that $Z = 1$ for $X = Y$ or for points connected by a null geodesic, $Z > 1$ for points connected by a timelike geodesic, $|Z| < 1$ for points connected by a spacelike geodesic, and $Z < -1$ for points which cannot be connected by any geodesic in real de Sitter space. In the latter case, the points are not causally related; see Fig. 3. Note that $Z > -1$ in the static patch. Thus, if points X and Y are in the static patch, then there is a geodesic connecting these two points.

On complex de Sitter space we may take $t = \sigma + i\tau$ in static coordinates to write

$$Z = \cos\theta_x \cos\theta_y [\cosh(\sigma_x - \sigma_y) \cos(\tau_x - \tau_y) - i \sinh(\sigma_x - \sigma_y) \sin(\tau_x - \tau_y)] + \sin\theta_x \sin\theta_y \cos\gamma^{D-2}, \tag{2.17}$$

$$F(a, b; c; x) = 1 + \sum_{n=1}^{\infty} \frac{a(a+1) \cdots (a+n-1) b(b+1) \cdots (b+n-1)}{n! c(c+1) \cdots (c+n-1)} x^n. \tag{2.21}$$

We will be interested in the analytic properties of (2.19) for general complex Z . The only singularities are branch points³ at $Z = 1$ and $Z = \infty$, and we take the branch cut to connect these points along the positive real axis. It will be particularly important to understand the singularity

³These are poles if D is even and if the scalar is conformally coupled and massless.




FIG. 3. Carter-Penrose diagram of de Sitter space with timelike geodesics from point O drawn with solid lines and spacelike geodesics from it drawn with dashed lines.

so that

$$|Z|^2 = |\cos\theta_x \cos\theta_y \cosh(\sigma_x - \sigma_y) \cos(\tau_x - \tau_y) + \sin\theta_x \sin\theta_y \cos\gamma^{D-2}|^2 + \cos^2\theta_x \cos^2\theta_y \sinh^2(\sigma_x - \sigma_y) \sin^2(\tau_x - \tau_y), \tag{2.18}$$

$\theta_x, \theta_y \in [0, \pi/2)$.

B. de Sitter propagators

Consider two points X, Y on Euclidean de Sitter S^D . In terms of $Z = X \cdot Y$, the scalar propagator on S^D is [34,35]

$$\Delta(X, Y) = \frac{\Gamma(a_+) \Gamma(a_-)}{2(2\pi)^{D/2} \Gamma(\frac{D}{2})} (1 - Z)^{(2-D)/2} \times F\left(\frac{D}{2} - a_+, \frac{D}{2} - a_-; \frac{D}{2}; \frac{1+Z}{2}\right), \tag{2.19}$$

where

$$a_{\pm} = \frac{1}{2}[D - 1 \pm \sqrt{(D - 1)^2 - 4m^2}]. \tag{2.20}$$

Here F is Gauss' hypergeometric function:

structure in terms of static coordinates (2.7). Careful inspection of (2.17) shows the following:

Observation. The Green's function for two points X, Y with static coordinates (t_x, θ_x) and (t_y, θ_y) with $\theta_x, \theta_y \in [0, \pi/2)$ is analytic for all complex t_x, t_y except when $t_x - t_y$ is real modulo $2\pi i$ (so that the two points lie on the same Lorentz-signature real section) and the two points obtained by replacing t_x and t_y by $\text{Re}t_x$ and $\text{Re}t_y$, respectively, are causally related. (within this real section).

It will be useful to regulate the divergences of (2.19) at $Z = 1$ using Pauli-Villars subtractions both for the internal and external propagators so that all propagators become bounded functions of Z . Because the unbounded nature of the external propagators needs to be taken into account only in the coincidence limit, where the vertex integral is convergent due to the small integration measure, it is in fact possible to show the equivalence of the Poincaré and Euclidean formalisms regulating only the internal propagators. However, since analyzing such issues in detail would make the argument more cumbersome, we choose to regulate the external propagators as well.

For each m, D we define a regulated propagator

$$\Delta^{\text{reg}}(X, Y) = \Delta(X, Y) + \sum_{i=1}^{[D/2]} C_i \Delta_{M_i}(X, Y), \quad (2.22)$$

where $[D/2]$ denotes the integer part of $D/2$, $\Delta_{M_i}(X, Y)$ is the propagator (2.19) for a particle of mass M_i , and C_i are constants. We will always assume $M_i \gg 1$ in units of the de Sitter scale, so that, in particular, the masses M_i correspond to principal series representations [36] of the de Sitter group. One may choose the coefficients C_i so that $\Delta^{\text{reg}}(X, Y)$ has a well-defined finite limit as $Z \rightarrow 1$ (see, e.g., [37] for $D = 4$). For $D = 2, 3$ we have $[D/2] = 1$ and one may take $C_1 = -1$ for any M_1 . For $D = 4, 5$ one may choose any C_1, C_2, M_1, M_2 which satisfy $C_1 + C_2 = -1$ and $C_1 M_1^2 + C_2 M_2^2 = -m^2$. Nevertheless, $\Delta^{\text{reg}}(X, Y)$ is not analytic at $Z = 1$. Instead, $Z = 1$ remains a branch point analogous to that of the function $x \ln x$ or $x^{1/2}$ at $x = 0$.

If desired, one can also make further subtractions to define regulated propagators with continuous (and thus bounded) derivatives to any specified order. Such additional subtractions are useful in treating theories with derivative interactions, or for consideration of field-renormalization counterterms. Below, we will focus on nonderivative interactions for which the above subtractions will suffice. But it will be clear from the argument that the same results hold for derivative interactions so long as an appropriate number of additional Pauli-Villars subtractions have been made.

Finally, it is useful to study $\Delta(X, Y)$ at large $|Z|$. There, Δ behaves either like $|Z|^{-a_-}$ (for $m^2 < (D-1)^2/4$) or $|Z|^{-(D-1)/2}$ (for $m^2 \geq (D-1)^2/4$). Hence, for given choices of regulator parameters C_i, M_i the modulus of the regularized propagator $|\Delta^{\text{reg}}(Z)|$ is bounded. It is useful to take each C_i, M_i to be a given function of the smallest regulator mass M , so that the regulator is removed as $M \rightarrow \infty$. We may then take the bound on $|\Delta^{\text{reg}}(Z)|$ to be $B(M)$, determined only by m and the lightest regulator mass M .

III. EUCLIDEAN CORRELATORS VS. THERMAL STATIC PATCH CORRELATORS

We now turn to step 1 of the argument sketched in the introduction. Our task here is to show that the analytic

continuation of Euclidean correlators is equivalent to those computed using in-in perturbation theory (defined using the propagator of the free Euclidean vacuum) in the so-called static patch of de Sitter. This essentially amounts to checking that conditions are right for the usual relation between Euclidean field theory and Lorentz-signature thermal field theory to hold; i.e., that the Hartle-Hawking correlators are indeed thermal correlators in the static patch. At a formal level, this follows from the fact that correlation functions $Tr[\phi(x_1) \dots \phi(x_n) e^{-\beta H}]$ in the canonical ensemble are given by an imaginary-time path integral; see e.g., [33]. However, in order not to miss any subtleties (perhaps due to IR divergences of the sort predicted in [14]) and because of the many controversies surrounding de Sitter quantum field theory, we will proceed slowly through an explicit perturbative argument. Below, we consider diagrams using the Pauli-Villars regularized propagators (2.22) so that $|\Delta^{\text{reg}}(Z)| \leq B(M)$. We restrict attention to connected diagrams since vacuum bubbles are automatically excluded both in the Euclidean and in-in formalisms. Because the desired result is trivial for the diagram with two external points connected by a single propagator, we also exclude this diagram from our discussion. Nonderivative interactions are assumed for simplicity, though the argument is readily extended to derivative interactions so long as additional Pauli-Villars subtractions are made as described in Sec. II B above.

Recall that in static coordinates (2.7) points of de Sitter space are labeled by a pair (t, \hat{X}) where \hat{X} is a point in the (open) northern hemisphere of S^{D-1} . We will use these coordinates for both the static patch of Lorentz signature dS_D (where $t \in \mathbb{R}$) and on Euclidean-signature de Sitter S^D (where $-it \in (-\pi, \pi)$.) We imagine that the integrals over the time coordinates t_i of the internal vertices will be performed first, followed later by the integrals over \hat{X}_i . So for the moment we consider the \hat{X}_i to be fixed. We also assume that all internal vertices and external points correspond to distinct spatial points \hat{X} ; i.e., $\hat{X}_i \neq \hat{X}_j$ for $i \neq j$. Because of our Pauli-Villars regularization, we can always recover information at coincidence by continuity.

Let us first review the general argument relating Euclidean correlators to in-in correlators (see e.g., [33]) using our de Sitter static patch notation. In the Euclidean approach the time integrals of the internal vertices are all from $i\pi$ to $-i\pi$. The external points are taken to lie on this contour and, at least for the moment, we take them to all lie close to (though not necessarily precisely at) $t = 0$. Since the \hat{X}_i are distinct, it follows from the *Observation* of Sec. II B that the integrand is analytic in all time coordinates t_i in a region containing the contour of integration. Thus, the contour can be deformed. In fact, taking all internal coordinates t_i to be integrated along the same contour C , we note that the contour can be freely deformed so long as (i) it begins at some $t = t_0 + i\pi$ with t_0 real and ends at $t = t_0 - i\pi$, (ii) the imaginary part of t is strictly

decreasing everywhere (so that no two points on the path have the same value of $\text{Im}t$) and (iii) the path continues to pass through the external points. In particular, we are free to take the limit $t_0 \rightarrow -\infty$.

Now, these rules allow us to choose the contour $C = A_1 + C_1 + B + C_2 + A_2$ to be as in Fig. 4. Here $\epsilon (< \pi)$ is a nonzero positive number. The imaginary part is increasing infinitesimally on the horizontal portions of the contour. This is equivalent to using the Feynman propagator when the two points are both on the upper horizontal portion and the Dyson (or anti-Feynman) propagator if the two points are both on the lower horizontal portion. In general, in the $\epsilon \rightarrow 0$ limit (and where the imaginary parts of the times for all external points are also taken to zero), one may say that the above contour computes correlators using the free path-ordered two-point function as the propagator, just as occurs in the in-in formalism. Furthermore, since all integrals converge after Pauli-Villars regularization, it is clear that the integral along B is of order ϵ and can be neglected in the limit $\epsilon \rightarrow 0$.

As a result, the Euclidean correlators (evaluated at $t = 0$) agree with the corresponding in-in correlators in the static patch (computed using the propagators of the free Euclidean vacuum) so long as a property called *factorization* [33] holds, which states that the A_1, A_2 pieces of the contour C can be neglected in the $t_0 \rightarrow -\infty$ limit. We now establish this property for our systems, diagram by diagram.⁴ For each Feynman diagram, let us choose one external point $X = (t_e, \hat{X}_e)$ and one internal point Y that lies on either segment A_1 or A_2 . To show that the integral of Y over the above segments can be neglected, we also choose a path through the diagram from X to Y ; i.e., a particular chain of propagators.

Now, recall from Sec. II B that at fixed Pauli-Villars regulator mass M all propagators are bounded by some $B(M)$. To establish a bound on the integrals, we may thus replace the integrand with its magnitude and replace all propagators *not* on the chosen path by $B(M)$. Next consider the propagators on the chosen path. For at least one such propagator, external or internal, the (static-patch) time coordinates of its two arguments have real parts differing by at least $(t_e - t_0)/K$, where K is the number of propagators in the chain. From (2.18) and the asymptotics of the propagators discussed in Sec. II B, this means that this propagator is of order $[\cos\theta_1 \cos\theta_2 e^{(t_e - t_0)/K}]^{-\nu}$ or smaller for some positive ν determined by the mass m of the quantum fields, where θ_1 and θ_2 are the θ coordinates of the two arguments of this propagator, if $t_e - t_0$ is large enough. Replacing all other propagators on this chain with $B(M)$, we integrate the time coordinate τ of Y along the segments from $t_0 + i\pi$ to $t_0 + i\epsilon$ and from $t_0 - i\epsilon$ to




FIG. 4. A deformation of the Euclidean contour.

$t_0 - i\pi$. We also perform all other t integrals at the vertices. The result is clearly bounded by

$$c_2 [B(M)]^{n_1} (t_e - t_0)^{n_2} [\cos\theta_1 \cos\theta_2 e^{(t_e - t_0)/K}]^{-\nu} \quad (3.1)$$

for some constants c_2, n_1, n_2 , where the factors of $(t_e - t_0)^{n_2}$ come from the measure. It is important to note that c_2, n_1, n_2 are independent of the positions of all vertices, as well as t_0 .

To complete the argument, we divide the integrals over θ_1, θ_2 (or, say, just θ_1 if the second point is external so that $\cos\theta_2$ is fixed and independent of t_0) into two regions. In the first, we take $\cos\theta_1, \cos\theta_2 > e^{-(t_e - t_0)/3K}$. The bound in (3.1) then shows that the integral over this region tends zero at least like $(t_e - t_0)^{n_2} e^{-2\nu(t_e - t_0)/3K}$ as $t_0 \rightarrow -\infty$. The remaining region of integration is small since one of the variables to be integrated (θ_1 and/or θ_2) satisfies $\cos\theta < e^{-(t_e - t_0)/3K}$ and the integration measure $\sin^{D-2}\theta \cos\theta d\theta$ contains a factor of $\cos\theta$. We note that the length of the interval on which θ is integrated is of order $e^{-(t_e - t_0)/3K}$ as well. We may therefore replace *all* propagators by the bound $B(M)$ and find that the contribution from this region is again bounded by a number of the form $c_3 (t_e - t_0)^{n_3} e^{-2(t_e - t_0)/3K}$, which of course tends to zero as $t_0 \rightarrow -\infty$. This establishes the fact that sections A_1 and A_2 can be neglected in the desired limit for any (finite or infinitesimal) choice of ϵ in Fig. 4. In particular, this demonstrates the agreement of Euclidean and static patch in-in correlators (computed using the propagator of the free Euclidean vacuum) when the external points are located at $t = 0$.

To demonstrate agreement for more general external points, we need only analytically continue the correlators as a function of the time coordinates of the external points. This is in fact the *definition* of the Euclidean correlators evaluated at more general times, and we will show that it also gives the static patch in-in correlators. For this step, it is convenient to take the external points to have distinct (and fixed) values of $\text{Im}t$. At the end of the argument we will take the limit where all of these imaginary parts vanish.

We first consider the analyticity of the integrand for some given diagram in the time coordinate t_1 of some external point with the time coordinates of all other points (both internal and external) held fixed and taken to lie on one of the contours C discussed above. We also take the spatial coordinates of all points to be fixed and distinct.

⁴For a general contour C , we will refer to the associated diagrams below as Feynman diagrams, even though they may sometimes involve Dyson (or other) propagators as noted above.

Because of the observation of Sec. II B, the singularities are then a finite distance from the contour C . For example, for the original Euclidean integral, if the external point with the time coordinate t_1 is connected to a vertex with time coordinate t which is also connected to two other vertices, the singularities and associated branch cuts on the complex t plane are similar to those shown in Fig. 5.

We may thus analytically continue t_1 to any complex value so long as we avoid the branch cuts. Let us do so holding $\text{Im}t_1$ fixed and distinct from the imaginary parts of all other external time coordinates. Then the only singularities which are of concern are those due to the vertex connected to t_1 with the same imaginary part of t ; i.e., for which $\text{Im}t = \text{Im}t_1$. As indicated in Fig. 5, for a fixed contour C this will in general allow only a finite range over which the integrand can be analytically continued in $\text{Re}t_1$. However, as noted earlier, we are also free to further deform the contour. For example, by shifting the contour for all vertices a bit to the right at $\text{Im}t_1$, we shift the allowed window for analytic continuation a bit to the right, and we do so without changing the size of this window. It is thus clear that, by dragging the contour along with the external point in this way, we may analytically continue the result of the time integrations to arbitrary values of $\text{Re}t_1$ for any given distinct set of spatial coordinates. But as before, our Pauli-Villars regularization scheme implies the same result holds for general spatial coordinates by continuity.⁵ It follows that the analytic continuation of Euclidean correlators can be computed via the usual Feynman diagrams associated with any contour which (i) begins at some $t = -\infty + i\epsilon$ with any real and positive ϵ ends at $t = -\infty - i\epsilon$, (ii) has the imaginary part of t strictly decreasing everywhere (so that no two points on the path have the same value of $\text{Im}t$) and (iii) passes through all external points.⁶

⁵Continuity of the integrand is clear from the regularization scheme. Continuity of the result of the time integrations follows from the fact that these integrals converge absolutely. This in turn follows from the same estimates used to show factorization above.

⁶The reader may ask if the analytic continuation of a full diagram (after all integrals, including space integrals, have been performed) coincides with the result described above (in which the integrand is first continued, before performing the spatial integrals). The potential obstacle is the fact that spatial coordinates will necessarily coincide somewhere during the integrals over space, and such coincidences shrink the windows (used to enact the analytic continuation above) between past- and future-branch cuts to zero size. One may show that this is not an issue by performing a further regularization in which all propagators $\Delta(Z)$ are replaced by $\Delta(Z - s)$ for some positive s . This regularization maintains windows of finite size even at coincidence. Furthermore, so long as one drags the contour along with the external point as described above, one finds that the resulting integral is analytic in the external time variables for all positive s on the domain where the external times have distinct imaginary parts. Then we find that the full diagram is analytic at $s = 0$, and its analytic continuation is given by the prescription above. The argument is very similar to that given in Section IV B to establish the analyticity of the Poincaré in-in correlators in the conformal-time variables.




FIG. 5. Singularities and branch cuts of propagators.




FIG. 6. Deformed contour for external points at t_i with finite imaginary parts.




FIG. 7. t -integration contour for the in-in formalism in the static patch. The open circles denote external points at times t_i .

An example is shown in Fig. 6. Taking the limit $\epsilon \rightarrow 0$ (and taking the limit in which all external time coordinates now become real) gives the usual closed-time-path representation of the static patch in-in correlators (defined using the propagators of the free Euclidean vacuum) just as described above for external points at $t = 0$; see Fig. 7.

IV. ANALYTICITY OF IN-IN CORRELATORS

Recall that our goal is to demonstrate the equivalence of the Poincaré and Euclidean formalisms for perturbation theory. We outlined a three-step argument in the introduction. As described there, it is clear that the in-in formalism in the static patch is a restriction of that in the Poincaré patch (step 2). Since we have now shown that the static patch in-in correlators agree with those of the Euclidean formalism (step 1), it remains only to show that Poincaré in-in correlators are appropriately analytic in their arguments (step 3). The desired result then follows since two analytic functions that agree in any nonempty open subset of a real section must in fact agree everywhere. In this section we will establish analyticity of Poincaré in-in correlators as functions of the conformal-time coordinates with space coordinates fixed. This will turn out to be sufficient for our purpose.

Recall that, for given external points $X_j = (\tilde{\lambda}_j, \mathbf{x}_j)$, in the coordinates of (2.11) any Poincaré correlator is a sum of terms of the form

$$\begin{aligned} \mathcal{A}_P &= c_1 \left(\prod_{k=1}^n \int d^{D-1} \mathbf{y}_k \int_C \frac{d\lambda_k}{\lambda_k^D} \right) F(Y_1, \dots, Y_n) \\ &\times \prod_{j=1}^m \Delta^{\text{reg}}(X_j, Y_k), \end{aligned} \quad (4.1)$$

where a typical contour C is shown in Fig. 8 and we have used the Pauli-Villars regulated propagators $\Delta^{\text{reg}}(X_j, Y_k)$. The contour is infinitesimally away from the real line, and the imaginary part of λ *increases* infinitesimally everywhere along the contour, even on the horizontal sections. Time-ordered correlators are obtained by putting the λ coordinates, $\tilde{\lambda}_j$, of the external points on the lower horizontal line, whereas anti-time-ordered correlators are obtained by putting them on the upper horizontal line. We will refer to any such \mathcal{A}_P as an amplitude, and we will again refer to the associated diagram as a Feynman diagram even though diagrams include Dyson (or other) propagators in computing \mathcal{A}_P .

Like the in-in amplitude in the static patch, the Poincaré in-in amplitude \mathcal{A}_P is obtained by first considering the corresponding amplitude with finite and distinct imaginary parts $\text{Im}\tilde{\lambda}_j$ of the conformal-time coordinates of the external points and then taking the limit $\text{Im}\tilde{\lambda}_j \rightarrow 0$. For this reason we let $\tilde{\lambda}_j$ satisfy $\text{Im}\tilde{\lambda}_1 < \text{Im}\tilde{\lambda}_2 < \dots < \text{Im}\tilde{\lambda}_m$ without loss of generality and use the contour analogous to that considered for the static patch. Figure 9 shows an example with $m = 4$ with finite imaginary parts, before taking the $\epsilon \rightarrow 0$ limit.

It is important to note that, in general, one must integrate over the conformal-time coordinates λ_k first in (4.1), before integrating over the spatial coordinates as the integrand may otherwise decay too slowly at large $\|\mathbf{y}_k\|$ for the \mathbf{y}_k integrals to converge if each λ_k is fixed on the contour. We will show below that, with our Pauli-Villars regulators, all integrals converge so long as the λ_k integrals are performed first. We then use this result to demonstrate the desired analyticity of \mathcal{A}_P .



FIG. 8. Typical λ contour for the in-in formalism in the Poincaré patch. All external points have $\lambda > \lambda_f$.




FIG. 9. Deformed contour for external points with finite imaginary parts in the Poincaré patch.

A. Convergence of \mathcal{A}_P

We now verify that integrals defining the amplitude \mathcal{A}_P converge with the contour C chosen as in Fig. 9 so long as we perform the λ_k integrations before the \mathbf{y}_k integrations. The general strategy is deform the λ contour at each vertex as much to the right as possible, while avoiding singularities, so that the regions of spacetime over which the vertices are integrated become small enough to guarantee absolute convergence.

The structure of singularities in the complex λ plane is directly analogous to that discussed in the complex t plane in Sec. III. We again fix the spatial coordinates of all points, both internal and external, and take them to be distinct. An example for the conformal time λ_1 of the vertex $Y_1 = (\lambda_1, \mathbf{y}_1)$ is shown in Fig. 10, where dashed lines again indicate branch cuts. Of the two singularities with the same imaginary part, we call the one with the larger (smaller) real part a past (future) singularity. For example, the singularities due to vertex $(\lambda_3, \mathbf{y}_3)$ are at

$$\lambda_{\pm} = \lambda_3 \pm \|\mathbf{y}_1 - \mathbf{y}_3\|. \quad (4.2)$$

The points λ_+ and λ_- are a past singularity and a future singularity, respectively. Notice that $(\text{Re}\lambda_+, \mathbf{y}_1)$ and $(\text{Re}\lambda_-, \mathbf{y}_1)$ are on the past and future light cones of $(\text{Re}\lambda_3, \mathbf{y}_3)$, respectively.

Also in the same way as in Sec. III, each λ_k contour can be deformed as we like so long as it encloses all past singularities and avoids all future singularities. In particular, for the given values of all spatial coordinates $\mathbf{x}_i, \mathbf{y}_k$, the portion of the contour to the left of a vertical line segment connecting two points on the contour can be replaced by this line segment provided that all past singularities lie to its right. For example, the λ_1 contour in Fig. 10 can be deformed as in Fig. 11. Note that this contour may no longer pass through certain λ values corresponding either to external points or to other contours which were not similarly deformed.

Using this observation, we deform the contours as follows. We begin with an integral where all λ_k are integrated over the same contour C of the form shown in Fig. 9 for some given values of the spatial coordinates $\mathbf{x}_j, \mathbf{y}_k$. We deform *all* of the λ_k contours *in the same way* as follows.




FIG. 10. Singularities in the complex λ_1 plane. Two external points lie at $\tilde{\lambda}_1, \tilde{\lambda}_3$ and two internal points lie at λ_2, λ_3 as indicated by the open circles. Filled circles are singularities and dashed lines are branch cuts.




FIG. 11. The λ_1 contour in Fig. 10 deformed as much to the right as possible for the given values of the spatial coordinates as described in the text.

We choose a vertical line segment connecting two points of the contour such that all past singularities on the complex λ_k plane for all k is to its right. Then we let this line segment replace the portion of the contour to its left. We keep deforming the contour in this manner by moving the vertical line segment to the right until it encounters a past singularity, say, on the λ_{k_1} plane, λ_{k_1} being the conformal time for Y_{k_1} , due to some external point, say X_{j_1} . We then stop deforming the contour for Y_{k_1} (since we cannot deform it beyond the singularity) and hold it fixed. We describe this relationship between Y_{k_1} and X_{j_1} by saying that Y_{k_1} is past-related to X_{j_1} (for the given values of all spatial coordinates) and writing⁷ $Y_{k_1} \rightarrow X_{j_1}$.

We then choose some value of λ_{k_1} on its fixed contour and deform the remaining contours by moving the vertical line segment to the right with λ_{k_1} fixed until one of them, say a contour for Y_{k_2} , hits a past singularity due to, say X_{j_2} , which is either an external point or the vertex Y_{k_1} whose contour is being held fixed. We write $Y_{k_2} \rightarrow X_{j_2}$ and hold the contour for Y_{k_2} fixed from now on. We continue in this manner until each vertex is past-related to another point, so that all contours have been fixed.⁸

To understand the resulting structure, we now use the above past-relations to decorate the Feynman diagram under discussion for each fixed set of spatial coordinates. Note that any pair (A, B) of vertices with A past-related to B must be connected by at least one line on the diagram.⁹ If there is one line from A to B , we decorate it with an arrow

⁷For certain spatial coordinates, our contour will encounter two singularities due to distinct external points X_{j_1} and X_{j_2} at the same time. Since this happens only on a set of spatial coordinates of measure zero, we will ignore such cases and assume below that Y_{k_1} is past-related to only one point, and similarly for other vertices in the diagram.

⁸It may be that $X_{j_2} = Y_{k_1}$ for some values of λ_{k_1} while for other values X_{j_2} is an external point. In this way, our definition of new past-relations can depend on the positions of integration variables along contours that have already been fixed. It is straightforward to deal with this seeming complication as discussed in footnote ¹¹ below.

⁹Otherwise the location of point B could not produce singularities in the propagators evaluated at A . In particular, our notion of past-relation is *not* transitive.




FIG. 12. A Feynman diagram in which arrows indicate past relations as described in the text, determined by some particular set of spatial coordinates.

pointing from A to B (i.e., toward the future). If there is more than one such line, we decorate only one of them. Once all past-relations have been indicated in this way, we replace all remaining undecorated propagators with dashed lines. An example is shown in Fig. 12.

In the deformation of contours described above, the contours are deformed until one of them encounters a past singularity. Although this procedure is sufficient to show the convergence of \mathcal{A}_p itself, we need to modify it slightly for proving convergence of the derivatives of \mathcal{A}_p with respect to the external coordinates, which diverge at past singularities. Here we briefly describe this modification. The main difference is that the modified deformation keeps the contours away from past singularities.

We choose the initial contour common to all λ_k as before. We define an *effective past singularity* as follows: if λ is a past singularity, then the corresponding effective past singularity is $\lambda - b$, where b is a small but positive constant. We deform the contours in the same way as before except that they are deformed until one of the contours encounters an effective past singularity rather than a true one. We define the past-relation as before. It may happen that some effective past singularities are outside the contour though the true ones must be inside. If the effective past singularity on the complex λ_{k_3} -plane due to a point X_{j_3} , external or internal, is outside the contour, we stop deforming the contour for λ_{k_3} , fix the value of λ_{k_3} on this contour, and let $Y_{k_3} \rightarrow X_{j_3}$. (If there are two or more effective past singularities outside the contour, we choose one to define the past-relation.) The rest is the same as the original deformation.¹⁰

As noted in the introduction, it is well known that in-in diagrams can be computed by integrating only over the

¹⁰Notice that, since the contour is separated from the past singularities due to external points by a finite distance for large $|\lambda|$, the Z in (2.16) for an external propagator is bounded away from 1 as $|\lambda| \rightarrow \infty$. This means that a differentiated external propagator is bounded on the contours and that the proof for convergence of \mathcal{A}_p below can be used virtually unaltered for the derivative of \mathcal{A}_p .

past light cones of external points. The choice of contours above gives a similar result, but one which is clearly valid for finite ϵ . To see the similarity, note that when A is past-related to B the real part of point A lies in the casual past of the real part of point B over most of the contour for A . The exception is a finite piece near the minimum value of $\text{Re}\lambda$ due to the use of effective past singularities in the modified contour deformation. Since any internal point is connected by some chain of arrows to some external point, except for a set of finite-sized pieces as noted above, the projection of the integration region onto the real λ axis lies in the causal past of at least one external point. We will find it useful below to break up the integration region into such past light cones and finite-sized protruding segments.

Now we establish convergence using the modified deformation of contour. Recall that each internal point A is past-related to precisely one point B (see footnote ⁷), which may be either internal or external. Also recall that, starting at any internal point, one may always follow a chain of arrows upwards until one arrives at an external point. As a result, deleting all dashed lines results in a set of disconnected subdiagrams for which each connected component is a tree whose root (which in this case means that future-most point) is an external point. As a result, if we replace every dashed-line propagator by the bound $B(M)$, our amplitude \mathcal{A}_p factorizes into a product of tree amplitudes in which all points are connected by a chain of past-relations.¹¹

But each such tree amplitude is easy to bound. We begin by bounding the integrals corresponding to some past-most vertex $Y = (\lambda, \mathbf{y})$ in a given tree (e.g., Y_2 , Y_4 or Y_6 in Fig. 12). Taking the magnitude of the integrand, this integral takes the form

$$I = \int d^{D-1}\mathbf{y} \int_{C_{\mathbf{y}}} \frac{|d\lambda|}{|\lambda|^D} |\Delta^{\text{reg}}(Y', Y)|, \quad (4.3)$$

where the notation indicates that the contour $C_{\mathbf{y}}$ over which we integrate λ can depend on the spatial coordinates \mathbf{y} . Recall that $\Delta^{\text{reg}}(Y', Y)$ behaves like $|Z|^{-\nu}$, $\nu > 0$, for large $Z = Y \cdot Y'$. As a result, $|\Delta^{\text{reg}}(Y', Y)|$ behaves at most like $Z_R^{-\nu}$, where $Z_R := \text{Re}Z$. Let us choose some Z_0 large enough that for $Z_R > Z_0$ our $|\Delta^{\text{reg}}(Y', Y)|$ is bounded by $\alpha Z_R^{-\nu}$ for some real constant α .

¹¹Since past-relations depend on both the spatial coordinates and the conformal-time coordinates of the previously fixed contours (see footnote ⁸), the tree structure exhibits a similar dependence. It would therefore be better to say that each amplitude can be written as a finite sum of products of tree amplitudes, where the amplitudes for any given term in the product are integrated only over some subset of the spacetime coordinates. But since we wish only to establish absolute convergence of the amplitude, it does no harm to extend the spatial integrations for each tree to the full space $\mathbb{R}^{n(D-1)}$ and each λ integrations over the whole of the appropriate contour and to then abuse language by referring to the amplitude as a ‘‘product’’ of tree amplitudes without mentioning the remaining sum explicitly.

It is now useful to break up the integration domain into several pieces. First consider the portion of $C_{\mathbf{y}}$ noted above that protrudes from the past light cone of $\text{Re}Y'$. The past singularity which has stopped this contour from being deformed further is at $\lambda_y^+ = \lambda_{y'} + \|\mathbf{y} - \mathbf{y}'\|$ and the corresponding effective past singularity is at $\lambda_y^{+, \text{eff}} = \lambda_y^+ - b$ on the complex λ_y plane. Hence, the length of this portion of the contour is bounded by a constant, which is larger than $2b$ because the contour has a finite width. We also find that $\text{Re}\lambda_y \geq \lambda_0 + c \|\mathbf{y} - \mathbf{y}'\|$, where λ_0 and c are some positive constants, on this portion of the contour. This is because $\text{Re}\lambda_y \geq \lambda_{\min}$, where λ_{\min} is the minimum of the real part of λ_y at $\mathbf{y} = \mathbf{y}'$, and that $\text{Re}\lambda_y^{+, \text{eff}} / \|\mathbf{y} - \mathbf{y}'\| \rightarrow 1$ as $\|\mathbf{y}\| \rightarrow \infty$. The contribution to I from the protruding portions is thus bounded by a constant times $B(M) \times \int d^{D-1}\mathbf{y} (\lambda_0 + c \|\mathbf{y}\|)^{-D}$.

Next consider the contribution to I from the region $0 < Z_R < Z_0$. This is bounded by $\beta B(M)$ times the total measure $\int d^{D-1}\mathbf{y} \int |d\lambda| |\lambda|^{-D}$ of this region, where β is a constant, assuming that this measure is finite. To see that this is so, consider any point X in the Poincaré patch of real de Sitter space and, furthermore, consider the part of its past light cone that is both within embedding distance Z_0 and which also lies to the future of the cosmological horizon. This region is compact and thus has finite volume. Since widening of the contour described above for large $\text{Re}\lambda$ has little effect at large Z , for fixed ϵ , we may therefore choose Z_0 large enough that the measure of the desired region in complex de Sitter is within, say, a factor of 2 of the volume of the region just discussed in real de Sitter space. Thus, this part of our integral is easily bounded.

We can similarly bound the contribution from the region $Z_R > Z_0$. For large enough Z_0 , this contribution is no more than, say, a factor of 2 times the integral of $\alpha Z_R^{-\nu}$ over the region of real de Sitter space lying to the future of the cosmological horizon but more than an embedding distance Z_0 to the past of the point $(\text{Re}\lambda_{y'}, \mathbf{y}')$. To proceed further, one should compute the volume of surfaces lying a constant embedding distance Z_R to the past of the given point but to the future of the cosmological horizon. In the limit of large Z_R , this volume turns out to approach the constant $1/(D-1)$. We also note that the proper time difference between the two surfaces at Z_R and $Z_R + dZ_R$ is dZ_R/Z_R for large Z_R . As a result, for large enough Z_0 the contribution from the region $\text{Re}Z > Z_0$ is bounded by, say, $4\alpha(D-1)^{-1} \int_{Z_0}^{\infty} dZ_R Z_R^{-1-\nu}$. Combining this with our observations above shows that (4.3) is bounded by some constant $B(I)$ which (for, say, $|\epsilon| < 1$) depends only on the mass of our field and which, in particular, is independent of both ϵ and the location of the point Y' .

As a result, we can bound the integral corresponding to any of the above tree diagrams by $B(I)$ times the integral corresponding to the diagram shortened by cutting off a

lowest line. We can clearly repeat this procedure and continue to remove the lowest lines until we are left with no lines at all. Thus, the integral corresponding to each arrowed tree diagram is bounded by $(B(I))^n$, where n is the number of lines in the given tree. Hence, the integral for the amplitude \mathcal{A}_P given by (4.1) is (absolutely) convergent after translating the contours appropriately.

B. Analyticity of the amplitude \mathcal{A}_P

To complete the argument for equivalence between the Euclidean and Poincaré in-in correlators, we now establish the desired analyticity property of the amplitude \mathcal{A}_P , which we have shown above to be well defined. Specifically, we will show that \mathcal{A}_P is analytic as a function of the conformal times $\tilde{\lambda}_i$ of the external points if $(\tilde{\lambda}_1, \dots, \tilde{\lambda}_m) \in U = \{(\mu_1, \dots, \mu_m) \in \mathbb{C}^m: \text{Im}\mu_i < \text{Im}\mu_{i+1}, i = 1, \dots, m-1\}$, or more generally if the imaginary parts of $\tilde{\lambda}_i$ are all distinct, for any given spatial coordinates \mathbf{x}_i . For this purpose we introduce an additional regulator defined by some $s > 0$ and show that the regulated correlators are analytic functions on U . We then show that this analyticity property persists in the $s \rightarrow 0$ limit.

Our choice of regulator is straightforward to introduce. We define the amplitude $\mathcal{A}_{P,s}$ for $s > 0$ by simply replacing each (already Pauli-Villars regulated) propagator $\Delta^{\text{reg}}(Z)$ with $\Delta_s^{\text{reg}}(Z) = \Delta^{\text{reg}}(Z - s)$, where these propagators are written as functions of the embedding distance Z defined by (2.16). Note that s is indeed a regulator in the sense that it widens the gap between any pair of past and future singularities such as those shown in Figs. 10 and 11. As a result, any contour that can also be used to compute the unregulated \mathcal{A}_P can be used to compute $\mathcal{A}_{P,s}$ for $s > 0$. Thus, contours similar to Fig. 9 are again allowed for $(\tilde{\lambda}_1, \dots, \tilde{\lambda}_m) \in U$. The (absolute) convergence of the integrals for $\mathcal{A}_{P,s}$ can be established in exactly the same way as in the $s = 0$ case.

Now consider complex $\tilde{\lambda}_i$ derivatives of $\mathcal{A}_{P,s}$ computed formally by differentiating the integrand, which is a product of propagators, and then integrating over the contours. Our s regularization makes the integrand analytic in an open neighborhood of $(\tilde{\lambda}_1, \dots, \tilde{\lambda}_m)$ with the contours fixed so that complex derivatives of the integrand are well defined. Furthermore, differentiated propagators are bounded at fixed s and their behavior as $Z \rightarrow \infty$ is not worse than that of undifferentiated propagators. Hence, the argument for the (absolute) convergence of the integrals defining $\mathcal{A}_{P,s}$ applies equally well to integrals of the differentiated integrands. But absolute convergence guarantees that these latter integrals do in fact give the complex $\tilde{\lambda}_i$ derivatives of $\mathcal{A}_{P,s}$. It follows that such integrals are well defined and that each $\mathcal{A}_{P,s}$ is analytic in U .

Now, since the integrals defining \mathcal{A}_P converge, it is clear that $\mathcal{A}_{P,s}$ tends to \mathcal{A}_P as $s \rightarrow 0$. As for the λ_i

derivative of $\mathcal{A}_{P,s}$, the integrand will be divergent in the $s \rightarrow 0$ limit only where the arguments of the differentiated external (regulated) propagator, become coincident. However, due to our Pauli-Villars regularization this divergence is very mild and does not spoil absolute convergence. It follows that the λ_i derivative of $\mathcal{A}_{P,s}$ has a finite limit as $s \rightarrow 0$ which gives the λ_i derivative of \mathcal{A}_P . In particular, these derivatives are well defined on U , so that \mathcal{A}_P is analytic in this domain. This completes our step 3.

Let us now assemble the facts demonstrated above to establish the equivalence of the Euclidean and Poincaré in-in correlators. The amplitude \mathcal{A}_P and the corresponding Euclidean amplitude, which we call \mathcal{A}_E , are both analytic functions of the conformal-time variables $\tilde{\lambda}_i$ of the external points $(\tilde{\lambda}_i, \mathbf{x}_i)$ if $(\tilde{\lambda}_1, \dots, \tilde{\lambda}_m) \in U$ (step 3). These amplitudes coincide in the limit where the imaginary parts of the conformal-time variables tend to zero if the limits of the external points all lie in the static patch of real de Sitter space. (This was established in two steps: In step 1 we established that \mathcal{A}_E agrees with the static in-in amplitude, and in step 2 we established (rather trivially) that the latter agrees with the Poincaré in-in amplitude if the limits of the external points are all in the static patch of real de Sitter space.) Hence, by uniqueness of analytic continuation,¹² $\mathcal{A}_P = \mathcal{A}_E$ for all $\tilde{\lambda}_i$ wherever these amplitudes are well defined. Then, \mathcal{A}_P and \mathcal{A}_E have, of course, the same limit as $\text{Im}\tilde{\lambda}_i \rightarrow 0$, producing the same physical amplitude for any points X_i in the Poincaré patch.

V. EXPLICIT CHECKS IN SIMPLE EXAMPLES

As a check on our arguments, we now explicitly compare the Euclidean and Poincaré in-in results for one-loop corrections to propagators from ϕ^4 and ϕ^3 interactions. As the Euclidean computations (including the analytic continuation to Lorentz-signature de Sitter) were performed in [23], we focus on the in-in calculations here.

A. ϕ^4 correction

Consider the 1-loop correction to the propagator due to an interaction term of the type $\mathcal{L}_{\text{int}}[\phi] = -\frac{\lambda}{4!}\phi(X)^4$. The relevant Feynman diagram is shown in Fig. 13(a). The in-in correlation function is given by

¹²Here, we are using the agreement of \mathcal{A}_P and \mathcal{A}_E on an open subset of a real section, $B = \{(\mu_1, \dots, \mu_m) \in \mathbb{C}^m: \text{Im}\mu_i = 0, i = 1, \dots, m\}$, on the boundary of the region of analyticity U to conclude $\mathcal{A}_P = \mathcal{A}_E$ in U . This is a simple corollary of Bogolubov's edge-of-the-wedge theorem (see, e.g., Theorem 2-17 in [38]).




FIG. 13. The 1-loop corrections to the propagator.

$$\begin{aligned} \langle T\phi(X_1)\phi(X_2) \rangle = & -\frac{\lambda i}{2} \int_Y \{ \Delta_{m_3}^{\text{reg}}(Y, Y) \Delta_{m_1}^{\text{reg}}(Y, X_1) \Delta_{m_2}^{\text{reg}}(Y, X_2) \\ & - \Delta_{m_3}^{\text{reg}*}(Y, Y) W_{m_1}^{\text{reg}}(Y, Y_1) W_{m_2}^{\text{reg}}(Y, X_2) \} \end{aligned} \quad (5.1)$$

Here $\int_Y \dots$ denotes an integral over the Poincaré patch, and we remind the reader that $\Delta_{m^2}(X, Y)$, $\Delta_{m^2}^*(X, Y)$, and $W_{m^2}(X, Y)$ are the time-ordered, anti-time-ordered, and Wightman 2-point functions of the Gaussian theory. It is convenient to let each line in the Feynman diagram have a distinct mass; one may take the limit of equal masses later. This expression has a UV divergence for $D \geq 4$ which we control by using Pauli-Villars regularization. For simplicity, we regulate only the internal lines, though we could of course also regulate the external lines as well.

To simplify (5.1) we first note that the regulated Feynman function $\Delta_{m^2}^{\text{reg}}(X, Y)$ evaluated at coincident points is real and independent of position, so $\Delta_{m^2}^{\text{reg}}(Y, Y) = \Delta_{m^2}^{\text{reg}*}(Y, Y) := \Delta_{m^2}^{\text{reg}}(1)$. After removing a common factor of $\Delta_{m_3}^{\text{reg}}(1)$ from the integrand the remaining integral is

$$\begin{aligned} I(X_1, X_2) := & i \int_Y \{ \Delta_{m_1}^{\text{reg}}(Y, X_1) \Delta_{m_2}^{\text{reg}}(Y, X_2) \\ & - W_{m_1}^{\text{reg}}(Y, X_1) W_{m_2}^{\text{reg}}(Y, X_2) \}. \end{aligned} \quad (5.2)$$

The integral (5.2) can be quickly performed as follows. Consider a theory of two free massive scalar fields $\Phi_{1,2}(X)$ with masses $M_1^2 \neq M_2^2$. We can rewrite this theory in terms of two new fields $\phi_{1,2}(X)$ by performing an $SO(2)$ rotation in field space:

$$\begin{aligned} \phi_1(X) &= \cos\omega \Phi_1(X) - \sin\omega \Phi_2(X), \\ \phi_2(X) &= \sin\omega \Phi_1(X) + \cos\omega \Phi_2(X). \end{aligned} \quad (5.3)$$

The fields $\phi_{1,2}(X)$ have masses $m_{1,2}^2$ that are functions of $M_{1,2}^2$ and ω , and also an interaction $-g\phi_1(X)\phi_2(X)$ in the Lagrangian with the coupling $g = (M_1^2 - M_2^2) \sin\omega \cos\omega$. Now consider the correlation function $\langle T\phi_1(X_1)\phi_2(X_2) \rangle$. We may compute this correlation function using standard in-in perturbation theory; the term at lowest order in g (or equivalently, in ω) is

$$\langle T\phi_1(X_1)\phi_2(X_2) \rangle = -gI(X_1, X_2) + O(g^3). \quad (5.4)$$

On the other hand, by simply using (5.3) we can compute $\langle T\phi_1(X_1)\phi_2(X_2) \rangle$ exactly¹³:

$$\begin{aligned} \langle T\phi_1(X_1)\phi_2(X_2) \rangle &= \sin\omega \cos\omega [\langle T\Phi_1(X_1)\Phi_1(X_2) \rangle - \langle T\Phi_2(X_1)\Phi_2(X_2) \rangle] \\ &= \sin\omega \cos\omega [\Delta_{M_1^2}(X_1, X_2) - \Delta_{M_2^2}(X_1, X_2)]. \end{aligned} \quad (5.5)$$

We can then write M_1^2 , M_2^2 and ω in terms of m_1^2 , m_2^2 and g , expand the right-hand side of (5.5) in a power series in g , and equate the $O(g)$ term with the right-hand side of (5.4). The result is the equality

$$I(X_1, X_2) = \frac{\Delta_{m_1^2}(X_1, X_2) - \Delta_{m_2^2}(X_1, X_2)}{m_2^2 - m_1^2}. \quad (5.6)$$

Returning to (5.1), we may use (5.6) to obtain

$$\begin{aligned} \langle T\phi(X_1)\phi(X_2) \rangle &= \frac{\lambda}{2} \Delta_{m_3}^{\text{reg}}(1) \left[\frac{\Delta_{m_1^2}(X_1, X_2) - \Delta_{m_2^2}(X_1, X_2)}{m_1^2 - m_2^2} \right]. \end{aligned} \quad (5.7)$$

It is clear that the same steps can be used to compute the Euclidean expression. The analogue of (5.2) then involves only Euclidean propagators, but these are just what are needed to arrive at the analogue of (5.6). After analytic continuation to real de Sitter space, the result is precisely (5.7).

We note that the above calculations could be performed equally well using dimensional regularization rather than the Pauli-Villars scheme. In dimensional regularization the computation is performed in an arbitrary real dimension which is sufficiently small such that there are no ultraviolet divergences. As in Pauli-Villars regularization, the values of de Sitter-invariant Green's functions $\Delta_{m^2}(1)$, etc., are divergent but de Sitter-invariant constants. By the usual arguments [39], the manipulations we performed to derive (5.6) and its Euclidean analogue are valid for arbitrary real dimension.

B. ϕ^3 correction

Next we turn to the 1-loop correction to the propagator that arises from the interaction $\mathcal{L}_{\text{int}}[\phi] = -\frac{g}{3!}\phi^3(X)$. The relevant Feynman diagram is shown in Fig 13(b). Once again it is convenient to let each leg of this diagram have a distinct mass. This correction has a UV divergence for spacetime dimension $D \geq 4$. Both to draw on results of

¹³A truly skeptical reader might ask whether (5.4) must necessarily give the vacuum correlator of the theory defined by (5.3). But at this order the result must be a Gaussian state invariant under translations, rotations, and the scaling symmetry of the Poincaré patch. This determines the state uniquely, assuming that the results are finite. Finiteness in turn can be shown by either a careful direct analysis or by using the results of [12] to expand both m_1^2 and m_2^2 about the conformal-coupling value $m_c^2 = \frac{1}{4}D(D-2)$ and then using the explicit calculations of that reference.

[23] and to simplify the arguments, we carry out the computations below using dimensional regularization. In particular, we will compute this correction in arbitrary $D < 2$, then analytically continue D to extend the result to higher dimensions. However, we also explain how similar results (with more complicated explicit forms) can be obtained via Pauli-Villars techniques.

It is useful to introduce a so-called linearization formula for the Green's functions $\Delta_{m^2}(X, Y)$, $\Delta_{m^2}^*(X, Y)$ and $W_{m^2}(X, Y)$. We use the variable $\alpha := (D - 1)/2$ to keep track of spacetime dimension and the mass variable σ defined by the equation $-\sigma(\sigma + 2\alpha) = m^2 \ell^2$. All three Green's functions are proportional to the Gegenbauer function $C_\sigma^\alpha(Z)$. The following linearization formula for the Gegenbauer function allows us to replace a product of Gegenbauer functions with an integral of a single Gegenbauer function [23]:

$$C_{\sigma_1}^\alpha(Z)C_{\sigma_2}^\alpha(Z) = -\frac{4\pi^\alpha}{\Gamma(\alpha)} \sin(\pi\sigma_1) \sin(\pi\sigma_2) \times \int_\mu \frac{\rho_{\sigma_1\sigma_2}^\alpha(\mu)}{\sin(\pi\mu)} C_\mu^\alpha(Z). \quad (5.8)$$

In this equation $C_\sigma^\alpha(Z)$ is the Gegenbauer function which is analytic in the complex Z plane cut along $Z \in (-\infty, -1]$. We assume $\text{Re}\sigma_1 < 0$ and $\text{Re}\sigma_2 < 0$, which is valid for $m_{1,2}^2 > 0$. The shorthand $\int_\mu \dots$ denotes a contour integral in the complex μ plane with measure $d\mu/2\pi i$. The integration contour runs from $-i\infty$ to $+i\infty$ within the strip $\text{Re}(\sigma_1 + \sigma_2) < \text{Re}\mu < 0$. Within this strip the integrand is analytic and the contour integral converges absolutely. From (5.8) we may write the following linearization

$$\begin{aligned} \langle T\phi(X_1)\phi(X_2) \rangle &= -g^2 \int_{Y_1} \int_{Y_2} \{ \Delta_{\sigma_1}(Y_1, X_1)\Delta_{\sigma_2}(Y_2, X_2)\Delta_{\sigma_3}(Y_1, Y_2)\Delta_{\sigma_4}(Y_1, Y_2) \\ &\quad - W_{\sigma_1}(Y_1, X_1)\Delta_{\sigma_2}(Y_2, X_2)W_{\sigma_3}(Y_1, Y_2)W_{\sigma_4}(Y_1, Y_2) + W_{\sigma_1}(Y_1, X_1)W_{\sigma_2}(Y_2, X_2)\Delta_{\sigma_3}^*(Y_1, Y_2)\Delta_{\sigma_4}^*(Y_1, Y_2) \\ &\quad - \Delta_{\sigma_1}(Y_1, X_1)W_{\sigma_2}(Y_2, X_2)W_{\sigma_3}(Y_2, Y_1)W_{\sigma_4}(Y_2, Y_1) \}. \end{aligned} \quad (5.11)$$

The first two terms in (5.11) contain the integral over Y_1 :

$$T_1 := \int_{Y_1} \{ \Delta_{\sigma_1}(Y_1, X_1)\Delta_{\sigma_3}(Y_1, Y_2)\Delta_{\sigma_4}(Y_1, Y_2) - W_{\sigma_1}(Y_1, X_1)W_{\sigma_3}(Y_1, Y_2)W_{\sigma_4}(Y_1, Y_2) \}. \quad (5.12)$$

To compute T_1 we first use the linearization formula (5.9) in each term, then use (5.6) to integrate over Y_1 :

$$\begin{aligned} T_1 &= \int_\mu \rho_{\sigma_3\sigma_4}^\alpha(\mu) \int_{Y_1} \{ \Delta_{\sigma_1}(Y_1, X_1)\Delta_\mu(Y_1, Y_2) - W_{\sigma_1}(Y_1, X_1)W_\mu(Y_1, Y_2) \} \\ &= \frac{1}{i} \int_\mu \rho_{\sigma_3\sigma_4}^\alpha(\mu) \left[\frac{\Delta_\mu(Y_2, X_1) - \Delta_{\sigma_1}(Y_2, X_1)}{m_1^2 - m_\mu^2} \right] \\ &= \frac{1}{i} \int_\mu \frac{\rho_{\sigma_3\sigma_4}^\alpha(\mu)}{m_1^2 - m_\mu^2} \Delta_\mu(Y_2, X_1). \end{aligned} \quad (5.13)$$

¹⁴The definition of $\rho_{\sigma_1\sigma_2}^\alpha(\mu)$ used here is (-2) times the $\rho_{\sigma_1\sigma_2}^\alpha(L)$ of that paper.

formula for the Green's functions, with $H_\sigma(X, Y)$ standing for $\Delta_\sigma(X, Y)$, $\Delta_\sigma^*(X, Y)$, or $W_\sigma(X, Y)$:

$$H_{\sigma_1}(X, Y)H_{\sigma_2}(X, Y) = \int_\mu \rho_{\sigma_1\sigma_2}^\alpha(\mu)H_\mu(X, Y). \quad (5.9)$$

Of course, most of the content of (5.9) is contained in the details of the function $\rho_{\sigma_1\sigma_2}^\alpha(\mu)$. The explicit form of $\rho_{\sigma_1\sigma_2}^\alpha(\mu)$ can be found in [23]¹⁴; we will not need the explicit form. We need only note that:

- (i) $\rho_{\sigma_1\sigma_2}^\alpha(\mu)$ is itself analytic in the region $\text{Re}\mu > \text{Re}(\sigma_1 + \sigma_2)$ and that in this region the function behaves at large $|\mu| \gg 1$ like $|\mu|^{2\alpha-3} \log(\mu)$. In particular, it follows that

$$\int_\mu \frac{\rho_{\sigma_1\sigma_2}^\alpha(\mu)}{m^2 + \mu(\mu + 2\alpha)} = 0 \quad \text{for } \alpha < 2 \quad (5.10)$$

for $m^2 > 0$ with the μ contour lying to the right of poles at $\mu = -\alpha \pm \sqrt{\alpha^2 - m^2}$ (both of which lie in the left half-plane).

- (ii) The function $\rho_{\sigma_1\sigma_2}^\alpha(\mu)$ is proportional to $\Gamma(2 - 2\alpha)$ and so has simple poles as a function of α at $\alpha = 1, 3/2, 2, \dots$. Of course, the left-hand sides of (5.8) and (5.9) are regular for these values of α ; the integral over μ cancels these poles. However, the integral of an arbitrary function of μ times $\rho_{\sigma_1\sigma_2}^\alpha(\mu)$ will generically not cancel this divergence and so will diverge at these values of α .

The $O(g^2)$ correction to the propagator in this theory is given in the in-in formalism by the expression

We compute with $\alpha < 3/2$, so the final equality follows from (5.10). The latter two terms in (5.11) contain the integral over Y_1 :

$$T_2 := \int_{Y_1} \{W_{\sigma_1}(Y_1, X_1)\Delta_{\sigma_3}^*(Y_1, Y_2)\Delta_{\sigma_4}^*(Y_1, Y_2) - \Delta_{\sigma_1}(Y_1, X_1)W_{\sigma_3}(Y_2, Y_1)W_{\sigma_4}(Y_2, Y_1)\}. \quad (5.14)$$

To compute T_2 we again use the linearization formula (5.9), then use the integral

$$\begin{aligned} J(X_1, X_2) &:= i \int_Y \{W_{\sigma_1}(X_1, Y)\Delta_{\sigma_2}(Y, X_2) \\ &\quad - \Delta_{\sigma_1}^*(X_1, Y)W_{\sigma_2}(Y, X_2)\} \\ &= \frac{W_{\sigma_1}(X_1, X_2) - W_{\sigma_2}(X_1, X_2)}{m_2^2 - m_1^2}. \end{aligned} \quad (5.15)$$

This integral may be derived in the same manner as $I(X_1, X_2)$ by examining the Wightman correlation function $\langle \phi_1(x_1)\phi_2(x_2) \rangle$ in the $SO(2)$ -rotated theory.

Inserting (5.15) into (5.14) yields

$$\begin{aligned} T_2 &= \int_{\mu} \rho_{\sigma_3\sigma_4}^{\alpha}(\mu) \int_{Y_1} \{W_{\sigma_1}(Y_1, X_1)\Delta_{\mu}^*(Y_1, Y_2) \\ &\quad - \Delta_{\sigma_1}(Y_1, X_1)W_{\mu}(Y_2, Y_1)\} \\ &= -\frac{1}{i} \int_{\mu} \rho_{\sigma_3\sigma_4}^{\alpha}(\mu) \left[\frac{W_{\mu}(Y_2, X_1) - W_{\sigma_1}(Y_2, X_1)}{m_1^2 - m_{\mu}^2} \right] \\ &= -\frac{1}{i} \int_{\mu} \frac{\rho_{\sigma_3\sigma_4}^{\alpha}(\mu)}{m_1^2 - m_{\mu}^2} W_{\mu}(Y_2, X_1). \end{aligned} \quad (5.16)$$

Once again the last equality follows from (5.10). Assembling (5.13) and (5.16) we may write the propagator correction as

$$\begin{aligned} \langle T\phi(X_1)\phi(X_2) \rangle &= -\frac{g^2}{i} \int_{\mu} \frac{\rho_{\sigma_3\sigma_4}^{\alpha}(\mu)}{m_1^2 - m_{\mu}^2} \int_{Y_2} \{\Delta_{\sigma_2}(Y_2, X_2)\Delta_{\mu}(Y_2, X_1) \\ &\quad - W_{\sigma_2}(Y_2, X_2)W_{\mu}(Y_2, X_1)\}. \end{aligned} \quad (5.17)$$

The remaining integral over Y_2 may be performed using (5.6):

$$\begin{aligned} \langle T\phi(X_1)\phi(X_2) \rangle &= g^2 \int_{\mu} \frac{\rho_{\sigma_3\sigma_4}^{\alpha}(\mu)}{m_1^2 - m_{\mu}^2} \left[\frac{\Delta_{\mu}(X_1, X_2) - \Delta_{\sigma_2}(X_1, X_2)}{m_2^2 - m_{\mu}^2} \right] \\ &= g^2 \int_{\mu} \frac{\rho_{\sigma_3\sigma_4}^{\alpha}(\mu)}{(m_1^2 - m_{\mu}^2)(m_2^2 - m_{\mu}^2)} \Delta_{\mu}(X_1, X_2). \end{aligned} \quad (5.18)$$

The expected UV divergence of this expression is in the factor $\Gamma(2 - 2\alpha)$ contained in $\rho_{\sigma_1\sigma_2}^{\alpha}(\mu)$.

The Euclidean computation is essentially identical, using the analogue of (5.15) involving only Euclidean propagators,¹⁵ so that the results agree under analytic continuation as desired. The details of the Euclidean calculation were given in [23], where it is also shown that both the final expression and the counterterms used to render a finite expression in higher dimensions agree with the standard flat-space results in the limit $\ell \rightarrow \infty$.

One can perform essentially the same computations using Pauli-Villars regularization instead of dimensional regularization. Note that the key steps above were the linearization formula (5.8), the property (5.10) of the form factor $\rho_{\sigma_1\sigma_2}^{\alpha}$, and the composition rules (5.6) and (5.15). But it is clear from the derivation in [23] that a similar linearization formula can be used to express the product of two Pauli-Villars regularized Green's functions as an integral over (unregularized) Gegenbauer functions. In this case, the corresponding form factor $\rho_{\sigma_1\sigma_2}^{\alpha, M}$ is manifestly finite for all α , but depends on the Pauli-Villars regulator mass M . While $\rho_{\sigma_1\sigma_2}^{\alpha, M}$ is analytic as above, it falls off faster at large μ so that the analogue of (5.10) is in fact satisfied for all α . Expanding any remaining regularized propagators as a sum of unregularized propagators then allows us to apply the composition rules (5.6) and (5.15) and to complete the calculation. The result is similar to that above with the replacement $\rho_{\sigma_1\sigma_2}^{\alpha} \rightarrow \rho_{\sigma_1\sigma_2}^{\alpha, M}$ and with extra terms coming from the regulators. The Euclidean Pauli-Villars computation proceeds in precisely the same way and again agrees after analytic continuation.

Finally, we note that the analogous 1-loop correction to the Wightman function $\langle \phi(X_1)\phi(X_2) \rangle$ of this theory was recently considered by Krotov and Polyakov (see Sec. 6 of [17]; the same correlation function is considered in Sec. 7, but with respect to a different state). Our result for this correlation function is simply the right-hand side of (5.18) with the replacement $\Delta_{\mu}(X_1, X_2) \rightarrow W_{\mu}(X_1, X_2)$. It is difficult to compare these two results exactly because the result of [17] has not been renormalized (our renormalized result is presented in [23]). However, we can safely compare the behavior of the two results in the infrared where the effect of renormalization is clear. To compare with [17] we set all masses to be equal. Using techniques presented in [23] we find the leading behavior at large $|Z_{12}| \gg 1$ to be

¹⁵The Euclidean analogue of (5.15) is identical to the Euclidean analogue of (5.2).

$$\begin{aligned}
& \langle \phi(X_1)\phi(X_2) \rangle \\
&= \frac{g^2 \rho_{\sigma\sigma}^\alpha(\sigma) - \delta m^2 + m^2 \delta\phi}{16\pi^{\alpha+1}(\sigma + \alpha)^2} \{ \Gamma(-\sigma)\Gamma(\sigma + \alpha) \\
&\quad \times (-2Z_{12})^\sigma \log Z_{12} + \Gamma(\sigma + 2\alpha)\Gamma(-\sigma - \alpha) \\
&\quad \times (-2Z_{12})^{-(\sigma+2\alpha)} \log Z_{12} \} [1 + O(Z_{12}^{-1})]. \quad (5.19)
\end{aligned}$$

Here δm^2 and $\delta\phi$ are the real, divergent, coefficients of the mass and field renormalization counterterms which cancel the divergent terms in $\rho_{\sigma\sigma}^\alpha(\sigma)$. We find the same asymptotic dependence on Z_{12} as [17]; in particular, while the Wightman function of the free theory has two asymptotic branches which decay like Z_{12}^σ and $Z_{12}^{-(\sigma+2\alpha)}$, the $O(g^2)$ correction has two asymptotic branches that each decay slower by a multiplicative factor of $\log Z_{12}$.

The authors of [17] interpret the appearance of the logarithm in the asymptotic behavior (5.19) as an indication of an ‘‘infrared correction’’ to the correlator. Indeed, the logarithm indicates that the 1-loop correction induces an $O(g^2)$ correction to the mass parameter σ ; as a result, the asymptotic expansion of the correlator is altered in perturbation theory like $(Z)^{\sigma+O(g^2)} = O(g^2)(Z_{12})^\sigma \times \log Z_{12} + O(g^4)$. The $O(g^2)$ correction to σ can be computed by performing the sum over 1PI diagrams of the form of Fig. 13(b). This analysis was performed in detail [23]. There it was found that, at least for scalar fields with bare masses belonging to the principal series of $SO(D, 1)$, the $O(g^2)$ correction to σ has a finite negative real part (equivalently, the correction introduces a finite negative imaginary part to the self-energy) which cannot be removed with a local Hermitian counterterm. Thus the $O(g^2)$ correction unambiguously *increases* the rate of decay of the 1PI-summed correlator so that this correlator decays *faster* than any free Wightman function. This agrees with the analogous computation in flat-space where the 1PI-summed correlator also enjoys an enhanced exponential rate of decay at large separations [40].

VI. DISCUSSION

We have shown that Euclidean techniques and in-in perturbation theory on the Poincaré (a.k.a. cosmological) patch of de Sitter yield identical correlation functions for scalar field theories with positive masses. This is in contrast with the situation for the in-in perturbation theory defined by global coordinates on de Sitter, where the corresponding factorization property fails [17] and the in-in scheme contains infrared divergences. Our equivalence holds diagram by diagram and for any finite value of appropriate Pauli-Villars regulator masses. It thus also holds for the fully renormalized diagrams. While we focussed on nonderivative interactions, interactions involving derivatives can be handled in precisely the same way so long as additional Pauli-Villars subtractions are made as described in Sec. II B. We used a 3-step argument in the

main text, though a more direct analytic continuation is described in the Appendix.

As a check on the above arguments, we also explicitly calculated the one-loop propagator corrections due to both ϕ^3 and ϕ^4 interactions for all masses and in all dimensions in Sec. V. The Poincaré in-in and Euclidean calculations agreed precisely.¹⁶ We suspect that methods similar to those used in Sec. V, perhaps combined with Mellin-Barnes techniques as in [24], could be used to give a rather direct diagram-by-diagram proof of the equivalence of Euclidean and Poincaré in-in techniques, but we have not explored the details.

A number of points merit further discussion. First, some physicists have conjectured that in-in calculations in the Poincaré patch lead to IR divergences, even for fields with $m^2 > 0$ due to contributions with vertices at large conformal time λ . But there are clearly no such divergences in Euclidean signature. So how can the two forms of perturbation theory agree diagram by diagram? We believe that, if there are such divergences, they are better classified as ultraviolet (UV) divergences and are associated with the fact that the limit $\lambda \rightarrow \infty$ defines a null surface (the cosmological horizon) so that light-cone singularities can arise even at what appear to be large separations between points.

To a certain extent, the classification of these divergences as UV or IR in the cosmological patch may be a matter of semantics. What is important is that any divergences may be cancelled using only local counterterms. This much is clear from our analysis: We have seen that adding a Pauli-Villars regulator M^2 removes all divergences, and that the in-in and Euclidean calculations agree at all finite values of M^2 . This means that they have the same divergence structure in the limit $M^2 \rightarrow \infty$, and that divergences can be removed using the same sets of counterterms. But all divergences for massive theories on S^D are clearly ultraviolet in nature and so are the same as on \mathbb{R}^d . Local counterterms suffice to remove them.

Second, the reader will recall that the argument given in Sec. III to show factorization (i.e., that the vertical sections of the contour at infinite past may be neglected) required the propagators to fall off at large timelike separations. Without such falloff, the two formalisms should not agree. Instead, analytic continuation of the Euclidean perturbation theory would give the terms of the in-in formalism, together with terms associated with integrals over some contour at infinity in the complex t plane. How then should

¹⁶We have also used a combination of analytic and numerical techniques to check the agreement of Poincaré in-in and Euclidean correlators for the tree-level 3-point function for $D = 4$ for $m^2 = 2$ (conformal coupling) and also for the one-loop correction to the 4-point function for $D = 3$ and $m^2 = 3/4$ (also conformal coupling) evaluated at two pairs of coincident points. Both of these diagrams are finite and require no regularization. Our numerics indicate agreement to at least one part in 10^7 . As these calculations do not yield significant insights, we have refrained from presenting the details.

we interpret this disagreement? If the propagators do not fall off at large times, then integrals over the contour at infinity will generally diverge. Thus, one would expect at most one formalism to give finite results. Let us suppose that the Euclidean formalism is well defined and finite. If one can establish the appropriate positivity properties, then analytic continuation will define a good quantum state. In this case, it would appear that any divergences of the in-in formalism are an unphysical artifact of this particular perturbative framework, and one might hope to better relate the two formalisms through an appropriate resummation of the divergent in-in formalism.

There is some potential for this scenario to hold in perturbative gravity. For example, the tree-level three-point correlator constructed by Maldacena in [41] in the momentum space is IR divergent when inverse Fourier-transformed to position space. On the other hand, the three-point function constructed on S^D using Euclidean propagators would have no IR divergences (see, e.g., [42,43] for $D = 4$). It may therefore be interesting to re-examine the three-point function in Euclidean gravity. However, we note that some physicists have raised objections to these propagators [44,45]. In addition, at least with generic gauge choices the Euclidean gravitational action is not bounded below (though see [46]). This means that one cannot rely on Osterwalder-Schröder arguments [28] to guarantee that analytic continuation of the Euclidean correlators defines a positive-definite Hilbert space, and positivity would need to be verified.

The other possibility when propagators do not fall off is that both forms of perturbation theory are ill-defined. This is the case for massless scalars on de Sitter. But even here the divergences can be an artifact of the particular scheme for perturbation theory. In [26], Rajaraman showed that, in the presence of a ϕ^4 interaction with positive coefficient in the Hamiltonian, the Euclidean scheme can be resummed to give a new well-defined perturbation theory. Since the Euclidean action is bounded below, the resulting Euclidean correlators will satisfy reflection-positivity and can be analytically continued to give a good state of the Lorentzian theory.

We close with a brief comment on other generalizations. Recall that our first step was to verify that the usual connection between Euclidean methods and thermal in-in field theory on a static spacetime holds in the context of the de Sitter static patch. It is clear that similar arguments will hold in the static regions of generic spacetimes with bifurcate Killing horizons, so long as the propagators again fall off sufficiently quickly at large separations. For a particularly amusing application, consider the standard Minkowski space correlators (in the Minkowski vacuum) for which the usual perturbation theory integrates the vertices of Feynman diagrams over all of Minkowski space. We now see that, so long as their arguments are taken to lie in, say, the right Rindler wedge, these

correlators can in fact be computed using in-in perturbation theory in the Rindler wedge, and thus by integrating vertices of the in-in diagrams only over this Rindler wedge. One would expect this fact to be well known, but we have been unable to find any discussions in the literature.

ACKNOWLEDGMENTS

We thank C. Fewster, S. Hollands, I. McIntosh, S. Polyakov, A. Rajaraman, A. Roura, M. Srednicki, and T. Tanaka for useful discussions and correspondence. D. M. and I. M. were supported in part by the U.S. National Science Foundation under NSF Grant No. PHY08-55415 and by funds from the University of California. A. H. thanks the Astro-Particle Theory and Cosmology Group and Department of Applied Mathematics at the University of Sheffield and the Physics Department at UCSB for kind hospitality, while part of this work was carried out. His work at UCSB was supported by a Royal Society International Travel Grant.

APPENDIX: DIRECT ANALYTIC CONTINUATION

If an analytic function $f(z_1, \dots, z_N)$ of N variables is integrated over a real N -dimensional compact surface \mathcal{S} with no boundary in \mathbb{C}^N as

$$I = \int_{\mathcal{S}} f(z_1, \dots, z_N) dz_1 \wedge \dots \wedge dz_N, \quad (\text{A1})$$

we have $I = 0$ as long as f has no singularities on or inside \mathcal{S} because the differential form $f(z_1, \dots, z_N) dz_1 \wedge \dots \wedge dz_N$ is closed. This generalization of Cauchy's theorem can be used for the analytic continuation of correlators in the Euclidean formalism to those in the Poincaré in-in formalism. In either formalism the integration is over a manifold of the form M^n , where M is a real D -dimensional surface in complexified sphere, \mathbb{S}^D , and where n is the number of internal vertices. We showed in Sec. III that we can take $M = C_E \times S_h^{D-1}$ where C_E is a contour similar to that shown in Fig. 6 and where S_h^{D-1} is a $D - 1$ dimensional half-sphere in the Euclidean formalism. On the other hand, we take $M = C_P \times \mathbb{R}^{D-1}$ where C_P is a contour on the complex λ plane with measure $d\lambda/\lambda^D$ (see Fig. 9) in the Poincaré in-in formalism. The generalized Cauchy's theorem together with the regularization of the propagator in Sec. IV B can be used to show that the amplitude, which is an integral over M^n , is analytically continued as an analytic function of the external points on \mathbb{S}^D if M can be deformed, with the external points moving and remaining on M , without letting it cross any singularities of the integrand.¹⁷ In this Appendix we demonstrate that this deformation of the surface M of integration from

¹⁷We expect that the integrals on all intermediate surfaces can be shown to converge by methods similar to those employed in Secs. III and IV.

$S_1 = C_E \times S_h^{D-1}$ (Euclidean formalism) to $S_2 = C_P \times \mathbb{R}^{D-1}$ (Poincaré in-in formalism) can indeed be achieved.

We start with the surface S_1 for the Euclidean formalism. It can be given in Poincaré coordinates as follows:

$$S_1 = \{(\Lambda e^{i\tau}, \mathbf{X} e^{i\tau}) : \tau \in (-\epsilon, \epsilon), \Lambda^2 - \|\mathbf{X}\|^2 = f(\tau) > 0, \Lambda > 0, \mathbf{X} \in \mathbb{R}^{D-1}\}, \quad (\text{A2})$$

where $f(\tau) \rightarrow \infty$ as $\tau \rightarrow \pm\epsilon$. This can be shown using the following relationship between the static and Poincaré coordinates, (t, θ, \hat{X}) with $\hat{X} \cdot \hat{X} = 1$, and (λ, \mathbf{x}) , respectively:

$$e^{-2t} = \lambda^2 - \mathbf{x} \cdot \mathbf{x}, \quad (\text{A3})$$

$$\hat{X}^i \sin\theta = x^i / \lambda. \quad (\text{A4})$$

On the other hand the contour in Fig. 9, which is the λ contour for the Poincaré in-in formalism before taking the limit $\text{Im}\lambda \rightarrow 0$, corresponds to

$$S_2 = \{([f(\tau)]^{1/2} + i\tau, \mathbf{X}) : \tau \in (-\epsilon, \epsilon), \mathbf{X} \in \mathbb{R}^{D-1}\}. \quad (\text{A5})$$

If the points $X_1 = (\lambda_1, \mathbf{x}_1)$ and $X_2 = (\lambda_2, \mathbf{x}_2)$ are the arguments of a propagator, then (2.16) shows that it is singular if and only if

$$(X_1 - X_2)^2 = -(\lambda_1 - \lambda_2)^2 + (\mathbf{x}_1 - \mathbf{x}_2) \cdot (\mathbf{x}_1 - \mathbf{x}_2) = 0. \quad (\text{A6})$$

It can readily be seen that this equation is not satisfied by any pair of distinct points on S_1 or S_2 . Since the integrand is a product of propagators with arguments on the surface of integration, what we need to show is that there is a continuous deformation from S_2 to S_1 such that no intermediate surfaces contain two distinct points satisfying (A6).¹⁸ We note that, if the vector $\text{Im}X_1 - \text{Im}X_2$ is timelike, then (A6) does not hold.

First consider the following one-parameter family of surfaces:

¹⁸We can show as in Sec. IV B that coincidence singularities do not spoil the analytic continuation argument here.

$$S_{2,\gamma} = \{(\Lambda + i\tau, \mathbf{X}) : \Lambda^2 - \gamma \|\mathbf{X}\|^2 = f(\tau)\}, \quad (\text{A7})$$

where $f(\tau)$ is the same positive function as in (A2) and where $0 \leq \gamma \leq 1$. Note that $S_{2,0} = S_2$. For any two points $X_j = (\Lambda_j + i\tau_j, \mathbf{X}_j)$, $j = 1, 2$, on $S_{2,\gamma}$, we have

$$\text{Im}X_1 - \text{Im}X_2 = (\tau_1 - \tau_2, \mathbf{0}), \quad (\text{A8})$$

which is timelike if $\tau_1 \neq \tau_2$. If $\tau_1 = \tau_2$, then $(X_1 - X_2)^2 > 0$ because $X_1 = (\Lambda_1, \mathbf{X}_1)$ and $X_2 = (\Lambda_2, \mathbf{X}_2)$ are both on the hyperboloid $\Lambda^2 - \gamma \|\mathbf{X}\|^2 = f(\tau_1)$ with $0 \leq \gamma \leq 1$. Thus, the deformation of S_2 to $S_{2,1}$ leads to analytic continuation of the integral.

Next we consider the following two-parameter family of surfaces:

$$S_{(\alpha,\beta)} = \{((\Lambda + i\alpha\tau)e^{i\beta\tau}, \mathbf{X}e^{i\beta\tau}) : \Lambda^2 - \|\mathbf{X}\|^2 = f(\tau)\}, \quad (\text{A9})$$

where $0 \leq \alpha, \beta \leq 1$. We note that $S_{2,1} = S_{(1,0)}$ and $S_1 = S_{(0,1)}$. Consider two points on $S_{(\alpha,\beta)}$:

$$X_1 = ((\Lambda_1 + i\alpha\tau_1)e^{i\beta\tau_1}, \mathbf{X}_1 e^{i\beta\tau_1}), \quad (\text{A10})$$

$$X_2 = ((\Lambda_2 + i\alpha\tau_2)e^{i\beta\tau_2}, \mathbf{X}_2 e^{i\beta\tau_2}). \quad (\text{A11})$$

Define $\tilde{X}_j := e^{-i\beta\tau_j} X_j$, $j = 1, 2$. Since (A6) is invariant under multiplication of X_1 and X_2 by a common factor, it is not satisfied if $\text{Im}\tilde{X}_1 - \text{Im}\tilde{X}_2$ is timelike. We find

$$\text{Im}\tilde{X}_1 - \text{Im}\tilde{X}_2 = (\Lambda_1 \sin\beta(\tau_1 - \tau_2) + \alpha\tau_1 \cos\beta(\tau_1 - \tau_2) - \alpha\tau_2, \mathbf{X}_1 \sin\beta(\tau_1 - \tau_2)). \quad (\text{A12})$$

If ϵ is sufficiently small—recall $|\tau_1|, |\tau_2| < \epsilon$ —then this vector is timelike for $0 \leq \alpha, \beta \leq 1$ provided that at least one of them is nonzero and that $\tau_1 \neq \tau_2$. If $\tau_1 = \tau_2$, then we have $(\tilde{X}_1 - \tilde{X}_2)^2 > 0$ because $\tilde{X}_1 = (\Lambda_1, \mathbf{X}_1)$ and $\tilde{X}_2 = (\Lambda_2, \mathbf{X}_2)$ are both on the hyperboloid $\Lambda^2 - \|\mathbf{X}\|^2 = f(\tau_1)$. Thus, the deformation of $S_{2,1}$ to S_1 gives analytic continuation of the integral if ϵ is sufficiently small, and, hence, so does the deformation of S_2 to S_1 . When combined with the various convergence and falloff arguments from the main text, this result implies that the correlators computed using the contour of Fig. 9 in the Poincaré in-in formalism is equal to the corresponding analytic continuation of the Euclidean correlators.

[1] B. Allen, *Phys. Rev. D* **32**, 3136 (1985).

[2] A. A. Starobinsky, *Pis'ma Zh. Eksp. Teor. Fiz.* **30**, 719 (1979) [*JETP Lett.* **30**, 682 (1979)].

[3] E. Mottola, *Phys. Rev. D* **31**, 754 (1985); *Physical Origins of Time Asymmetry*, edited by J. J. Halliwell *et al.* (Cambridge University Press, Cambridge, UK, 1993),

p. 504; I. Antoniadis, P. O. Mazur, and E. Mottola, *New J. Phys.* **9**, 11 (2007); E. Mottola, arXiv:1008.5006.

[4] B. L. Hu and D. J. O'Connor, *Phys. Rev. Lett.* **56**, 1613 (1986).

[5] B. L. Hu and D. J. O'Connor, *Phys. Rev. D* **36**, 1701 (1987).

- [6] N. C. Tsamis and R. P. Woodard, *Phys. Lett. B* **301**, 351 (1993); N. C. Tsamis and R. P. Woodard, *Ann. Phys. (N.Y.)* **238**, 1 (1995); *Nucl. Phys.* **B474**, 235 (1996); *Ann. Phys. (N.Y.)* **253**, 1 (1997); *Nucl. Phys.* **B724**, 295 (2005).
- [7] A. M. Polyakov, *Nucl. Phys.* **B797**, 199 (2008).
- [8] G. Perez-Nadal, A. Roura, and E. Verdaguier, *Classical Quantum Gravity* **25**, 154013 (2008).
- [9] M. Faizal and A. Higuchi, *Phys. Rev. D* **78**, 067502 (2008).
- [10] E. T. Akhmedov and P. V. Buividovich, *Phys. Rev. D* **78**, 104005 (2008).
- [11] A. Higuchi, *Classical Quantum Gravity* **26**, 072001 (2009).
- [12] A. Higuchi and Y. C. Lee, *Classical Quantum Gravity* **26**, 135019 (2009).
- [13] E. T. Akhmedov, *Mod. Phys. Lett. A* **25**, 2815 (2010).
- [14] A. M. Polyakov, *Nucl. Phys.* **B834**, 316 (2010).
- [15] C. P. Burgess, R. Holman, L. Leblond, and S. Shandera, *J. Cosmol. Astropart. Phys.* **10** (2010) 017.
- [16] S. B. Giddings and M. S. Sloth, *J. Cosmol. Astropart. Phys.* **01** (2011) 023.
- [17] D. Krotov and A. M. Polyakov, [arXiv:1012.2107](https://arxiv.org/abs/1012.2107).
- [18] P. Hájíček, “A New Generating Functional for Expectation Values of Field Operators,” 1978 (unpublished).
- [19] B. S. Kay, *Commun. Math. Phys.* **71**, 29 (1980).
- [20] R. D. Jordan, *Phys. Rev. D* **33**, 444 (1986).
- [21] E. Calzetta and B. L. Hu, *Phys. Rev. D* **35**, 495 (1987).
- [22] J. B. Hartle and S. W. Hawking, *Phys. Rev. D* **13**, 2188 (1976).
- [23] D. Marolf and I. A. Morrison, *Phys. Rev. D* **82**, 105032 (2010).
- [24] D. Marolf and I. A. Morrison, [arXiv:1010.5327](https://arxiv.org/abs/1010.5327).
- [25] S. Hollands, [arXiv:1010.5367](https://arxiv.org/abs/1010.5367).
- [26] A. Rajaraman, *Phys. Rev. D* **82**, 123522 (2010).
- [27] D. Schlingemann, [arXiv:hep-th/9912235](https://arxiv.org/abs/hep-th/9912235).
- [28] J. Glimm and A. Jaffe, *Quantum Physics* (Springer-Verlag, New York, 1987), Secs. 6.1 and 10.4.
- [29] G. W. Gibbons and M. J. Perry, *Proc. R. Soc. A* **358**, 467 (1978).
- [30] G. W. Gibbons and S. W. Hawking, *Phys. Rev. D* **15**, 2738 (1977).
- [31] J. S. Schwinger, *J. Math. Phys. (N.Y.)* **2**, 407 (1961).
- [32] L. V. Keldysh, *Zh. Eksp. Teor. Fiz.* **47**, 1515 (1964).
- [33] N. P. Landsman and C. G. van Weert, *Phys. Rep.* **145**, 141 (1987).
- [34] T. S. Bunch and P. C. W. Davies, *Proc. R. Soc. A* **360**, 117 (1978).
- [35] B. Allen and T. Jacobson, *Commun. Math. Phys.* **103**, 669 (1986).
- [36] N. Ya. Vilenken and A. U. Klimyk, *Representations of Lie Groups and Special Functions* Simplest Lie Groups, Special Functions and Integral Transforms (Mathematics and its Applications) (Kluwer Academic Publishers, Dordrecht, 1991), Vol. 1; *Representations of Lie Groups and Special Functions* Vol. 2 (Kluwer Academic Publishers, Dordrecht, 1992); *Representations of Lie Groups and Special Functions* Vol. 3 (Kluwer Academic Publishers, Dordrecht, 1993).
- [37] R. Camporesi and A. Higuchi, *Phys. Rev. D* **45**, 3591 (1992).
- [38] R. F. Streater and A. S. Wightman, *PCT, Spin and Statistics, and All That*, Advanced Book Classics, (Addison-Wesley, Redwood City, CA, 1989), p. 207 .
- [39] J. C. Collins, *Renormalization. An Introduction To Renormalization, The Renormalization Group, And The Operator Product Expansion* (Cambridge University Press, Cambridge UK, 1984), p. 380.
- [40] M. Srednicki, *Quantum Field Theory* (Cambridge University Press, Cambridge, UK, 2007), p. 641.
- [41] J. M. Maldacena, *J. High Energy Phys.* **05** (2003) 013.
- [42] A. Higuchi and S. S. Kouris, *Classical Quantum Gravity* **18**, 2933 (2001).
- [43] A. Higuchi and S. S. Kouris, *Classical Quantum Gravity* **18**, 4317 (2001).
- [44] S. P. Miao, N. C. Tsamis, and R. P. Woodard, *J. Math. Phys. (N.Y.)* **50**, 122502 (2009).
- [45] S. P. Miao, N. C. Tsamis, and R. P. Woodard, *J. Math. Phys. (N.Y.)* **51**, 072503 (2010).
- [46] A. Dasgupta and R. Loll, *Nucl. Phys.* **B606**, 357 (2001).