

Ghosts of critical gravity

Massimo Porrati and Matthew M. Roberts

*Center for Cosmology and Particle Physics Department of Physics New York University 4 Washington Place,
New York, New York 10003, USA*

(Received 20 April 2011; published 8 July 2011)

Recently proposed “critical” higher-derivative gravities in AdS_D $D > 3$ are expected to carry logarithmic representation of the anti-de Sitter isometry group. In this article, we quantize linear fluctuations of these critical gravities, which are known to be either identical with linear fluctuations of Einstein’s gravity or satisfy logarithmic boundary conditions at spacial infinity. We identify the scalar product uniquely defined by the symplectic structure implied by the classical action, and show that it does not possess null vectors. Instead, we show that the scalar product between any two Einstein modes vanishes, while the scalar product of an Einstein mode with a logarithmic mode is generically nonzero. This is the basic property of logarithmic representation that makes them neither unitary nor unitarizable.

DOI: 10.1103/PhysRevD.84.024013

PACS numbers: 04.50.Kd

I. INTRODUCTION

It has been known for many years that power-counting renormalizable theories of gravity can be obtained by adding to the Einstein-Hilbert action appropriate terms, quadratic in the Ricci and Weyl tensors. In the absence of a cosmological constant, these theories admit Minkowski space as background, but they are also perturbatively non-unitary [1]. Since the entire point of having a power-counting renormalizable theory of gravity is to make perturbative calculations possible, these theories have been justly abandoned long since. Recently, quadratic-curvature actions *with* cosmological constant were re-examined in four [2] and D [3] dimensions. In either case, it was found that there exist a choice of parameters for which these theories possess one anti-de Sitter (AdS) background on which neither massive fields, nor massless scalars or vectors propagate. Moreover, on the AdS background, the standard graviton, i.e. the massless tensor mode of Einstein-Hilbert gravity, also propagates and has vanishing energy (the energy of course depends on the action, not just on the form of the mode) [2,3].

Besides those that satisfy the homogenous Einstein equations on AdS_D , other tensor modes propagate in the “critical” theory [4–6]. Their asymptotic behavior at spacelike infinity differs from standard Einstein-Hilbert modes by terms logarithmic in the AdS radial coordinate. A complete set of propagating modes for critical D -dimensional gravity was presented in [6].

In this paper, we show that there exists an unambiguous manner to define the energy and the norm of all modes of log gravity. With that definition, the scalar product of two modes that solve the homogenous Einstein equation vanishes.

Next we address the main result: the scalar product of a homogenous mode with some of the logarithmic modes is nonzero. In other words, homogenous modes *are not* null vectors and cannot be factored out to yield a (positive-norm) Hilbert space, except if we restrict the physical

space to homogenous modes only, and then factor them out. This procedure leaves a profoundly uninteresting theory made only by the vacuum state. This picture should be compared to the case of $D = 3$, where CFTs possess two copies of the Virasoro algebra. There, in topologically massive gravity (TMG) at the critical point [7], restriction to homogenous modes—which can be promoted to a bona fide nonperturbative constraint on the Hilbert space—selects the vacuum of one such algebra, but allows for nontrivial states of the other [7,8].

This result shows that the critical (a.k.a. log) theory is neither unitary nor does it contain a unitary subspace other than the vacuum.¹ Though the lack of unitarity proven here is bad news for log gravity to give a viable quantum theory of gravity in AdS_D , it is consistent with (and indeed required by) it being dual to a logarithmic conformal field theory in $D - 1$ dimensions. Such a theory could be of interest in statistical mechanics; its basic properties are summarized in the next section.

II. KINEMATICS OF LOGARITHMIC CFTS

The isometry group of AdS_D is $SO(2, D - 1)$, which is also the conformal group in $D - 1$ space-time dimensions. So, the Hilbert space² of a consistent quantum gravity in AdS_D decomposes into a direct sum of representations of $SO(2, D - 1)$.

If we do not demand that the representation be unitary, then the Hilbert space, H , can contain a reducible but *indecomposable* representation. Let us consider in detail the case of AdS_4 . Its isometry group, $SO(2, 3)$, admits a Cartan decomposition into four positive roots E^{α_a} , $a = 1$,

¹The ghost pole in one-particle exchange amplitudes between physical sources may cancel [9]; this is not enough to rescue unitarity as we shall discuss in the last section.

²By Hilbert space we mean here a vector space H , endowed with a nondegenerate but not necessarily positive bilinear form $\langle u, v \rangle$ such that $u \in H, v \in H \mapsto \langle u, v \rangle \in \mathbb{C}$.

2, 3, 4, four negative roots $E^{-\alpha_a}$, and two Cartan generators H_1, H_2 .³ In the Cartan basis the $SO(2, 3)$ algebra is

$$\begin{aligned} [H_i, H_j] &= 0, & [H_i, E^{\alpha_a}] &= \alpha_a^i E^{\alpha_a}, \\ [E^{-\alpha_a}, E^{\alpha_a}] &= \frac{2}{|\alpha_a|^2} \alpha_a \cdot H, \end{aligned} \quad (1)$$

with $|\alpha_a|^2 = \sum_{i=1}^2 (\alpha_a^i)^2$.

$SO(2, 3)$ representations with energy H_1 bounded below possess a ground state ψ , annihilated by all $E^{\alpha_a} \psi = 0$. Lowest-weight vectors can also define logarithmic representations if they are not eigenstates of H_1 , but obey instead⁴

$$H_1 \psi = E_0 \psi + \phi, \quad H_1 \phi = E_0 \phi. \quad (2)$$

When the generators H_i, E^α , are self-adjoint with respect to a scalar product, \langle, \rangle , not necessarily positive definite, then the vector ϕ has zero norm:

$$\begin{aligned} 0 &= \langle \psi, (H_1 - E_0)^2 \psi \rangle = \langle (H_1 - E_0) \psi, (H_1 - E_0) \psi \rangle \\ &= \langle \phi, \phi \rangle. \end{aligned} \quad (3)$$

The standard procedure to obtain a nondegenerate scalar product is to identify vectors modulo null vectors. So, if the vector ϕ is null, i.e. if $\langle \chi, \phi \rangle = 0$ for all χ in the representation V , Eqs. (2) actually define a standard lowest-weight representation on the quotient space. So, to obtain a truly new representation, the scalar product of ϕ with some vector $\chi \in V$ (such that $\langle \chi, \chi \rangle \neq 0$) must be nonvanishing.

Of course, the latter property is incompatible with unitarity, i.e. with the scalar product being positive definite. For if $\langle \phi, \chi \rangle = A \neq 0$, the norm of $z\phi + \chi$ is $\langle \chi, \chi \rangle + 2\text{Re}Az$, which can have either sign when z ranges over the complex plane.

Two dimensional logarithmic conformal field theories too are characterized by having zero norm non-null vectors [10] (for a review see [11]). Topologically massive gravity at the critical point, which is conjectured to be dual to a logarithmic CFT, indeed contains such vectors [12–14].

Four-dimensional critical gravity is in many ways the higher-dimensional analog of TMG at the critical point; so, the next question to ask is does critical gravity also contain logarithmic representations of $SO(2, 3)$?

III. THE EXAMPLE OF $D = 4$ CRITICAL GRAVITY

Reference [6] gives a complete set of modes for critical gravity. With some obvious changes of notations and simplifications, the four-dimensional action of [6] is

$$\begin{aligned} S &= \frac{1}{16\pi G} \int d^4x \sqrt{-g} \left[R - 2\Lambda - \frac{1}{2} f^{\mu\nu} G_{\mu\nu} \right. \\ &\quad \left. + \frac{m^2}{8} (f^{\mu\nu} f_{\mu\nu} - f^2) \right], \end{aligned} \quad (4)$$

with $G_{\mu\nu}$ the Einstein tensor and $f_{\mu\nu}$ an auxiliary symmetric tensor field. Elimination of $f_{\mu\nu}$ through its algebraic equations of motion gives an action quadratic in curvatures. Critical gravity is obtained when the cosmological constant is

$$\Lambda = -3m^2. \quad (5)$$

In four-dimensions, Λ is the usual cosmological constant and action (4) admits an anti-de Sitter background $\bar{g}_{\mu\nu}$ with $R^{\mu\nu}{}_{\rho\sigma} = (\Lambda/3)(\delta_\rho^\mu \delta_\sigma^\nu - \mu \leftrightarrow \nu)$.

Expanding $g_{\mu\nu}$ and $f_{\mu\nu}$ around the AdS background as

$$\begin{aligned} g_{\mu\nu} &= \bar{g}_{\mu\nu} + h_{\mu\nu}, \\ f_{\mu\nu} &= -2(\bar{g}_{\mu\nu} + h_{\mu\nu}) + \frac{2}{3m^2} k_{\mu\nu}, \end{aligned} \quad (6)$$

Action (4) reduces to a constant term plus the quadratic action [6]

$$6m^2 S_2 = \int d^4x \sqrt{-\bar{g}} \left[2h^{\mu\nu} \mathcal{G}_{\mu\nu}(k) - \frac{1}{3} (k^{\mu\nu} k_{\mu\nu} - k^2) \right]. \quad (7)$$

The linearized Einstein operator $\mathcal{G}_{\mu\nu}$ reduces to $-(1/2) \times (\square + 2m^2)$ on transverse-traceless modes. The equations of motion following from action (7) are

$$\mathcal{G}_{\mu\nu}(h) = \frac{1}{3}(k_{\mu\nu} - \bar{g}_{\mu\nu} k), \quad \mathcal{G}_{\mu\nu}(k) = 0. \quad (8)$$

Thanks to the Bianchi identity, $k_{\mu\nu}$ is transverse and traceless. In the gauge $D^\mu h_{\mu\nu} - D_\nu h = 0$ Eqs. (8) become

$$-\frac{1}{2}(\square + 2m^2)h_{\mu\nu} = \frac{1}{3}k_{\mu\nu}, \quad -\frac{1}{2}(\square + 2m^2)k_{\mu\nu} = 0. \quad (9)$$

In global coordinates (colatitude θ , longitude ϕ , radius ρ , and time t) the AdS₄ metric is

$$\begin{aligned} m^2 ds^2 &= -\cosh^2(\rho) dt^2 + d\rho^2 + \sinh^2(\rho) \\ &\quad \times [d\theta^2 + \sin^2(\theta) d\phi^2]. \end{aligned} \quad (10)$$

The Cartan generators of $SO(2, 3)$ are $H_1 = i\partial_t$, $H_2 = -i\partial_\phi$ [6]. The explicit expressions for the E^α generators are also given in [6]. Imposing $E^{\alpha_a} \psi = 0$ one finds two particularly interesting classes of solutions to (9).

One, $\psi_{\mu\nu}^E$, solves the homogeneous linearized Einstein equation $\mathcal{G}_{\mu\nu}(\psi^E) = 0$. In global coordinates it has the form $\psi_{\mu\nu}^E = e^{-iE_0 t + 2i\phi} F(\rho, \theta)_{\mu\nu}$. For large ρ one finds

$$\begin{aligned} F(\rho, \theta)_{\rho\rho} &\sim e^{-(E_0+2)\rho}, \\ F(\rho, \theta)_{\rho*} &\sim e^{-E_0\rho}, \\ F(\rho, \theta)_{**} &\sim e^{-(E_0-2)\rho}. \end{aligned} \quad (11)$$

³The basis used in Ref. [6] is $\alpha_1 = (-1, 1)$, $\alpha_2 = (0, 1)$, $\alpha_3 = (-1, -1)$, $\alpha_4 = (-1, 0)$.

⁴One could consider in principle also polylogarithmic representations defined by $(H_1 - E_0)\psi^k = \psi^{k+1}$, $k = 0, \dots, n$.

Here $*$ denotes coordinates other than ρ , and we did not spell out the θ dependence in F . Normalizability of ψ^E for $\rho \rightarrow \infty$ gives $E_0 = 3$ [6].

The other one is

$$f(t, \rho)\psi_{\mu\nu}^E, \quad f(t, \rho) = it + \log \sinh \rho. \quad (12)$$

It obeys Eq. (8) with $k_{\mu\nu} = -(9m^2/2)\psi_{\mu\nu}^E$.⁵

After this brief review of the result of [6] we come to the definition of the scalar product and energy for linearized critical gravity.

IV. THE INNER PRODUCT OF QUADRATIC THEORIES

Consider a general quadratic action

$$S = \int dt \frac{1}{2} (-\dot{q}^T L \dot{q} + \dot{q}^T Q q + q^T K q), \quad (13)$$

where L and K are symmetric matrices while Q is anti-symmetric and commuting with L : $[L, Q] = 0$. They are defined in terms of a matrix Ω obeying

$$L\Omega - \Omega^T L = 2iQ, \quad \Omega^T L \Omega = K. \quad (14)$$

Reality of K follows from $[L, Q] = 0$; together with the first of the equations above, it implies that $L\Omega = S + iQ$, with S a real symmetric matrix. If Ω satisfies Eqs. (14) so does $-\Omega^*$. When some of the eigenvalues of the matrix Ω coincide, Ω may be nondiagonalizable but it can always be put in the Jordan form. We choose Ω to have ‘‘positive frequency’’ by demanding that its eigenvalues are positive definite.

Next we split the vector q into $q = 2^{-1/2}(A + A^*)$ with $i\dot{A} = \Omega A$ ($i\dot{A}^* = -\Omega^* A^*$). Equations (14) guarantee that A solves the equations of motion $L\ddot{A} - 2Q\dot{A} + KA$. The canonical momentum p conjugate to q is $p = L\dot{q} - Qq$. The canonical momenta conjugate to A, A^* are

$$\begin{aligned} P &= L\dot{A}^* - QA^* = iL\Omega^* A^* - QA^*, \\ P^* &= L\dot{A} - QA = -iL\Omega A - QA. \end{aligned} \quad (15)$$

By using Eqs. (14) and (15) it is straightforward to find that the conserved energy is

$$H = \frac{1}{2}(\dot{q}^T L \dot{q} + q^T K q) = \frac{1}{2}(P^T \dot{A} + P^{*T} \dot{A}^*) \quad (16)$$

In canonical quantization, A^* is replaced by the Hermitian conjugate A^\dagger and the nonzero canonical commutation relations are $[A^I, P_J] = i\delta^I_J$, $[A^{\dagger I}, P_J^\dagger] = i\delta^I_J$. The standard Fock vacuum obeys $A|0\rangle = 0$, so a classical positive-frequency solution of the equations of motion, $i\dot{\Phi}(t) = \Omega\Phi(t)$, defines a one-particle state $|\Phi\rangle = -i\Phi^I P_I|0\rangle$, which obeys $A(t)|\Phi\rangle = \Phi(t)|0\rangle$. The scalar product of two states $|\Phi\rangle, |\Psi\rangle$ is then

⁵Our inhomogeneous mode equals that in [6] plus a homogeneous mode.

$$\langle \Psi | \Phi \rangle = i(\Psi^{*T} L \dot{\Phi} - \Psi^{*T} Q \Phi). \quad (17)$$

An example directly related to action (7) is

$$\begin{aligned} L &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & \Omega &= \begin{pmatrix} \omega & 0 \\ 1/2\omega & \omega \end{pmatrix}, \\ A(t) &= \begin{pmatrix} 2\omega i \alpha e^{-i\omega t} \\ (\alpha t + \beta) e^{-i\omega t} \end{pmatrix} \end{aligned} \quad (18)$$

Notice that, though $A(t)$ contains terms linear in t , the scalar product is time-independent. Explicitly, on two solutions defined by constants $\alpha, \beta, \alpha', \beta'$, the scalar product formula (17) reduces to $\langle \Psi | \Phi \rangle = 2\omega \alpha'^* \alpha + 2\omega^2 i(\beta'^* \alpha - \alpha'^* \beta)$. States with $\alpha = \alpha' = 0$ have vanishing norm, but these states are not null: the norm of state $\alpha = 0, \beta \neq 0$ with an $\alpha' \neq 0$ state does not vanish.

Action (7) has the form (13). To see that, we can use the fact that $k_{\mu\nu}$ is transverse-traceless and integrate by part in time⁶

$$\begin{aligned} 6m^2 S_2 &= \int d^4 x \sqrt{-\bar{g}} \left[\bar{g}^{00} D_0 h^{\mu\nu} D_0 k_{\mu\nu} \right. \\ &\quad \left. - h^{\mu\nu} \left(\sum_{i=1}^3 D_i D^i + 2m^2 \right) k_{\mu\nu} - \frac{1}{3} (k^{\mu\nu} k_{\mu\nu} - k^2) \right]. \end{aligned} \quad (19)$$

Now, a straightforward application of formula (17) gives the scalar product of any two positive-frequency modes of log gravity as

$$\begin{aligned} \langle \psi | \phi \rangle &= \frac{3i}{4} \int d^3 x \sqrt{-\bar{g}} \bar{g}^{00} [(\square + 2m^2) \psi_{\mu\nu}^* D_0 \phi^{\mu\nu} \\ &\quad + \psi_{\mu\nu}^* D_0 (\square + 2m^2) \phi^{\mu\nu}]. \end{aligned} \quad (20)$$

Evidently, the scalar product between two solutions of the homogeneous Einstein equations, $\phi = \phi^E, \psi = \psi^E$ vanishes. Also, the potentially dangerous term $\propto t$ in the scalar product of two ‘‘log’’ modes $\phi^s = f(t, \rho)\phi^E, \psi = f(t, \rho)\psi^E$ (see definition in Eq. (12)) vanishes.

Crucially, the homogeneous modes ψ^E are *not* null vectors, since their scalar product with the log mode constructed from $\psi^E, \phi^s = f(t, \rho)\psi^E$ is nonzero:

$$\langle \psi^E | \phi^s \rangle = -\frac{9im^2}{4} \int d^3 x \sqrt{-\bar{g}} \bar{g}^{00} \psi_{\mu\nu}^{E*} D_0 \psi^{E\mu\nu} \neq 0. \quad (21)$$

Nonvanishing of Eq. (21) can be proven by a simple direct calculation or by noticing that (21) is proportional to the

⁶Initial time and final time configurations are held fixed when varying the action, so we can always add a total *time* derivative to the action without changing the equations of motion. Adding a total divergence of space coordinate, instead, changes the boundary conditions at the AdS boundary, so in general it changes the equations of motion by modifying the boundary behavior of the fields.

norm of the transverse-traceless mode $\psi_{\mu\nu}^E$ in standard Einstein gravity.⁷

The Einstein modes do have zero scalar product with special logarithmic modes: the spin-1 ‘‘Proca’’ modes [6], which have the form $\psi_{\mu\nu}^s = f\psi^E$, $\psi^E = D_{(\mu}A_{\nu)}$. Transversality and tracelessness of $\psi_{\mu\nu}^E$ hold when A_μ obeys the massive spin-1 Proca equation

$$D^\nu F_{\nu\mu} = 6m^2 A_\mu. \quad (22)$$

Vanishing of $\langle \psi^E | f D_{(\mu} A_{\nu)}$ follows immediately from the fact that this quantity is proportional to the Einstein gravity scalar product $\int d^3x \sqrt{-\bar{g}} \bar{g}^{00} \psi_{\mu\nu}^{E*} D_0 D^\mu A^\nu$, which vanishes because in Einstein gravity transverse modes are orthogonal to pure gauge modes.

In three-dimensions, the Proca modes are the only inhomogeneous solutions of Eq. (8), hence the Einstein modes are true null vectors that can be modded out to yield a positive-metric Hilbert space. Actually, the normalizable Proca modes decay so rapidly at infinity that they too are null in the norm induced by the NMG action. So, the modding out by these null vectors yields a trivial theory in three-dimensions too. Of course, one can also define the norm of the Proca fields using the Proca action; this is what makes new massive gravity [15] at the critical point potentially nontrivial and consistent, at least at linear order. In $D > 3$, the Proca modes [6] and the truly transverse-traceless spin-2 logarithmic modes mix under the action of $SO(D-1) \subset SO(2, D-1)$; therefore, one cannot consistently keep the Proca modes, which are transverse to the Einstein modes, without also keeping the spin-2 modes, which are not.

V. LINEARIZED ENERGY

A reasonable way to define the energy of a solution which is asymptotically AdS is to construct the linearized stress tensor of a metric perturbation, and integrate the tt component over a spacial slice, which by the Gauss’ law constraint reduces to a boundary term. Including the non-linear terms schematically in the right-hand side of the equations of motion, we find the equations

$$\mathcal{G}(k_{\mu\nu}) = \Theta_{\mu\nu}^{(NL)}, \quad \mathcal{G}(h_{\mu\nu}) - \frac{1}{3}(k_{\mu\nu} - \bar{g}_{\mu\nu}k) = \Xi_{\mu\nu}^{(NL)}. \quad (23)$$

Note that thanks to the Bianchi identity $\Theta_{\mu\nu}^{(NL)}$ is automatically covariantly conserved, while $\Xi_{\mu\nu}^{(NL)}$ is not. We therefore can use $\Theta = \mathcal{G}(k)$ to construct conserved charges following the method of [16] and replacing the metric perturbation $h_{\mu\nu}$ with $k_{\mu\nu} = 3\mathcal{G}_{\mu\nu}(h) - \bar{g}_{\mu\nu}\mathcal{G}_\sigma^\sigma(h)$, which gives that the conserved charge for a killing vector ξ is

$$E(\bar{\xi}) = \frac{1}{8\pi G} \oint dS_i \sqrt{-\bar{g}} [D_\beta K^{0i\nu\beta} - K^{0j\nu i} D_j] \bar{\xi}_\nu, \quad (24)$$

⁷See e.g. Ref. [13] for the analogous calculation in three-dimensional chiral gravity.

where

$$K^{\mu\alpha\nu\beta} = \frac{1}{2}[\bar{g}^{\mu\beta}H^{\nu\alpha} + \bar{g}^{\nu\alpha}H^{\mu\beta} - \bar{g}^{\mu\nu}H^{\alpha\beta} - \bar{g}^{\alpha\beta}H^{\mu\nu}], \\ H^{\mu\nu} = k^{\mu\nu} - \frac{1}{2}\bar{g}^{\mu\nu}k. \quad (25)$$

It is clear from this formulation that any solution that is purely an Einstein mode will have $k_{\mu\nu} = 0$ and therefore will have all conserved charges vanish, in agreement with other calculations of the energy in critical gravity. To find a nonzero energy we must turn on a nonzero $k_{\mu\nu}$ mode. At large radius the only static, spherically symmetric solution to (8) is simple in terms of $k_{\mu\nu} = \tilde{k}_{\mu\nu} + D_{(\mu}\xi_{\nu)}$, the Coulomb tail of a mass in AdS ($\tilde{k}_{\mu\nu}$), along with a vector mode⁸ ($D_{(\mu}\xi_{\nu)}$). The Coulomb tail of a mass in AdS₄ behaves as $\tilde{k}_{tt} = -\bar{g}_{tt}\tilde{k}_{\rho\rho} = \tilde{M}/\sinh\rho$. We must include a vector mode to ensure the consistency of the equation for h , because $D^\mu \mathcal{G}_{\mu\nu} = 0$ and we must therefore require that $D^\mu(k_{\mu\nu} - \bar{g}_{\mu\nu}k) = 0$ so that the equation of motion is covariantly conserved. This is ensured with $\xi^r = \tilde{M}/6\sinh^2\rho \cosh\rho$, which gives

$$k_{tt} = -k_{\rho\rho}\sinh^2\rho\cosh^2\rho = 2k_{\theta\theta} = 2k_{\phi\phi}/\sin^2\theta \\ = M/\sinh\rho, \quad (26)$$

where $M = 2\tilde{M}/3$. It is easy to check that the mass as defined in (24) is simply $M/2G$. We also note that this nonzero $k_{\mu\nu}$ sources a logarithmic falloff in h ,

$$h_{tt} = \frac{M \log\cosh\rho}{3 \sinh\rho} + \frac{M \cosh^2\rho}{3} \\ \times \left(\pi - 4 \arctan[\tanh(\rho/2)] - \frac{1 + \tanh^2\rho}{\sinh\rho} \right), \\ h_{\rho\rho} = \frac{M \log\cosh\rho}{3 \cosh^2\rho \sinh\rho}. \quad (27)$$

We can also turn on a homogeneous mode in h , that is one satisfying $\mathcal{G}(h) = 0$, but this will clearly not contribute to the energy.

VI. MISCELLANEOUS REMARKS

The one-particle Hilbert space obtained by quantizing critical gravity on its AdS background splits into a sum of Einstein modes and log modes: $H = H^s \oplus H^E$. The norm N on such space vanishes on H^E but it is off diagonal. Schematically, for $|\psi\rangle = |\psi^s\rangle \oplus |\psi^E\rangle$, $|\phi\rangle = |\phi^s\rangle \oplus |\phi^E\rangle$, one has

$$\langle \psi | N | \phi \rangle = (\langle \psi^s |, \langle \psi^E |) \begin{pmatrix} 1 & \alpha \\ \alpha & 0 \end{pmatrix} \begin{pmatrix} |\phi^s\rangle \\ |\phi^E\rangle \end{pmatrix}, \quad \alpha \neq 0. \quad (28)$$

A computation of the energy using formula (16) shows that, on a mode of frequency ω , its *unnormalized*

⁸This is of course not an honest linear diffeomorphism, as by Eq. (8) and the diff invariance of \mathcal{G} we see that k does not transform.

expectation value is proportional to the norm N : $\langle \psi | H | \phi \rangle = \omega \langle \psi | N | \phi \rangle$. The Fock space contains only states with positive frequency, so that whenever the norm is nonzero, the normalized energy, $\langle \psi | H | \phi \rangle / \langle \psi | N | \phi \rangle$, is positive and equal to the frequency ω , as expected. When the norm vanishes, by consistency one must define the energy so that it also vanishes. Restricting the multiparticle Hilbert space to the Einstein modes is thus equivalent to selecting the zero-energy subsector of critical gravity, in perfect analogy with three-dimensional chiral gravity [8]. To obtain a properly defined Hilbert space, the restriction to Einstein modes must be followed by modding out by null states.

Unlike three-dimensional chiral gravity, here the end result of this procedure leaves only one state in the theory, the Fock vacuum.

The results described so far were obtained by studying linearized critical gravity. Yet, the general structure we found can be promoted to a full nonlinear analysis. In particular, the vanishing of energy in solutions which asymptotically become Einstein modes has been shown to hold for black holes too in [2,3]. A general proof that all asymptotically-Einstein solutions have zero energy should be possible using the general definition of energy in quadratic-curvature gravities given in [17]. Conversely, in the previous section we just proved that nonvanishing energy requires the asymptotic behavior of log modes.

It was conjectured in [18] and proven in [19] that TMG in three-dimensions is perturbatively stable around certain warped AdS_3 vacua, even for noncritical values of the Chern-Simons coupling constant. Stability is achieved by restricting the asymptotic boundary conditions in such a way that only boundary gravitons of one chirality propagate. This is in perfect analogy with the behavior of TMG at the critical point on nonwarped AdS_3 . In this paper, we argued that in $D > 3$, restriction to the Einstein asymptotics can give us only a trivial Hilbert space made only of the vacuum. So, if an analog to the results of [18,19] could be found for $D > 3$ higher-derivative gravity, it would probably require asymptotics on fields that leave no physical state beyond the vacuum.

It may happen that negative-norm ghost poles cancel against some positive-norm poles in one-particle exchange

diagrams between physical sources [9]. This is not enough to ensure unitarity, since in the full nonlinear theory defined by Eq. (4) ghosts can obviously also be pair produced. Moreover, even at the level of one-particle diagrams, one must still mod out by null states, arriving again at an empty theory.

Finally, we must point out explicitly that the Hilbert space of Einstein modes is the only physically meaningful subspace without negative-norm states (unfortunately, it also has no positive-norm states). Obviously, by diagonalizing the metric N in Eq. (28) one can obtain a positive-norm subspace. Unfortunately, that subspace is not closed under $SO(2,3)$ transformations, since logarithmic representations are indecomposable. It is amusing to check explicitly this property by verifying that a positive-norm subspace is not even closed under time evolution. In fact, its vectors must be linear combinations of log modes and gravity modes. Consider, in particular, a mode of frequency ω : $|\psi\rangle = |\psi^s\rangle + |\phi^E\rangle$, $\psi^s = f(t, \rho)\psi^E$. Under a time translation $t \rightarrow t + \tau$, it transforms into $\psi(t) \rightarrow \psi(t + \tau) = \exp(-i\omega\tau)(\psi + i\tau\psi^E)$. Equation (21) says that $\langle \psi^s | \psi^E \rangle = c \langle \psi^E | \psi^E \rangle_E$, where $\langle \cdot \rangle_E$ is the scalar product in Einstein gravity and c is a positive constant; so the scalar product $\langle \psi(t + \tau) | N | \psi(t + \tau) \rangle$ is independent of τ . Now consider the linear combination $\psi(t) + C\psi(t + \tau)$. Its norm is

$$\begin{aligned} & \langle \psi(t) + C\psi(t + \tau) | N | \psi(t) + C\psi(t + \tau) \rangle \\ &= (CC^* + 1)\langle \psi | \psi \rangle + 2\langle \psi | \psi \rangle \text{Re}(Ce^{-i\omega\tau}) \\ & \quad + -2c\langle \psi^E | \psi^E \rangle_E \text{Im}(\tau Ce^{-i\omega\tau}). \end{aligned} \quad (29)$$

The right-hand side in this equation becomes negative for $C \neq 0$ and τ large.

ACKNOWLEDGMENTS

It is a pleasure to thank O. Hohm, C. Pope, and E. Bergshoeff for useful correspondence and discussions. M.P. is supported in part by the NSF under Grant No. PHY-0758032, and by ERC Advanced Investigator Grant No. 226455 *Supersymmetry, Quantum Gravity and Gauge Fields (Superfields)*.

-
- [1] K. S. Stelle, *Phys. Rev. D* **16**, 953 (1977); *Gen. Relativ. Gravit.* **9**, 353 (1978).
 - [2] H. Lu and C. N. Pope, *Phys. Rev. Lett.* **106**, 181302 (2011).
 - [3] S. Deser, H. Liu, H. Lu, C. N. Pope, T. C. Sisman, and B. Tekin, *Phys. Rev. D* **83**, 061502 (2011).
 - [4] M. Alishahiha and R. Fareghbal, *Phys. Rev. D* **83**, 084052 (2011).
 - [5] I. Gullu, M. Gurses, T. C. Sisman, and B. Tekin, *Phys. Rev. D* **83**, 084015 (2011).
 - [6] E. A. Bergshoeff, O. Hohm, J. Rosseel, and P. K. Townsend, *Phys. Rev. D* **83**, 104038 (2011).
 - [7] W. Li, W. Song, and A. Strominger, *J. High Energy Phys.* **04** (2008) 082.
 - [8] A. Maloney, W. Song, and A. Strominger, *Phys. Rev. D* **81**, 064007 (2010).
 - [9] T. C. Sisman, I. Gullu, and B. Tekin, arXiv:1103.2307.
 - [10] V. Gurarie, *Nucl. Phys.* **B410**, 535 (1993).
 - [11] M. Flohr, *Int. J. Mod. Phys. A* **18**, 4497 (2003).

- [12] K. Skenderis, M. Taylor, and B.C. van Rees, *J. High Energy Phys.* **09** (2009) 045; [arXiv:0909.5617](#).
- [13] D. Grumiller and I. Sachs, *J. High Energy Phys.* **03** (2010) 012.
- [14] D. Grumiller and N. Johansson, *J. Phys. Conf. Ser.* **222**, 012047 (2010).
- [15] E.A. Bergshoeff, O. Hohm, and P.K. Townsend, *Phys. Rev. Lett.* **102**, 201301 (2009).
- [16] L.F. Abbott and S. Deser, *Nucl. Phys.* **B195**, 76 (1982).
- [17] S. Deser and B. Tekin, *Phys. Rev. Lett.* **89**, 101101 (2002); *Phys. Rev. D* **67**, 084009 (2003).
- [18] D. Anninos, W. Li, M. Padi, W. Song, and A. Strominger, *J. High Energy Phys.* **03** (2009) 130.
- [19] D. Anninos, M. Esole, and M. Guica, *J. High Energy Phys.* **10** (2009) 083.