

Supersymmetric Higgs sector and $B - \bar{B}$ mixing for large $\tan\beta$

Martin Gorbahn,¹ Sebastian Jäger,² Ulrich Nierste,³ and Stéphanie Trine³

¹*Technische Universität München, Institute for Advanced Study, Lichtenbergstraße 2 a, 85748 Garching, Germany,
& Technische Universität München, Excellence Cluster “Universe,” Boltzmannstraße 2, D-85748 Garching, Germany*

²*University of Sussex, Department of Physics and Astronomy, Falmer, Brighton BN1 9QH, United Kingdom*

³*Institut für Theoretische Teilchenphysik, Universität Karlsruhe, Karlsruhe Institute of Technology,
Engesserstraße 7, D-76128 Karlsruhe, Germany*

(Received 10 May 2011; published 18 August 2011)

We match the Higgs sector of the most general flavor-breaking and CP -violating minimal supersymmetric standard model (MSSM) onto a generic two-Higgs-doublet model, paying special attention to the definition of $\tan\beta$ in the effective theory. In particular, no $\tan\beta$ -enhanced loop corrections appear in the relation to $\tan\beta$ defined in the $\overline{\text{DR}}$ scheme in the MSSM. The corrections to the Higgs-mediated flavor-changing amplitudes, which result from this matching, are especially relevant for the B_d and B_s mass differences $\Delta M_{d,s}$ for minimal flavor violation, where the superficially leading contribution vanishes. We give a symmetry argument to explain this cancellation and perform a systematic study of all Higgs-mediated effects, including Higgs loops. The corrections to ΔM_s are at most 7% for $\mu > 0$ and $M_A < 600$ GeV if constraints from other observables are taken into account. For $\mu < 0$ they can be larger, but are always less than about 20%. Contrary to recent claims, we do not find numerically large contributions here, nor do we find any $\tan\beta$ -enhanced contributions from loop corrections to the Higgs potential in $B^+ \rightarrow \tau^+ \nu$ or $B \rightarrow X_s \gamma$. We further update supersymmetric loop corrections to the Yukawa couplings, where we include all possible CP -violating phases and correct errors in the literature. The possible presence of CP -violating phases generated by Higgs exchange diagrams is briefly discussed as well. Finally, we provide improved values for the bag factors P_1^{VLL} , P_2^{LR} , and P_1^{SLL} at the electroweak scale.

DOI: 10.1103/PhysRevD.84.034030

PACS numbers: 13.25.Hw, 12.60.Jv, 14.80.Da

I. INTRODUCTION

Supersymmetry constrains the structure of the Yukawa couplings of the minimal supersymmetric standard-model (MSSM) to those of a special two-Higgs-doublet model (2HDM). In this 2HDM of type II one Higgs doublet, H_u , only couples to up-type fermions, while the other one, H_d , only couples to down-type fermions. As a consequence, there are no dangerous tree-level flavor-changing neutral current (FCNC) couplings of the neutral-Higgs bosons. However, the presence of supersymmetry-breaking terms destroys this pattern at the one-loop level, permitting couplings of both Higgs doublets to all fermions. Thus the resulting Higgs sector is that of a general 2HDM, often called 2HDM of type III. As pointed out first by Hall, Rattazzi and Sarid, the loop-induced Yukawa couplings can compete with the tree-level ones in the limit of a large $\tan\beta = v_u/v_d$, which is the ratio of the vacuum expectation values (vevs) of H_u and H_d [1]: in the relationship between $H_{u,d}$ -couplings and observed masses of the down-type fermions the loop-suppression factor ~ 0.02 is offset by a factor of $\tan\beta$, so that $\mathcal{O}(1)$ corrections to the type II 2HDM are possible for $\tan\beta \sim 50$. In such scenarios, also $\mathcal{O}(1)$ loop-induced FCNC couplings of neutral-Higgs bosons appear [2], which allow the branching fractions of (yet unobserved) leptonic B decays to exceed their standard model values by more than 2 orders of magnitude [3]. This observation has stimulated a large activity in flavor physics and powerful constraints on the MSSM Higgs

sector in scenarios with large $\tan\beta$ have been derived from B factory data [3–6]. These Higgs-induced effects in flavor physics are very transparent in the limit

$$M_{\text{SUSY}} \gg M_A \sim v, \quad (1)$$

where M_{SUSY} denotes the generic mass scale of the superpartners and the masses M_A , M_H , M_h , and $M_{H^\pm}$ of the five physical Higgs bosons are taken to be of the order of the electroweak scale $v \equiv \sqrt{v_u^2 + v_d^2} = 246$ GeV. All low-energy observables can be computed in the type III 2HDM, which emerges as the effective theory in the limit of Eq. (1). The new couplings can be calculated from finite one-loop diagrams with supersymmetric particles and thus become functions of the MSSM parameters, so that the desired constraints on the supersymmetric parameter space can be derived. The effective 2HDM Lagrangian efficiently incorporates all large- $\tan\beta$ effects, equivalent to a perturbative all-order resummation of those radiative corrections which are enhanced by a factor of $\tan\beta$ [7].

$B_q - \bar{B}_q$ mixing (with $q = d$ or s) plays a special role among the FCNC transitions of B mesons. Here the leading new effect stems from effective tree-level diagrams with neutral-Higgs bosons (see Fig. 1). *A priori* the dominant contribution is expected from Yukawa couplings to right-handed b quarks, generating the effective $\Delta B = 2$ operator

$$Q_1^{\text{SLL}} \equiv (\bar{b}_R q_L)(\bar{b}_R q_L). \quad (2)$$




FIG. 1. Leading contributions to $B_q - \bar{B}_q$ mixing from supersymmetric Higgs bosons. The FCNC couplings are induced by supersymmetric loops. The coefficient of $Q_1^{\text{SLL}} = (\bar{b}_R q_L)(\bar{b}_R q_L)$ vanishes, if the tree-level relations between Higgs masses and mixing angles are used.

However, the corresponding coefficient C_1^{SLL} vanishes exactly if one employs the tree-level relations between the Higgs masses and mixing angles [2]. Nevertheless, sizeable effects in $B_s - \bar{B}_s$ mixing are possible even in scenarios with minimal flavor-violation (MFV) [8–16], in which the Cabibbo-Kobayashi-Maskawa (CKM) matrix [17] is the only source of flavor-violation: keeping the strange Yukawa coupling nonzero one finds a nonvanishing contribution to the coefficient of

$$Q_2^{\text{LR}} \equiv (\bar{b}_R q_L)(\bar{b}_L q_R), \quad (3)$$

which depletes the $B_s - \bar{B}_s$ mass difference ΔM_s [5]. The tree-level vanishing of C_1^{SLL} calls for a systematic analysis of all subleading effects. In particular, the contribution that stems from Q_1^{SLL} can *a priori* compete with the contribution of the operator Q_2^{LR} above if the one-loop corrections to the MSSM Higgs potential [18–24] are taken into account. While a lot of work has been devoted to the analysis of the Yukawa sector [2,3,5–7,25], little attention has been given to effects from the Higgs potential. An exception is Ref. [26], which finds large contributions. We revisit these effects in the present paper and perform a systematic matching of the MSSM Higgs sector onto the type III 2HDM. The result is not only relevant for the calculation of C_1^{SLL} , it also clarifies the relationship between the definitions of $\tan\beta$ in the MSSM and the effective 2HDM. This is important to link the constraints from flavor physics to other fields of MSSM phenomenology, in particular, Higgs physics. Our paper is organized as follows. We derive the corrected $B - \bar{B}$ mixing amplitude in Sec. II, including all relevant subleading contributions. The renormalization of $\tan\beta$ and some further technical issues are the subject of Sec. III. In Sec. IV, we apply our new formulas to the phenomenology of $B - \bar{B}$ mixing, analyzing the mass differences ΔM_d and ΔM_s as well as CP violation. Our results are summarized in Sec. V. We list our notation and our technical results in four appendices. Parts of our results were previously presented by one of us at a conference [27].

II. HIGGS-MEDIATED EFFECTS IN $B - \bar{B}$ MIXING

The quantity governing the $B_q - \bar{B}_q$ mass difference is the off-diagonal element of the $B_q - \bar{B}_q$ meson mass matrix: $\Delta M_q = 2|M_{21}^q|$, with

$$M_{21}^q = \frac{\langle \bar{B}_q | \mathcal{H}_{\text{eff}}^{\Delta B=2} | B_q \rangle}{2M_{B_q}}. \quad (4)$$

The $\Delta B = 2$ effective weak Hamiltonian $\mathcal{H}_{\text{eff}}^{\Delta B=2}$ consists in general of eight dimension-six operators:

$$\mathcal{H}_{\text{eff}}^{\Delta B=2} = \frac{G_F^2 M_W^2 \lambda_{qb}^2}{16\pi^2} \sum_{i=1}^8 C_i(\mu_h) Q_i(\mu_h), \quad (5)$$

with $\lambda_{qb} \equiv V_{tq} V_{tb}^*$. The set of operators in Eq. (5) comprises the standard-model operator,

$$Q_1^{\text{VLL}} = (\bar{b}_L \gamma_\mu q_L)(\bar{b}_L \gamma^\mu q_L), \quad (6)$$

the two scalar operators defined in Eqs. (2) and (3), the operator

$$Q_1^{\text{SRR}} \equiv (\bar{b}_L q_R)(\bar{b}_L q_R), \quad (7)$$

and four other operators. The complete list of operators plus the relevant evanescent operators is given in Eq. (C4) and (C5) of Appendix C. We express our results in terms of matrix elements at the high-scale μ_h , which we choose equal to the top mass $\bar{m}_t(\bar{m}_t) = 164$ GeV. In this way, the other four operators do not appear in our formulas. However, some of them are needed to connect $Q_i(\mu_h)$ with $Q_i(\mu_b)$ at the low scale $\mu_b \sim m_b$, at which their matrix elements are computed, because they mix with Q_1^{SLL} , Q_1^{SRR} , or Q_2^{LR} under renormalization. We follow the conventions of Refs. [28,29] for operators and matrix elements. In particular, we parametrize the hadronic matrix elements as

$$\langle \bar{B}_q | Q_i(\mu_h) | B_q \rangle = \frac{2}{3} M_{B_q}^2 f_{B_q}^2 P_i. \quad (8)$$

The P_i 's are obtained [29] by renormalization-group evolution from the conventional bag factors B_i computed at the low scale μ_b . We calculate the P_i 's from up-to-date lattice QCD results in Appendix C, where we fully exploit constraints from heavy-quark relations. This is a new feature of our analysis compared to previous studies of new-physics effects in $B - \bar{B}$ mixing.

A. Effective tree-level Higgs exchange

The Higgs sector of the MSSM contains two $SU(2)$ doublets H_u and H_d ,

$$H_u = \begin{pmatrix} h_u^+ \\ h_u^0 \end{pmatrix}, \quad H_d = \epsilon \begin{pmatrix} h_d^+ \\ h_d^0 \end{pmatrix}^*, \quad \epsilon = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad (9)$$

of hypercharge $+1/2$ and $-1/2$, respectively, with vacuum expectation values (vevs) $\langle h_{u,d}^0 \rangle = v_{u,d}/\sqrt{2}$ of relative size $\tan\beta = v_u/v_d$. Integrating out supersymmetric particles, the Lagrangian of the resulting effective 2HDM is no longer restricted to be of type II, and is constrained only by the electroweak symmetry. Neither will it be renormalizable, with operators of dimension greater than four

encoding effects that decouple at least as v/M_{SUSY} for heavy superpartners. We begin with a short review of some pertinent aspects of the general 2HDM.

Defining

$$\begin{pmatrix} \Phi \\ \Phi' \end{pmatrix} = \begin{pmatrix} \cos\beta & \sin\beta \\ -\sin\beta & \cos\beta \end{pmatrix} \begin{pmatrix} -\epsilon H_d^* \\ H_u \end{pmatrix}, \quad (10)$$

the most general fermion-Higgs interactions up to dimension-four read

$$\begin{aligned} \mathcal{L}_Y = & -\frac{\sqrt{2}}{v} \bar{d}_{Ri} M_{dij} \Phi^\dagger Q_{Lj} - \bar{d}_{Ri} \kappa_{ij} \Phi'^\dagger Q_{Lj} \\ & -\frac{\sqrt{2}}{v} \bar{u}_{Ri} M_{u_{ij}} Q_{Lj} \cdot \Phi - \bar{u}_{Ri} \tilde{\kappa}_{ij} Q_{Lj} \cdot \Phi' + \text{h.c.}, \end{aligned} \quad (11)$$

where we have employed the notation $a \cdot b \equiv a^T \epsilon b$. By construction, the vev of Φ' vanishes, whereas Φ has $\langle \Phi \rangle = (0, v/\sqrt{2})^T$ and contains all three Goldstone bosons. Hence, only Φ can contribute to the fermion masses and only Φ' can have flavor-violating neutral couplings. The flavor basis is defined such that the down-quark mass matrix M_d is diagonal. In this basis, the FCNC Higgs couplings to b -quarks are governed by κ_{bq} or κ_{qb} ($q = d$ or s).

The renormalizable Higgs self-interactions are comprised in the most general gauge-invariant dimension-four two-Higgs-doublet potential [30],

$$\begin{aligned} V = & m_{11}^2 H_d^\dagger H_d + m_{22}^2 H_u^\dagger H_u + \{m_{12}^2 H_u \cdot H_d + \text{H.c.}\} \\ & + \frac{\lambda_1}{2} (H_d^\dagger H_d)^2 + \frac{\lambda_2}{2} (H_u^\dagger H_u)^2 + \lambda_3 (H_u^\dagger H_u) (H_d^\dagger H_d) \\ & + \lambda_4 (H_u^\dagger H_d) (H_d^\dagger H_u) + \left\{ \frac{\lambda_5}{2} (H_u \cdot H_d)^2 \right. \\ & \left. - \lambda_6 (H_d^\dagger H_d) (H_u \cdot H_d) - \lambda_7 (H_u^\dagger H_u) (H_u \cdot H_d) + \text{H.c.} \right\}. \end{aligned} \quad (12)$$

The couplings m_{12}^2 , λ_5 , λ_6 , and λ_7 are in general complex, yet the vevs $v_{u,d}$ can be made real by a $U(1)$ transformation on the Higgs fields. The definitions of m_{ij}^2 and λ_i in Eq. (12) coincide with Ref. [30], except for λ_3 and λ_4 : we associate a different operator with λ_4 to eliminate it from tree-level neutral-Higgs phenomenology and have instead $\lambda_3 = \lambda_3^{[30]} + \lambda_4^{[30]}$ and $\lambda_4 = -\lambda_4^{[30]}$.

Shifting the fields in Eq. (12) by their vevs, which minimize V at tree-level,¹

$$h_{u,d}^0 = \frac{1}{\sqrt{2}} (v_{u,d} + \phi_{u,d} + i\chi_{u,d}), \quad (13)$$

determines the physical Higgs-boson mass matrices and interactions. We write the neutral-Higgs mass matrix in the basis $(\phi_d, \phi_u, \chi_d, \chi_u)$ in terms of 2×2 blocks,

¹“Tree-level” here refers to the 2HDM. We defer a discussion of quantum corrections to v_u and v_d to Sec. III.

$$M_0^2 = \begin{pmatrix} M_R^2 & M_{RI}^2 \\ M_{RI}^{2T} & M_I^2 \end{pmatrix}, \quad (14)$$

with M_R^2 , M_{RI}^2 , and M_I^2 given in Eqs. (23)–(26) below. In the CP -conserving case, $M_{RI}^2 = 0$, and M_R^2 and M_I^2 are diagonalized by rotating the CP -even and CP -odd Higgs fields through angles α and β , respectively:

$$\begin{pmatrix} \phi_d \\ \phi_u \end{pmatrix} = \begin{pmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{pmatrix} \begin{pmatrix} H^0 \\ h^0 \end{pmatrix}, \quad \begin{pmatrix} \chi_d \\ \chi_u \end{pmatrix} = \begin{pmatrix} \cos\beta & -\sin\beta \\ \sin\beta & \cos\beta \end{pmatrix} \begin{pmatrix} G^0 \\ A^0 \end{pmatrix}. \quad (15)$$

The same angle $\beta = \arctan v_u/v_d$ as defined above appears because (and only when) v_u, v_d minimize V . If CP violation is present, four physical mixing angles $\alpha_{1,2,3}$ and β are required to diagonalize M_0^2 . The charged-Higgs mass matrix $M_\pm^2$ is always diagonalized by β ,

$$\begin{pmatrix} h_d^+ \\ h_u^+ \end{pmatrix} = \begin{pmatrix} \cos\beta & -\sin\beta \\ \sin\beta & \cos\beta \end{pmatrix} \begin{pmatrix} G^+ \\ H^+ \end{pmatrix}. \quad (16)$$

The nonstandard $\Delta B = 2$ effective operators Q_2^{LR} , Q_1^{SLL} , and Q_1^{SRR} are generated at tree-level via the exchange of neutral-Higgs bosons (see Fig. 1) with the Wilson coefficients

$$\begin{aligned} C_2^{\text{LR}} &= -\frac{8\pi^2}{G_F^2 M_W^2 \lambda_{qb}^2} (\kappa_{qb}^* \kappa_{bq}) \mathcal{F}^+, \\ C_1^{\text{SLL}} &= -\frac{4\pi^2}{G_F^2 M_W^2 \lambda_{qb}^2} (\kappa_{bq})^2 \mathcal{F}^-, \end{aligned} \quad (17)$$

and C_1^{SRR} obtained from C_1^{SLL} through the replacement $(\kappa_{bq})^2 \mathcal{F}^- \rightarrow (\kappa_{qb}^*)^2 \mathcal{F}^{*-}$. We find that, in the general case, the Higgs propagation factors can be expressed as follows:

$$\mathcal{F}^+ = \frac{\det(M_R^2 + M_I^2 + iM_{RI}^2 - iM_{RI}^{2T})}{m_1^2 m_2^2 m_3^2} \equiv \frac{\det B}{m_1^2 m_2^2 m_3^2}, \quad (18)$$

$$\mathcal{F}^- = -\frac{\det(M_R^2 - M_I^2 - iM_{RI}^2 - iM_{RI}^{2T})}{m_1^2 m_2^2 m_3^2} \equiv -\frac{\det A}{m_1^2 m_2^2 m_3^2}, \quad (19)$$

where the denominators contain the product of the three nonzero eigenvalues of M_0^2 . In the CP -conserving case, Eqs. (18) and (19) reduce to the well-known expressions

$$\mathcal{F}^\pm = \frac{\sin^2(\alpha - \beta)}{M_{H_h}^2} + \frac{\cos^2(\alpha - \beta)}{M_{H_h}^2} \pm \frac{1}{M_A^2}, \quad (20)$$

where $M_{H,h}$ and M_A denote the CP -even and CP -odd Higgs-boson masses, respectively.

The discussion so far has been completely general. Particularizing to the MSSM, a perturbative matching calculation relates the two theories. At tree-level, this trivially results in

$$\begin{aligned}
M_d^{(0)} &= \frac{v}{\sqrt{2}} \cos \beta Y_d, & M_u^{(0)} &= \frac{v}{\sqrt{2}} \sin \beta Y_u, & \kappa^{(0)} &= -\sin \beta Y_d, & \tilde{\kappa}^{(0)} &= \cos \beta Y_u, & m_{11}^{2(0)} &= |\mu|^2 + m_{H_d}^2 \equiv m_1^2, \\
\lambda_1^{(0)} &= \lambda_2^{(0)} = -\lambda_3^{(0)} = (g^2 + g'^2)/4 \equiv \tilde{g}^2/4, & m_{22}^{2(0)} &= |\mu|^2 + m_{H_u}^2 \equiv m_2^2, & \lambda_4^{(0)} &= g^2/2, & m_{12}^{2(0)} &= B\mu, & \lambda_5^{(0)} &= \lambda_6^{(0)} = \lambda_7^{(0)} = 0.
\end{aligned}
\tag{21}$$

At this order, $\kappa^{(0)}$ and $\tilde{\kappa}^{(0)}$ are aligned with $M_d^{(0)}$ and $M_u^{(0)}$, respectively, so that no FCNC are induced, as it must be in a model II. At one-loop, all couplings in Eq. (12) are generated. Moreover, the corrections to the Yukawa couplings have the more general form

$$\begin{aligned}
M_d^{(1)} &= \frac{v}{\sqrt{2}} \cos \beta [\Delta Y_d + \tan \beta \Delta K], \\
\kappa^{(1)} &= -\sin \beta [\Delta Y_d - \cot \beta \Delta K],
\end{aligned}
\tag{22}$$

where ΔY_{dij} and ΔK_{ij} parametrize the one-loop vertices $\bar{d}_{Ri} H_d \cdot Q_{Lj}$ and $\bar{d}_{Ri} H_u^\dagger Q_{Lj}$, respectively. Diagonalizing M_d rotates $\kappa^{(0)}$, giving rise to a flavor-violating coupling $\propto Y_d \tan \beta / (16\pi^2)$, which can be of $\mathcal{O}(1)$ for $\tan \beta \sim 50$.

The origin of this explicit $\tan \beta$ enhancement (in addition to the mere presence of large down-type Yukawa couplings), which can compensate the loop factor $1/(16\pi^2)$, is the replacement of v_d by $v_u \gg v_d$ in the contribution of ΔK to M_d [1].² This removal of a v_d suppression can happen only in dimensionful quantities. In the fermion mass terms, only one power of $\tan \beta$ can appear because there was only one power of v_d to begin with. This is in agreement with the findings in [7]. Our approach using unshifted Higgs fields (“unbroken-theory”) makes particularly evident that this result holds to all orders, as the Yukawa Lagrangian only involves dimensionless couplings and there are no hidden factors of $\tan \beta$. Although we have integrated out only the

sparticles—as we assume a hierarchy $v, M_A \ll M_{\text{SUSY}}$ —the argument continues to hold if we also integrate over the Higgs fields, keeping only constant background values of Φ, Φ' (spurions). The reason is that for determining the mass matrices, the relevant external four-momenta are of $\mathcal{O}(m_q)$, providing an expansion parameter m_q/v or m_q/M_A . Hence, the Higgs contributions to the effective potential (which on general grounds respects the electro-weak symmetry) can be organized into a (local) effective Lagrangian, with m_q -suppressed corrections to the form Eq. (11) encoded in higher-dimensional operators with additional derivatives acting on d_{Ri} or Q_{Lj} . The contribution from both Higgs and sparticle loops to M_d is then simply obtained upon substituting for Φ, Φ' their vacuum expectation values. This mass matrix is to be identified with a short-distance (such as $\overline{\text{MS}}$) mass in the effective QCD $\times$ QED at low energies, where the dependence on the chosen scheme cancels against the explicit form of the matching (of the 2HDM onto QCD $\times$ QED).

There is only one other place where a similar $\tan \beta$ enhancement can occur, namely, in the dimensionful self-couplings of the (shifted) Higgs fields, that is, their masses and trilinear couplings. Indeed, at dimension two it is exhibited in the neutral-Higgs mass matrix Eq. (14). Explicitly, one has (with $s_\beta \equiv \sin \beta$, $c_\beta \equiv \cos \beta$, and $\lambda_k^r \equiv \text{Re} \lambda_k$)

$$M_R^2 = v^2 \begin{pmatrix} \lambda_5^r s_\beta^2 + 2\lambda_6^r s_\beta c_\beta + \lambda_1 c_\beta^2 & \lambda_7^r s_\beta^2 + \lambda_3 s_\beta c_\beta + \lambda_6^r c_\beta^2 \\ \lambda_7^r s_\beta^2 + \lambda_3 s_\beta c_\beta + \lambda_6^r c_\beta^2 & \lambda_2 s_\beta^2 + 2\lambda_7^r s_\beta c_\beta + \lambda_5^r c_\beta^2 \end{pmatrix} + M_I^2,
\tag{23}$$

$$M_I^2 = M_A^2 \begin{pmatrix} s_\beta^2 & -s_\beta c_\beta \\ -s_\beta c_\beta & c_\beta^2 \end{pmatrix},
\tag{24}$$

where $(m_{12}^2)^r$ has been traded for M_A^2 , with

$$s_\beta c_\beta M_A^2 = (m_{12}^2)^r - \frac{v^2}{2} (\lambda_7^r s_\beta^2 + 2\lambda_5^r s_\beta c_\beta + \lambda_6^r c_\beta^2).
\tag{25}$$

If CP is conserved, in the limit of infinite $\tan \beta$ ($c_\beta \rightarrow 0$) the leading mass splitting $M_H^2 - M_A^2 = \lambda_5 v^2$, and the leading correction to the tree-level result $\alpha = 0$ is deter-

mined by λ_7 . In the former case, an enhancement by two powers of $\tan \beta$ occurs ($M_H^2 - M_A^2 = \mathcal{O}(\cos^2 \beta)$ at tree-level), while the loop correction to α is enhanced by a single power of $\tan \beta$ with respect to its tree-level value. Either effect is sufficient to remove the cancellation in $\mathcal{F}^-$ in Eq. (20) observed at tree-level. Moreover, a $1/\tan \beta$ -unsuppressed CP -violating contribution proportional to λ_5^i and λ_7^i appears to occur:

$$M_{RI}^2 = \frac{v^2}{2} \begin{pmatrix} \lambda_5^i s_\beta^2 + 2\lambda_6^i s_\beta c_\beta & -\lambda_5^i s_\beta c_\beta - 2\lambda_6^i c_\beta^2 \\ 2\lambda_7^i s_\beta^2 + \lambda_5^i s_\beta c_\beta & -2\lambda_7^i s_\beta c_\beta - \lambda_5^i c_\beta^2 \end{pmatrix},
\tag{26}$$

where $\lambda_k^i \equiv \text{Im} \lambda_k$. However, as we show in Sec. IIc below, the individual phases of λ_5 and λ_7 become unphysical in the limit $\tan \beta \rightarrow \infty$, and mixing between the CP -even and CP -odd sectors is described by a single

²We tacitly assume that the fermion kinetic terms in the effective 2HDM have been made canonical. Such a field renormalization does not contribute factors of $\tan \beta$ because it is determined by dimensionless couplings. Cf. Sec. III for a discussion of field renormalization. Our ΔK and ΔY_d correspond to $\Delta_u Y_d$ and $-\Delta_d Y_d$, respectively, in the first paper of Ref. [5].

angle α' , determined by the relative phase of λ_5 and λ_7' . Finally, the charged-Higgs mass matrix is given by

$$M_{\pm}^2 = \left(1 + \frac{v^2(\lambda_4 + \lambda_5')}{2M_A^2}\right) M_{\pm}^2. \quad (27)$$

Here, no $\tan\beta$ enhancement due to loop-induced couplings occurs.

Unlike the case of the fermion mass matrix, the typical momentum flowing through the effective Lagrangian Eq. (12) for an on-shell Higgs is itself of $\mathcal{O}(v)$ or $\mathcal{O}(M_A)$. Hence Higgs-loop contributions to the Higgs masses cannot be included in Eq. (12), but rather the full effective action would be needed. Higgs-loop effects in κ_{bq} and κ_{qb} multiplying $\mathcal{F}^{\pm}$ could, however, be included via Eq. (11), since again the momenta flowing through the vertices are much smaller than v, M_A . This is not possible in Higgs boxes, where large momenta flow through the FCNC vertices. We will present a systematic method to include all Higgs-loop contributions in Sec. IIc.

It is instructive to consider the explicit form of the numerator in Eq. (19), which is

$$\begin{aligned} \det A = v^4 [& (\lambda_2 \lambda_5^* - \lambda_7'^2) s_{\beta}^4 + 2(\lambda_2 \lambda_6^* - \lambda_3 \lambda_7' + \lambda_5^* \lambda_7) s_{\beta}^3 c_{\beta} \\ & + (\lambda_1 \lambda_2 - \lambda_3^2 + |\lambda_5|^2 - 2\lambda_6 \lambda_7^* + 4\lambda_6^* \lambda_7) s_{\beta}^2 c_{\beta}^2 \\ & + 2(\lambda_5 \lambda_6^* - \lambda_3 \lambda_6 + \lambda_1 \lambda_7) s_{\beta} c_{\beta}^3 + (\lambda_1 \lambda_5 - \lambda_6^2) c_{\beta}^4]. \end{aligned} \quad (28)$$

With Eq. (21), $\det A = v^4 (\lambda_1 \lambda_2 - \lambda_3^2) s_{\beta}^2 c_{\beta}^2 = 0$, reproducing the known vanishing of $\mathcal{F}^-$ employing the tree-level MSSM Higgs sector. The cancellation is removed already at the leading-logarithmic level. For instance, λ_2 alone receives a large additive correction $\propto y_t^4$ due to top-quark loops, which is also responsible for the most important correction to the tree-level mass of h . The corresponding corrections could be computed by RG-evolving the tree-level couplings in the effective 2HDM. However, as we are considering large $\tan\beta$, we expect (and find below) the most important effect to be due to λ_5 and λ_7 , which remove the $\mathcal{O}(c_{\beta}^2)$ suppression of the leading-log result, as anticipated above.

B. The case of minimal flavor-violation

From the discussion so far, it follows that $|\mathcal{F}^+| = \mathcal{O}(1/M_A^2) \gg |\mathcal{F}^-| = \mathcal{O}(1/(16\pi^2 M_A^2))$, implying $|C_2^{\text{LR}}| \gg |C_1^{\text{SL}}|$ for generic κ_{ij} ,³ such that the motivation to consider $\mathcal{F}^-$ at all is not very strong. The situation is fundamentally different for MFV because then the contribution propor-

tional to $\mathcal{F}^+$ turns out to be suppressed by a light quark mass, introducing a further small parameter m_q/m_b comparable to $1/(16\pi^2)$ or $1/\tan\beta$ for $q = s$ (and negligible for $q = d$). For simplicity, in this paper we consider the simplest version of MFV, assuming flavor-universal soft-breaking terms $\tilde{m}_Q^2$, $\tilde{m}_u^2$, and $\tilde{m}_d^2$ and trilinear SUSY-breaking terms $T_{u_{ij}}$, $T_{d_{ij}}$ which are proportional to the Yukawa matrices and therefore diagonal in the super-CKM basis (denoted with a hat): $\hat{T}_{u_{ij}} = a_t y_{u_i} \delta_{ij}$ and $\hat{T}_{d_{ij}} = a_b y_{d_i} \delta_{ij}$, see Appendix A for details of our notation. The structure of our results, however, does not depend on these additional assumptions. The $\tan\beta$ -enhanced loop-induced FCNC couplings of the neutral-Higgs bosons in Eq. (11) can be expressed as:

$$\kappa_{bq} = \epsilon_Y y_t^2 \lambda_{qb} \frac{\sqrt{2} m_b}{v \cos^2 \beta} \frac{1}{1 + \tilde{\epsilon}_3 \tan \beta} \frac{1}{1 + \epsilon_0 \tan \beta}, \quad (29)$$

$$\kappa_{qb} = \epsilon_Y y_t^2 \lambda_{qb}^* \frac{\sqrt{2} m_q}{v \cos^2 \beta} \frac{1}{1 + \tilde{\epsilon}_3 \tan \beta} \frac{1}{1 + \epsilon_0 \tan \beta}, \quad (30)$$

with $y_t = \sqrt{2} m_t / (v \sin \beta)$ and $\lambda_{qb} = V_{tq} V_{tb}^*$. The effective couplings ϵ_Y , ϵ_0 , and $\tilde{\epsilon}_3$, which depend on the MSSM parameters, have been analyzed in the decoupling limit $M_{\text{SUSY}} \gg v$ in the limit $g = g' = 0$ in Refs. [2–4] for the case that ϵ_Y , ϵ_0 and $\tilde{\epsilon}_3$ are real and in Ref. [25] for the maximally CP -violating MFV scenario. We consider the CP -violating case allowing μ , the universal trilinear term a_t and the gaugino mass parameters to be complex. Effects from nonzero g, g' have been taken into account in Ref. [5], where also effects beyond the decoupling limit were considered. The corresponding expressions for $M_{\text{SUSY}} \gg v$, suited for our analysis, were derived in Ref. [26]. We have recalculated the FCNC couplings of neutral-Higgs bosons for $g, g' \neq 0$ including all CP -violating phases and found agreement with the results for the FCNC self-energies given in Ref. [5], but encountered a significant discrepancy with Ref. [26]. In our results, the phase conventions of μ , a_t , and M_2 can be inferred from Eqs. (A1) and (A3) of Appendix A. The phase convention for $M_{1,3}$ complies with that of M_2 and the gluino mass equals $M_{\tilde{g}} = |M_3|$. Of course, one can choose one of these parameters (e.g. M_3) real. Now the effective couplings of Eqs. (29) and (30) read:

$$\begin{aligned} \epsilon_0 = & \frac{-2\alpha_s}{3\pi} \frac{\mu^*}{M_3} H_2 \left(\frac{M_{\tilde{b}L}^2}{|M_3|^2}, \frac{M_{\tilde{b}R}^2}{|M_3|^2} \right) + \frac{g'^2}{96\pi^2} \frac{\mu^*}{M_1} \\ & \times \left[H_2 \left(\frac{M_{\tilde{b}L}^2}{|M_1|^2}, \frac{|\mu|^2}{|M_1|^2} \right) + 2H_2 \left(\frac{M_{\tilde{b}R}^2}{|M_1|^2}, \frac{|\mu|^2}{|M_1|^2} \right) \right] \\ & + \frac{g'^2}{144\pi^2} \frac{\mu^*}{M_1} H_2 \left(\frac{M_{\tilde{b}L}^2}{|M_1|^2}, \frac{M_{\tilde{b}R}^2}{|M_1|^2} \right) \\ & + \frac{3g^2}{32\pi^2} \frac{\mu^*}{M_2} H_2 \left(\frac{M_{\tilde{b}L}^2}{|M_2|^2}, \frac{|\mu|^2}{|M_2|^2} \right), \end{aligned} \quad (31)$$

³In Ref. [13], an argument based on $SU(2) \times U(1)$ gauge invariance was used to infer that (in the present notation) $\mathcal{F}^- = \mathcal{O}(v^2/M_A^4)$. This statement, which clearly is respected by our Eq. (19) in conjunction with Eq. (28) (recall $M_h^2 = \mathcal{O}(v^2)$), is about the asymptotic behavior as $v/M_A \rightarrow 0$. The latter is not necessarily a small number in practice. Indeed, many of the analyses in the literature have dealt with the case $M_A \sim 200$ GeV.

$$\epsilon_Y = \frac{-1}{16\pi^2} \frac{a_t^*}{\mu} H_2\left(\frac{M_{\tilde{t}L}^2}{|\mu|^2}, \frac{M_{\tilde{t}R}^2}{|\mu|^2}\right) + \epsilon_{Y,v/M}, \quad (32)$$

$$\tilde{\epsilon}_3 = \epsilon_0 + y_t^2 \epsilon_Y. \quad (33)$$

Here

$$H_2(x, y) = \frac{x \log x}{(1-x)(x-y)} + \frac{y \log y}{(1-y)(y-x)}. \quad (34)$$

Numerically, the electroweak contributions in ϵ_0 can be of $\mathcal{O}(10\%)$. They improve the comparison with the results computed with full chargino and squark mass matrices (see Eq. (5.1) in the second paper in Ref. [5]).

Ref. [5] also discusses threshold corrections to the fermion kinetic operators (wave-function renormalizations). While these terms are not $\tan\beta$ -enhanced, the flavor-diagonal quark wave-function-renormalization constants receive sizable contributions from squark-gluino loops. One can parametrize these loops in terms of a new quantity $\epsilon_0|_{\text{kin}}$ which will add to ϵ_0 in the relation between the MSSM Yukawa coupling y_{d_i} and the physical quark mass m_{d_i} (see Eq. (A6) for the case of the bottom Yukawa coupling). $\epsilon_0|_{\text{kin}}$ will likewise appear in the relation between κ_{ij} and y_{d_i} , but it drops out once κ_{ij} is expressed in terms of m_{d_i} , so that it does not appear in Eqs. (29) and (30). This cancellation of the flavor-diagonal quark wave-function-renormalization can be verified by inserting Eq. (2.29) into Eq. (2.26) of the second paper in Ref. [5]. This feature can be traced back to the fact that the wave-function-renormalization affects both the tree-level and the loop-induced Yukawa couplings with the same multiplicative factor.

Comparing our result with Ref. [26], we find different results for ϵ_0 and ϵ_Y : In Ref. [26], the chargino-stop contribution proportional to g^2 is erroneously assigned to ϵ_Y rather than ϵ_0 . Since this piece does not contain any up-type Yukawa couplings, all three generations contribute in the same way and the resulting overall CKM structure combines to $V_{ub}^* V_{uq} + V_{cb}^* V_{cq} + V_{tb}^* V_{tq}$, which is zero for $q \neq b$ and equal to one for $q = b$. This GIM cancellation eliminates the wino-stop loop from ϵ_Y , while this loop contributes to ϵ_0 twice as much as the corresponding loop with a neutral winolike neutralino and a sbottom. The two terms are combined into the last term in Eq. (31). Omitting the chargino loop here would violate $SU(2)$ gauge symmetry, which also enforces $M_{\tilde{t}L} = M_{\tilde{b}L}$ in the limit $M_{\text{SUSY}} \gg v$. Since ϵ_Y normalizes all Higgs-induced FCNC couplings, one should verify the accuracy of this limit: It is easy to include the $\tan\beta$ -enhanced contributions to ϵ_Y to all orders in v/M_{SUSY} . To this end, one merely has to calculate the FCNC $\bar{b}_R q_L$ self-energy using the exact chargino and up-squark mass eigenstates. This self-energy renormalizes the off-diagonal pieces of the quark mass matrix and causes the mismatch between the flavor structures of the latter with the Yukawa couplings leading to $\epsilon_Y \neq 0$. In

higher loop-orders, $\tan\beta$ -enhanced contributions are suppressed by products of small CKM elements (and are negligible) or are flavor-conserving and therefore contribute to ϵ_0 rather than to ϵ_Y . Using the $\bar{b}_R q_L$ self-energy $(\Sigma_{mL}^d)^{3i}$ (with $q = d_i$) from Ref. [5], one finds

$$\epsilon_{Y,v/M} = \frac{1}{16\pi^2} \frac{a_t^*}{\mu} H_2\left(\frac{M_{\tilde{t}L}^2}{|\mu|^2}, \frac{M_{\tilde{t}R}^2}{|\mu|^2}\right) + \frac{\sqrt{2}}{vy_t^2 \lambda_{bd_i}} \frac{(\Sigma_{mL}^d)^{3i}}{y_b} \quad (35)$$

(Note that $(\Sigma_{mL}^d)^{3i} \propto y_b$ and be aware of the different sign conventions for y_b in Eq. (A6) and Ref. [5].) We stress that Eq. (35) must be evaluated for $i \neq 3$, so that the GIM cancellation of the above-mentioned wino-stop loop takes place. Numerically, one finds a marginal impact of $\epsilon_{Y,v/M}$: Setting all supersymmetric massive parameters equal to a common value M_{SUSY} , one finds that $\epsilon_{Y,v/M}$ amounts to a mere 1.4% correction to ϵ_Y for $M_{\text{SUSY}} = 400$ GeV. Even for $M_{\text{SUSY}} = 150$ GeV, for which the expansion in v/M_{SUSY} formally breaks down, $\epsilon_{Y,v/M}$ depletes ϵ_Y by as little as 8%. $\epsilon_{Y,v/M}$ also enters $\tilde{\epsilon}_3$ through Eq. (33). It can be inferred from Ref. [7] that this procedure indeed leads to the correct all-order resummation of the $\tan\beta$ -enhanced corrections involving y_t . Corrections to $\tilde{\epsilon}_3$ beyond the $M_{\text{SUSY}} \gg v$ limit from α_s , g , g' , and y_b are considered in Refs. [5,7]. We remark that no terms proportional to y_b^2 occur in Eqs. (31)–(33), because the corresponding loops violate hypercharge and involve a suppression factor of v^2/M_{SUSY}^2 .

We verify from Eqs. (29) and (30) that $\kappa_{qb}^* \kappa_{bq}$ multiplying $\mathcal{F}^+$ in Eq. (17) is suppressed by a factor m_q/m_b relative to κ_{bq}^2 , which multiplies $\mathcal{F}^-$. Hence, C_1^{SLL} is naively leading (over C_2^{LR}) from the point of view of MFV alone, and a meaningful analysis of $B_q - \bar{B}_q$ mixing requires a systematic investigation of all leading corrections to its vanishing “tree” value. (The coefficient C_1^{SRR} both undergoes a strong m_q^2/m_b^2 suppression and involves $\mathcal{F}^{*-}$, and can thus be disregarded.) It is then useful to think of the $\Delta B = 2$ amplitude as being a function of the four small parameters identified so far:

$$l \equiv \frac{1}{(4\pi)^2}, \quad \omega \equiv \frac{m_q}{m_b}, \quad \frac{1}{\tan\beta}, \quad \nu \equiv \frac{v}{M_{\text{SUSY}}}. \quad (36)$$

The vanishing 2HDM tree diagram for $\mathcal{F}^-$ is (superficially) $\mathcal{O}((\cot\beta)^{-2} l^2 \nu^0 \omega^0)$, i.e. $\mathcal{O}(1)$ when treating all expansion parameters on the same footing. Conversely, $\mathcal{F}^+$ is nonzero at the tree-level but is suppressed by one power of ω , which is non-negligible only for $q = s$. We have already seen that $\mathcal{F}^-$ vanishes exactly for tree-level matching (or up to $\mathcal{O}(1/\tan^2\beta)$ when including leading logs), so there are no $\mathcal{O}(1/\tan\beta)$ corrections at first sub-leading order. This leaves loop corrections (via sparticle corrections to the λ_i as well as loops in the effective 2HDM) and possible corrections due to higher-dimensional operators, not written in Eqs. (11) and (12). We now discuss these contributions in turn.

Sparticle loops One-loop contributions from higgsinos, gauginos, and sfermions correct the values of $\lambda_{1,2,3,4}$ in Eq. (12) and induce nonzero couplings $\lambda_{5,6,7}$. As a technical result of our paper, we have computed the λ_i for general sparticle masses and flavor structure. These results are reported in Appendix B. At tree-level in the effective theory and in the leading-order of $1/\tan\beta$ the quantity $\mathcal{F}^-$ receives only contributions from λ_2 , λ_5 , and λ_7 , cf. Equation (28). The general results of Eqs. (71), (B1), (B5), (B6), (B8), (B10), (B11), and (B13), for the MFV case read

$$\begin{aligned} \lambda_7 &= \bar{\lambda}_7 \\ &= \frac{1}{16\pi^2} \left\{ \frac{1}{4} \mu a_b |y_b|^2 ((3g^2 + g'^2) C_0(\tilde{m}_d, \tilde{m}_Q, \tilde{m}_Q) + 2g'^2 C_0(\tilde{m}_Q, \tilde{m}_d, \tilde{m}_d)) + \mu a_t |y_t|^4 (3|a_t|^2 D_0(\tilde{m}_Q, \tilde{m}_Q, \tilde{m}_u, \tilde{m}_u) \right. \\ &\quad + 3C_0(\tilde{m}_Q, \tilde{m}_Q, \tilde{m}_u) + 3C_0(\tilde{m}_Q, \tilde{m}_u, \tilde{m}_u)) + 3\mu a_b |\mu|^2 |y_b|^4 D_0(\tilde{m}_d, \tilde{m}_d, \tilde{m}_Q, \tilde{m}_Q) \\ &\quad - \frac{1}{4} \mu a_t |y_t|^2 ((3g^2 - g'^2) C_0(\tilde{m}_Q, \tilde{m}_Q, \tilde{m}_u) + 4g'^2 C_0(\tilde{m}_Q, \tilde{m}_u, \tilde{m}_u)) + \frac{1}{4} \mu a_\tau |y_\tau|^2 (2g'^2 C_0(\tilde{m}_l, \tilde{m}_e, \tilde{m}_e) \\ &\quad + (g^2 - g'^2) C_0(\tilde{m}_e, \tilde{m}_l, \tilde{m}_l)) + \mu a_\tau |\mu|^2 |y_\tau|^4 D_0(\tilde{m}_e, \tilde{m}_e, \tilde{m}_l, \tilde{m}_l) - \frac{1}{4} \tilde{g}^2 \mu (3a_b |y_b|^2 B'_0(\tilde{m}_d, \tilde{m}_Q) \\ &\quad + 3a_t |y_t|^2 B'_0(\tilde{m}_u, \tilde{m}_Q) + a_\tau |y_\tau|^2 B'_0(\tilde{m}_e, \tilde{m}_l)) + g^4 \left(3\mu M_2 \tilde{D}_2(|M_2|, |M_2|, |\mu|, |\mu|) - \frac{3}{4} \mu M_2 B'_0(|M_2|, |\mu|) \right) \\ &\quad - \frac{1}{4} g^2 g'^2 \mu (M_1 B'_0(|M_1|, |\mu|) + 3M_2 B'_0(|M_2|, |\mu|) - 4(M_1 + M_2) \tilde{D}_2(|M_1|, |M_2|, |\mu|, |\mu|)) \\ &\quad \left. + g'^4 (\mu M_1 \tilde{D}_2(|M_1|, |M_1|, |\mu|, |\mu|) - \frac{1}{4} \mu M_1 B'_0(|M_1|, |\mu|)) \right\}, \end{aligned} \quad (37)$$

$$\begin{aligned} \lambda_5 &= \bar{\lambda}_5 \\ &= -\frac{1}{16\pi^2} \mu^2 \{ 3a_b^2 |y_b|^4 D_0(\tilde{m}_d, \tilde{m}_d, \tilde{m}_Q, \tilde{m}_Q) + 3a_t^2 |y_t|^4 D_0(\tilde{m}_Q, \tilde{m}_Q, \tilde{m}_u, \tilde{m}_u) + a_\tau^2 |y_\tau|^4 D_0(\tilde{m}_e, \tilde{m}_e, \tilde{m}_l, \tilde{m}_l) \\ &\quad - 3g^4 M_2^2 D_0(|M_2|, |M_2|, |\mu|, |\mu|) - 2g^2 g'^2 M_1 M_2 D_0(|M_1|, |M_2|, |\mu|, |\mu|) - g'^4 M_1^2 D_0(|M_1|, |M_1|, |\mu|, |\mu|) \}, \end{aligned} \quad (38)$$

and

$$\begin{aligned} \lambda_2 &= \bar{\lambda}_2 \\ &= \frac{\tilde{g}^2}{4} + \frac{1}{16\pi^2} \left\{ -\frac{3}{4} g'^4 B_0(\tilde{m}_e, \tilde{m}_e) - \frac{3}{8} (g^4 + g'^4) B_0(\tilde{m}_l, \tilde{m}_l) + \frac{1}{2} (g'^2 - g^2) |\mu y_\tau|^2 C_0(\tilde{m}_e, \tilde{m}_l, \tilde{m}_l) - g'^2 |\mu y_\tau|^2 C_0(\tilde{m}_l, \tilde{m}_e, \tilde{m}_e) \right. \\ &\quad - |\mu y_\tau|^4 D_0(\tilde{m}_e, \tilde{m}_e, \tilde{m}_l, \tilde{m}_l) - \frac{1}{4} g'^4 B_0(\tilde{m}_d, \tilde{m}_d) + (-3|y_t|^4 + 2g^2 |y_t|^2 - g'^4) B_0(\tilde{m}_u, \tilde{m}_u) + \frac{1}{8} (-9g^4 - 24|y_t|^4 - g'^4 \\ &\quad - 4|y_t|^2 (g'^2 - 3g^2)) B_0(\tilde{m}_Q, \tilde{m}_Q) + \frac{1}{2} (3g^2 - 12|y_t|^2 - g'^2) |a_t y_t|^2 C_0(\tilde{m}_Q, \tilde{m}_Q, \tilde{m}_u) + 2(g'^2 - 3|y_t|^2) |a_t y_t|^2 C_0(\tilde{m}_Q, \tilde{m}_u, \tilde{m}_u) \\ &\quad - g'^2 |\mu y_b|^2 C_0(\tilde{m}_Q, \tilde{m}_d, \tilde{m}_d) - \frac{1}{2} (3g^2 + g'^2) |\mu y_b|^2 C_0(\tilde{m}_d, \tilde{m}_Q, \tilde{m}_Q) - 3|a_t y_t|^4 D_0(\tilde{m}_Q, \tilde{m}_Q, \tilde{m}_u, \tilde{m}_u) \\ &\quad - 3|\mu y_b|^4 D_0(\tilde{m}_d, \tilde{m}_d, \tilde{m}_Q, \tilde{m}_Q) + \frac{1}{2} \tilde{g}^2 (3|\mu y_b|^2 B'_0(\tilde{m}_d, \tilde{m}_Q) + |\mu y_\tau|^2 B'_0(\tilde{m}_e, \tilde{m}_l) + 3|a_t y_t|^2 B'_0(\tilde{m}_u, \tilde{m}_Q)) \\ &\quad + \frac{1}{24} \left[-2 \log \frac{\tilde{m}_d^2}{\mu_0^2} g'^4 - 6 \log \frac{\tilde{m}_e^2}{\mu_0^2} g'^4 - 8 \log \frac{\tilde{m}_u^2}{\mu_0^2} g'^4 - 3 \log \frac{\tilde{m}_l^2}{\mu_0^2} (g^4 + g'^4) - \log \frac{\tilde{m}_Q^2}{\mu_0^2} (9g^4 + g'^4) \right] \\ &\quad - \frac{1}{24} g^4 \left[-12 \tilde{D}_2(|M_2|, |M_2|, |\mu|, |\mu|) |M_2|^2 - 60 \tilde{D}_4(|M_2|, |M_2|, |\mu|, |\mu|) + 9W(|M_2|, |\mu|) \right. \\ &\quad \left. + 4 \log \frac{|\mu|^2}{\mu_0^2} + 8 \log \frac{M_2^2}{\mu_0^2} + 14 \right] - \frac{1}{8} g^2 g'^2 [-8 \text{Re}(M_1 M_2^*) \tilde{D}_2(|M_1|, |M_2|, |\mu|, |\mu|) - 8 \tilde{D}_4(|M_1|, |M_2|, |\mu|, |\mu|) \\ &\quad + W(|M_1|, |\mu|) + 3W(|M_2|, |\mu|) + 4] - \frac{1}{24} g'^4 \left[-12 \tilde{D}_2(|M_1|, |M_1|, |\mu|, |\mu|) |M_1|^2 - 12 \tilde{D}_4(|\mu|, |\mu|, |M_1|, |M_1|) \right. \\ &\quad \left. + 3W(|M_1|, |\mu|) + 4 \log \left(\frac{|\mu|^2}{\mu_0^2} + 6 \right) \right] \right\}, \end{aligned} \quad (39)$$

where the loop functions $B_0, C_0, D_0, B'_0, \tilde{D}_2, \tilde{D}_4$, and W are defined in Appendix B d, and the notation $\bar{\lambda}_i$ refers to the matching scheme as explained in Sec. III. Inspecting Eq. (28), λ_7 enters quadratically, which formally is of higher loop order. Nevertheless, it can be seen that $\lambda_7^2 \propto y_t^8$, as opposed to $\lambda_2 \lambda_5^* \propto \tilde{g}^2 y_t^4$, which can partly offset the additional loop-suppression. Indeed we find that, numerically, neglecting λ_7 is not always a good approximation (Sec. IV).

The form of the matching result depends on the renormalization schemes of both the full theory, i.e., the MSSM, and the effective theory, i.e., the 2HDM. The latter cancels in physical quantities, while explicit MSSM scheme dependence cancels against the one implicit in the MSSM parameters, to ensure that the couplings in the effective theory are independent of the renormalization of the MSSM at any given order of perturbation theory. The residual scheme dependence in both cases may, however, be important as we are considering a leading effect. We will discuss scheme issues in Sec. IIIa, paying special attention to the definitions of $\tan\beta$.

Higgs loops There is a considerable number of one-loop diagrams in the effective 2HDM that can contribute to $B - \bar{B}$ mixing amplitudes (Fig. 2, upper row). These give the following contributions to the Wilson coefficients multiplying Q_1^{VLL} and Q_1^{VRR} :

$$C_1^{\text{VLL}}|_{\text{Higgsloops}} = -\frac{1}{4} \frac{m_b^2}{v^2 \cos^2 \beta (1 + \tilde{\epsilon}_3^* \tan \beta)^2} \times \frac{\kappa_{bq}^2}{G_F^2 M_W^2 \lambda_{qb}^2} C_0(M_A^2, M_A^2, 0), \quad (40)$$

$$C_1^{\text{VRR}}|_{\text{Higgsloops}} = -\frac{1}{4} \frac{m_b^2}{v^2 \cos^2 \beta (1 + \tilde{\epsilon}_3 \tan \beta)^2} \times \frac{\kappa_{qb}^{*2}}{G_F^2 M_W^2 \lambda_{qb}^2} C_0(M_A^2, M_A^2, 0). \quad (41)$$

In these expressions, we have neglected the small Yukawa coupling y_q and employed tree-level MSSM mass relations, in agreement with our approximation of working to leading-order in small parameters (in the present case, the loop factor $1/(16\pi^2)$). C_1^{VRR} is suppressed by two powers of m_q/m_b inside κ_{qb}^{*2} in the MFV case, hence beyond our accuracy. The results Eqs. (40) and (41) involve a great deal of cancellations, which can be understood in terms of symmetry arguments, as explained in Sec. IIc below. We note the absence of charged-Higgs contributions in the approximation considered here.

v/M -suppressed effects All of the couplings given in Eq. (11) correspond to the zeroth order in the v/M_{SUSY} expansion, or equivalently to the level of dimension-four operators. Gauge invariance forbids dimension-five operators built from quark and Higgs fields, so the leading higher-dimensional operators have dimension six. This can lead to more general Higgs-fermion couplings than those deriving from Eq. (11) and, in consequence, the cancellation leading to $C_1^{\text{SLL}} = 0$ might be broken. To see that this is indeed the case, consider the operator

$$Q^{(6)} = \frac{1}{M_{\text{SUSY}}^2} (H_u^\dagger H_u) (\bar{b}_R H_u^\dagger Q_L), \quad (42)$$

which gives rise, *inter alia*, to effective dimension-three and -four couplings




FIG. 2. Upper row: A subset of one-loop diagrams for $B_q - \bar{B}_q$ mixing in the effective two-Higgs-doublet model. Lower row: Tree and one-loop diagrams contributing at large $\tan\beta$ when employing the Lagrangian $\mathcal{L}_{\text{lb}}$ and tree-level couplings. The crosses denote the flavor-changing neutral-Higgs couplings and [in diagrams (f) and (g)] loop-suppressed Higgs mass terms. On the lower row, arrows designate the flow of the conserved $U(1)$ charge discussed in Sec. IIc.

$$\frac{2\sqrt{2}v_u^3}{M_{\text{SUSY}}^2} \bar{b}_R s_L + \frac{2v_u^2}{M_{\text{SUSY}}^2} (\bar{b}_R s_L h_u^0 + 2\bar{b}_R s_L h_u^{0*}). \quad (43)$$

The first term is removed by a re-diagonalization of the quark mass matrices, but the two remaining terms, in general, are not. The appearance of h_u^0 in addition to h_u^{0*} leads to a contribution to C_1^{SLL} proportional to $\kappa_{bq} C^{(6)}$. However, because of R -parity, SUSY particles do not contribute to tree graphs with external standard particles only, such that $Q^{(6)}$ (or any other higher-dimension operator) is only induced at the loop level, and this loop-suppression factor is not compensated by factors of $\tan\beta$. (Recall that the $\mathcal{O}(1)$ FCNC couplings at dimension-four are nothing but rotated tree-level Yukawa couplings.) Hence any v/M_{SUSY} corrections that break the cancellation in $\mathcal{F}^-$ involve an additional loop-suppression, and can be neglected for the present analysis. On the other hand, as Eq. (43) shows, the higher-dimensional operators do have an impact on the re-diagonalization of the quark mass matrices and, consequently, on the size of the FCNC couplings κ_{bq} . These effects preserve the cancellations in $\mathcal{F}^-$ discussed above but have a mild impact on the FCNC couplings multiplying $\mathcal{F}^+$ in C_2^{LR} (cf. Eq. (35) and the discussion around it).

C. $U(1)_{\text{PQ}}$ and effective Lagrangian for large $\tan\beta$

To better understand the various types of cancellations in $\mathcal{F}^-$ and in the Higgs-loop contributions to C_1^{VLL} , as well as the suppression of the $\mathcal{F}^+$ contribution, we now introduce an effective 2HDM Lagrangian at large $\tan\beta$. This will allow us, on the basis of simple symmetry arguments, to clarify the role of the parameters λ_5 and λ_7 , the structure of Eqs. (18), (19), and (28), as well as the vanishing of $\mathcal{F}^-$ for tree-level Higgs couplings at leading-order in $1/\tan\beta$. It also provides a tool for computing loop diagrams involving Higgs bosons efficiently and consistently, which may be useful in other contexts such as collider processes with Higgses in the initial or final state.

As before, we eliminate m_{11}^2 , m_{22}^2 , and $(m_{12}^2)^i$ by the minimization conditions and trade $(m_{12}^2)^r$ for M_A^2 via Eq. (25). We then take the limit

$$v_d \rightarrow 0, \quad v_u \rightarrow v, \quad M_A^2 \text{ fixed}, \quad \lambda_i \text{ fixed}, \quad (44)$$

of the Lagrangian (12) in the broken phase.⁴ We also keep the Yukawa couplings fixed when considering the couplings to fermions. In this limit, we have $\Phi = H_u$, $\Phi' = \epsilon H_d^*$, and

$$\begin{aligned} h_u^0 &= \frac{1}{\sqrt{2}}(v + \phi_u + iG^0), & h_d^0 &= \frac{1}{\sqrt{2}}(\phi_d - iA^0), \\ h_u^+ &= G^+, & h_d^+ &= H^+. \end{aligned} \quad (45)$$

⁴This procedure will be justified in Sec. IIIb

If there were no mixing among neutral Higgses, we would have $\phi_u = h^0$ and $\phi_d = H^0$, and A^0 would be a mass eigenstate. The mass matrices are compactly expressed by the quadratic potential

$$\begin{aligned} V_{\text{lib}}^{(2)} &= \left[m_A^2 + \frac{\lambda_5^r}{2} v^2 \right] H_d^\dagger H_d + \frac{\lambda_4}{2} v^2 |h_d^+|^2 + \frac{\lambda_2}{2} v^2 \phi_u^2 \\ &+ \left[\frac{\lambda_5}{4} (h_d^{0*})^2 + \frac{\lambda_7}{\sqrt{2}} \phi_u h_d^{0*} + \text{H.c.} \right] v^2, \end{aligned} \quad (46)$$

valid up to corrections of order $\cos\beta \sim 1/\tan\beta \ll 1$. The trilinear terms are given in Appendix D; the quartic terms follow trivially from those in the symmetric Lagrangian Eq. (12). Note that the first line of Eq. (46) is symmetric under the $U(1)$ Peccei-Quinn (PQ) transformation

$$\begin{aligned} h_d^0 &\rightarrow e^{-i\delta} h_d^0, & h_d^+ &\rightarrow e^{-i\delta} h_d^+, \\ \text{or equivalently, } H_d &\rightarrow e^{i\delta} H_d, \end{aligned} \quad (47)$$

while the second line is not. In the MSSM, the noninvariant terms appear only at the loop level. We note that the $U(1)$ symmetry is not spontaneously broken in the large- $\tan\beta$ limit, so there is no massless boson, in agreement with our keeping M_A^2 fixed.⁵ Next, a PQ transformation makes λ_5 real, such that the first term on the second line of Eq. (46) contributes with opposite sign to the mass terms for ϕ_d and $\chi_d = -A^0 + \mathcal{O}(\cos\beta)$, splitting the two. There are only two independent mixing angles that do not vanish: they can be identified with the CP -conserving angle $\alpha = \mathcal{O}(\lambda_7^r)$ and a CP -violating $\alpha' = \mathcal{O}(\lambda_7^i)$; a third angle present in the general 2HDM is suppressed by $\mathcal{O}(\cot\beta; v/M)$. All of these are symmetry-breaking effects. To lowest order in the PQ-breaking couplings, the mass matrices are diagonalized by

$$\begin{pmatrix} H_1 \\ H_2 \\ H_3 \end{pmatrix} = \begin{pmatrix} 1 & -\frac{\lambda_7^r v^2}{M_A^2 - \lambda_2 v^2} & \frac{\lambda_7^i v^2}{M_A^2 - \lambda_2 v^2} \\ \frac{\lambda_7^r v^2}{M_A^2 - \lambda_2 v^2} & 1 & 0 \\ -\frac{\lambda_7^i v^2}{M_A^2 - \lambda_2 v^2} & 0 & 1 \end{pmatrix} \begin{pmatrix} \phi_u \\ \phi_d \\ A^0 \end{pmatrix}, \quad (48)$$

$$m_1^2 = \lambda_2 v^2, \quad m_2^2 = M_A^2 + |\lambda_5| v^2, \quad m_3^2 = M_A^2. \quad (49)$$

In a general basis, CP -violating Higgs mixing is present if and only if λ_7^i/λ_5 is complex. Note that there is no mixing for the charged scalars according to Eq. (46), i.e. no mixing between charged-Higgs and Goldstone bosons due to sparticles in the large- $\tan\beta$ limit.

These considerations can be extended to the Higgs-fermion interactions. The operators up to dimension-four follow from (11), which, in the limit of infinite $\tan\beta$, becomes

⁵Also at finite (but large) $\tan\beta$, there is no (pseudo-) Goldstone boson, as $m_{11}^2 \sim M_A^2 > 0$ contributes to the mass terms of both ϕ_d and A^0 (see also Sec. IIIb).

$$\begin{aligned} \mathcal{L}_{\text{lib}}^Y = & -\frac{\sqrt{2}}{v} \bar{d}_{Ri} M_{d_{ij}} H_u^\dagger Q_{Lj} - \bar{d}_{Ri} \kappa_{ij} Q_{Lj} \cdot H_d \\ & -\frac{\sqrt{2}}{v} \bar{u}_{Ri} M_{u_{ij}} Q_{Lj} \cdot H_u + \bar{u}_{Ri} \tilde{\kappa}_{ij} H_d^\dagger Q_{Lj} + \text{H.c.} \end{aligned} \quad (50)$$

This can be made approximately invariant by extending the symmetry transformation (47) to fermions. One judicious PQ charge assignment is

$$d_{Ri} \rightarrow e^{i\delta} d_{Ri}, \quad Q_{Lj} \rightarrow Q_{Lj}, \quad u_{Rk} \rightarrow u_{Rk}, \quad (51)$$

which commutes with the SM gauge group, implying that neutral and charged gauge-boson couplings respect the symmetry. It has been previously used in [13] to classify the Higgs-fermion couplings in MFV. However, since for MFV one has one more small parameter $\kappa_{qb}/\kappa_{bq} \propto m_q/m_b$ for $q = s$ or d , it is useful to consider the following variant of Eq. (51):

$$b_R \rightarrow e^{i\delta} b_R, \quad q_R \rightarrow q_R, \quad Q_{Lj} \rightarrow Q_{Lj}, \quad u_{Rk} \rightarrow u_{Rk}. \quad (52)$$

Now $\kappa_{ij} \bar{d}_{Ri} Q_{Lj} \cdot H_d$ in Eq. (50) breaks the symmetry unless $d_{Ri} = d_{R3} = b_R$. However, all $U(1)_{\text{PQ}}$ breaking is still proportional to one of the small parameters of Eq. (36):

$\kappa_{qj} = \mathcal{O}(\omega)$ and $\tilde{\kappa}_{ij} = \mathcal{O}(l)$. The modified symmetry Eq. (52) forbids all operators in the weak Hamiltonian Eq. (4) (Table I), including the would-be leading one, Q_1^{SLL} , except for the standard-model operator Q_1^{VLL} and for $Q_{1,2}^{\text{SRR}}$. The last two are, however, forbidden by the original charge assignment in Eq. (51). Hence, the Wilson coefficients of these operators are suppressed by $\omega = m_q/m_b$ or by factors of loop-induced effective couplings, respectively.

At the tree-level (in the 2HDM), $\mathcal{F}^+$, which induces Q_2^{LR} , is multiplied by a factor κ_{qb}^* , which is a PQ-breaking coupling. On the other hand, $\mathcal{F}^-$, which induces Q_1^{SLL} , is multiplied by the unsuppressed factor κ_{qb}^2 . Hence, $\mathcal{F}^-$ must be proportional to PQ-breaking couplings in the Higgs potential (up to $1/\tan\beta$ -suppressed terms). This is also seen from the fact that in the infinite $\tan\beta$ limit, it is given by

$$\int d^4x \langle T(h_d^0(x) h_d^0(0)) \rangle,$$

which vanishes if the PQ symmetry is unbroken. Explicitly, in the large $\tan\beta$ limit one has:

$$\mathcal{F}^+ = \frac{2\lambda_2 M_A^2 + (\lambda_2 \lambda_5^r - |\lambda_7|^2) v^2}{\lambda_2 M_A^4 + (\lambda_2 \lambda_5^r - |\lambda_7|^2) v^2 M_A^2 - (\lambda_7^i \text{Im}(\lambda_5^* \lambda_7) + \frac{1}{4} \lambda_2 \lambda_5^{i2}) v^4} \simeq \frac{2}{M_A^2}, \quad (53)$$

$$\mathcal{F}^- = \frac{-(\lambda_2 \lambda_5^* - \lambda_7^{*2}) v^2}{\lambda_2 M_A^4 + (\lambda_2 \lambda_5^r - |\lambda_7|^2) v^2 M_A^2 - (\lambda_7^i \text{Im}(\lambda_5^* \lambda_7) + \frac{1}{4} \lambda_2 \lambda_5^{i2}) v^4} \simeq \frac{-(\lambda_2 \lambda_5^* - \lambda_7^{*2}) v^2}{\lambda_2 M_A^4}, \quad (54)$$

where the rightmost expressions hold up to higher orders of small couplings. For $\mathcal{F}^-$, this is identical to the sum of the two leading diagrams in a “mass-insertion approximation,” where the PQ-breaking contributions to the Higgs mass terms are treated as interactions [Fig. 2(f) and 2(g)].

At the loop level (in the 2HDM), up to doubly suppressed contributions one can employ the PQ-conserving parts of Eqs. (50) and (46), i.e. set $\lambda_5 = \lambda_6 = \lambda_7 = 0$, as well as ignore κ_{qb} and $\tilde{\kappa}_{ij}$. The matching onto the weak Hamiltonian can be organized according to one-light-particle-irreducible chirality amplitudes. There are three amplitudes:

$$\mathcal{A}_{RR} = \langle T(b_R(x_1) b_R(x_2) \bar{s}_L(x_3) \bar{s}_L(x_4)) \rangle, \quad (55)$$

$$\mathcal{A}_{RL} = \langle T(b_R(x_1) b_L(x_2) \bar{s}_L(x_3) \bar{s}_R(x_4)) \rangle, \quad (56)$$

$$\mathcal{A}_{VLL} = \langle T(b_L(x_1) b_L(x_2) \bar{s}_L(x_3) \bar{s}_L(x_4)) \rangle, \quad (57)$$

plus the parity conjugates of $\mathcal{A}_{RR}$ and $\mathcal{A}_{VLL}$. (We have omitted amplitudes that cannot match onto Lorentz-invariant local dimension-six operators.) Only $\mathcal{A}_{VLL}$ is invariant under $U(1)_{\text{PQ}}$ (both versions) and can be generated from a symmetric Lagrangian. It matches onto

the standard-model operator Q_1^{VLL} . There is a single diagram contributing, see Fig. 2(h). (Diagram (i) matches onto Q_1^{VRR} and would be allowed for the unmodified PQ assignment of Eq. (51).)

The present discussion could be extended to other processes, and to higher loop-orders, by systematically treating the PQ-breaking couplings as interactions and working to a fixed total order in the small parameters; in practice, at such higher precision, one might want to extend the effective 2HDM by higher-dimensional operators to account for v/M_{SUSY} corrections.

Finally, let us remark that because our choice of shift parameters v_u and v_d minimize the potential V in the potential of our effective theory and not necessarily the full effective potential, the one-point functions for the (shifted) Higgs fields $\langle 0|h_i|0 \rangle$ ($h_i = \phi_u, \phi_d, A^0$) will, in general, not vanish. Hence, also “tadpole” diagrams involving quark or Higgs loops would have to be considered at the outset [Fig. 2(e)]. That they cancel in $B - \bar{B}$ mixing in our approximation follows from the fact that no such diagrams are present when working with a complex h_d^0 field and the Lagrangian V_{lib} . Tadpoles may, however, be relevant in other contexts. We discuss our renormalization of v_u , v_d , and $\tan\beta$ in detail in the following section.

TABLE I. Charges of the operators in the weak Hamiltonian under the approximate $U(1)$ symmetry discussed in the text, see Eq. (52). The number in brackets denotes the charge under the “unmodified” charge assignment of Eq. (51).

Operator [field content]	$U(1)$ charge	Suppression of leading Higgs-mediated contribution	Remark
$Q_{1,2}^{\text{SL}}[\bar{b}_R q_L \bar{b}_R q_L]$	2	λ_5 (sparticle loop)	new
$Q_{1,2}^{\text{LR}}[\bar{b}_R q_L \bar{b}_L q_R]$	1 [0]	ω	known
$Q_1^{\text{VLL}}[\bar{b}_L q_L \bar{b}_L q_L]$	0	2HDM loop	SM operator
$Q_1^{\text{VRR}}[\bar{b}_R q_R \bar{b}_R q_R]$	2 [0]	$\omega^2 \times$ 2HDM loop	tiny
$Q_{1,2}^{\text{SRR}}[\bar{b}_L q_R \bar{b}_L q_R]$	0 [-2]	$\omega^2 \times$ sparticle loop	tiny

III. SYSTEMATICS OF THE LARGE- $\tan\beta$ MSSM

The present section is devoted to certain technical aspects of the large $\tan\beta$ limit. The first concerns the definition (i.e. renormalization) of $\tan\beta$ in the MSSM and in the effective two-Higgs-doublet-model description of low-energy (i.e., Higgs, electroweak, and flavor) phenomenology, and the matching between the two. This is of phenomenological importance, as $\tan\beta$ definitions used in the literature on the MSSM are known to differ by parametrically large expressions $\mathcal{O}(\tan\beta \times \text{loop factor})$. This can lead to ambiguities in the value of $\tan\beta$ of 10–15 in certain regions of the MSSM parameter space between schemes that have been extensively used in the study of radiative corrections to the MSSM Higgs sector [31]. Having clarified the connection between our “full” and “effective” $\tan\beta$, we justify the systematic expansion in $1/\tan\beta$ at the Lagrangian level employed in Sec. IIc.

A. Renormalization of $\tan\beta$

In the MSSM, $\tan\beta = v_u/v_d$ is defined as a ratio of vacuum expectation values. This is an unambiguous notion at tree-level, because a preferred basis is provided by the chiral Higgs supermultiplets of definite hypercharges $\pm 1/2$. Beyond tree-level, a scheme dependence arises as the bare parameters p_i^0 ($p_i = m_1^2, m_2^2, B\mu, g, g'$, etc.) are renormalized, $p_i^0 = p_i + \delta p_i$, as well as in the normalization of the fields and in defining renormalized shift parameters v_d, v_u . To formalize the renormalization program, we first define bare shifts that minimize the bare effective potential including radiative corrections, which is equivalent to requiring vanishing one-point functions for the shifted fields, i.e.,

$$\left\langle h_i^{0,\text{bare}} - \frac{1}{\sqrt{2}} v_i^0 \right\rangle \stackrel{!}{=} 0, \quad (58)$$

such that the v_i^0 are indeed vacuum expectation values. Identifying (for any definition of renormalized shift parameters)

$$v_i^0 = Z_i^{1/2}(v_i - \delta v_i), \quad i = d, u, \quad (59)$$

scheme dependence arises through, and only through, field renormalization and the counterterms δv_i . Reference [32] argued that for a stable perturbation expansion, it is desirable to define the renormalized v_i such as to mini-

mize the renormalized effective potential, i.e. $\delta v_i = 0$, and implemented this proposal for $\overline{\text{DR}}$ field renormalization and Landau gauge. The same condition and gauge fixing was imposed in the computation of one-loop corrections to the MSSM Higgs masses in [18–21]. References [22,23] chose to work with on-shell fields and in R_ξ gauge instead, and their shifts do not strictly minimize the one-loop effective potential. In fact, in general gauges, for $\delta v_i = 0$ the effective action is not finite and the v_i are both divergent [22,33] and gauge-dependent [33,34] (as are the bare vevs v_i^0).⁶ Hence, to have finite renormalized v_i and $\tan\beta$, $\delta v_i \neq 0$, containing a gauge-dependent divergence, is required. For $\tan\beta$, we have

$$\begin{aligned} \tan\beta^0 &\equiv \frac{v_u^0}{v_d^0} \stackrel{1\text{-loop}}{=} \tan\beta \left(1 + \frac{1}{2} \delta Z_u - \frac{1}{2} \delta Z_d - \frac{\delta v_u}{v_u} + \frac{\delta v_d}{v_d} \right) \\ &\equiv \tan\beta + \delta \tan\beta. \end{aligned} \quad (60)$$

Minimal subtraction for $Z_u, Z_d, \delta v_u/v_u, \delta v_d/v_d$ defines $\tan\beta^{\overline{\text{DR}}}$ [21]. It also follows from Eq. (60) that a change between two schemes R and R' can be calculated from

$$\tan\beta^R - \tan\beta^{R'} = \delta \tan\beta^{R'} - \delta \tan\beta^R,$$

hence any scheme where $\delta \tan\beta$ is a pure divergence has $\tan\beta = \tan\beta^{\overline{\text{DR}}}$ regardless of any nonminimal field renormalizations as those employed in [31]. In the latter case, however, $\delta v_u, \delta v_d$ are nonminimal and the counterterm for $\tan\beta$ has no simple relation to the field renormalization constants.

$\tan\beta^{\overline{\text{DR}}}$ is gauge-dependent [36], but to one-loop order, the gauge-dependence drops out for the R_ξ gauges. In spite of its gauge-dependence, the $\overline{\text{DR}}$ scheme for $\tan\beta$ has been shown to lead to a well-behaved perturbation expansion [31] and is also used in the most recent version of the

⁶This is, in particular, true in R_ξ gauges if $\xi \neq 0$. The apparent contradiction to the results in [35], whose authors are able to renormalize the effective action with purely “symmetric” counterterms, is resolved by noticing that in the Lorenz gauge employed in [35] the gauge-fixed Lagrangian still respects an invariance under constant (“global”) gauge transformations. This is sufficient to forbid divergences that cannot be removed by symmetric counterterms. Conversely, the R_ξ gauges break also this global invariance, for instance through Goldstone and ghost mass terms, which are indeed responsible for the “non-symmetric” divergences at one-loop [33]. The exception is the Landau gauge $\xi = 0$, which has the invariance.

publicly available computer programs FeynHiggs [37] and CPsuperH [38].

A second issue is that a fully minimal subtraction scheme, where in particular $\delta v_i^{\text{finite}} = 0$, generally entails v_i that do not minimize the (renormalized) tree potential, such that the renormalized Lagrangian contains linear terms

$$\mathcal{L} \supset t_d \phi_d + t_u \phi_u \quad (61)$$

for the shifted (real parts of the) Higgs fields. On the other hand, from Eq. (58) and (59) it follows that

$$\Gamma_i^{\text{ren}} = t_i + \Gamma_i^{(1)} + \delta t_i = 0 \quad (62)$$

always holds, if only δv_u and δv_d are included in δt_i . The presence of t_u, t_d is perfectly fine, but tadpole diagrams then have to be retained in the calculation. (In particular, they appear in the expressions relating Higgs and gauge-boson mass parameters to the Lagrangian parameters. If all renormalization constants are minimal, Eq. (62) determines t_i in terms of the bare proper one-point functions $\Gamma_i^{(1)}$ [21].) Yet it may be more convenient to perform additional finite renormalizations to work in a scheme where $t_i = 0$. This can be achieved either by suitable finite terms in δv_i or by finite renormalizations of the mass and coupling parameters. The former shifts $\tan\beta$ from its $\overline{\text{DR}}$ value according to

$$\tan\beta^{\text{tad}} = \tan\beta^{\overline{\text{DR}}} \left(1 - \frac{\delta v_d^{\text{tad}}}{v_d} + \frac{\delta v_u^{\text{tad}}}{v_u} \right). \quad (63)$$

The latter option does not modify $\tan\beta$.

Going from the MSSM to a general 2HDM, $\tan\beta$ becomes—strictly speaking—an ill-defined notion, as there is no preferred basis. Identifying $H_1 = -\epsilon H_d^*$ and $H_2 = H_u$, an $SU(2)$ rotation $H_i \rightarrow U_{ij} H_j$ removes the vacuum expectation value of one doublet; this corresponds to the (Φ, Φ') basis introduced in Sec. II. Only Φ receives a vev, provides for the Higgs mechanism, and has flavor-conserving couplings, while Φ' is an ordinary scalar with FCNC couplings. To make contact with MSSM phenomenology, however, it is useful to keep the notion of $\tan\beta$ in the effective theory. In principle, we could fix a basis to enforce $\tan\beta^{\text{EFT}} \equiv \tan\beta^{\overline{\text{DR}}}$, but find it technically simpler to allow for a parametrically small (i.e. not $\tan\beta$ -enhanced) shift, as we discuss in the following.

In complete analogy with the MSSM case discussed above, if we employ a general gauge and $\overline{\text{MS}}$ everywhere in the effective theory, v_1 and v_2 will not minimize the tree-level (nor the effective) potential.⁷ This would require a modification of the formalism in Sec. II. In particular, in writing the mass matrices Eqs. (23)–(25) and the flavor structure of the scalar-fermion couplings in Eq. (11) we assumed the minimization conditions $t_1 = t_2 = 0$. To

avoid such modifications, as well as changed expressions for neutral meson mixing, we can either perform renormalizations on the parameters m_{11}^2 and m_{22}^2 such that v_1 and v_2 minimize the 2HDM potential, or achieve this through nonminimal $\delta v_{1,2}$. We pursue the latter option, keeping the symmetric parameters of the 2HDM minimally subtracted. This has the added virtue that the $\tan\beta$ such defined is gauge-independent at the order considered, as it is fully determined by $\overline{\text{MS}}$ mass and coupling parameters. These are gauge-invariant at one-loop, which is clear from our explicit matching calculation. We presume this to hold also at higher orders, at least if the appropriate wave-function renormalization is employed. The δv_i are determined entirely in terms of “light”-particle loops and, at least at one-loop, do not lead to parametrically large shifts $\propto \tan^2\beta \frac{1}{16\pi^2}$, as can be verified from the explicit expressions for the tadpoles in [23] or by considering tadpole diagrams in the large- $\tan\beta$ effective Lagrangian.

To find the precise connection between $\tan\beta^{\overline{\text{DR}}}$ and our effective $\tan\beta$, consider the total tree plus one-loop contribution of the superpartners to the (MSSM) effective action for the gauge and Higgs fields,

$$\begin{aligned} S_{gh} = \int d^4x & \left[(1 + \Delta Z_W) \left(-\frac{1}{4} \right) W_{\mu\nu}^A W^{\mu\nu A} + (1 + \Delta Z_B) \right. \\ & \times \left(-\frac{1}{4} \right) B_{\mu\nu} B^{\mu\nu} + (\delta_{ij} + \Delta Z_{ij}) (D_\mu H_i)^\dagger (D^\mu H_j) \\ & \left. - \hat{m}_{ij}^2 H_i^\dagger H_j - \sum_{k=1}^7 \hat{\lambda}_k O_k + \dots \right]. \quad (64) \end{aligned}$$

Here, O_i are the quartic terms constructed from the Higgs fields appearing in Eq. (12), and the dots denote higher-dimensional local terms, and H_1, H_2 are related to H_d, H_u as above. The precise values for the coefficients depend on the MSSM renormalization scheme. We assume the MSSM has been regularized by dimensional reduction while the couplings, Higgs fields and $\tan\beta$ are minimally subtracted ($\overline{\text{DR}}$). The corresponding expressions $\hat{\lambda}_k$ are reported in Eq. (21) (tree-level) and in Appendix B2 and 3 (one-loop).

Equation (64) can be identified with the classical action (ignoring 2HDM loops) for an effective two-Higgs-doublet model with noncanonically normalized fields. To obtain from this the $\overline{\text{MS}}$ -renormalized Lagrangian in the presence of light-particle loops, one simply has to add the contributions (which are local) due to loops of 2ϵ scalars present in DRED⁸ and subsequently rescale the fields,

$$\begin{pmatrix} -\epsilon H_d^{\overline{\text{DR}*}} \\ H_u^{\overline{\text{DR}}} \end{pmatrix} = \begin{pmatrix} Z_{dd} & Z_{du} \\ Z_{ud} & Z_{uu} \end{pmatrix} \begin{pmatrix} H_1^{\overline{\text{eff}}} \\ H_2^{\overline{\text{eff}}} \end{pmatrix}, \quad (65)$$

⁷In this section, the renormalized 2HDM vevs are denoted by $v_{1,2}$ instead of $v_{d,u}$ in Sec. II in order to avoid confusion with the renormalized MSSM vevs. Correspondingly, the 2HDM tadpoles are denoted by $t_{1,2}$.

⁸Integrating over the 2ϵ scalars leaves a path integral over light fields that is identical to that in the DREG-regularized effective theory, including the $1/\epsilon$ divergence structure. We recall that the 2ϵ scalars should be thought of as having a nonzero mass of $\mathcal{O}(M_{\text{SUSY}})$ [39].

subject to the condition $Z^\dagger(1 + \Delta Z)Z = \mathbf{1}$. This provides the relation between the $\overline{\text{DR}}$ fields of the MSSM and one out of an infinite choice of $\overline{\text{MS}}$ fields in the effective theory, labeled “eff”. We fix the freedom to choose the Higgs basis in the effective theory by setting $Z_{du} = 0$ and $Z_{uu}^i = Z_{dd}^i = 0$.⁹ The relation between the shifts and $\tan\beta$ of the MSSM and of the 2HDM are now determined according to

$$\begin{aligned}\bar{v}_2(\mu) &\equiv v_2(\mu)^{\text{eff}} = v_u^{\overline{\text{DR}}} - \delta Z_{ud}v_d - \delta Z_{uu}v_u + \delta v_2^{\text{tad}} \\ \bar{v}_1(\mu) &\equiv v_1(\mu)^{\text{eff}} = v_d^{\overline{\text{DR}}} - \delta Z_{dd}v_d + \delta v_1^{\text{tad}} \\ \tan\beta(\mu)^{\text{eff}} &= \tan\beta^{\overline{\text{DR}}}\left(1 - \frac{\delta v_1^{\text{tad}}}{v_1} + \frac{\delta v_2^{\text{tad}}}{v_2} + \delta Z_{dd} \right. \\ &\quad \left. - \delta Z_{uu} - \delta Z_{ud}\cot\beta\right).\end{aligned}\quad (66)$$

Here, we have expanded $Z_{uu/dd} = \mathbf{1} + \delta Z_{uu/dd}$ and $Z_{ud} = \delta Z_{ud}$, and the δZ_{ij} are related to the ΔZ_{ij} via $\Delta Z_{11/22} = -2\delta Z_{uu/dd}$ and $\Delta Z_{12} = -\delta Z_{ud}^*$, with the explicit expressions given in Appendix B1. The shifts $\delta v_{1,2}^{\text{tad}}$ are defined implicitly as discussed above. In summary, we have constructed a $\tan\beta$ which is appropriate for effective weak interactions, gauge-independent and, up to an ordinary (i.e., not $\tan\beta$ -enhanced) loop correction, coincides with the widely used $\tan\beta^{\overline{\text{DR}}}$. It means that the $\tan\beta$ measured in flavor physics, for instance through $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$, and employed in our analysis, can be identified with the corresponding $\overline{\text{DR}}$ parameter at large $\tan\beta$, up to small corrections.

We note that our framework leads to a transparent expression for the relation between the $\overline{\text{DR}}$ scheme and the so-called DCPR scheme employed in [22,23] in the limit $v \ll M_{\text{SUSY}}$. In the latter scheme, finite but, unlike in our effective 2HDM, “diagonal” wave-function renormalizations of H_u, H_d are performed, i.e., in our notation, $\delta Z_{ud} = \delta Z_{du} = 0$, as well as $\delta Z_{uu/dd} = 2\delta Z_{u/d}^{\text{finite}}$. Moreover, the renormalization conditions include

$$\frac{\delta v_u}{v_u} = \frac{\delta v_d}{v_d}, \quad \text{Re}\Sigma_{A^0 Z^0}(M_A^2) = 0, \quad (67)$$

where $\Sigma_{A^0 Z^0}(k^2)$ parameterizes the A^0 - Z^0 mixing according to $\Sigma_{A^0 Z^0}^\mu(k) = k^\mu \Sigma_{A^0 Z^0}(k^2)$. Now, the sparticle contribution to $\Sigma_{A^0 Z^0}(k^2)$ reads

$$\begin{aligned}\Sigma_{A^0 Z^0}(k^2) &= M_Z \sin^2 \beta \text{Re}\Delta Z_{12} \\ &\quad + M_Z \sin\beta \cos\beta (\delta Z_d^{\text{finite}} - \delta Z_u^{\text{finite}}) + \dots,\end{aligned}\quad (68)$$

where the dots denote terms proportional to $\cos\beta$ but not involving the wave-function renormalization constants, or terms suppressed in the limit where v and $k \ll M_{\text{SUSY}}$. This follows either by considering the mixed gauge-boson-Higgs-boson bilinear terms resulting from the covariant kinetic operator for the Higgs fields in Eq. (64), or via the Ward identity

$$k_\mu \Sigma_{A^0 Z^0}^\mu(k^2) + M_Z \Sigma_{G^0 A^0}(k^2) = \mathcal{O}(k^2 - M_A^2) \quad (69)$$

⁹As before, superscripts “r” and “i” denote real and imaginary parts.

(which is trivially satisfied in our $SU(2)$ -invariant formalism) from the terms bilinear in the gauge fields in the same term. The two conditions in Eq. (67) then determine

$$\begin{aligned}(\delta \tan\beta^{\text{DCPR}})^{\text{finite}} &= \frac{\tan\beta}{2} (\delta Z_u^{\text{finite}} - \delta Z_d^{\text{finite}}) \\ &= \frac{\tan^2 \beta}{2} \text{Re}\Delta Z_{12} + \dots,\end{aligned}\quad (70)$$

where the omitted terms are not $\tan\beta$ -enhanced. This explains the large numerical differences between $\tan\beta^{\overline{\text{DR}}}$ and $\tan\beta^{\text{DCPR}}$ found in [31] as a parametrically large effect. Hence, $\tan\beta$ measured in flavor physics should not be identified with the corresponding DCPR parameter at large $\tan\beta$.¹⁰

As with the Higgs fields, we explicitly decouple the contributions of heavy particles to the gauge field wave

¹⁰The renormalization of $\tan\beta$ in the $M_{\text{SUSY}} \gg v$ limit was recently discussed in [40], where a different conclusion was reached, namely, that the shift between the $\overline{\text{DR}}$ and DCPR schemes is not $\tan\beta$ -enhanced. As we do, they introduce separate sets of renormalized Higgs fields in the MSSM and in an effective 2HDM and proceed to assert that the 2HDM field renormalization does not contribute $\tan\beta$ -enhanced terms in Higgs-mediated processes. Next, within their effective 2HDM, the authors write (with left-hand sides corresponding to our notation)

$$\begin{aligned}\frac{1}{\sqrt{2}}v_i^0 &= v_i + \Delta v_i, \\ h_d^{0,\text{bare}} &= v_d + \Delta v_d + \left(1 - \frac{z_{11}}{2}\right)H_d^0 + \frac{1+a}{2}z_{12}H_u^{0*}, \\ h_u^{0,\text{bare}} &= v_u + \Delta v_u + \left(1 - \frac{z_{22}}{2}\right)H_u^0 + \frac{1-a}{2}z_{12}H_d^{0*}, \\ \tan\beta^0 &= \tan\beta + \frac{\Delta v_u}{v_u} - \frac{\Delta v_d}{v_d}.\end{aligned}$$

the fields H_u^0, H_d^0 have vanishing vev. Δv_i can be identified with $1/\sqrt{2}(\delta Z_{ii}v_i - \delta v_i)$, while z_{ij} is equivalent to ΔZ_{ij} , in our notation. Differently to our treatment, wave-function renormalizations are only performed on the shifted fields. For the latter, they correspond to our 2HDM field basis if $a = -1$. The above field redefinitions then contribute a counterterm mixing A^0 and Z^0 , which is adjusted to remove the loop-induced A^0 - Z^0 mixing. (An $A_0 Z_0$ counterterm could also be generated from Δv_i but vanishes if v_i minimize the renormalized tree-level potential, as assumed both here and in [40].) However, off-diagonal renormalizations are not present in a manifestly supersymmetric renormalization scheme such as DCPR, hence the renormalized A^0 field in the 2HDM of [40] cannot be identified with the DCPR A^0 field for any value of a or Δv_i , even in the large- $\tan\beta$ limit. If MSSM wave-function renormalization is taken into account, its effect on v_i could be absorbed into Δv_i , with DCPR then requiring $\Delta v_u/v_u - \Delta v_d/v_d = 1/2(\delta Z_u - \delta Z_d)$. Our discussion above shows that this expression is $\tan\beta$ -enhanced when the other DCPR requirement – vanishing $A^0 - Z^0$ mixing – is enforced. One should remark in this context that one can be misled by an erroneous zero in Eq. (A.8) in [23], which also disagrees with [22]. We acknowledge and enjoyed extensive discussions with M. Beneke on this issue.

functions (hence to $g^{(l)}$) by a finite renormalization $g^{(l)\overline{\text{DR}}} = g^{(l)\text{eff}} + g^{(l)\text{eff}} \delta g^{(l)}$, cancelling the terms ΔZ_B and ΔZ_W in Eq. (64) of the gauge fields, $B_\mu^{\overline{\text{DR}}} = Z_B^{1/2} B_\mu^{\text{eff}}$ and $W_\mu^{\overline{\text{DR}}} = Z_W^{1/2} W_\mu^{\text{eff}}$. For $\overline{\text{DR}}$ -subtracted MSSM couplings, this gives $\overline{\text{MS}}$ -renormalized 2HDM gauge couplings.

We denote the quartic couplings in our 2HDM scheme by $\bar{\lambda}_i$. The finite renormalizations leave λ_5 invariant, $\bar{\lambda}_5 = \hat{\lambda}_5$, while the other quartic coupling constants transform like

$$\begin{aligned}\bar{\lambda}_1 &= \hat{\lambda}_1 + \tilde{g}^2 \delta Z_{dd}^r + \frac{1}{2}(g^2 \delta g + g'^2 \delta g'), \\ \bar{\lambda}_2 &= \hat{\lambda}_2 + \tilde{g}^2 \delta Z_{uu}^r + \frac{1}{2}(g^2 \delta g + g'^2 \delta g'), \\ \bar{\lambda}_3 &= \hat{\lambda}_3 - \frac{\tilde{g}^2}{2}(\delta Z_{dd}^r + \delta Z_{uu}^r) - \frac{1}{2}(g^2 \delta g + g'^2 \delta g'), \\ \bar{\lambda}_4 &= \hat{\lambda}_4 + g^2(\delta Z_{dd}^r + \delta Z_{uu}^r) + g^2 \delta g, \\ \bar{\lambda}_6 &= \hat{\lambda}_6 - \frac{\tilde{g}^2}{4} \delta Z_{ud}^*, \quad \bar{\lambda}_7 = \hat{\lambda}_7 + \frac{\tilde{g}^2}{4} \delta Z_{ud}^*,\end{aligned}\quad (71)$$

where x^r and x^i denote the real and imaginary part of x , respectively. The couplings $\bar{\lambda}_i$ and $g^{(l)} = g^{(l)\text{eff}}$ are $\overline{\text{MS}}$ couplings from the viewpoint of the effective theory.

The modification of the dimensionless couplings by the finite wave-function renormalizations affects the $B - \bar{B}$ mixing amplitudes as a formally higher-order effect, as does the scheme dependence of $\tan\beta$. Unlike the latter, however, the former is never $\tan\beta$ enhanced unless the wave-function-renormalization constants themselves are.

Invariance of $B - \bar{B}$ mixing under field renormalization

The effects of an arbitrary linear field redefinition in the 2HDM such as Eq. (65) on the Higgs-mediated FCNC Eq. (11) are twofold: (i) the values for $\cos\beta$ and $\sin\beta$ in Eq. (10) are modified. This cancels the contributions to $\mathcal{F}^\pm$ from the redefinition of the mass matrices up to a global factor:

$$\begin{aligned}\mathcal{F}^+(\lambda_i, v_{u,d}, M_A) &\rightarrow \mathcal{F}^+(\lambda'_i, v'_{u,d}, M'_A) \\ &= \frac{v^2}{v'^2} |\det Z|^2 \mathcal{F}^+(\lambda_i, v_{u,d}, M_A),\end{aligned}\quad (72)$$

$$\begin{aligned}\mathcal{F}^-(\lambda_i, v_{u,d}, M_A) &\rightarrow \mathcal{F}^-(\lambda'_i, v'_{u,d}, M'_A) \\ &= \frac{v^2}{v'^2} (\det Z^*)^2 \mathcal{F}^-(\lambda_i, v_{u,d}, M_A).\end{aligned}\quad (73)$$

(ii) the $v_{u,d}$ redefinition in (i) comes with a modification of κ_{ij} :

$$\kappa_{ij} \rightarrow \kappa'_{ij} = \frac{v'}{v} (\det Z^*)^{-1} \kappa_{ij}.\quad (74)$$

The above factors cancel each other out in the products $\kappa_{bq}^2 \mathcal{F}^-$ and $\kappa_{qb}^* \kappa_{bq} \mathcal{F}^+$, as they should. In particular, our

choice of wave-function renormalization acting on the leading FCNC coupling Eq. (29) produces an extra term:

$$\delta \kappa_{bq} = -\kappa_{bq}(s_\beta^2 \delta Z_{dd}^r - s_\beta c_\beta \delta Z_{ud}^r + c_\beta^2 \delta Z_{uu}^r). \quad (75)$$

Considering Eqs. (29), (30), (71), and (75), gives the same Wilson coefficients C_2^{LR} and C_1^{SLL} as does considering Eqs. (29), (30), and (71), with the finite parts of δZ_{ij} set to zero. While in practice, wave-function renormalization has to be performed to relate the parameter M_A to the physical Higgs-boson masses and to take $v = (\sqrt{2}G_F)^{-1/2} \simeq 246$ GeV beyond leading-order precision, such renormalizations are not the source of a nonvanishing of the Q_1^{SLL} amplitude, to be found instead in the corrections to Higgs masses and mixings (via the self-couplings λ_i , in particular λ_5); wave-function-renormalization effects enter that amplitude only at higher orders (as might have been expected). In this regard our findings disagree with the conclusions of [26].

B. Health of the large- $\tan\beta$ limit and fine-tuning

In Sec. IIc we took the limit $\tan\beta \rightarrow \infty$ ($v_d \rightarrow 0$, $M_A^2 = \text{const}$, $v_u^2 + v_d^2 = \text{const}$, $\lambda_i = \text{const}$, v_u and v_d defined as minima of the tree potential) at the Lagrangian level. One might wonder whether this procedure is valid at the quantum level. To justify it, we show that the $v_d = 0$ case and the $v_d \neq 0$ case are analytically connected, i.e. one can be reached from the other without a phase transition. It then follows that amplitudes are (in some neighborhood of a parameter point with $v_d = 0$) analytic functions of the parameters (either symmetric or broken). The renormalizability of the effective potential V_{lb} then follows by standard arguments from the fact that it is equivalent to the symmetric potential Eq. (12) (for a certain choice of parameters), which is renormalizable.

We first note that the number of independent minimization conditions is unchanged in the $v_d = 0$ limit. First, for general values of the parameters, out of the four real (two complex) minimization conditions, at most three are independent. This follows from the $U(1)_Y$ invariance but is easy to verify explicitly. Fixing v_u to be real and positive, three polynomials of degree three determining three unknowns v_u , v_d^r , v_d^i remain. The system has a solution $v_d = 0$ if

$$\lambda_2 m_{12}^2 + \lambda_7 m_{22}^2 = 0, \quad v_u^2 = -\frac{2m_{22}^2}{\lambda_2}.\quad (76)$$

Here, the second equation determines $v_u = v$ as a function of m_{22}^2 and λ_2 similarly to the case of a single doublet, while the first equation can be viewed as a fine-tuning condition between m_{12}^2 and λ_7 . The dimensionless, complex parameter

$$\epsilon = \frac{m_{12}^2}{m_{11}^2} + \frac{\lambda_7}{\lambda_2} \frac{m_{22}^2}{m_{11}^2}\quad (77)$$

parameterizes the deviation from the fine-tuning limit; we may trade m_{12}^2 in favor of ϵ . Clearly, at the limiting point

$\epsilon = 0$, we indeed have three independent equations. Now, it is easy to verify that, writing the four real minimization conditions in the form

$$f_i(\lambda_i, m_{ij}^2, v_u, v_d) = 0, \quad (78)$$

the Jacobian matrix

$$\frac{\partial(f_1, f_2, f_3, f_4)}{\partial(v_u, v_d^r, v_d^i)} \quad (79)$$

has maximal rank (3) at any point with $v_d = 0$. (Physically, this just means that the neutral-Higgs mass matrix has three nonzero eigenvalues there.) Hence, by the implicit function theorem, we may solve for (v_u, v_d) in a neighborhood of it, where the solutions will be (real-)analytic functions of ϵ . In particular, v_u behaves analytically (and is strictly positive) around $\epsilon = 0$, i.e. no phase boundary is encountered. Explicitly and to linear order, the real and imaginary parts of v_d are determined by

$$\begin{pmatrix} 1 + \frac{\lambda_3 + \lambda_5^r}{2} \frac{v_u^2}{m_{11}^2} & \frac{\lambda_5^i}{2} \frac{v_u^2}{m_{11}^2} \\ \frac{\lambda_5^i}{2} \frac{v_u^2}{m_{11}^2} & 1 + \frac{\lambda_3 - \lambda_5^r}{2} \frac{v_u^2}{m_{11}^2} \end{pmatrix} \begin{pmatrix} v_d^r \\ v_d^i \end{pmatrix} = v_u \begin{pmatrix} \epsilon^r \\ \epsilon^i \end{pmatrix} + \mathcal{O}(\epsilon^2), \quad (80)$$

such that $\tan\beta = \mathcal{O}(1/|\epsilon|)$. The nonsingular linear term allows us to change variables from m_{12}^2 to a complex v_d . Of course, we may always perform a field redefinition of H_d such that v_d is real. Then, the mass parameters besides m_{11}^2 are power series in $1/\tan\beta$, which read

$$\begin{aligned} m_{12}^{2,i} &= \frac{1}{2} \lambda_6^i v_d^2 + \frac{1}{2} v_u \lambda_5^i v_d + \frac{1}{2} v_u^2 \lambda_7^i, \\ m_{22}^2 &= \frac{v_d m_{11}^2}{v_u} + \frac{1}{2} v_u v_d (\lambda_3 + \lambda_5^r) + \frac{1}{2} v_u^2 \lambda_7^r + \mathcal{O}(v_d^2/v_u^2), \\ M_A^2 &= m_{11}^2 + \frac{\lambda_3 - \lambda_5^r}{2} v_u^2 + \mathcal{O}(v_d/v_u), \\ M_h^2 &= \lambda_2 v^2 + \mathcal{O}(\lambda_7^2; \epsilon), \quad M_H^2 = m_{11}^2 + \frac{\lambda_3 + \lambda_5^r}{2} v^2 + \mathcal{O}(\lambda_7^2; \epsilon), \\ M_{H^\pm}^2 &= m_{11}^2 + \frac{\lambda_3 + \lambda_4}{2} v^2 + \mathcal{O}(v_d/v_u). \end{aligned} \quad (81)$$

We see explicitly that we can continuously change the dimensionful parameters in the Higgs potential from a situation where $v_d \neq 0$ to one where $v_d = 0$, keeping M_A^2 (and the dimensionless couplings) fixed, as was assumed in Sec. IIc. The last three equations illustrate that the large- $\tan\beta$ case is characterized by a “primary” doublet H_u which receives a large vev v_u and a “secondary” doublet H_d with a positive gauge-invariant mass m_{11}^2 that receives corrections of $\mathcal{O}(v^2)$ and $\mathcal{O}(\epsilon)$, respectively, due to its dimensionless and dimensionful couplings to H_u . Those corrections differ among the physical components of the doublet, approximately to be identified with H^0 , A^0 , $H^\pm$, due to electroweak symmetry-breaking. In principle, m_{11}^2 could be negative, but in that case, $v_d \approx 0$ will typically not be the global minimum of the potential.

We close this section by considering the fine-tuning which is necessary to obtain a large $\tan\beta$ while keeping the mass M_A fixed.¹¹ For v_d real, Eq. (80) implies

$$m_{12}^{2,r} = -\frac{\lambda_7^r}{\lambda_2} m_{22}^2 + \cot\beta \left(m_{11}^2 + \frac{\lambda_3 + \lambda_5^r}{2} v_u^2 \right), \quad (82)$$

which illustrates the tuning that is known to be necessary to have large $\tan\beta$ in the MSSM. For the generic situation $m_{11}^2 \sim M_{\text{SUSY}}^2 \gg M_Z^2$, the right-hand side is dominated by the m_{11}^2 term: λ_7^r is down by a loop factor relative to λ_2 , and $m_{22}^2 \sim v^2 \ll M_{\text{SUSY}}^2$ (the little hierarchy). Hence,

$$m_{12}^{2,r}/M_{\text{SUSY}}^2 \sim 1/\tan\beta, \quad (83)$$

which implies an extra tuning beyond the one to achieve the correct weak scale. For smaller $m_{11}^2 \sim M_A^2 \sim M_Z^2$, which is interesting from the point of view of B -physics phenomenology, the required tuning gets even worse—unless, of course, the whole SUSY scale is lowered to the weak scale, which is, however, problematic since then $M_h^2 \approx \lambda_2 v^2$ is generally below the experimental lower limit. On the other hand, as we have seen, $M_A^2 \sim m_{11}^2$, such that no extra tuning is required to keep M_A^2 finite, while one might have expected otherwise from the well-known tree-level formula

$$M_A^2 = (\tan\beta + \cot\beta) m_{12}^2, \quad (84)$$

which is generalized by Eq. (25). Also, while a small m_{12}^2 is indeed sensitive to radiative corrections, those are automatically correlated with shifts of v_d and in consequence of $\tan\beta$ in such a way that M_A^2 receives only mild corrections.

IV. PHENOMENOLOGY

In Sec. II, we performed a detailed study of the supersymmetric contributions to ΔM_d and ΔM_s in the generic framework of an effective 2HDM. The corresponding matching coefficients were computed at the one-loop level in Sec. III and Appendix B. In this section, we assess the maximal size of the various types of effects identified in the MFV case taking into account the existing constraints on the supersymmetric parameter space, in particular, from the $B_s \rightarrow \mu^+ \mu^-$, $B^+ \rightarrow \tau^+ \nu$, and $b \rightarrow s \gamma$ branching fractions. For convenience, we start with a compendium of the formulas derived in Sec. II:

$$\begin{aligned} \Delta M_q &= |\Delta M_q^{\text{SM}} + \Delta M_q^{\text{LR}} + \Delta M_q^{\text{LL}} + \Delta M_q^{\text{HL}}| \\ &\equiv |1 + h_q| \Delta M_q^{\text{SM}}, \end{aligned} \quad (85)$$

where the standard-model, the left-right and left-left Higgs-pole (cf. Equation (17)), and the neutral-Higgs-loop (cf. Equation (40)) contributions read

¹¹This aspect has been considered before, and recently in an EFT framework in Ref. [40].

TABLE II. Input values. Here M_{SUSY} stands for any of the supersymmetric parameters $|\mu|$, $M_{\tilde{L}}$, $M_{\tilde{R}}$, $M_{\tilde{B}}$, $M_{\tilde{W}}$, $M_{\tilde{Z}}$, $|a_t|$, $|a_b|$, $|a_\tau|$, M_1 , M_2 , M_3 . The renormalized parameters M_A and $\tan\beta$ are identical in the MSSM and effective 2HDM for $M_{\text{SUSY}} \gg v$, in the scheme introduced in Sec. IIIa. The quark masses and α_s are defined in the $\overline{\text{MS}}$ scheme at the scale m_t . The bag factors, defined at the scale m_t as well, are discussed in Appendix C.

Quark masses and α_s	Bag factors	SUSY parameters
$m_t = 164 \text{ GeV}$	$P_1^{\text{VLL}} \in [0.66, 0.76]$	$\tan\beta \in [10, 60]$
$m_b = 2.75 \text{ GeV}$	$P_2^{\text{LR}} m_s \in [0.12, 0.22] \text{ GeV}$	$M_A \in [120, 600 \text{ GeV}]$
$m_d/m_s = 1/19$	$P_1^{\text{SLL}} \in [-1.48, -1.24]$	$M_{\text{SUSY}} \in [600, 1800 \text{ GeV}]$
$\alpha_s = 0.108$		

$$\begin{aligned}
\Delta M_q^{\text{SM}} &= |V_{tq} V_{tb}^*|^2 f_{B_q}^2 M_{B_q} P_1^{\text{VLL}} \left[\frac{G_F^2 M_W^2}{6\pi^2} S_0(m_t^2/M_W^2) \right], & h_s &= \left\{ -2.40 \left[\frac{m_s/m_b}{0.053/2.75} \right] \left[\frac{P_2^{\text{LR}}/P_1^{\text{VLL}}}{3.2/0.71} \right] \right. \\
\Delta M_q^{\text{LR}} &= |V_{tq} V_{tb}^*|^2 f_{B_q}^2 M_{B_q} P_2^{\text{LR}} \left[\frac{-1}{3} \frac{m_b m_q}{v^2} |\bar{\kappa}|^2 \mathcal{F}^+ \right], & & + 0.35 \frac{16\pi^2 (-\lambda_5^* + \lambda_7^{*2}/\lambda_2) (120 \text{ GeV})^2 e^{2i\phi_{\bar{\kappa}}}}{M_A^2} \\
\Delta M_q^{\text{LL}} &= |V_{tq} V_{tb}^*|^2 f_{B_q}^2 M_{B_q} P_1^{\text{SLL}} \left[\frac{-1}{6} \frac{m_b^2}{v^2} \bar{\kappa}^2 \mathcal{F}^- \right], & & \times \left[\frac{P_1^{\text{SLL}}/P_1^{\text{VLL}}}{-1.36/0.71} \right] + 0.01 \frac{e^{2i\phi_{\bar{\kappa}}}}{(1 + \tilde{\epsilon}_3 t_\beta)^2} \left[\frac{t_\beta}{40} \right]^2 \left[\frac{m_b}{2.75} \right]^2 \Big\} \\
\Delta M_q^{\text{HL}} &= |V_{tq} V_{tb}^*|^2 f_{B_q}^2 M_{B_q} P_1^{\text{VLL}} \left[\frac{1}{12} \frac{y_b^{*2}}{16\pi^2} \frac{m_b^2}{v^2} \bar{\kappa}^2 \frac{1}{M_A^2} \right], & & \times \frac{|16\pi^2 \epsilon_Y|^2 (120 \text{ GeV})^2}{|1 + \tilde{\epsilon}_3 t_\beta|^2 |1 + \epsilon_0 t_\beta|^2 M_A^2} \left[\frac{t_\beta}{40} \right]^4 \left[\frac{m_b}{2.75} \right]^2, \quad (89)
\end{aligned}$$

respectively. The Inami-Lim function S_0 is given by $S_0(x) = (x - 11x^2/4 + x^3/4)(1-x)^{-2} - (3x^3 \log(x)/2) \times (1-x)^{-3}$ and $v = (\sqrt{2}G_F)^{-1/2} = 246 \text{ GeV}$. The flavor-changing and flavor-conserving quark-Higgs couplings were defined in Secs. II A and II B:

$$\begin{aligned}
\bar{\kappa} &\equiv \frac{\kappa_{bq} v}{\lambda_{qb} m_b} = \frac{\kappa_{qb} v}{\lambda_{qb}^* m_q} = \frac{y_t^2 \sqrt{2}}{c_\beta^2} \frac{\epsilon_Y}{(1 + \tilde{\epsilon}_3 t_\beta)(1 + \epsilon_0 t_\beta)}, \\
y_b &= \frac{\sqrt{2} m_b}{v c_\beta} \frac{1}{1 + \tilde{\epsilon}_3 t_\beta}, \quad (87)
\end{aligned}$$

with $\epsilon_{0,Y}$ and $\tilde{\epsilon}_3$ given in Eqs. (31)–(33) and y_t in Appendix A. The $\mathcal{F}^\pm$ factors describing the propagation of the neutral Higgses were defined in Eqs. (18) and (19), with the effective couplings λ_i entering the neutral-Higgs mass matrix computed in Sec. IIIa and Appendix B. For large $\tan\beta$, we have in very good approximation:

$$\mathcal{F}^+ \simeq \frac{2}{M_A^2}, \quad \mathcal{F}^- \simeq \frac{(-\lambda_5^* + \lambda_7^{*2}/\lambda_2) v^2}{M_A^4} \quad (88)$$

(exact formulas were used in our numerical analysis though). For simplicity, we have omitted the Higgs wavefunction-renormalization terms in Eqs. (87) and (88), which drop out in physical observables, as discussed at the end of Sec. IIIa. Explicit expressions for λ_5 , λ_7 , and λ_2 in the MFV case were given in Eqs. (39) and (37). Altogether, counting ϵ_Y , $\mathcal{F}^- \sim (16\pi^2)^{-1}$ and $M_A \sim 120 \text{ GeV}$ to get an idea of the naive size of the various effects in the absence of constraints, we obtain:

where m_b is in GeV and $\phi_{\bar{\kappa}} \equiv \arg(\bar{\kappa})$. h_d is given by the same expression with m_s replaced by m_d , so that the first term becomes subleading.

A first obvious remark is that ΔM_q^{HL} cannot compete with ΔM_q^{LR} or ΔM_q^{LL} unless y_b becomes nonperturbative. This is rather accidental (notice the small loop factor in Eq. (86) as well as the smallness of P_1^{VLL} with respect to P_2^{LR} and P_1^{SLL}). Further, the contribution of ΔM_q^{LL} seems somewhat limited. However, the loop functions λ_5 and λ_7 could be enhanced for large μ or $a_{t,b}$, see Eqs. (38) and (37). A more quantitative analysis is thus desirable. In the next two sections, we perform a random scan of the MFV-MSSM parameter space to find the maximal ΔM_q^{LL} and ΔM_q^{LR} values allowed by current experimental data. Equations (85)–(89) do allow for new CP -violating phases,¹² yet these will be set to zero in the scan. CP -violating effects within the MFV scenario will be shortly discussed in Sec. IVc.

A. Scan of the parameter space

The values of the various input parameters used in the scan are collected in Table II. Note that only the products $P_2^{\text{LR}} m_s$ and $P_2^{\text{LR}} m_d$, or alternatively $P_2^{\text{LR}} m_s$ and m_d/m_s , are needed, see Eq. (86). We scan over $P_2^{\text{LR}} m_s$ but keep m_d/m_s fixed as ΔM_d^{LR} is doomed to be small anyway. The decay constants f_{B_q} and CKM factors $|V_{tq} V_{tb}^*|$ are not specified.

¹²Let us recall that in that case M_A^2 , defined as the nonzero eigenvalue of the CP -odd mass matrix M_J^2 in Eq. (14), is no more an eigenvalue of the full Higgs mass matrix.

Instead, outputs are formulated in terms of ratios free from these rather poorly known parameters. Finally, we take $\alpha = 1/127.9$, $\sin^2\theta_W = 0.231$, and $M_Z = 91.1876$ GeV.

For simplicity, the gaugino mass parameters are assumed to have the same sign (which we can choose positive), as well as the trilinear terms (positive or negative). Note that the absolute scale of M_{SUSY} plays no role as supersymmetric parameters enter $\epsilon_{0,Y}$ and λ_i by means of ratios. Only the spread of the interval chosen for M_{SUSY} matters. Still, M_{SUSY} should not be taken too large to help satisfy the $b \rightarrow s\gamma$ constraint in the case $\mu < 0$. We will come back to this point later. We allow for rather large values of M_A , close to the lower end of the interval chosen for M_{SUSY} . Still, the matching performed in Sec. III and Appendix B remains valid as the corrections from higher-dimension operators at the electroweak scale are ruled by the ratio v/M_{SUSY} and not M_A/M_{SUSY} . The formulas for the various observables at the B mass scale are thus unaffected.

The constraints imposed on the points generated inside the above ranges are summarized in Table III. We now discuss them in turn:

- (i) The bottom Yukawa coupling y_b is maintained small enough, say, $y_b < 2$, to guarantee the validity of perturbation theory. This condition removes possible fine-tuned points in parameter space for which the denominators in Eq. (87) are close to zero.
- (ii) The lightest Higgs-boson mass M_h has to come out large enough to comply with the LEP II experimental lower bound. M_h is obtained from the CP -even Higgs mass matrix in Eq. (23), with the effective couplings λ_i computed at the one-loop level. Higher-order corrections to λ_2 are known to be important [18–20]. However, h^0 comes up in the FCNC vertices κ_{ij} of Eq. (11), along with a $\cot\beta$ suppression factor. The $\tan\beta$ -enhanced effects considered here are thus largely uncorrelated with M_h . For this reason we do not correct the one-loop formulas and simply impose $M_h > 115$ GeV.
- (iii) The following bounds are imposed on a_t and a_b to avoid the occurrence of color symmetry-breaking vacua at tree-level [41]:

$$\begin{aligned} |a_t|^2 &< 3(M_{\tilde{t}_L}^2 + M_{\tilde{t}_R}^2 + m_{22}^2), \\ |a_b|^2 &< 3(M_{\tilde{t}_L}^2 + M_{\tilde{b}_R}^2 + m_{11}^2). \end{aligned} \quad (90)$$

The corresponding bound for a_τ is not imposed as slepton parameters have very little impact on the quark FCNC considered here anyway.

- (iv) The most stringent constraint on the FCNC coupling $\bar{\kappa}$ comes from the $B_s \rightarrow \mu^+ \mu^-$ branching fraction [3–5,42], which we normalize to ΔM_s to avoid the occurrence of the parameters f_{B_s} and $V_{ts} V_{tb}^*$. This time the Higgs-pole contribution overcomes the standard-model and Higgs-loop pieces.

In addition, these last two interfere destructively, so we will neglect them. The counterpart of Eq. (86) then reads (with $m_\mu^2/M_{B_q}^2 = 0$ for simplicity):

$$\begin{aligned} \mathcal{B}(B_q \rightarrow \mu^+ \mu^-) &= \tau_{B_q} |V_{tq} V_{tb}^*|^2 f_{B_q}^2 M_{B_q}^5 \frac{m_\mu^2}{64\pi v^4} \frac{|\bar{\kappa}|^2 [|\mathcal{F}_P|^2 + |\mathcal{F}_S|^2]}{\cos^2\beta |1 + \epsilon_\mu t_\beta|^2} \\ &\equiv R_q \mathcal{B}(B_q \rightarrow \mu^+ \mu^-)^{\text{SM}}, \end{aligned} \quad (91)$$

where $\mathcal{F}_P$ and $\mathcal{F}_S$ refer to the Wilson coefficients of the effective operators $Q_P = (\bar{b}_R s_L)(\bar{\ell} \gamma_5 \ell)$ and $Q_S = (\bar{b}_R s_L)(\bar{\ell} \ell)$ arising from neutral-Higgs exchanges and

$$\begin{aligned} \epsilon_\mu &= \frac{g'^2}{16\pi^2} \frac{\mu^*}{M_1} \left[-\frac{1}{2} H_2 \left(\frac{M_{\tilde{\mu}L}^2}{|M_1|^2}, \frac{|\mu|^2}{|M_1|^2} \right) \right. \\ &\quad \left. + H_2 \left(\frac{M_{\tilde{\mu}R}^2}{|M_1|^2}, \frac{|\mu|^2}{|M_1|^2} \right) \right] - \frac{g'^2}{16\pi^2} \frac{\mu^*}{M_1} \\ &\quad \times H_2 \left(\frac{M_{\tilde{\mu}L}^2}{|M_1|^2}, \frac{M_{\tilde{\mu}R}^2}{|M_1|^2} \right) + \frac{3g^2}{32\pi^2} \frac{\mu^*}{M_2} \\ &\quad \times H_2 \left(\frac{M_{\tilde{\mu}L}^2}{|M_2|^2}, \frac{|\mu|^2}{|M_2|^2} \right). \end{aligned} \quad (92)$$

This result agrees with [43] but disagrees with [44]. The loop function H_2 was defined in Eq. (34) and $M_{\tilde{\mu}L(R)} = M_{\tilde{\tau}L(R)}$ in our MFV scenario. In the large $\tan\beta$ limit and at tree-level in the Higgs potential, we have: $\mathcal{F}_P = -\mathcal{F}_S = \mathcal{F}^+/2 = 1/M_A^2$, so that $\mathcal{B}(B_q \rightarrow \mu^+ \mu^-)$ is tightly correlated with ΔM_q^{LR} [5]. Going beyond the tree-level and large $\tan\beta$ approximations we obtain: $\mathcal{F}_P = s_\beta(\mathcal{F}^+ - \mathcal{F}^-)/2$, with $\mathcal{F}^\pm$ given in Eqs. (18) and (19). This formula is actually valid in any 2HDM, including arbitrary CP -violating phases (in the CP -conserving case it reduces to the usual identity $\mathcal{F}_P = s_\beta/M_A^2$). We did not find such a general and simple form for $\mathcal{F}_S$, yet it is straightforward to write it in terms of M_A , $\tan\beta$, and the λ_i 's (alternatively one can of course express it in terms of the neutral-Higgs masses and mixing angles). Note that one still has $\mathcal{F}_S = -(\mathcal{F}^+ + \mathcal{F}^-)/2$ up to $\cot\beta$ -suppressed terms. Sparticle loop corrections to the Higgs self-energies turn out to be relevant in the case of $\mathcal{F}_S$: they can be as large as 15% for small M_A after all constraints are taken into account, as we will see in Sec. IVb. Numerically, Higgs-mediated effects can easily be very large:

$$\begin{aligned} R_s &= 9930 \left[1 + \frac{(-\lambda_5^+ + |\lambda_7|^2/\lambda_2)v^2}{M_A^2} \right] \\ &\quad \times \frac{|16\pi^2 \epsilon_Y|^2 (120 \text{ GeV})^4}{|1 + \tilde{\epsilon}_3 t_\beta|^2 |1 + \epsilon_0 t_\beta|^2 |1 + \epsilon_\mu t_\beta|^2 M_A^4} \left[\frac{t_\beta}{40} \right]^6 \end{aligned} \quad (93)$$

and $R_d \simeq R_s$. The first correction factor above captures the bulk of the effects from the Higgs self-energies, yet the exact formula for $\mathcal{F}_S$ should be used for better precision. In practice, the looser constraint $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)/\Delta M_s < 5.7 \times 10^{-9}$ ps, obtained from $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)^{\text{exp}} < 10^{-7}$ [45] and $\Delta M_s^{\text{exp}} = 17.77 \pm 0.1 \pm 0.07 \text{ ps}^{-1}$ [46], is built-in in the scan procedure, then the current 95% C.L. bound $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)/\Delta M_s < 3.3 \times 10^{-9}$ ps corresponding to $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)^{\text{exp}} < 5.8 \times 10^{-8}$ [47] is imposed.¹³ We also checked the bound $\mathcal{B}(B_d \rightarrow \mu^+ \mu^-)/\Delta M_d < 3.6 \times 10^{-8}$ ps, corresponding to $\mathcal{B}(B_d \rightarrow \mu^+ \mu^-)^{\text{exp}} < 1.8 \times 10^{-8}$ [47] and $\Delta M_d^{\text{exp}} = 0.507 \pm 0.005 \text{ ps}^{-1}$ [49]. This provides no additional constraint. Neither do $\mathcal{B}(B_{s,d} \rightarrow \mu^+ \mu^-)$ and $\Delta M_{s,d}$ taken separately due to the large parametric uncertainties from f_{B_q} .

- (v) The $b \rightarrow s\gamma$ branching fraction with the energy cut $E_\gamma > 1.6 \text{ GeV}$ is computed using the fortran code SusyBSG [50]. Higgs-mediated effects now appear at loop level with smaller powers of $\tan\beta$, so that purely supersymmetric loop corrections (scaling as $1/M_{\text{SUSY}}$) are comparatively more important. For $a_t \mu < 0$ and relatively light M_{SUSY} , chargino and charged-Higgs loops can interfere destructively and more room is left for New Physics. This interplay is welcome when $\mu < 0$ as the charged-Higgs contribution then tends to overshoot the experimental branching fraction. On the other hand, in that case, the discrepancy between the $(g-2)_\mu$ standard-model prediction and its present measurement [51] increases (for a recent discussion, see e.g. [43] and references therein). The significance of this discrepancy, however, is questioned by the new $e^+e^- \rightarrow \pi^+\pi^-\gamma$ BABAR data [52]. We, therefore, still include the situation $\mu < 0$ in our considerations. The $\mathcal{B}(b \rightarrow s\gamma)$ experimental world average reads: $\mathcal{B}(b \rightarrow s\gamma)^{\text{exp}} = (3.52 \pm 0.23 \pm 0.09) \times 10^{-4}$ [53]. The standard-model central value of the SusyBSG program agrees well with the next-to-next-to-leading-order prediction $\mathcal{B}(b \rightarrow s\gamma) = (3.15 \pm 0.23) \times 10^{-4}$ [54]. We combine the experimental error with the uncertainties discussed in Ref. [50] and obtain the following two-sigma range: $2.71 \times 10^{-4} < \mathcal{B}(b \rightarrow s\gamma) < 4.33 \times 10^{-4}$.
- (vi) The $B^+ \rightarrow \tau^+ \nu$ branching fraction is given by

$$\mathcal{B}(B^+ \rightarrow \tau^+ \nu) = \frac{G_F^2}{8\pi} \tau_{B^+} |V_{ub}|^2 f_{B_d}^2 \times M_{B^+} m_\tau^2 \left(1 - \frac{m_\tau^2}{M_{B^+}^2}\right)^2 |1 - g_P|^2, \quad (94)$$

¹³A newer preliminary result is $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)^{\text{exp}} < 4.3 \times 10^{-8}$ at 95% C.L. [48].

where

$$g_P = \frac{M_{B^+}^2 t_\beta^2}{(1 + \epsilon_0 t_\beta)(1 + \epsilon_\tau^* t_\beta) M_{H^+}^2} \quad (95)$$

parametrizes Higgs-mediated effects. ϵ_τ is obtained from ϵ_μ in Eq. (92) by the replacement $M_{\tilde{\mu}L(R)} \rightarrow M_{\tilde{\tau}L(R)}$. Corrections to the Higgs potential merely change the value of M_{H^+} , which becomes a function of M_A , $\tan\beta$, and the various supersymmetric parameters. Again, we include these corrections in our numerical analysis. Given the large theoretical and experimental uncertainties, we impose: $g_P < 0.36 \cup 1.64 < g_P < 2.73$. The constraint from $\mathcal{B}(B \rightarrow D\tau\nu)$ allows to reduce the second interval, and we end up with $g_P < 0.36 \cup 1.64 < g_P < 1.79$ [55].

B. Size of the new contributions

The various Higgs-mediated contributions ΔM_s^{LR} , ΔM_q^{LL} and ΔM_q^{HL} normalized to the standard-model prediction ΔM_q^{SM} are displayed in Fig. 3(a)–3(c) as a function of the FCNC coupling $\bar{\kappa}$. As expected from Eq. (89), Higgs-loop effects are very small (the bottom Yukawa coupling actually does not reach its upper bound $y_b = 2$ in the presence of the other constraints, see Fig. 3(d). The upper and lower branches correspond to $\mu < 0$ and $\mu > 0$, respectively). Further, the contribution of ΔM_q^{LL} appears to be much smaller than that of ΔM_s^{LR} despite the fact that $m_b \mathcal{F}^-$ can compete with $m_s \mathcal{F}^+$, see Fig. 3(e) and 3(f). This suppression is a consequence of the $B_s \rightarrow \mu^+ \mu^-$ constraint. Indeed, large values of $\mathcal{F}^-$ are obtained for small values of M_A , to which $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ is particularly sensitive. As a result, the recent CDF bound [47] only leaves room for very small $\bar{\kappa}$ couplings, killing practically all effects in ΔM_q^{LL} (and actually also in ΔM_s^{LR} for such small M_A values). In Fig. 3(g), we illustrate this decrease of the maximal $\bar{\kappa}$ value allowed by the $B_s \rightarrow \mu^+ \mu^-$ constraint with M_A . Blue/magenta/red (dark grey/light grey/grey) points correspond to $M_A = 550/350/150 \text{ GeV}$ (the constraints in the right column of Table III were not imposed to keep the focus on $B_s \rightarrow \mu^+ \mu^-$). As one can see, for M_A fixed, the largest possible $\bar{\kappa}$ first increases with $\tan\beta^2$, as expected from Eq. (87), saturates the $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ experimental upper bound for some $\tan\beta$ value, and is then forced to decrease. For smaller M_A , the $B_s \rightarrow \mu^+ \mu^-$ constraint is more stringent and only a smaller $\bar{\kappa}_{\text{max}}$ can be achieved. This growth of $\bar{\kappa}_{\text{max}}$ with M_A is sufficient to overcome the $1/M_A^4$ suppression factor in ΔM_s^{LR} but not the $1/M_A^4$ one in ΔM_q^{LL} . Overall, Higgs-mediated effects in ΔM_q are of the LR type and the room for such effects increases with M_A . The correlation between ΔM_s and $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ pointed out in Ref. [5] is thus preserved, up to the relatively small Higgs self-energy corrections to $B_s \rightarrow \mu^+ \mu^-$ mentioned above Eq. (93). These are only




FIG. 3 (color online). Study of Higgs-mediated contributions to ΔM_q (see text). Black dots denote the points in parameter space that satisfy all constraints, while gray dots refer to those that only satisfy the initial constraints (see Table III). In plots (g) and (h), blue/magenta/red (dark grey/light grey/grey) points correspond to $M_A = 550/350/150$ GeV, respectively. Plain lines indicate the $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ constraint. In plot (h), the dashed line corresponds to the more stringent $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ constraint in the right column of Table III.

relevant for $\mu > 0$, $\tan\beta \lesssim 25$, small M_A , and large λ_5 , though (see Fig. 4). The mass difference in the B_d system, on the other hand, remains unaffected. These results seem to contradict those of Ref. [56], where large LL-type

effects were claimed. Without attempting a close numerical comparison (the sign of $\Delta M_s^{\text{LL}} / \Delta M_s^{\text{LR}}$ in [56] is actually reversed with respect to ours), let us point out that, as shown by Figs. 3(e) and 3(f), a large $\Delta M_s^{\text{LL}} / \Delta M_s^{\text{LR}}$ ratio

TABLE III. Constraints built-in in the scan procedure (left) and imposed afterwards (right).

Built-in constraints	Additional constraints
$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)/\Delta M_s < 5.7 \times 10^{-9}$ ps	$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)/\Delta M_s < 3.3 \times 10^{-9}$ ps
$y_b < 2$	$2.71 \times 10^{-4} < \mathcal{B}(b \rightarrow s\gamma) < 4.33 \times 10^{-4}$
$M_h > 115$ GeV	$g_P < 0.36$ or $1.64 < g_P < 1.79$
Stability bounds, see Eq. (90)	

does not automatically lead to large nonstandard effects in ΔM_s due to the $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ constraint.

Being of the LR type, the maximal effect allowed in ΔM_s is essentially determined by the current $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ experimental upper bound for a fixed (but large enough) value of the ratio $\tan\beta/M_A$. This is illustrated in Fig. 3(h) for a slightly larger bound (cf. left-hand side column in Table III). The correlation itself is displayed in Fig. 5, where each diagonal corresponds to a fixed value of the ratio $\tan\beta/M_A$. We distinguish the cases $\mu > 0$ and $\mu < 0$ as the latter leads to larger effects due to smaller denominators in Eq. (87) but is disfavored by the measurement of $(g - 2)_\mu$, as mentioned previously. The sign of the various a -terms, on the other hand, has only little impact. Still, in the case $\mu < 0$, $a_t > 0$ helps satisfy the $b \rightarrow s\gamma$ constraint. Note that the effect of the $B^+ \rightarrow \tau^+ \nu$ constraint is particularly transparent on Fig. 5: it removes the points with large $\tan\beta/M_A$ ratios, i.e., the steepest diagonals. Altogether, for $M_A < 600$ GeV, Higgs-mediated effects in ΔM_s can reach $\sim 7\%$ ($\sim 20\%$) for $\mu > 0$ ($\mu < 0$). These findings agree with those of Ref. [57]. They merely follow from the $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ constraint, as one can see from Figs. 3(g), 3(h), and 5.

Finally, for completeness (or out of curiosity), we display in Fig. 6 the typical size of various quantities and their dependence on effective couplings or supersymmetric parameters. In particular, in the last four plots, we illustrate how the loop functions ε_0 , ε_Y , ε_τ , and λ_5 increase with the range chosen for M_{SUSY} (more precisely, they increase with the trilinear and μ terms and decrease with the squark and slepton mass parameters $M_{\tilde{f}_L}$ and $M_{\tilde{f}_R}$ with $f = t, b, \tau$).

C. CP-violating effects

The Higgs-mediated $B - \bar{B}$ mixing amplitudes studied here can in principle generate new contributions to the CP-violating phases measured in the $B_d \rightarrow J/\psi K_S$ time-dependent CP asymmetry and the $B_s \rightarrow J/\psi \phi$ time-dependent angular distribution. The coefficients of the $\sin(\Delta M_q t)$ terms are

$$\begin{aligned} S_{J/\psi K_S} &= \sin(2\beta + \phi_d^\Delta), \\ S_{J/\psi \phi} &= -\sin(-2\beta_s + \phi_s^\Delta), \end{aligned} \quad (96)$$

where $\beta \equiv \arg[-(V_{td}^* V_{tb})/(V_{cd}^* V_{cb})]$, $\beta_s \equiv -\arg[-(V_{ts}^* V_{tb})/(V_{cs}^* V_{cb})]$, and

$$\phi_q^\Delta = \arg(M_{12}^q/M_{12}^{q,\text{SM}}) \equiv \arg(1 + h_q). \quad (97)$$

In $B_s \rightarrow J/\psi \phi$, an angular analysis separates the different CP components, the sign quoted for $S_{J/\psi \phi}$ in Eq. (96) refers to the dominant CP-even component. These phases have received a lot of attention recently. In particular, the new measurements of $-2\beta_s + \phi_s^\Delta$ by the CDF and D0 collaborations [58], both more than 1.5 sigma above its SM prediction [59], have triggered speculations about the validity of the SM [60]. A possible tension between the value of $\sin 2\beta$ obtained from $S_{J/\psi K_S}$ and the amount of CP violation in the kaon system was also pointed out [61].

Looking back at (17)–(19), it is clear that the new phases ϕ_q^Δ , when associated with the Q_2^{LR} effective operator, have to be brought up by the quark-Higgs couplings κ_{ij} as $\mathcal{F}^+$ cannot develop an imaginary part. When associated with Q_1^{SLL} or Q_1^{SRR} , on the other hand, they can arise from both




FIG. 4. Correction factor to $\mathcal{B}(B_q \rightarrow \mu^+ \mu^-)$ arising from supersymmetric loop effects in the Higgs self-energies as a function of the effective coupling λ_5 (left) and the total Higgs-mediated effects in $B_s \rightarrow \mu^+ \mu^-$ (right), for $\mu > 0$. On the left-hand side, the upper (lower) line corresponds to $M_A = 120(600)$ GeV, assuming the approximate formula of Eq. (93) for the correction factor.




FIG. 5. Correlation between ΔM_q and $\mathcal{B}(B_q \rightarrow \mu^+ \mu^-)$: (a) $q = s$, $\mu < 0$; (b) $q = s$, $\mu > 0$; (c) $q = d$, $\mu < 0$; (d) $q = d$, $\mu > 0$. The descending lines correspond to a fixed value of the ratio $\tan\beta/M_A$. From left to right: $\tan\beta/M_A = 0.05, 0.075, 0.10, 0.13, 0.21 \text{ GeV}^{-1}$. The ascending lines refer to the $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)/\Delta M_s$ constraints, see Table III. These lines do not take into account the uncertainties on the quark masses and bag factors, nor the effects from sparticle loop corrections to the Higgs potential in ΔM_s , $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$, and to the lepton Yukawa couplings in $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$, so that the actual points do not follow them exactly but are somewhat scattered.

the Yukawa sector and the Higgs potential via $\mathcal{F}^-$. Within MFV, $\kappa_{qb}^* \kappa_{bq} = |\bar{\kappa}|^2 \lambda_{qb}^2 m_q m_b / v^2$ and only C_1^{SLL} can produce a new phase (via $\epsilon_{0,Y}$ or $\lambda_{5,7}$). However, the $B_s \rightarrow \mu^+ \mu^-$ branching fraction is barely affected by CP -violating effects, so that its constraints on $|\bar{\kappa}|$ are still very well approximated by the plain lines in Fig. 3(g) (for some representative M_A values). As a result, just like in the CP -conserving case, the net effect of the suppression of $|\bar{\kappa}|$ and enhancement of $\mathcal{F}^-$ for small M_A is quite small. The MSSM with large $\tan\beta$ and MFV is thus not able to account for a large nonstandard phase in $B_s - \bar{B}_s$ (or $B_d - \bar{B}_d$) mixing, if the evidence for such a phase were confirmed. Let us emphasize, however, that the formulation of MFV adopted here does not coincide exactly with the full symmetry-based definition of Ref. [13]. In the formalism of Ref. [13], it was shown recently that new phases could appear in the $\delta_{LL}^{13,23}$ sector, in addition to those in the $(\delta_{LR}^d)^{13,23}$ sector [16]. The possible impact of these MFV phases via $\kappa_{qb}^* \kappa_{bq}$ in C_2^{LR} is *a priori* rather limited due to the $B_s \rightarrow \mu^+ \mu^-$ constraint, yet a more quantitative analysis is desirable.

Beyond MFV, the Q_2^{LR} contribution is expected to dominate. As said before, supersymmetric loop corrections to the Higgs propagator $\mathcal{F}^+$ do not bring in any new phases. These can only enter via the quark-Higgs couplings κ_{qb}^*

and κ_{bq} . The possible size of CP -violating effects generated in this way without violating the existing constraints deserves a study on its own. We will not discuss this further here.

V. CONCLUSIONS

We have studied supersymmetric loop corrections to the MSSM Higgs sector. While the tree-level Higgs sector of the MSSM is a 2HDM of type II, the soft supersymmetry-breaking terms lead to new loop-induced couplings which result in a generic 2HDM with FCNC couplings of neutral-Higgs bosons to quarks, even if the supersymmetry-breaking sector is minimally flavor-violating. The strength of these couplings to d -type quarks grows with $\tan\beta$ and precision observables of flavor physics are known to severely constrain large- $\tan\beta$ scenarios of the MSSM. The appropriate tool for such studies is an effective Lagrangian, which is derived by integrating out the heavy supersymmetric particles. The abundant literature on the subject has primarily focused on the flavor-changing Yukawa couplings [1–7]. Among the FCNC quantities, $B - \bar{B}$ mixing plays a special role, because the apparently dominant contribution of Fig. 1 vanishes. Therefore, $B - \bar{B}$ mixing is sensitive to subleading effects, whose systematic study was the main motivation for this paper. Pursuing this goal we




FIG. 6. Dependence of various quantities on effective couplings or supersymmetric parameters.

have derived several conceptual and analytic results which can be applied well beyond this topic. They can be classified into three categories:

1. *MSSM Higgs sector.* We have matched the complete MSSM Higgs sector, i.e. both the Yukawa interactions and the Higgs potential, onto an effective 2HDM. Our results for the effective Yukawa couplings are valid for arbitrary

CP phases of μ , a_t , and the gaugino masses; and (31) and (32) correct the gaugino contributions to ϵ_0 and ϵ_Y quoted in Ref. [26]. The complete one-loop matching corrections for the quartic Higgs couplings for the most general MSSM are explicitly listed in one place for the first time. This result goes beyond minimal flavor-violation and beyond the large- $\tan\beta$ limit. It is well-known that improper

choices of the MSSM renormalization scheme can lead to radiative corrections which grow with $\tan\beta$ rendering perturbative results unreliable [31]. At the heart of this problem is the feature that $\tan\beta$ is an ill-defined parameter in the general 2HDM, which permits arbitrary rotations among the two Higgs doublets. In the matching of the MSSM onto the 2HDM this feature enters through the wave-function-renormalization, and we propose an $\overline{\text{MS}}$ renormalization of $\tan\beta$ in the 2HDM which is stable in the limit of large $\tan\beta$. The relation to a $\overline{\text{DR}}$ -renormalized $\tan\beta$ in the MSSM is discussed including electroweak corrections. We identify the places in the effective Higgs potential where physical $\tan\beta$ -enhanced effects occur. The coefficients λ_2 , λ_5 and λ_7 , which are important for $B - \bar{B}$ mixing, are explicitly specified for the MFV case in (39), (38), and (37). Certain loop corrections to the Higgs potential (λ_5 , λ_6 , λ_7) and their impact on $\tan\beta$ and the Higgs-fermion couplings have also been considered in an effective-field-theory framework in Ref. [40], which appeared during completion of this paper. Their results for the λ 's agree with ours for the parameter values considered. As far as the renormalization of $\tan\beta$ is concerned, in Sec. III of the present paper we critically compared with Ref. [40]. In particular, we do find a $\tan\beta$ -enhanced term in the relation of the $\overline{\text{DR}}$ and DCPR $\tan\beta$ parameters. We stress that, in general, only the former is numerically close to the $\tan\beta$ parameter extracted from B -physics observables.

2. *Large $\tan\beta$ phenomenology.* The prime application of our results is $B - \bar{B}$ mixing. We have identified a global $U(1)$ symmetry of the $\bar{b}_R q_L$ Higgs-mediated FCNC transitions and the tree-level Higgs potential in the large- $\tan\beta$ limit which suppresses the superficially leading contribution of Fig. 1. A systematic study of $B - \bar{B}$ mixing has required the analyses of four subleading contributions, which are governed by the small parameters $m_{d,s}/m_b$, $1/\tan\beta$, v/M_{SUSY} and the loop factor $1/(4\pi)^2$. These parameters either provide a breaking of the $U(1)$ symmetry or allow for a contribution proportional to the $U(1)$ -conserving standard-model effective operator. Prior to this work, only corrections involving $m_{d,s}/m_b$ had been studied [5] (with the exception of Ref. [26]). The v/M_{SUSY} corrections are found to be suppressed. The new loop contributions include all non-decoupling SUSY corrections to the quartic Higgs interactions λ_1 – λ_7 and the contribution of neutral-Higgs box diagrams in the effective theory. In the complex MSSM the results for $\mathcal{F}^\pm$ comprising the neutral-Higgs propagators become cumbersome. We have expressed $\mathcal{F}^\pm$ in terms of subdeterminants of the neutral-Higgs mass matrix. These expressions are easy to implement and clearly reveal the invariance of the Higgs-mediated amplitudes under rotations of the basis $(-\epsilon H_d^*, H_u)$. The results for the Higgs sector are also used to refine the MSSM predictions for the $B^+ \rightarrow \tau^+ \nu$ and the $B_{s,d} \rightarrow \mu^+ \mu^-$

branching ratios. In this context we stress that loop corrections to the Higgs potential do not give rise to additional $\tan\beta$ -enhanced contributions to the charged-Higgs-fermion couplings beyond those known before Ref. [26] appeared. Hence no parametrically large modification of the charged-Higgs contributions to $B^+ \rightarrow \tau^+ \nu$ or $B \rightarrow X_s \gamma$ relative to Ref. [5] occurs.

While the MSSM corrections to $B - \bar{B}$ mixing in the large $\tan\beta$ scenario could be dominated by the contribution of λ_5 and λ_7 , the size of this piece is limited by the experimental upper bound on $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$. After performing an exhaustive analysis of this quantity, $\mathcal{B}(B \rightarrow X_s \gamma)$, $\mathcal{B}(B^+ \rightarrow \tau^+ \nu)$ and the mass of the lightest neutral-Higgs boson M_h , we find that the impact of the corrections to the Higgs potential on ΔM_s is always weaker than that of the m_s/m_b correction identified in Ref. [5]. Assessing the total Higgs-mediated MSSM corrections to ΔM_s we find an upper limit of 7% of the SM contribution for $\mu > 0$ and $M_A < 600$ GeV. If μ is negative, the upper bound is around 20%. This is in contrast with Refs. [26,56], which claim large effects of the Higgs potential on $B - \bar{B}$ mixing. The corrections to $\mathcal{B}(B_q \rightarrow \mu^+ \mu^-)$ from the Higgs potential are typically also small, but can reach 15% in some corners of the parameter space. In summary, the correlation between an enhancement of $\mathcal{B}(B_q \rightarrow \mu^+ \mu^-)$ and a (moderate) depletion of ΔM_s found in Ref. [5] remains essentially intact.

We finally note that our new contributions can alter the CP phase of the $B - \bar{B}$ mixing amplitude, while the previously known Higgs contribution proportional to $m_s m_b \mathcal{F}^+$ has the same phase as the SM term (in MFV scenarios). While the maximal possible CP phase is clearly below the sensitivity of the current Tevatron experiments, it is an open question whether future $B - \bar{B}$ mixing experiments can help to unravel the CP structure of the MSSM Higgs potential.

3. *Heavy-quark relations and bag parameters.* We have transformed the NLO anomalous dimensions computed in Ref. [28] to an operator basis and a renormalization scheme typically used in lattice calculations. The corresponding anomalous dimensions are needed to evaluate the “bag” parameters, which parametrise the hadronic matrix elements, at the electroweak scale. We further employed a heavy-quark relation to improve the numerical prediction of the bag parameter B_1^{SLL} entering the SUSY contributions to $B - \bar{B}$ mixing. The heavy-quark relation essentially determines B_1^{SLL} in terms of the bag parameter B_1^{VLL} , which is needed for the SM prediction [59]. We found $p_1^{\text{SLL}} = -\frac{5}{8} B_1^{\text{SLL}}(m_t) = -1.36 \pm 0.12$.

ACKNOWLEDGMENTS

M.G. and S.J. appreciate helpful conversations with Dominik Stöckinger, Martin Beneke, John Ellis, Gian Giudice, Uli Haisch, Janusz Rosiek, Pietro Slavich, Jure Zupan, and other members of the CERN theory group on

various aspects of this work. U.N. appreciates fruitful discussions with Lars Hofer und Dominik Scherer. S.J. acknowledges the hospitality of the University of Karlsruhe. We are grateful to A.J. Buras for comments on the manuscript, and to J.S. Lee and A. Pilaftsis for communications. This work was supported by the DFG Grant No. NI 1105/1-1, by project C6 of the DFG Research Unit SFB-TR 9 *Computergestützte Theoretische Teilchenphysik* by the European Union Contract No. MRTN-CT-2006-035482, “FLAVIANet”, the DFG cluster of excellence “Origin and Structure of the Universe”, and the RTN European Program Grant No. MRTN-CT-2004-503369.

APPENDIX A: NOTATIONS AND CONVENTIONS

To state our phase conventions for μ and M_2 we quote the chargino mass matrix:

$$\mathcal{M}_{\chi^+} = \begin{pmatrix} M_2 & \frac{gv \sin \beta}{\sqrt{2}} \\ \frac{gv \cos \beta}{\sqrt{2}} & \mu \end{pmatrix} \quad (\text{A1})$$

with $v = 246$ GeV and the chargino mass term in the Lagrangian

$$\mathcal{L}_{\chi^+}^{\text{mass}} = -(\lambda^-, \tilde{h}_d^2) \mathcal{M}_{\chi^+} (\lambda^+, \tilde{h}_u^1)^T. \quad (\text{A2})$$

For the case of a general flavor structure of the squark mass matrices we define the trilinear couplings $\hat{T}_{u_{ij}}$ and $\hat{T}_{d_{ij}}$ (with flavor indices i, j) such that the squark mass matrices read

$$\begin{aligned} \hat{\mathcal{M}}_{\tilde{u}}^2 &= \begin{pmatrix} \hat{M}_{\tilde{u}_L}^2 & \frac{v \sin \beta}{\sqrt{2}} [\hat{T}_u^\dagger - \mu \hat{Y}_u^\dagger \cot \beta] \\ \frac{v \sin \beta}{\sqrt{2}} [\hat{T}_u - \mu^* \hat{Y}_u \cot \beta] & \hat{M}_{\tilde{u}_R}^2 \end{pmatrix}, \\ \hat{\mathcal{M}}_{\tilde{d}}^2 &= \begin{pmatrix} \hat{M}_{\tilde{d}_L}^2 & \frac{v \cos \beta}{\sqrt{2}} [\hat{T}_d^\dagger - \mu \hat{Y}_d^\dagger \tan \beta] \\ \frac{v \cos \beta}{\sqrt{2}} [\hat{T}_d - \mu^* \hat{Y}_d \tan \beta] & \hat{M}_{\tilde{d}_R}^2 \end{pmatrix}, \end{aligned} \quad (\text{A3})$$

in the super-CKM basis, where the ($\overline{\text{DR}}$ -renormalized) Yukawa matrices are diagonal: $\hat{Y}_q = \text{diag}(y_{q_1}, y_{q_2}, y_{q_3})$, $q = u, d$. The mass matrices in (A3) correspond to the squark mass term

$$\mathcal{L}_{\tilde{q}}^{\text{mass}} = -\Phi_{\tilde{u}}^\dagger \hat{\mathcal{M}}_{\tilde{u}}^2 \Phi_{\tilde{u}} - \Phi_{\tilde{d}}^\dagger \hat{\mathcal{M}}_{\tilde{d}}^2 \Phi_{\tilde{d}} \quad (\text{A4})$$

with $\Phi_{\tilde{u}} = (\tilde{u}_L, \tilde{c}_L, \tilde{t}_L, \tilde{u}_R, \tilde{c}_R, \tilde{t}_R)^T$ and $\Phi_{\tilde{d}} = (\tilde{d}_L, \tilde{s}_L, \tilde{b}_L, \tilde{d}_R, \tilde{s}_R, \tilde{b}_R)^T$.

Our sign convention for the MSSM Yukawa couplings implies the following relations between 2HDM quark masses and MSSM Yukawa couplings in the up sector:

$$y_{u_i} = \frac{\sqrt{2}}{v \sin \beta} m_{u_i}. \quad (\text{A5})$$

In general, the analogous relations in the down sector involve complex phases associated with the $\tan \beta$ -enhanced threshold corrections in (22). In particular, Ref. [24], which discusses the complex MSSM for the case without flavor mixing, relates the b Yukawa coupling to the b quark mass as

$$y_b = \frac{\sqrt{2}}{v \cos \beta} \frac{m_b}{1 + \tilde{\epsilon}_3 \tan \beta}, \quad (\text{A6})$$

rendering y_b complex for complex $\tilde{\epsilon}_3$ in (33). Our approach of matching the MSSM to an effective 2HDM permits different phase conventions, because the quark fields in the MSSM and the 2HDM can be chosen to differ by a phase factor. We can rephase the b_R superfield of the MSSM in such a way that y_b is real and positive and $1 + \tilde{\epsilon}_3 \tan \beta$ in (A6) is replaced by $|1 + \tilde{\epsilon}_3 \tan \beta|$. The (physical) phase of $1 + \tilde{\epsilon}_3 \tan \beta$ will then, however, appear explicitly in the Higgs and higgsino couplings to bottom (s)quarks. Introducing 3×3 flavor mixing, the relation between $\hat{Y}_d$ and m_d, m_s, m_b is found from (22). Now the quark fields in the 2HDM differ from those in the MSSM by a complex rotation in flavor space and a particular choice for the phases of MSSM fields appears less obvious. In particular, one could render all y_{d_i} real and positive by suitable rephasings of the right-handed superfields. Note that within MFV y_b is still related to m_b via (A6) in good approximation without such rephasings. The analogous relation for the first two generations reads

$$y_{d,s} = \frac{\sqrt{2}}{v \cos \beta} \frac{m_{d,s}}{1 + \epsilon_0 \tan \beta}, \quad (\text{A7})$$

while in the lepton sector, we have

$$y_\ell = \frac{\sqrt{2}}{v \cos \beta} \frac{m_\ell}{1 + \epsilon_\ell \tan \beta} \quad \text{with } \ell = e, \mu, \tau. \quad (\text{A8})$$

In (most of) the paper we express our results in terms of fermion masses (i.e. avoiding $y_{d_i, \ell}$) to achieve formulas which are independent of such phase conventions. Note that the phases of ϵ_0 and ϵ_Y are physical and no phase convention other than that of the CKM matrix matters for κ_{ij} in (29) and (30). While the phase convention of y_{q_i} enters the phases in $\hat{T}_q$, it drops out from the MFV parameters a_q in (A12).

Finally, the quadratic squark soft-breaking terms are defined as follows:

$$\begin{aligned}
(\hat{M}_{\tilde{u}_L}^2)_{ij} &= (V_{\text{CKM}}^0 \tilde{m}_Q^2 V_{\text{CKM}}^{0\dagger})_{ij} + \frac{v^2 \sin^2 \beta}{2} \delta_{ij} |y_{u_i}|^2 + \delta_{ij} M_Z^2 \cos 2\beta (1/2 - 2\sin^2 \theta_W/3), \\
(\hat{M}_{\tilde{d}_L}^2)_{ij} &= (\tilde{m}_Q^2)_{ij} + \frac{v^2 \cos^2 \beta}{2} \delta_{ij} |y_{d_i}|^2 + \delta_{ij} M_Z^2 \cos 2\beta (-1/2 + \sin^2 \theta_W/3), \\
(\hat{M}_{\tilde{u}_R}^2)_{ij} &= (\tilde{m}_u^2)_{ij} + \frac{v^2 \sin^2 \beta}{2} \delta_{ij} |y_{u_i}|^2 + 2\delta_{ij} M_Z^2 \cos 2\beta \sin^2 \theta_W/3, \\
(\hat{M}_{\tilde{d}_R}^2)_{ij} &= (\tilde{m}_d^2)_{ij} + \frac{v^2 \cos^2 \beta}{2} \delta_{ij} |y_{d_i}|^2 - \delta_{ij} M_Z^2 \cos 2\beta \sin^2 \theta_W/3,
\end{aligned} \tag{A9}$$

where V_{CKM}^0 corresponds to the relative rotation of left-handed u -type and d -type quark fields performed when diagonalizing the Yukawa matrices. It differs from the actual CKM matrix, defined by the rotations that diagonalize the 2HDM mass matrices rather than the MSSM Yukawa couplings, by loop-suppressed (but $\tan\beta$ -enhanced) corrections. In particular, within MFV, we have:

$$V_{\text{CKM}_{ij}} = \begin{cases} \left| \frac{1 + \epsilon_0 \tan\beta}{1 + \tilde{\epsilon}_3 \tan\beta} \right| V_{\text{CKM}_{ij}}^0 & \text{for } (i, j) = (u, b), (c, b), (t, d), (t, s), \\ V_{\text{CKM}_{ij}}^0 & \text{otherwise.} \end{cases} \tag{A10}$$

The relations (A10) take a particularly compact form in the (exact) Wolfenstein parametrization, where one has

$$A = \left| \frac{1 + \epsilon_0 \tan\beta}{1 + \tilde{\epsilon}_3 \tan\beta} \right| A^0, \quad \lambda = \lambda^0, \quad \bar{\rho} = \bar{\rho}^0, \quad \bar{\eta} = \bar{\eta}^0. \tag{A11}$$

Whenever we consider the case of MFV we write

$$\begin{aligned}
\hat{T}_{u_{ij}} &= a_t y_{u_i} \delta_{ij}, & \hat{T}_{d_{ij}} &= a_b y_{d_i} \delta_{ij}, & \tilde{m}_{Q_{ij}}^2 &= \tilde{m}_Q^2 \delta_{ij}, \\
\tilde{m}_{u_{ij}}^2 &= \tilde{m}_u^2 \delta_{ij}, & \tilde{m}_{d_{ij}}^2 &= \tilde{m}_d^2 \delta_{ij}.
\end{aligned} \tag{A12}$$

The $SU(2)$ relation between $\hat{M}_{\tilde{d}_L}^2$ and $\hat{M}_{\tilde{u}_L}^2$ then implies for the third generation:

$$M_{\tilde{t}_L}^2 = M_{\tilde{b}_L}^2 + m_t^2 - \frac{m_b^2}{|1 + \tilde{\epsilon}_3 \tan\beta|^2} + M_Z^2 \cos 2\beta (1 - \sin^2 \theta_W). \tag{A13}$$

In the strict $SU(2)$ limit (i.e., $v/M_{\text{SUSY}} \rightarrow 0$) one has $M_{\tilde{t}_L}^2 = M_{\tilde{b}_L}^2$, but for small $M_{\tilde{b}_L}^2$, the term involving m_t^2 can be relevant. Also, FCNC $\tilde{W}-\tilde{u}_{Li}$ loops vanish (for universal $M_{\tilde{u}_L}^2$) by the GIM mechanism up to the m_t^2 term in (A13).

Finally, it is convenient to define the so-called superflavor basis, obtained from a generic electroweak interaction eigenstate basis by rotating the supermultiplets Q_L , u_R and d_R such that the quadratic squark soft-breaking terms are diagonal. We denote the corresponding entries by $\tilde{m}_{Q_i}^2$, $\tilde{m}_{u_i}^2$, and $\tilde{m}_{d_i}^2$. For $M_{\text{SUSY}} \gg v$, these are just the squark masses, and the computation of the effective couplings λ_i induced by heavy squark loops for arbitrary flavor and CP structure is greatly simplified. The Yukawa matrices and trilinear terms in this basis are simply written $Y_{u,d}$ and $T_{u,d}$, respectively. They are given in terms of $\hat{Y}_{u,d}$ and $\hat{T}_{u,d}$ as follows:

$$\begin{aligned}
Y_u^T &= U_u \hat{Y}_u V_{\text{CKM}}^0 V_d^\dagger, & Y_d^T &= U_d \hat{Y}_d V_d^\dagger, \\
T_u^T &= U_u \hat{T}_u V_{\text{CKM}}^0 V_d^\dagger, & T_d^T &= U_d \hat{T}_d V_d^\dagger,
\end{aligned} \tag{A14}$$

where the matrices V_d , U_u and U_d are defined such that

$$\begin{aligned}
\text{diag}(\tilde{m}_{Q_i}^2) &= V_d \tilde{m}_Q^2 V_d^\dagger, & \text{diag}(\tilde{m}_{u_i}^2) &= U_u \tilde{m}_u^2 U_u^\dagger, \\
\text{diag}(\tilde{m}_{d_i}^2) &= U_d \tilde{m}_d^2 U_d^\dagger.
\end{aligned} \tag{A15}$$

Assuming MFV, one is allowed to choose $V_d = U_u = U_d = 1$.

Our conventions comply with the Les Houches accord [62]. In particular, our $Y_{u,d}$ and $T_{u,d}$ matrices correspond to a particular choice of the generic $Y_{u,d}$ and $T_{u,d}$ matrices of Ref. [62]. Our conventions also agree with those of Ref. [63], except that the sign convention of our $\hat{Y}_d$ in (A6) is opposite. Besides, our $\hat{T}_u$ equals $-A_u$ and our $\hat{T}_d$ equals A_d of Ref. [63], respectively.

APPENDIX B: MATCHING OF THE MSSM ONTO A 2HDM

The notation of Sec. III distinguishes between the coefficients $\hat{\lambda}_i$ and $\bar{\lambda}_i$. The former quantities contain the results from the supersymmetric loop corrections to the quadrilinear Higgs couplings, whose tree-level values are given in (21). The latter coefficients also include the effect of the wave-function and gauge coupling renormalization constants in Appendix B1.

In the following, we summarize the one-loop matching corrections for the quartic Higgs-coupling constants in the general MSSM. While the calculation of loop corrections to the Higgs sector of the MSSM has a long history, see for example [18–24], and a determined reader could extract part of the matching coefficients below from these works, the results collected in this appendix as a service to the

reader are more complete than those in the literature, capturing the effects of the full set of mass, flavor-violation, and CP -violation parameters of the most general MSSM. The effective potential V in (12) must be used with $\lambda_i = \bar{\lambda}_i$ and the relation between $\bar{\lambda}_i$ and $\hat{\lambda}_i$ is given in (71); the renormalization constants needed in this relation are given in Appendix B1. In the following subsections, we quote the results for $\hat{\lambda}_i$ in the general MSSM and decompose $\hat{\lambda}_{1-7}$ as

$$\hat{\lambda}_i = \lambda_i^{\text{tree}} + \frac{\lambda_i^{\text{ino}} + \lambda_i^{\text{sferm}}}{16\pi^2}. \quad (\text{B1})$$

The tree-level values λ_i^{tree} are given in (21). λ_i^{ino} and λ_i^{sferm} , given in Appendices B2, and B3, contain the contributions from higgsino and gaugino loops and from sfermion loops, respectively. Finally we also list the relevant loop functions in Appendix B4. All these results are given in the superflavor basis including the most general soft-breaking terms.

1. Renormalization constants

The renormalization of $g^{(i)}$ is related only to the field renormalization of W and B , $Z_{W,B} = 1 + \delta Z_{W,B}$, if we

decouple the sfermionic, higgsino and gaugino contributions: $\delta g' = -\delta Z_B/2$ and $\delta g = -\delta Z_W/2$. The finite part of the one-loop wave-function renormalization constants of the gauge bosons are

$$\begin{aligned} \delta Z_W &= \frac{g^2}{16\pi^2} \frac{1}{6} \left[4 \log \frac{|\mu|^2}{\mu_0^2} + 8 \log \frac{M_2^2}{\mu_0^2} \right. \\ &\quad \left. + \sum_{i=1}^3 \left(\log \frac{\tilde{m}_{l_i}^2}{\mu_0^2} + N_C \log \frac{\tilde{m}_{Q_i}^2}{\mu_0^2} \right) - 4 \right] \\ \delta Z_B &= \frac{g'^2}{16\pi^2} \frac{1}{3} \left[2 \log \frac{|\mu|^2}{\mu_0^2} + \sum_{i=1}^3 \left(\log \frac{\tilde{m}_{e_i}^2}{\mu_0^2} + \frac{1}{2} \log \frac{\tilde{m}_{l_i}^2}{\mu_0^2} \right. \right. \\ &\quad \left. \left. + \frac{4N_C}{9} \log \frac{\tilde{m}_{u_i}^2}{\mu_0^2} + \frac{N_C}{9} \log \frac{\tilde{m}_{d_i}^2}{\mu_0^2} + \frac{N_C}{18} \log \frac{\tilde{m}_{Q_i}^2}{\mu_0^2} \right) \right], \quad (\text{B2}) \end{aligned}$$

where μ_0 is the renormalization scale and the soft-breaking terms are written in the superflavor basis (see Appendix A).

The sfermionic contributions to the wave-function renormalization constants of the Higgs bosons are

$$\begin{aligned} \delta Z_{dd} &= \frac{1}{32\pi^2} \sum_{ij} [3B'_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}) T_{d_{ji}} T_{d_{ji}}^* + 3|\mu|^2 B'_0(\tilde{m}_{u_i}, \tilde{m}_{Q_j}) Y_{u_{ji}} Y_{u_{ji}}^* + B'_0(\tilde{m}_{e_i}, \tilde{m}_{l_j}) T_{e_{ji}} T_{e_{ji}}^*] \\ \delta Z_{ud} &= -\frac{1}{16\pi^2} \sum_{ij} [3\mu^* B'_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}) T_{d_{ji}}^* Y_{d_{ji}} + 3\mu^* B'_0(\tilde{m}_{u_i}, \tilde{m}_{Q_j}) T_{u_{ji}}^* Y_{u_{ji}} + \mu^* B'_0(\tilde{m}_{e_i}, \tilde{m}_{l_j}) T_{e_{ji}}^* Y_{e_{ji}}] \\ \delta Z_{uu} &= \frac{1}{32\pi^2} \sum_{ij} [3B'_0(\tilde{m}_{u_i}, \tilde{m}_{Q_j}) T_{u_{ji}} T_{u_{ji}}^* + 3|\mu|^2 B'_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}) Y_{d_{ji}} Y_{d_{ji}}^* + |\mu|^2 B'_0(\tilde{m}_{e_i}, \tilde{m}_{l_j}) Y_{e_{ji}}^* Y_{e_{ji}}], \quad (\text{B3}) \end{aligned}$$

while the respective contributions of the gaugino and higgsino loops read:

$$\begin{aligned} \delta Z_{dd} &= -\frac{1}{16\pi^2} \frac{1}{8} (g'^2 W(|M_1|, |\mu|) + 3g^2 W(|M_2|, |\mu|)) \\ \delta Z_{ud} &= -\frac{1}{16\pi^2} \mu^* (g'^2 M_1^* B'_0(|M_1|, |\mu|) + 3g^2 M_2^* B'_0(|M_2|, |\mu|)) \\ \delta Z_{uu} &= -\frac{1}{16\pi^2} \frac{1}{8} (g'^2 W(|M_1|, |\mu|) + 3g^2 W(|M_2|, |\mu|)). \quad (\text{B4}) \end{aligned}$$

2. Higgsino-gaugino contributions to λ_1 – λ_7

The situation of $\bar{\lambda}_5 = \hat{\lambda}_5$ is particularly simple: The matching correction only involves the box function and λ_5^{ino} can be written in a compact form:

$$\lambda_5^{\text{ino}} = 3g^4 \mu^2 M_2^2 D_0(|M_2|, |M_2|, |\mu|, |\mu|) + 2g^2 g'^2 \mu^2 M_1 M_2 D_0(|M_1|, |M_2|, |\mu|, |\mu|) + g'^4 \mu^2 M_1^2 D_0(|M_1|, |M_1|, |\mu|, |\mu|), \quad (\text{B5})$$

if we use the loop functions defined in Appendix B4.

We find for $\lambda^{\text{ino}} = \lambda_1^{\text{ino}}, \dots, \lambda_4^{\text{ino}}, \lambda_6^{\text{ino}}, \lambda_7^{\text{ino}}$:

$$\begin{aligned} \lambda^{\text{ino}} &= g^4 (a_s + a_2 \tilde{D}_2(|M_2|, |M_2|, |\mu|, |\mu|) + a_4 \tilde{D}_4(|M_2|, |M_2|, |\mu|, |\mu|)) + g^2 g'^2 (a'_s + a'_2 \tilde{D}_2(|M_1|, |M_2|, |\mu|, |\mu|) \\ &\quad + a'_4 \tilde{D}_4(|M_1|, |M_2|, |\mu|, |\mu|)) + g'^4 (a''_s + a''_2 \tilde{D}_2(|M_1|, |M_1|, |\mu|, |\mu|) + a''_4 \tilde{D}_4(|M_1|, |M_1|, |\mu|, |\mu|)), \quad (\text{B6}) \end{aligned}$$

where the coefficients $a_s, \dots, a''_4$ depend on the index i labeling λ_i in (B1), which we suppress throughout this appendix. The coefficients $a_{2,4}^{(i)}$ are given in Table IV, while

TABLE IV. Coefficients entering $\lambda_1^{\text{ino}} - \lambda_4^{\text{ino}}$, λ_6^{ino} , and λ_7^{ino} in Eq. (B6).

λ	a_2	a_4	a'_2	a'_4	a''_2	a''_4
λ_1	$\frac{1}{2} M_2 ^2$	$\frac{5}{2}$	$\frac{1}{2}(M_1 M_2^* + M_1^* M_2)$	1	$\frac{1}{2} M_1 ^2$	$\frac{1}{2}$
λ_2	$\frac{1}{2} M_2 ^2$	$\frac{5}{2}$	$\frac{1}{2}(M_1 M_2^* + M_1^* M_2)$	1	$\frac{1}{2} M_1 ^2$	$\frac{1}{2}$
λ_3	$3 \mu ^2 + \frac{5}{2} M_2 ^2$	$\frac{1}{2}$	$2 \mu ^2 + \frac{1}{2}(M_1 M_2^* + M_1^* M_2)$	1	$ \mu ^2 + \frac{1}{2} M_1 ^2$	$\frac{1}{2}$
λ_4	$-3 \mu ^2 - 2 M_2 ^2$	2	$2 \mu ^2 - M_1 M_2^* - M_1^* M_2$	-2	$- \mu ^2$	0
λ_6	$3\mu M_2$	0	$\mu(M_1 + M_2)$	0	μM_1	0
λ_7	$3\mu M_2$	0	$\mu(M_1 + M_2)$	0	μM_1	0

$$a_s = \begin{cases} -\frac{3}{4} \\ 0 \\ 0 \end{cases}, \quad a'_s = \begin{cases} -\frac{1}{2} \\ 1 \\ 0 \end{cases}, \quad a''_s = \begin{cases} -\frac{1}{4} \\ 0 \\ 0 \end{cases} \quad \text{for } \begin{cases} \lambda_1 \text{ to } \lambda_3 \\ \lambda_4 \\ \lambda_5 \text{ to } \lambda_7 \end{cases} \quad (\text{B7})$$

3. Sfermion contributions to $\lambda_1 - \lambda_7$

The sfermion contribution to λ_{1-7} are products of loop functions and flavor dependent coefficients if we sum over the generation index of the internal sfermions. For $\hat{\lambda}_5$, our results then take the simple form

$$\lambda_5^{\text{sferm}} = d_1^{ijkl} D_0(\tilde{m}_{e_i}, \tilde{m}_{e_j}, \tilde{m}_{l_k}, \tilde{m}_{l_l}) + d_2^{ijkl} D_0(\tilde{m}_{d_i}, \tilde{m}_{d_j}, \tilde{m}_{Q_k}, \tilde{m}_{Q_l}) + d_3^{ijkl} D_0(\tilde{m}_{Q_i}, \tilde{m}_{Q_j}, \tilde{m}_{u_k}, \tilde{m}_{u_l}), \quad (\text{B8})$$

TABLE V. Slepton contribution to $\lambda_1^{\text{sferm}} - \lambda_7^{\text{sferm}}$ in Eqs. (B8), (B10), and (B13).

	d_1^{ijkl}
λ_1	$-T_{e_{ki}} T_{e_{lj}} T_{e_{kj}}^* T_{e_{li}}^*$
λ_2	$- \mu ^4 Y_{e_{kj}} Y_{e_{li}} Y_{e_{ki}}^* Y_{e_{lj}}^*$
λ_3	$- \mu ^2 T_{e_{li}} Y_{e_{kj}} (T_{e_{lj}}^* Y_{e_{ki}} + T_{e_{ki}}^* Y_{e_{lj}})$
λ_4	$ \mu ^2 T_{e_{li}} T_{e_{kj}}^* Y_{e_{kj}} Y_{e_{li}}^*$
λ_5	$-\mu^2 T_{e_{kj}} T_{e_{li}} Y_{e_{ki}}^* Y_{e_{lj}}^*$
λ_6	$\mu T_{e_{ki}} T_{e_{lj}} T_{e_{li}}^* Y_{e_{kj}}^*$
λ_7	$\mu \mu ^2 T_{e_{lj}} Y_{e_{ki}} Y_{e_{kj}}^* Y_{e_{li}}^*$

where the slepton contribution is contained in d_1^{ijkl} listed in Table V and d_{2-4}^{ijkl} comprises the squark contribution (Table VI). Only λ_4 receives a contribution from d_4^{ijkl} :

$$d_4^{ijkl} = -3(T_{d_{ki}} T_{u_{kl}}^* - |\mu|^2 Y_{d_{ki}} Y_{u_{kl}}^*) \times (T_{u_{jl}} T_{d_{ji}}^* - |\mu|^2 Y_{u_{jl}} Y_{d_{ji}}^*) \quad \text{for } \lambda_4, \quad (\text{B9})$$

while $d_4^{ijkl} = 0$ for λ_i with $i \neq 4$. The contributions to the matching coefficients depend on the Yukawa couplings $Y_{e,u,d}$ of the charged leptons, the up-type quarks, and the down-type quarks as well as on the trilinear soft-breaking terms $T_{e,u,d}$, defined in the superflavor basis (see Appendix A).

We write $\lambda_{1-4}^{\text{sferm}} = \lambda_{1-4}^{\text{sl}} + \lambda_{1-4}^{\text{sq}}$, and find for the slepton contribution

$$\lambda_{1-4}^{\text{sl}} = (b_1 \delta_{ij} + b_2 Y_{eei} \delta_{ij} + b_3 Y_{eeij} Y_{eeji}) B_0(\tilde{m}_{e_i}, \tilde{m}_{e_j}) + (b_4 \delta_{ij} + b_5 \bar{Y}_{eei} \delta_{ij} + b_6 \bar{Y}_{eeij} \bar{Y}_{eeji}) B_0(\tilde{m}_{l_i}, \tilde{m}_{l_j}) + (c_1 |\mu|^2 Y_{e_{ki}} Y_{e_{kj}}^* \delta_{ij} + c_2 T_{e_{ki}} T_{e_{kj}}^* \delta_{ij} + c_3 |\mu|^2 Y_{e_{ki}} Y_{e_{kj}}^* Y_{eeij} + c_4 T_{e_{ki}} T_{e_{kj}}^* Y_{eeij}) C_0(\tilde{m}_{e_i}, \tilde{m}_{e_j}, \tilde{m}_{l_k}) + (c_5 |\mu|^2 Y_{e_{ji}} Y_{e_{ji}}^* \delta_{jk} + c_6 T_{e_{ji}} T_{e_{ji}}^* \delta_{jk} + c_7 |\mu|^2 Y_{e_{ji}} Y_{e_{ki}}^* \bar{Y}_{eekj} + c_8 T_{e_{ji}} T_{e_{ki}}^* \bar{Y}_{eekj}) C_0(\tilde{m}_{e_i}, \tilde{m}_{l_j}, \tilde{m}_{l_k}) + d_1^{ijkl} D_0(\tilde{m}_{e_i}, \tilde{m}_{e_j}, \tilde{m}_{l_k}, \tilde{m}_{l_l}), \quad (\text{B10})$$

while the squark contribution reads

TABLE VI. D_0 squark contribution to $\lambda_1^{\text{sferm}} - \lambda_7^{\text{sferm}}$ in Eqs. (B8) and (B13).

	d_2^{ijkl}	d_3^{ijkl}
λ_1	$-3T_{d_{ki}} T_{d_{lj}} T_{d_{kj}}^* T_{d_{li}}^*$	$-3 \mu ^4 Y_{u_{il}} Y_{u_{jk}} Y_{u_{ik}}^* Y_{u_{jl}}^*$
λ_2	$-3 \mu ^4 Y_{d_{kj}} Y_{d_{li}} Y_{d_{ki}}^* Y_{d_{lj}}^*$	$-3T_{u_{ik}} T_{u_{jl}} T_{u_{il}}^* T_{u_{jk}}^*$
λ_3	$-3 \mu ^2 T_{d_{li}} Y_{d_{kj}} (T_{d_{lj}}^* Y_{d_{ki}} + T_{d_{ki}}^* Y_{d_{lj}})$	$-3 \mu ^2 T_{u_{jl}} Y_{u_{ik}} (T_{u_{jk}}^* Y_{u_{il}} + T_{u_{il}}^* Y_{u_{jk}})$
λ_4	$3 \mu ^2 T_{d_{li}} T_{d_{kj}}^* Y_{d_{kj}} Y_{d_{li}}^*$	$3 \mu ^2 T_{u_{jl}} T_{u_{il}}^* Y_{u_{ik}} Y_{u_{jk}}^*$
λ_5	$-3\mu^2 T_{d_{kj}} T_{d_{li}} Y_{d_{ki}}^* Y_{d_{lj}}^*$	$-3\mu^2 T_{u_{ik}} T_{u_{jl}} Y_{u_{il}}^* Y_{u_{jk}}^*$
λ_6	$3\mu T_{d_{ki}} T_{d_{lj}} T_{d_{li}}^* Y_{d_{kj}}^*$	$3\mu \mu ^2 T_{u_{il}} Y_{u_{jk}} Y_{u_{ik}}^* Y_{u_{jl}}^*$
λ_7	$3\mu \mu ^2 T_{d_{lj}} Y_{d_{ki}} Y_{d_{kj}}^* Y_{d_{li}}^*$	$3\mu T_{u_{il}} T_{u_{jk}} T_{u_{jl}}^* Y_{u_{ik}}^*$

$$\begin{aligned}
\lambda_{1-4}^{\text{sq}} = & (b_7 \delta_{ij} + b_8 Y_{ddii} \delta_{ij} + b_9 Y_{ddij} Y_{ddji}) B_0(\tilde{m}_{d_i}, \tilde{m}_{d_j}) + b_{10} Y_{duij} Y_{udji} B_0(\tilde{m}_{d_i}, \tilde{m}_{u_j}) \\
& + (b_{11} \delta_{ij} + b_{12} Y_{uuui} \delta_{ij} + b_{13} Y_{uuij} Y_{uuji}) B_0(\tilde{m}_{u_i}, \tilde{m}_{u_j}) + (b_{14} \delta_{ij} + b_{15} \bar{Y}_{ddii} \delta_{ij} + b_{16} \bar{Y}_{uuui} \delta_{ij} + b_{17} \bar{Y}_{ddij} \bar{Y}_{ddji} \\
& + b_{18} \bar{Y}_{uuij} \bar{Y}_{uuji} + b_{19} \bar{Y}_{ddij} \bar{Y}_{uuji}) B_0(\tilde{m}_{Q_i}, \tilde{m}_{Q_j}) + (c_9 |\mu|^2 Y_{d_{ki}} Y_{d_{ki}}^* \delta_{ij} + c_{10} T_{d_{ki}} T_{d_{ki}}^* \delta_{ij} + c_{11} |\mu|^2 Y_{d_{ki}} Y_{d_{kj}}^* Y_{ddij} \\
& + c_{12} T_{d_{ki}} T_{d_{kj}}^* Y_{ddij}) C_0(\tilde{m}_{d_i}, \tilde{m}_{d_j}, \tilde{m}_{Q_k}) + (c_{13} |\mu|^2 Y_{d_{ji}} Y_{d_{ji}}^* \delta_{jk} + c_{14} |\mu|^2 Y_{d_{ji}} Y_{d_{ki}}^* \bar{Y}_{ddkj} + c_{15} T_{d_{ji}} T_{d_{ji}}^* \delta_{jk} \\
& + c_{16} T_{d_{ji}} T_{d_{ki}}^* \bar{Y}_{ddkj} + c_{17} T_{d_{ji}} T_{d_{ki}}^* \bar{Y}_{uuuj}) C_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}, \tilde{m}_{Q_k}) + (-c_{18} Y_{d_{ji}} Y_{u_{jk}}^* Y_{duik} |\mu|^2 - c_{18} Y_{u_{jk}} Y_{d_{ji}}^* Y_{udki} |\mu|^2 \\
& + c_{18} T_{u_{jk}} T_{d_{ji}}^* Y_{udki} + c_{18} T_{d_{ji}} T_{u_{jk}}^* Y_{duik}) C_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}, \tilde{m}_{u_k}) + (c_{19} |\mu|^2 Y_{u_{ik}} Y_{u_{ik}}^* \delta_{ij} + c_{20} |\mu|^2 Y_{u_{jk}} Y_{u_{ik}}^* \bar{Y}_{uuji} \\
& + c_{21} T_{u_{ik}} T_{u_{ik}}^* \delta_{ij} + c_{22} T_{u_{ik}} T_{u_{jk}}^* \bar{Y}_{uuji} + c_{23} T_{u_{ik}} T_{u_{jk}}^* \bar{Y}_{ddji}) C_0(\tilde{m}_{Q_i}, \tilde{m}_{Q_j}, \tilde{m}_{u_k}) + (c_{24} |\mu|^2 Y_{u_{ij}} Y_{u_{ij}}^* \delta_{jk} \\
& + c_{25} |\mu|^2 Y_{u_{ik}} Y_{u_{ij}}^* Y_{uu_{kj}} + c_{26} T_{u_{ij}} T_{u_{ij}}^* \delta_{jk} + c_{27} T_{u_{ij}} T_{u_{ik}}^* Y_{uu_{jk}}) C_0(\tilde{m}_{Q_i}, \tilde{m}_{u_j}, \tilde{m}_{u_k}) + d_2^{ijkl} D_0(\tilde{m}_{d_i}, \tilde{m}_{d_j}, \tilde{m}_{Q_k}, \tilde{m}_{Q_l}) \\
& + d_3^{ijkl} D_0(\tilde{m}_{Q_i}, \tilde{m}_{Q_j}, \tilde{m}_{u_k}, \tilde{m}_{u_l}) + d_4^{ijkl} D_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}, \tilde{m}_{Q_k}, \tilde{m}_{u_l}). \tag{B11}
\end{aligned}$$

Here, we introduced shorthand notations for the products of two Yukawa coupling matrices:

$$Y_{xyij} \equiv Y_{xil}^\dagger Y_{yij}, \quad \bar{Y}_{xyij} \equiv Y_{xil} Y_{yij}^\dagger, \tag{B12}$$

where $x, y = e, u, d$ and we sum over the internal index l . The coefficients b_n , c_n , and d_n^{ijkl} for each $\lambda_1, \dots, \lambda_4$ are given in Tables V, VI, VIII, and IX.

We finally give the slepton and squark contributions to $\lambda_{6,7}$:

$$\begin{aligned}
\lambda_{6,7}^{\text{ferm}} = & (c'_1 \mu T_{e_{ki}} Y_{e_{ki}}^* \delta_{ij} + c'_2 \mu T_{e_{ki}} Y_{e_{kj}}^* Y_{eeij}) C_0(\tilde{m}_{e_i}, \tilde{m}_{e_j}, \tilde{m}_{l_k}) + (c'_3 \mu T_{e_{ji}} Y_{e_{ji}}^* \delta_{jk} + c'_4 \mu T_{e_{ji}} Y_{e_{kj}}^* \bar{Y}_{eeij}) C_0(\tilde{m}_{e_i}, \tilde{m}_{l_j}, \tilde{m}_{l_k}) \\
& + (c'_5 \mu T_{d_{ki}} Y_{d_{ki}}^* \delta_{ij} + c'_6 \mu T_{d_{ki}} Y_{d_{kj}}^* Y_{ddij}) C_0(\tilde{m}_{d_i}, \tilde{m}_{d_j}, \tilde{m}_{Q_k}) + (c'_7 \mu T_{d_{ji}} Y_{d_{ji}}^* \delta_{jk} + c'_8 \mu T_{d_{ji}} Y_{d_{ki}}^* \bar{Y}_{ddkj}) C_0(\tilde{m}_{d_i}, \tilde{m}_{Q_j}, \tilde{m}_{Q_k}) \\
& + (c'_9 \mu T_{u_{ik}} Y_{u_{ik}}^* \delta_{ij} + c'_{10} \mu T_{u_{ik}} Y_{u_{jk}}^* \bar{Y}_{uuji}) C_0(\tilde{m}_{Q_i}, \tilde{m}_{Q_j}, \tilde{m}_{u_k}) + (c'_{11} \mu T_{u_{ij}} Y_{u_{ij}}^* \delta_{jk} + c'_{12} \mu T_{u_{ij}} Y_{u_{ik}}^* Y_{uu_{jk}}) C_0(\tilde{m}_{Q_i}, \tilde{m}_{u_j}, \tilde{m}_{u_k}) \\
& + d_1^{ijkl} D_0(\tilde{m}_{e_i}, \tilde{m}_{e_j}, \tilde{m}_{l_k}, \tilde{m}_{l_l}) + d_2^{ijkl} D_0(\tilde{m}_{d_i}, \tilde{m}_{d_j}, \tilde{m}_{Q_k}, \tilde{m}_{Q_l}) + d_3^{ijkl} D_0(\tilde{m}_{Q_i}, \tilde{m}_{Q_j}, \tilde{m}_{u_k}, \tilde{m}_{u_l}) \tag{B13}
\end{aligned}$$

where the coefficients c'_n and d_n^{ijkl} are given in Tables V, VI, and VII.

4. Loop functions

In the UV-divergent loop functions we set $\epsilon = (4 - D)/2$. The loop functions are defined as

$$\begin{aligned}
\frac{i}{(4\pi)^2} A_0(m_1) \left(\frac{4\pi}{\mu_0^2} e^{-\gamma_E} \right)^\epsilon &= \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2 - m_1^2}, \\
\frac{i}{(4\pi)^2} B_0(k^2; m_1, m_2) \left(\frac{4\pi}{\mu_0^2} e^{-\gamma_E} \right)^\epsilon &= \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q+k)^2 - m_1^2} \frac{1}{q^2 - m_2^2}, \\
\frac{i}{(4\pi)^2} C_0(m_1, m_2, m_3) \left(\frac{4\pi}{\mu_0^2} e^{-\gamma_E} \right)^\epsilon &= \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2 - m_1^2} \frac{1}{q^2 - m_2^2} \frac{1}{q^2 - m_3^2}, \\
\frac{i}{(4\pi)^2} D_0(m_1, m_2, m_3, m_4) \left(\frac{4\pi}{\mu_0^2} e^{-\gamma_E} \right)^\epsilon &= \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2 - m_1^2} \frac{1}{q^2 - m_2^2} \frac{1}{q^2 - m_3^2} \frac{1}{q^2 - m_4^2}, \\
\frac{i}{(4\pi)^2} W(m_1, m_2) \left(\frac{4\pi}{\mu_0^2} e^{-\gamma_E} \right)^\epsilon &= -\frac{d}{dk^2} \bigg|_{k^2=0} \int \frac{d^D q}{(2\pi)^D} \frac{\text{Tr}[(\not{q} - \not{k})\not{q}]}{((q-k)^2 - m_1^2)(q^2 - m_2^2)}, \tag{B14}
\end{aligned}$$

and we also write

$$B_0(m_1^2, m_2^2) \equiv B_0(0; m_1^2, m_2^2), \quad B'_0(m_1^2, m_2^2) \equiv -\frac{d}{dk^2} \bigg|_{k^2=0} B_0(k^2; m_1^2, m_2^2). \tag{B15}$$

These functions read:

$$\begin{aligned}
A_0(m_1) &= \frac{m_1^2}{\epsilon} + m_1^2 + m_1^2 \log\left(\frac{\mu_0^2}{m_1^2}\right) & B_0(m_1, m_2) &= \frac{1}{\epsilon} + 1 + \frac{m_1^2 \log\left(\frac{\mu_0^2}{m_1^2}\right) - m_2^2 \log\left(\frac{\mu_0^2}{m_2^2}\right)}{m_1^2 - m_2^2} \\
B'_0(m_1, m_2) &= \frac{m_2^4 - m_1^4 + 2m_1^2 m_2^2 \log\left(\frac{m_2^2}{m_1^2}\right)}{2(m_1^2 - m_2^2)^3} & C_0(m_1, m_2, m_3) &= \frac{m_1^2 m_2^2 \log\left(\frac{m_2^2}{m_1^2}\right) + m_3^2 m_2^2 \log\left(\frac{m_3^2}{m_2^2}\right) + m_1^2 m_3^2 \log\left(\frac{m_1^2}{m_3^2}\right)}{(m_1^2 - m_2^2)(m_1^2 - m_3^2)(m_2^2 - m_3^2)} \\
D_0(m_1, m_2, m_3, m_4) &= \sum_{\substack{\{m_1^2, m_2^2, m_3^2, m_4^2\} + \text{cyclic permutations} \\ \{a, b, c, d\}}} \frac{a^2 b c \log\left(\frac{b}{c}\right) + a b^2 c \log\left(\frac{c}{a}\right) + a b c^2 \log\left(\frac{a}{b}\right)}{(a-b)(a-c)(a-d)(b-c)(b-d)(c-d)} \\
W(m_1, m_2) &= \frac{2}{\epsilon} + 2 \log\left(\frac{\mu_0^2}{m_1^2}\right) + \log\left(\frac{m_2^2}{m_1^2}\right) \frac{(2m_2^6 - 6m_1^2 m_2^4)}{(m_1^2 - m_2^2)^3} + \frac{m_1^4 - 6m_2^2 m_1^2 + m_2^4}{(m_1^2 - m_2^2)^2}
\end{aligned} \tag{B16}$$

$$\begin{aligned}
\tilde{D}_2(m_1, m_2, m_3, m_4) &= C_0(m_2, m_3, m_4) + m_1^2 D_0(m_1, m_2, m_3, m_4) \\
\tilde{D}_4(m_1, m_2, m_3, m_4) &= B_0(m_3, m_4) + (m_1^2 + m_2^2) C_0(m_2, m_3, m_4) + m_1^4 D_0(m_1, m_2, m_3, m_4)
\end{aligned} \tag{B17}$$

A further loop function, H_2 , is defined in (34).

APPENDIX C: RENORMALIZATION-GROUP AND BAG PARAMETERS

The standard-model contribution to $B - \bar{B}$ mixing involves the operator $Q_1^{\text{VLL}} = (\bar{b}_L \gamma_\mu q_L)(\bar{b}_L \gamma^\mu q_L)$ of (6). The main new supersymmetric contribution to $B - \bar{B}$ mixing presented in this paper comes with the four-quark operator $Q_1^{\text{SLL}} = (\bar{b}_R q_L)(\bar{b}_R q_L)$ with $q = d$ or $q = s$, see (2). Q_1^{SLL} mixes under renormalization with

$$\tilde{Q}_1^{\text{SLL}} = (\bar{b}_R^i q_L^j)(\bar{b}_R^j q_L^i) \tag{C1}$$

where i, j are color indices. The operators Q_1^{SLL} and $\tilde{Q}_1^{\text{SLL}}$ are widely studied in the context of the width difference $\Delta\Gamma$ among the two mass eigenstates in the $B - \bar{B}$ mixing system and the CP asymmetry a_{fs} in flavor-specific decays [64,65].

Yet the next-to-leading-order (NLO) anomalous dimensions have been calculated for an equivalent operator basis in Ref. [28]. These operators,

TABLE VII. Coefficients of λ_6^{sferm} and λ_7^{sferm} in Eq. (B13).

	λ_6	λ_7
c'_1	$-\frac{g'^2}{2}$	$\frac{g'^2}{2}$
c'_2	1	0
c'_3	$\frac{1}{4}(g'^2 - g'^2)$	$\frac{1}{4}(g'^2 - g'^2)$
c'_4	1	0
c'_5	$-\frac{g'^2}{2}$	$\frac{g'^2}{2}$
c'_6	3	0
c'_7	$\frac{1}{4}(-g'^2 - 3g'^2)$	$\frac{1}{4}(g'^2 + 3g'^2)$
c'_8	3	0
c'_9	$\frac{1}{4}(3g'^2 - g'^2)$	$\frac{1}{4}(g'^2 - 3g'^2)$
c'_{10}	0	3
c'_{11}	g'^2	$-g'^2$
c'_{12}	0	3

$$\begin{aligned}
\bar{Q}_1^{\text{VLL}} &= (\bar{b}_L \gamma_\mu q_L)(\bar{b}_L \gamma^\mu q_L), & \bar{Q}_1^{\text{LR}} &= (\bar{b}_L \gamma_\mu q_L)(\bar{b}_R \gamma^\mu q_R), \\
\bar{Q}_2^{\text{LR}} &= (\bar{b}_R q_L)(\bar{b}_L q_R), & \bar{Q}_1^{\text{SLL}} &= (\bar{b}_R q_L)(\bar{b}_R q_L), \\
\bar{Q}_2^{\text{SLL}} &= -(\bar{b}_R \sigma_{\mu\nu} q_L)(\bar{b}_R \sigma^{\mu\nu} q_L), \\
\bar{Q}_1^{\text{VRR}} &= (\bar{b}_R \gamma_\mu q_R)(\bar{b}_R \gamma^\mu q_R), & \bar{Q}_1^{\text{SRR}} &= (\bar{b}_L q_R)(\bar{b}_L q_R), \\
\bar{Q}_2^{\text{SRR}} &= -(\bar{b}_L \sigma_{\mu\nu} q_R)(\bar{b}_L \sigma^{\mu\nu} q_R),
\end{aligned} \tag{C2}$$

are split into five sectors (VLL, LR, SLL, VRR, SRR), which separately mix under renormalization—note that we define $\sigma_{\mu\nu} = \frac{i}{2}[\gamma_\mu, \gamma_\nu]$. The anomalous dimensions of the VRR and SRR sectors are the same as those of the VLL and SLL sectors, respectively. To define the renormalization scheme for the NLO we first note that we use the $\overline{\text{MS}}$ scheme with anticommuting γ_5 as in [28]. Then we must specify the definition of the evanescent operators which

TABLE VIII. Slepton loop contributions to $\lambda_1^{\text{sferm}} \dots \lambda_4^{\text{sferm}}$ in Eq. (B10).

	λ_1	λ_2	λ_3	λ_4
b_1	$-\frac{g'^4}{4}$	$-\frac{g'^4}{4}$	$\frac{g'^4}{4}$	0
b_2	g'^2	0	$-\frac{g'^2}{2}$	0
b_3	-1	0	0	0
b_4	$\frac{1}{8}(-g'^4 - g'^4)$	$\frac{1}{8}(-g'^4 - g'^4)$	$\frac{1}{8}(g'^4 + g'^4)$	$-\frac{g'^4}{4}$
b_5	$\frac{1}{2}(g'^2 - g'^2)$	0	$\frac{1}{4}(g'^2 - g'^2)$	$\frac{g'^2}{2}$
b_6	-1	0	0	0
c_1	0	$-g'^2$	$\frac{g'^2}{2}$	0
c_2	g'^2	0	$-\frac{g'^2}{2}$	0
c_3	0	0	-1	0
c_4	-2	0	0	0
c_5	0	$\frac{1}{2}(g'^2 - g'^2)$	$\frac{1}{4}(g'^2 - g'^2)$	$-\frac{g'^2}{2}$
c_6	$\frac{1}{2}(g'^2 - g'^2)$	0	$\frac{1}{4}(g'^2 - g'^2)$	$\frac{g'^2}{2}$
c_7	0	0	-1	1
c_8	-2	0	0	0

TABLE IX. Squark loop contributions to $\lambda_1^{\text{sferm}} \dots \lambda_4^{\text{sferm}}$ in Eq. (B11).

	λ_1	λ_2	λ_3	λ_4
b_7	$-\frac{g^4}{12}$	$-\frac{g^4}{12}$	$\frac{g^4}{12}$	0
b_8	g^2	0	$-\frac{g^2}{2}$	0
b_9	-3	0	0	0
b_{10}	0	0	0	-3
b_{11}	$-\frac{g^4}{3}$	$-\frac{g^4}{3}$	$\frac{g^4}{3}$	0
b_{12}	0	$2g^2$	$-g^2$	0
b_{13}	0	-3	0	0
b_{14}	$\frac{1}{24}(-9g^4 - g^4)$	$\frac{1}{24}(-9g^4 - g^4)$	$\frac{1}{24}(9g^4 + g^4)$	$-\frac{3}{4}g^4$
b_{15}	$\frac{1}{2}(3g^2 + g^2)$	0	$\frac{1}{4}(-3g^2 - g^2)$	$\frac{3}{2}g^2$
b_{16}	0	$\frac{1}{2}(3g^2 - g^2)$	$\frac{1}{4}(g^2 - 3g^2)$	$\frac{3}{2}g^2$
b_{17}	-3	0	0	0
b_{18}	0	-3	0	0
b_{19}	0	0	0	-3
c_9	0	$-g^2$	$\frac{g^2}{2}$	0
c_{10}	g^2	0	$-\frac{g^2}{2}$	0
c_{11}	0	0	-3	0
c_{12}	-6	0	0	0
c_{13}	0	$-\frac{1}{2}(3g^2 + g^2)$	$\frac{1}{4}(3g^2 + g^2)$	$-\frac{3}{2}g^2$
c_{14}	0	0	-3	3
c_{15}	$\frac{1}{2}(3g^2 + g^2)$	0	$\frac{1}{4}(-3g^2 - g^2)$	$\frac{3}{2}g^2$
c_{16}	-6	0	0	0
c_{17}	0	0	0	-3
c_{18}	0	0	0	-3
c_{19}	$\frac{1}{2}(g^2 - 3g^2)$	0	$\frac{1}{4}(3g^2 - g^2)$	$-\frac{3}{2}g^2$
c_{20}	0	0	-3	3
c_{21}	0	$\frac{1}{2}(3g^2 - g^2)$	$\frac{1}{4}(g^2 - 3g^2)$	$\frac{3}{2}g^2$
c_{22}	0	-6	0	0
c_{23}	0	0	0	-3
c_{24}	$-2g^2$	0	g^2	0
c_{25}	0	0	-3	0
c_{26}	0	$2g^2$	$-g^2$	0
c_{27}	0	-6	0	0

enter the NLO results as counterterms. In particular for the SLL sector, the evanescent operators of Ref. [28] read:

$$\begin{aligned}
\bar{E}_1^{\text{SLL}} &= (\bar{b}_R^i q_L^j)(\bar{b}_R^j q_L^i) + \frac{1}{2}\bar{Q}_1^{\text{SLL}} - \frac{1}{8}\bar{Q}_2^{\text{SLL}}, \\
\bar{E}_2^{\text{SLL}} &= -(\bar{b}_R^i \sigma_{\mu\nu} q_L^j)(\bar{b}_R^j \sigma^{\mu\nu} q_L^i) - 6\bar{Q}_1^{\text{SLL}} - \frac{1}{2}\bar{Q}_2^{\text{SLL}}, \\
\bar{E}_3^{\text{SLL}} &= (\bar{b}_R^i \gamma_\mu \gamma_\nu \gamma_\rho \gamma_\sigma q_L^j)(\bar{b}_R^j \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma q_L^i) \\
&\quad + (-64 + 96\epsilon)\bar{Q}_1^{\text{SLL}} + (-16 + 8\epsilon)\bar{Q}_2^{\text{SLL}}, \\
\bar{E}_4^{\text{SLL}} &= (\bar{b}_R^i \gamma_\mu \gamma_\nu \gamma_\rho \gamma_\sigma q_L^j)(\bar{b}_R^j \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma q_L^i) \\
&\quad - 64\bar{Q}_1^{\text{SLL}} + (-16 + 16\epsilon)\bar{Q}_2^{\text{SLL}}, \tag{C3}
\end{aligned}$$

where we use $\epsilon \equiv (4 - D)/2$.

The operator basis

$$\begin{aligned}
Q_1^{\text{VLL}} &= \bar{Q}_1^{\text{VLL}}, & Q_1^{\text{LR}} &= \bar{Q}_1^{\text{LR}}, & Q_2^{\text{LR}} &= \bar{Q}_2^{\text{LR}}, \\
Q_1^{\text{SLL}} &= \bar{Q}_1^{\text{SLL}}, & Q_2^{\text{SLL}} &= \bar{Q}_1^{\text{SLL}} = (\bar{b}_R^i q_L^j)(\bar{b}_R^j q_L^i), \\
Q_1^{\text{VRR}} &= \bar{Q}_1^{\text{VRR}}, & Q_1^{\text{SRR}} &= \bar{Q}_1^{\text{SRR}}, \\
Q_2^{\text{SRR}} &= \bar{Q}_1^{\text{SRR}} = (\bar{b}_L^i q_R^j)(\bar{b}_L^j q_R^i), \tag{C4}
\end{aligned}$$

which we adopt in this work agrees with the one of (C2) except for the SLL sector and the SRR sector. The evanescent operators are defined as in Refs. [59,64]:

$$\begin{aligned}
E_1^{\text{SLL}} &= (\bar{b}_R^i \gamma_\mu \gamma_\nu q_L^j)(\bar{b}_R^j \gamma^\mu \gamma^\nu q_L^i) + 8(1 - \epsilon)\bar{Q}_1^{\text{SLL}}, \\
E_2^{\text{SLL}} &= (\bar{b}_R^i \gamma_\mu \gamma_\nu q_L^j)(\bar{b}_R^j \gamma^\mu \gamma^\nu q_L^i) + 8(1 - \epsilon)Q_1^{\text{SLL}}. \tag{C5}
\end{aligned}$$

The hadronic matrix elements in this basis are parametrized in terms of “bag” parameters B_1^{VLL} , $B_1^{\text{SLL'}}$, and $\tilde{B}_1^{\text{SLL'}}$ defined as

$$\begin{aligned}
\langle \bar{B}_q | Q_1^{\text{VLL}}(\mu) | B_q \rangle &= \frac{2}{3} M_{B_q}^2 f_{B_q}^2 B_1^{\text{VLL}}(\mu), \\
\langle \bar{B}_q | Q_1^{\text{SLL}}(\mu) | B_q \rangle &= -\frac{5}{12} M_{B_q}^2 f_{B_q}^2 B_1^{\text{SLL'}}(\mu), \tag{C6} \\
\langle \bar{B}_q | \tilde{Q}_1^{\text{SLL}}(\mu) | B_q \rangle &= \frac{1}{12} M_{B_q}^2 f_{B_q}^2 \tilde{B}_1^{\text{SLL'}}(\mu).
\end{aligned}$$

Here, μ is the renormalization scale at which the matrix element is computed and f_{B_q} is the B_q meson decay constant. While f_{B_s} exceeds f_{B_d} by 10–30%, no nonperturbative calculation finds any dependence of a bag parameter on the flavor of the light valence quark. In the vacuum insertion approximation the bag parameters equal $B_1^{\text{VLL}}(\mu) = 1$ and $B_1^{\text{SLL'}}(\mu) = \tilde{B}_1^{\text{SLL'}}(\mu) = M_{B_q}^2/[m_b(\mu) + m_q(\mu)]^2$. Lattice computations determine the matrix elements at a low scale around 1 GeV and results are quoted for $\mu = \bar{m}_b(\bar{m}_b)$. In order to use the lattice results in our calculation, we need the renormalization-group (RG) evolution of the bag parameters to the high-scale μ_h which is set by the masses of the Higgs bosons exchanged in our $B - \bar{B}$ mixing diagrams. The matrix elements computed on a finite lattice are converted to continuum QCD by a matching calculation. This lattice-continuum matching is only meaningful beyond the leading-order of perturbative QCD. Thus, the dependence of the bag parameters on the chosen (continuum) renormalization scheme must be addressed: The NLO anomalous dimension matrices entering the RG evolution must be defined in the same renormalization scheme as the bag parameters, so that the scheme dependence properly cancels from physical observables. The NLO anomalous dimensions have been calculated for Q_1^{VLL} in Ref. [66]. As said previously, in the case of $(Q_1^{\text{SLL}}, \tilde{Q}_1^{\text{SLL}})$ the NLO anomalous dimensions have been calculated for the equivalent operator basis $(\bar{Q}_1^{\text{SLL}}, \bar{Q}_2^{\text{SLL}})$ with the evanescent operators of (C3) [28].

The purpose of this section is twofold: First, we present the transformation of the results of Ref. [28] to the

$(Q_1^{\text{SLL}}, \tilde{Q}_1^{\text{SLL}})$ basis and the scheme corresponding to the evanescent operators of Eq. (C5), for which lattice groups quote their results. These formulas are useful beyond the need to evolve the bag parameters given at $\mu = m_b$ up to $\mu = \mu_h$: In particular, lattice groups need to evolve $B_1^{\text{SLL}}(\mu)$ and $\tilde{B}_1^{\text{SLL}}(\mu)$ from a scale around 1 GeV up to $\mu = m_b$. Second, we exploit a heavy-quark relation among the bag factors in Eq. (C6) to sharpen the numerical prediction for $B_1^{\text{SLL}}(\mu_h)$ entering the SUSY contribution to $B - \bar{B}$ mixing. While constraints from the heavy-quark limit of QCD have been used to improve the predictions for $\Delta\Gamma$ and a_{fs} [59,64,65], they had escaped attention in studies of new-physics contributions to B physics observables so far.

1. NLO scheme transformation formulas

We decompose the anomalous dimension matrix in the usual way as

$$\gamma = \frac{\alpha_s(\mu)}{4\pi} \gamma^{(0)} + \left(\frac{\alpha_s(\mu)}{4\pi} \right)^2 \gamma^{(1)} + \mathcal{O}(\alpha_s^3). \quad (\text{C7})$$

The NLO correction $\gamma^{(1)}$ has been computed for the basis $(\tilde{Q}_1^{\text{SLL}}, \tilde{Q}_2^{\text{SLL}})$ in Ref. [28]. In four dimensions it is related to the basis (C4) by a simple Fierz identity:

$$\tilde{Q} = \left(\frac{Q_1^{\text{SLL}}}{\tilde{Q}_1^{\text{SLL}}} \right)^{D=4} \hat{R} \left(\frac{\tilde{Q}_1^{\text{SLL}}}{\tilde{Q}_2^{\text{SLL}}} \right) = \hat{R} \tilde{Q}, \quad (\text{C8})$$

where $\hat{R}$ is given in Eq. (C10) below.

Yet in D dimensions our change of basis involves a rotation of the operator basis—including the evanescent operators $\tilde{E} = (\tilde{E}_1^{\text{SLL}}, \tilde{E}_2^{\text{SLL}})^T$ —and a change of the renormalization scheme. We follow Ref. [67] and write the rotation as¹⁴

$$\tilde{Q} = \hat{R}(\tilde{Q} + \hat{W}\tilde{E}), \quad \tilde{E} = \hat{M}(\epsilon\hat{U}\tilde{Q} + [\hat{1} + \epsilon\hat{U}\hat{W}]\tilde{E}), \quad (\text{C9})$$

with

$$\begin{aligned} \hat{R} &= \begin{pmatrix} 1 & 0 \\ -\frac{1}{2} & \frac{1}{8} \end{pmatrix}, & \hat{W} &= \begin{pmatrix} 0 & 0 \\ 8 & 0 \end{pmatrix}, \\ \hat{U} &= \begin{pmatrix} \frac{1}{4} & -\frac{1}{8} \\ 8 & -\frac{1}{4} \end{pmatrix}, & \hat{M} &= \begin{pmatrix} -8 & 0 \\ -4 & 1 \end{pmatrix}. \end{aligned} \quad (\text{C10})$$

The information on the definition of the evanescent operators in Eqs. (C3) and (C4) is contained in the matrices $\hat{U}$ and $\hat{M}$. Now Eq. (C9) corresponds to a finite renormalization with renormalization constants [67]

¹⁴Two more evanescent operators (called $\tilde{E}_3^{\text{SLL}}$ and $\tilde{E}_4^{\text{SLL}}$ in Ref. [28]) must be specified to fully define the scheme of the calculated $\tilde{\gamma}^{(1)}$. This information enters the matrix $\hat{U}$ in Eq. (C10). We choose to add $\tilde{E}_3^{\text{SLL}}$ and $\tilde{E}_4^{\text{SLL}}$ also to the evanescent operators of Eq. (C3), so that we can in practice work with the change of basis defined in Eqs. (C9) and (C10).

$$\hat{Z}_{QQ}^{(1,0)} = \hat{R}[\hat{W}\hat{Z}_{EQ}^{(1,0)} - (\hat{Z}_{QE}^{(1,1)} + \hat{W}\hat{Z}_{EE}^{(1,1)} - \frac{1}{2}\tilde{\gamma}^{(0)}\hat{W})\hat{U}]\hat{R}^{-1}. \quad (\text{C11})$$

While the one-loop anomalous dimension matrix is just rotated, the two-loop anomalous dimension matrix undergoes an additional scheme transformation:

$$\begin{aligned} \gamma^{(0)} &= \hat{R}\tilde{\gamma}^{(0)}\hat{R}^{-1}, \\ \gamma^{(1)} &= \hat{R}\tilde{\gamma}^{(1)}\hat{R}^{-1} - [\hat{Z}_{QQ}^{(1,0)}, \gamma^{(0)}] - 2\beta^{(0)}\hat{Z}_{QQ}^{(1,0)}, \end{aligned} \quad (\text{C12})$$

with the one-loop operator renormalization constants

$$\begin{aligned} \hat{Z}_{QE}^{(1,1)} &= \begin{pmatrix} 0 & \frac{1}{2} \\ -8 & -8 \end{pmatrix}, & \hat{Z}_{EE}^{(1,1)} &= \begin{pmatrix} 2 & \frac{11}{6} \\ -\frac{16}{3} & -\frac{44}{3} \end{pmatrix}, \\ \hat{Z}_{EQ}^{(1,0)} &= \begin{pmatrix} \frac{37}{12} & \frac{29}{48} \\ -\frac{73}{3} & -\frac{1}{12} \end{pmatrix}. \end{aligned} \quad (\text{C13})$$

We can now calculate the new two-loop anomalous dimension matrix γ from the NLO anomalous dimension matrix $\tilde{\gamma}$ of Ref. [28],

$$\begin{aligned} \tilde{\gamma}^{(0)} &= \begin{pmatrix} -10 & \frac{1}{6} \\ -40 & \frac{34}{3} \end{pmatrix}, \\ \tilde{\gamma}_{[27]}^{(1)} &= \begin{pmatrix} -\frac{1459}{9} + \frac{74}{9}f & -\frac{35}{36} - \frac{1}{54}f \\ -\frac{6332}{9} + \frac{584}{9}f & \frac{2065}{9} - \frac{394}{27}f \end{pmatrix} \end{aligned} \quad (\text{C14})$$

We obtain

$$\gamma^{(0)} = \begin{pmatrix} -\frac{28}{3} & \frac{4}{3} \\ \frac{16}{3} & \frac{32}{3} \end{pmatrix}, \quad \gamma^{(1)} = \begin{pmatrix} -\frac{260}{3} + \frac{88}{27}f & -\frac{44}{3} + \frac{8}{27}f \\ \frac{242}{3} - \frac{76}{27}f & 198 - \frac{332}{27}f \end{pmatrix}. \quad (\text{C15})$$

Here, f denotes the number of active flavors and $\gamma^{(0)}$ coincides with the result in [64]. As a check, we have calculated the result of Eq. (C15) also in a different way: It is possible to define evanescent operators such that the Fierz identity holds for the one-loop matrix elements. This choice fixes the definitions of both E_1^{SLL} and E_2^{SLL} in Eq. (C5) and of the evanescent operators on the $(\tilde{Q}_1^{\text{SLL}}, \tilde{Q}_2^{\text{SLL}})$ basis. (One of the latter operators equals ϵ times a physical operator. Its impact is equivalent to a finite multiplicative renormalization of $\tilde{Q}_1^{\text{SLL}}$.) In this approach, one can simply rotate $\gamma^{(1)}$ in the same way as $\gamma^{(0)}$ in Eq. (C12). Finally, the result is transformed to the scheme of Ref. [28] using the scheme transformation formula of Ref. [68].

Next, we calculate the matrices governing the RG evolution in the $(Q_1^{\text{SLL}}, \tilde{Q}_1^{\text{SLL}})$ basis. The bag factors at the scale μ_h are obtained from those at the low scale $\mu_b = \mathcal{O}(m_b)$ via

$$\begin{pmatrix} -5B_1^{\text{SLL}}(\mu_h) \\ \tilde{B}_1^{\text{SLL}}(\mu_h) \end{pmatrix} = U(\mu_b, \mu_h)^T \begin{pmatrix} -5B_1^{\text{SLL}}(\mu_b) \\ \tilde{B}_1^{\text{SLL}}(\mu_b) \end{pmatrix} \quad (\text{C16})$$

In the spirit of [29], we write the evolution matrix as

$$U(\mu_b, \mu_h) = U^{(0)} \left(\frac{\alpha_s(\mu_h)}{\alpha_s(\mu_b)} \right) + \frac{\alpha_s(\mu_b)}{4\pi} \Delta U \left(\frac{\alpha_s(\mu_h)}{\alpha_s(\mu_b)} \right), \quad (\text{C17})$$

where $U^{(0)}$ is the LO evolution matrix and the NLO correction reads

$$\Delta U(\eta) = J_f U^{(0)}(\eta) - \eta U^{(0)}(\eta) J_f. \quad (\text{C18})$$

The 2×2 matrix J_f is calculated from the anomalous dimension matrix γ [69]. We only need J_5 , since we run with 5 active flavors to the scale μ_h . For applications in kaon physics, one also involves J_4 and J_3 . We quote all three matrices here, so that the formulas of Ref. [29] can be easily extended to the $(Q_1^{\text{SLL}}, \tilde{Q}_1^{\text{SLL}})$ basis:

$$J_5 = \begin{pmatrix} 1.474 & 0.707 \\ 0.306 & -5.350 \end{pmatrix}, \quad J_4 = \begin{pmatrix} 0.964 & 1.452 \\ 0.375 & -4.982 \end{pmatrix},$$

$$J_3 = \begin{pmatrix} 0.652 & 2.597 \\ 0.421 & -4.804 \end{pmatrix}. \quad (\text{C19})$$

We quote handy formulas for the five-flavor evolution matrix, similarly to Ref. [29]:

$$U_{f=5}^{(0)}(\eta) = \begin{pmatrix} 0.9831 & -0.2577 \\ -0.0644 & 0.0169 \end{pmatrix} \eta^{-0.6315} + \begin{pmatrix} 0.0169 & 0.2577 \\ 0.0644 & 0.9831 \end{pmatrix} \eta^{0.7184}. \quad (\text{C20})$$

The NLO correction reads:

$$\Delta U_{f=5}(\eta) = \begin{pmatrix} 1.4040 - 1.3707\eta & -0.3680 - 2.0731\eta \\ 0.6454 + 0.0898\eta & -0.1692 + 0.1358\eta \end{pmatrix} \eta^{-0.6315} + \begin{pmatrix} 0.0704 - 0.1037\eta & 1.0746 + 1.3665\eta \\ -0.3395 - 0.3958\eta & -5.1807 + 5.2141\eta \end{pmatrix} \eta^{0.7184}. \quad (\text{C21})$$

In our numerical analysis we drop the terms which are linear in η in the two matrices in Eq. (C21), because they are scheme-dependent. The scheme dependence of these terms cancels with that of the NLO QCD corrections to the $B - \bar{B}$ mixing diagrams with SUSY Higgs exchange. Yet these QCD corrections are unknown.

2. Hadronic matrix elements and heavy-quark relations

The three bag factors B_1^{VLL} , $B_1^{\text{SLL}}(\mu_b)$, and $\tilde{B}_1^{\text{SLL}}(\mu_b)$ obey a heavy-quark relation [65]:

$$B_1^{\text{SLL}}(\mu_b) = \frac{4}{5} \alpha_2(\mu_b) B_1^{\text{VLL}} + \frac{1}{5} \alpha_1(\mu_b) \tilde{B}_1^{\text{SLL}} + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}}{m_b}\right). \quad (\text{C22})$$

Here $\alpha_1(\mu)$ and $\alpha_2(\mu)$ comprise NLO QCD corrections [59,64]:

$$\alpha_1(\mu_b) = 1 + \frac{\alpha_s(\mu_b)}{4\pi} \left(16 \log \frac{\mu_b}{m_b} + 8 \right),$$

$$\alpha_2(\mu_b) = 1 + \frac{\alpha_s(\mu_b)}{4\pi} \left(8 \log \frac{\mu_b}{m_b} + \frac{26}{3} \right). \quad (\text{C23})$$

These values are specific to the definition of the evanescent operators as in Eq. (C5). As mentioned in Appendix C1, this definition allows to maintain the validity of Fierz identities at the loop level. Such a definition is preferred, if the bag factors are meant to parametrize the deviation of matrix elements from the vacuum insertion approximation (VIA), because the calculation of matrix elements in VIA approximation involves a Fierz transformation. In particular the choice in Eq. (C5) is crucial for Eq. (C22) to hold in the limit of a large number N_C of colors [64].

The bag factor B_1^{VLL} is very well studied in lattice QCD, so that it is worthwhile to study the constraint on the other bag factors when Eq. (C22) is combined with lattice results for B_1^{VLL} . Indeed, one can use Eq. (C22) to pinpoint the ratio

$$\frac{B_1^{\text{SLL}}(\mu_b)}{B_1^{\text{VLL}}(\mu_b)} = 0.93 + 0.23 \frac{\tilde{B}_1^{\text{SLL}}(\mu_b)}{B_1^{\text{VLL}}(\mu_b)} + (0.23 \pm 0.05) \frac{1}{B_1^{\text{VLL}}(\mu_b)} \quad (\text{C24})$$

quite precisely, even if $\tilde{B}_1^{\text{SLL}}$ is only poorly known, because its coefficient in Eq. (C24) is small. The last term in Eq. (C24) quantifies the Λ_{QCD}/m_b corrections, see [59] for details. The lattice results of [70] have been combined in Ref. [59] to

$$B_1^{\text{VLL}}(\mu_b) = 0.85 \pm 0.06 \quad \text{and} \quad \tilde{B}_1^{\text{SLL}}(\mu_b) = 1.41 \pm 0.12. \quad (\text{C25})$$

Inserting these values into Eq. (C24) yields

$$\frac{B_1^{\text{SLL}}(\mu_b)}{B_1^{\text{VLL}}(\mu_b)} = 1.57 \pm 0.08, \quad (\text{C26})$$

which is consistent with the direct determination

$$B_1^{\text{SLL}}(\mu_b) = 1.34 \pm 0.12 \quad (\text{C27})$$

from the lattice [70].

We are now in the position to accurately predict the bag factors at the high-scale μ_h . Choosing $\mu_h = \bar{m}_t(\bar{m}_t) = 164$ GeV, $\alpha_s(M_Z) = 0.1189$ and $\bar{m}_b(\bar{m}_b) = 4.2$ GeV and using Eqs. (C26) and (C27) we find

$$\begin{aligned}
B_1^{\text{SLL}'}(m_t) &= 1.62B_1^{\text{SLL}'}(m_b) + 0.01\tilde{B}_1^{\text{SLL}'}(m_b) \\
&= (2.54 \pm 0.13)B_1^{\text{VLL}}(m_b) + 0.01 \\
\tilde{B}_1^{\text{SLL}'}(m_t) &= 1.29B_1^{\text{SLL}'}(m_b) + 0.54\tilde{B}_1^{\text{SLL}'}(m_b) \\
&= (2.03 \pm 0.10)B_1^{\text{VLL}}(m_b) + 0.77 \pm 0.07
\end{aligned} \tag{C28}$$

Here, we have omitted the scheme-dependent terms proportional to η in Eq. (C21). The small (2, 1) element of $U_{f=5}^{(0)}$ in Eq. (C20) ensures that $\tilde{B}_1^{\text{SLL}'}(m_b)$ is inessential for $B_1^{\text{SLL}'}(m_t)$. One realises from Eq. (C28) that the uncertainty of the high-scale bag factors stems almost completely from the error of the lattice result for $B_1^{\text{VLL}}(m_b)$.

Switching finally to the P_i 's defined in Eq. (8) we get

$$\begin{aligned}
P_1^{\text{SLL}} &= -\frac{5}{8}B_1^{\text{SLL}'}(m_t) \\
&= -(1.59 \pm 0.08)B_1^{\text{VLL}}(m_b) - 0.01 \\
&= -1.36 \pm 0.12 \\
P_1^{\text{VLL}} &= B_1^{\text{VLL}}(m_t) = 0.83B_1^{\text{VLL}}(m_b) = 0.71 \pm 0.05.
\end{aligned} \tag{C29}$$

In the last line, the full NLO result of [66] has been used. We do not need $\tilde{P}_1^{\text{SLL}} = \tilde{B}_1^{\text{SLL}'}(m_t)/8$ for our analysis. Parity ensures that Q_1^{SLL} and the chirality-flipped operator Q_1^{SRR} defined in Eq. (7) have the same matrix element, i.e. $P_1^{\text{SRR}} = P_1^{\text{SLL}}$.

Finally, we compute P_2^{LR} using the formulas of Ref. [29] with the bag factors of Bećirević *et al.* [70]. This time the conversion between the bases of Ref. [29,70] is straightforward, since the renormalization scheme used in

Refs. [28,29] respects the Fierz symmetry and lattice results are already quoted for this scheme. The result is

$$P_2^{\text{LR}} = 3.2 \pm 0.2. \tag{C30}$$

The number in Eq. (C30) is significantly larger than $P_2^{\text{LR}} = 2.46$ quoted in Ref. [29], because our value for m_b is smaller and the lattice bag factors are larger than 1. The error in Eq. (C30) does not include the systematic error from the quenching approximation.

APPENDIX D: TRILINEAR HIGGS COUPLINGS

The trilinear terms of the effective Lagrangian at $\tan\beta = \infty$ introduced in Sec. IIc read

$$\begin{aligned}
V_{\text{tb}}^{(3)} &= \frac{v}{\sqrt{2}} \left\{ \frac{\lambda_2}{\sqrt{2}} r_u (r_u^2 + (G^0)^2 + 2|h_u^+|^2) + \sqrt{2}\lambda_3 r_u H_d^\dagger H_d \right. \\
&\quad + \lambda_4 (\sqrt{2}r_u |h_d^-|^2 + [h_d^0 h_d^- h_u^+ + \text{H.c.}]) \\
&\quad + \left[\lambda_5 h_d^{0*} \left(\frac{1}{\sqrt{2}} h_d^{0*} (r_u + iG^0) - h_d^- h_u^+ \right) + \lambda_6 (H_d^\dagger H_d) h_d^{0*} \right. \\
&\quad + \lambda_7 \left(h_d^{0*} \left[\frac{3}{2} r_u^2 + i r_u G^0 + \frac{1}{2} (G^0)^2 + |h_u^+|^2 \right] \right. \\
&\quad \left. \left. - \sqrt{2} r_u h_u^+ h_d^- \right) + \text{H.c.} \right] \left. \right\}. \tag{D1}
\end{aligned}$$

Again, the first two lines respect the $U(1)$ symmetry introduced in Sec. IIc, while the last two lines break it, and the breaking is proportional to loop-induced couplings. Finally, the quartic Lagrangian is obtained from the quartic terms in Eq. (12) by substituting $H_u \rightarrow (h_u^+, \frac{1}{\sqrt{2}}\phi_u^0)$ and $H_d \rightarrow (h_d^{0*}, -h_d^-)$. Also there, only λ_5 , λ_6 and λ_7 break the symmetry.

-
- [1] L. J. Hall, R. Rattazzi, and U. Sarid, *Phys. Rev. D* **50**, 7048 (1994); R. Hempfling, *Phys. Rev. D* **49**, 6168 (1994); M. Carena, M. Olechowski, S. Pokorski, and C. E. M. Wagner, *Nucl. Phys. B* **426**, 269 (1994); T. Blazek, S. Raby, and S. Pokorski, *Phys. Rev. D* **52**, 4151 (1995).
 - [2] C. Hamzaoui, M. Pospelov, and M. Toharia, *Phys. Rev. D* **59**, 095005 (1999).
 - [3] K. S. Babu and C. F. Kolda, *Phys. Rev. Lett.* **84**, 228 (2000).
 - [4] G. Isidori and A. Retico, *J. High Energy Phys.* **11** (2001) 001.
 - [5] A. J. Buras, P. H. Chankowski, J. Rosiek, and L. Slawianowska, *Phys. Lett. B* **546**, 96 (2002); *Nucl. Phys. B* **659**, 3 (2003).
 - [6] G. Isidori and P. Paradisi, *Phys. Lett. B* **639**, 499 (2006).
 - [7] M. Carena, D. Garcia, U. Nierste, and C. E. M. Wagner, *Nucl. Phys. B* **577**, 88 (2000).
 - [8] L. J. Hall and L. Randall, *Phys. Rev. Lett.* **65**, 2939 (1990).
 - [9] M. Ciuchini, G. Degrossi, P. Gambino, and G. F. Giudice, *Nucl. Phys. B* **534**, 3 (1998).
 - [10] A. Ali and D. London, *Eur. Phys. J. C* **9**, 687 (1999).
 - [11] A. J. Buras, P. Gambino, M. Gorbahn, S. Jager, and L. Silvestrini, *Nucl. Phys. B* **592**, 55 (2001).
 - [12] A. J. Buras, P. Gambino, M. Gorbahn, S. Jager, and L. Silvestrini, *Phys. Lett. B* **500**, 161 (2001).
 - [13] G. D'Ambrosio, G. F. Giudice, G. Isidori, and A. Strumia, *Nucl. Phys. B* **645**, 155 (2002).
 - [14] G. Isidori, F. Mescia, P. Paradisi, C. Smith, and S. Trine, *J. High Energy Phys.* **08** (2006) 064.
 - [15] W. Altmannshofer, A. J. Buras, and D. Guadagnoli, *J. High Energy Phys.* **11** (2007) 065.
 - [16] G. Colangelo, E. Nikolidakis, and C. Smith, *Eur. Phys. J. C* **59**, 75 (2008).
 - [17] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963); M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).

- [18] Y. Okada, M. Yamaguchi, and T. Yanagida, *Prog. Theor. Phys.* **85**, 1 (1991).
- [19] H. E. Haber and R. Hempfling, *Phys. Rev. Lett.* **66**, 1815 (1991).
- [20] J. R. Ellis, G. Ridolfi, and F. Zwirner, *Phys. Lett. B* **257**, 83 (1991).
- [21] A. Brignole, *Phys. Lett. B* **281**, 284 (1992).
- [22] P. H. Chankowski, S. Pokorski, and J. Rosiek, *Nucl. Phys. B* **423**, 437 (1994).
- [23] A. Dabelstein, *Z. Phys. C* **67**, 495 (1995).
- [24] A. Pilaftsis and C. E. M. Wagner, *Nucl. Phys. B* **553**, 3 (1999); M. S. Carena, J. R. Ellis, A. Pilaftsis, and C. E. M. Wagner, *Nucl. Phys. B* **586**, 92 (2000); **625**, 345 (2002).
- [25] J. R. Ellis, J. S. Lee, and A. Pilaftsis, *Phys. Rev. D* **76**, 115011 (2007).
- [26] A. Freitas, E. Gasser, and U. Haisch, *Phys. Rev. D* **76**, 014016 (2007).
- [27] S. Trine, [arXiv:0710.4955](https://arxiv.org/abs/0710.4955).
- [28] A. J. Buras, M. Misiak, and J. Urban, *Nucl. Phys. B* **586**, 397 (2000).
- [29] A. J. Buras, S. Jager, and J. Urban, *Nucl. Phys. B* **605**, 600 (2001).
- [30] see e.g., H. E. Haber, in *Perspectives on Higgs physics II*, edited by Gordon L. Kane (World Scientific, Singapore, 1997), and references therein.
- [31] A. Freitas and D. Stockinger, *Phys. Rev. D* **66**, 095014 (2002).
- [32] G. Gamberini, G. Ridolfi, and F. Zwirner, *Nucl. Phys. B* **331**, 331 (1990).
- [33] S. Weinberg, *Phys. Rev. D* **7**, 2887 (1973).
- [34] N. K. Nielsen, *Nucl. Phys. B* **101**, 173 (1975).
- [35] B. W. Lee and J. Zinn-Justin, *Phys. Rev. D* **5**, 3137 (1972); **8**, 4654(E) (1973).
- [36] Y. Yamada, *Phys. Lett. B* **530**, 174 (2002).
- [37] M. Frank, T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, *J. High Energy Phys.* **02** (2007) 047.
- [38] J. S. Lee, M. Carena, J. Ellis, A. Pilaftsis, and C. E. M. Wagner, *Comput. Phys. Commun.* **180**, 312 (2009).
- [39] I. Jack, D. R. T. Jones, S. P. Martin, M. T. Vaughn, and Y. Yamada, *Phys. Rev. D* **50**, R5481 (1994).
- [40] M. Beneke, P. Ruiz-Femenia, and M. Spinrath, *J. High Energy Phys.* **01**, (2009) 031.
- [41] J. A. Casas and S. Dimopoulos, *Phys. Lett. B* **387**, 107 (1996).
- [42] S. R. Choudhury and N. Gaur, *Phys. Lett. B* **451**, 86 (1999); P. H. Chankowski and L. Slawianowska, *Phys. Rev. D* **63**, 054012 (2001); C. S. Huang, W. Liao, Q. S. Yan, and S. H. Zhu, *Phys. Rev. D* **63**, 114021 (2001); **64**, 059902(E) (2001); C. Bobeth, T. Ewerth, F. Kruger, and J. Urban, *Phys. Rev. D* **64**, 074014 (2001).
- [43] S. Marchetti, S. Mertens, U. Nierste, and D. Stockinger, *Phys. Rev. D* **79**, 013010 (2009).
- [44] H. Itoh, S. Komine, and Y. Okada, *Prog. Theor. Phys.* **114**, 179 (2005).
- [45] A. Abulencia *et al.* (CDF Collaboration), CDF Public Note 8176.
- [46] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **97**, 242003 (2006).
- [47] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **100**, 101802 (2008).
- [48] CDF Collaboration, CDF Public Note 9892.
- [49] C. Amsler *et al.* (Particle Data Group), *Phys. Lett. B* **667**, 1 (2008).
- [50] G. Degross, P. Gambino, and P. Slavich, *Comput. Phys. Commun.* **179**, 759 (2008).
- [51] G. W. Bennett *et al.* (Muon G-2 Collaboration), *Phys. Rev. D* **73**, 072003 (2006).
- [52] M. Davier, "Final report of the E821 muon anomalous magnetic moment measurement at BNL", TAU08, <http://tau08.inp.nsk.su/prog.php>.
- [53] E. Barberio *et al.* (Heavy Flavor Averaging Group), [arXiv:0808.1297](https://arxiv.org/abs/0808.1297) and online update at <http://www.slac.stanford.edu/xorg/hfag/>.
- [54] M. Misiak *et al.*, *Phys. Rev. Lett.* **98**, 022002 (2007).
- [55] S. Trine, [arXiv:0810.3633](https://arxiv.org/abs/0810.3633); U. Nierste, S. Trine, and S. Westhoff, *Phys. Rev. D* **78**, 015006 (2008).
- [56] J. K. Parry, *Mod. Phys. Lett. A* **21**, 2853 (2006).
- [57] W. Altmannshofer, A. J. Buras, D. Guadagnoli, and M. Wick, *J. High Energy Phys.* **12** (2007) 096.
- [58] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **100**, 161802 (2008); V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **101**, 241801 (2008).
- [59] A. Lenz and U. Nierste, *J. High Energy Phys.* **06** (2007) 072.
- [60] M. Bona *et al.* (UTfit Collaboration), *PMC Phys. A* **3**, 6 (2009).
- [61] E. Lunghi and A. Soni, *Phys. Lett. B* **666**, 162 (2008); A. J. Buras and D. Guadagnoli, *Phys. Rev. D* **78**, 033005 (2008).
- [62] P. Skands *et al.*, *J. High Energy Phys.* **07** (2004) 036; *Comput. Phys. Commun.* **180**, 8 (2009).
- [63] J. Rosiek, *Phys. Rev. D* **41**, 3464 (1990), updated at <http://www.fuw.edu.pl/~rosiek/physics/prd41.html>
- [64] M. Beneke, G. Buchalla, C. Greub, A. Lenz, and U. Nierste, *Phys. Lett. B* **459**, 631 (1999).
- [65] M. Beneke, G. Buchalla, and I. Dunietz, *Phys. Rev. D* **54**, 4419 (1996); M. Ciuchini, E. Franco, V. Lubicz, F. Mescia, and C. Tarantino, *J. High Energy Phys.* **08** (2003) 031; M. Beneke, G. Buchalla, A. Lenz, and U. Nierste, *Phys. Lett. B* **576**, 173 (2003).
- [66] A. J. Buras, M. Jamin, and P. H. Weisz, *Nucl. Phys. B* **347**, 491 (1990).
- [67] M. Gorbahn and U. Haisch, *Nucl. Phys. B* **713**, 291 (2005).
- [68] S. Herrlich and U. Nierste, *Nucl. Phys. B* **455**, 39 (1995); **476**, 27 (1996).
- [69] A. J. Buras, M. Jamin, and M. E. Lautenbacher, *Nucl. Phys. B* **408**, 209 (1993).
- [70] D. Becirevic, V. Gimenez, G. Martinelli, M. Papinutto, and J. Reyes, *J. High Energy Phys.* **04** (2002) 025; S. Aoki *et al.* (JLQCD Collaboration), *Phys. Rev. Lett.* **91**, 212001 (2003); N. Yamada *et al.* (JLQCD Collaboration), *Nucl. Phys. B, Proc. Suppl.* **106-107**, 397 (2002).