

**D-foam-induced flavor condensates and breaking of supersymmetry in free Wess-Zumino fluids**

Nick E. Mavromatos\*

*CERN, Theory Division, CH-1211 Geneva 23, Switzerland*

Sarben Sarkar and Walter Tarantino

*King's College London, Department of Physics, Strand, London WC2R 2LS, United Kingdom*

(Received 23 February 2011; published 19 August 2011)

Recently [N.E. Mavromatos and S. Sarkar, *New J. Phys.* **10**, 073009 (2008) ; N.E. Mavromatos, S. Sarkar, and W. Tarantino, *Phys. Rev. D* **80**, 084046 (2009)], we argued that a particular model of string-inspired quantum space-time foam (D-foam) may induce oscillations and mixing among flavored particles. As a result, rather than the mass-eigenstate vacuum, the correct ground state to describe the underlying dynamics is the *flavor vacuum*, proposed some time ago by Blasone and Vitiello as a description of quantum field theories with mixing. At the microscopic level, the breaking of target-space supersymmetry is induced in our space-time foam model by the relative transverse motion of brane defects. Motivated by these results, we show that the flavor vacuum, introduced through an inequivalent representation of the canonical (anti-) commutation relations, provides a vehicle for the breaking of supersymmetry at a low-energy effective field-theory level; on considering the flavor-vacuum expectation value of the energy-momentum tensor and comparing with the form of a perfect relativistic fluid, it is found that the bosonic sector contributes as dark energy while the fermion contribution is like dust. This indicates a strong and novel breaking of supersymmetry, of a nonperturbative nature, which may characterize the low-energy field theory of certain quantum-gravity models.

DOI: [10.1103/PhysRevD.84.044050](https://doi.org/10.1103/PhysRevD.84.044050)

PACS numbers: 04.62.+v, 14.60.Pq

**I. INTRODUCTION**

We have, in an earlier work [1,2], investigated the behavior of an effective vacuum condensate in a field-theory arising from a brane model of space-time foam (*D-Particle Foam Model*). Within a braneworld context (cf. [1] and citations therein), this condensate is consistent with a flavor vacuum [3] of a string model (in a low-energy limit) that includes string excitations corresponding to particles of different flavors (on a D3 brane Universe) and a foam of D-particles (in the bulk); such a condensate was described through the formalism of quantum field theory (QFT) on a curved (classical) background [4] and the Blasone-Vitiello (BV) formalism for *flavor mixing* [3].

In our earliest investigation [1], a theory of two real scalar fields with flavor mixing, on a specific expanding  $1 + 1$ -dimensional universe, was considered. The equation of state  $p = w\rho$  (with  $p$  pressure and  $\rho$  energy density), associated with the flavor vacuum was found to have  $w = -1$  (on using a suitable regularization). In [2] we claimed that such a behavior would characterize also the  $3 + 1$ -dimensional case, in a generic Friedmann-Robertson-Walker (FRW) universe. Moreover, we analyzed a theory with two Dirac fields (complex spin  $1/2$ ) on a  $3 + 1$ -dimensional FRW universe. In this context, the  $w$  for the equation of state of the flavor vacuum was found to be in the range  $-1/3 < w < 0$  (the precise value being

determined by the cutoff for the momenta imposed on the QFT theory), leading to a contracting universe. Therefore, it has been advocated that this vacuum condensate in a realistic theory might contribute to the Dark Energy fluid that has been suggested as an explanation of the current acceleration of the Universe. This is in agreement with the general arguments of Ref. [5], except that the analysis of the model of [1,2] was done more consistently in a curved background, entailing particle creation and a dynamically generated cutoff scale. However the bosonic and fermionic particles were not considered as part of the same supersymmetric multiplet. Hence for an observer on the D3 brane a detailed signature of any supersymmetry (SUSY)-breaking needs to be sought. This will be effected by considering properties of a flavor vacuum state in a supersymmetric context.

In our microscopic brane model of [1,2], the seeds of microscopic Lorentz violation are provided by the recoil of D-particles, interacting with flavor matter states. An effective low-energy QFT description of this stringy picture needs to reflect this. Hence it is important to note at this point that a QFT, based on the flavor Fock space of the BV formalism, is necessarily Lorentz-violating. As remarked in [2], this distinct behavior in the collective properties between fermions and bosons is a novel nonperturbative measure of SUSY-breaking, which is consistent with the Lorentz-violating nature of the BV formalism.

In this article we embark on a detailed investigation of this point, considering as a simple, but instructive, example a *supersymmetric fluid* free Wess-Zumino (WZ) model

---

\*On leave from King's College London, Department of Physics, Strand, London WC2R 2LS, UK.

with flavor mixing.<sup>1</sup> The structure of the article is as follows: in Sec. II we introduce the reader to the concept of the medium of the D-particle space-time foam, which constitutes a background space-time whose interactions with open-string flavored excitations in the brane world necessitate the introduction of the flavor Fock states as the physically correct excitation spectrum. This illustrates that the interactions of a model can pick up the flavor vacuum as the physically correct vacuum, despite the fact that the latter may not be the minimal energy state in an effective theory. In the BV formalism this state has been associated with the vacuum state associated with the Fock space of an inequivalent representation of the field canonical commutation relations.

In this work, the two-flavor generalization of the free supersymmetric Wess-Zumino model will be viewed as an “effective” field theory of “free” fields, however now viewed as (Fock) excitations of the “flavored” vacuum. In Sec. II we motivate the use of the flavored supersymmetric WZ model as a prototype effective low-energy field-theory, describing flavor mixing induced by quantum-gravity space-time foam, in particular, stringy D-particle foam [7]. In Sec. III the model is introduced, with flavor mixing implemented through the BV formalism. In Sec. IV we outline the steps of the calculations of the expectation value of the energy-momentum tensor in the flavor vacuum. Further calculational details appear in Sec. IVA, and we present our results and comments in Secs. IV B and IV C. Not only a positive flavor-vacuum energy is demonstrated, but the equations of state for the fermion and boson parts are different, the bosons resembling a dark energy fluid (of cosmological type), while the fermions behave like dust. This indicates a strong breaking of SUSY, of a novel type, going beyond the conventional supersymmetry breaking situations characterized only by a nonzero vacuum energy.

Other scenarios, following from nonstandard model physics, which may lead to flavor condensates (and possible SUSY-breaking) will be considered in a forthcoming publication [8].

<sup>1</sup>A work [6] has appeared contemporaneously with the first draft of this article, available in the archives, where a free flavored WZ model has also been analyzed from the point of view of SUSY-breaking by the flavor condensate. However, the authors of that work, in contrast to our analysis here, examined only the energy part of the stress tensor (with results similar to ours) but did not analyze the pressure part of the fluid and thus the relevant equation of state. The latter requires a technically much more involved and highly nontrivial operator analysis, and, in fact, is directly relevant for a proper classification of the type of SUSY-breaking involved, as well as the phenomenology and, in particular, the cosmology of such models, in which context their true significance appears. Moreover, in view of our motivation outlined in Sec. II, the philosophy of our approach and the reasoning for the selection of the flavor vacuum are entirely different. This further differentiates our approach from theirs.

## II. MOTIVATION: D-PARTICLE FOAM AS A MEDIUM NECESSITATING FLAVOR FOCK STATES

### A. Features of the model

The supersymmetric (string) model illustrated in the upper panel of Fig. 1 [7] will serve as a concrete framework for D-foam phenomenology. In it, our Universe, after appropriate compactification, is represented as a Dirichlet three-brane (D3-brane), propagating in a 10-dimensional bulk space-time punctured by D-particle defects.<sup>2</sup> As the D3-braneworld moves through the bulk, the D-particles cross it. To an observer on the D3-brane the model looks like “space-time foam” with defects “flashing” on and off as the D-particles cross it: “D-foam” denotes this structure. *If no relative motion of branes takes place in the bulk target-space supersymmetry*, it implies the vanishing of the ground-state energy of the configuration. However, if there is relative motion in the bulk, SUSY is broken and there are nontrivial forces among the D-particles as well as between the D-particles, the braneworld and orientifolds [7]. The resulting nonzero contribution to the energy is proportional to  $v^2$  for *transverse* relative motion of branes with different dimensionalities, and to  $v^4$  for branes of the same dimensionality (there is *no* contribution to the energy of a p-braneworld from motion of other branes in directions *parallel* to its longitudinal directions). There is also a dependence on the relative distance of the various branes. We shall now give a quite heuristic analysis of the effects of a gas of D-particles on the flavored string state on the braneworld. (A more string-theoretic calculation will be given later which will model more rigorously the flavor changing features of the interaction of D-particles.) In particular, the interaction of a single D-particle, that lies far away from the D3 brane (i.e., a compactified D8) world, and moves adiabatically with a small velocity  $v_{\perp}$  in a direction transverse to the brane, results in the following potential<sup>3</sup> [7]

$$\mathcal{V}_{D0-D8}^{\text{long}} = + \frac{r(v_{\perp}^{\text{long}})^2}{8\pi\alpha'}, \quad r \gg \sqrt{\alpha'}. \quad (1)$$

On the other hand, a D-particle close to the D3-brane (compactified D8), at a distance  $r' \ll \sqrt{\alpha'}$ , moving adiabatically in the perpendicular direction with a velocity  $v_{\perp}^{\text{short}}$  will induce the following potential to it:

<sup>2</sup>Since an isolated D-particle cannot exist [9], because of string gauge flux conservation, the presence of a D-brane is essential.

<sup>3</sup>For brevity, in what follows we ignore potential contributions induced by compactification of the D8 braneworlds to D3 branes, stating only the expressions for the induced potential on the uncompactified braneworld as a result of a stretched string between the latter and the D-particle—the compactification does not affect our arguments on the negative energy contributions to the brane vacuum energy.

$$\mathcal{V}_{D0-D8}^{\text{short}} = -\frac{\pi\alpha'(v_{\perp}^{\text{short}})^2}{12r^3}. \quad (2)$$

D-foam involves different configurations of D-particles, and so one has to average over appropriate populations and quantum fluctuations [2,10] [due to recoiling D-particles with (nine-dimensional) recoil velocities  $v_i$ ]. Recoil fluctuations *along the D3-brane universe* represent a stochastic fraction of the incident momentum of the D-particle [10]. Because of the D-particle fluctuations in the transverse direction of the braneworlds, there would also be contributions to the potential energy of the brane [cf. (1) and (2), where now the quantities involving  $v_{\perp}^{\text{short,long}}$  should be averaged over populations of D-particles and quantum fluctuations]. We can plausibly use a general parametrization of the foam fluctuations as follows:

$$\begin{aligned} \delta v_A &= g_s \frac{r_{\parallel}}{M_s} p_A, \quad A = 1, 2, 3 \Rightarrow \langle r_{\parallel} \rangle = 0, \\ \langle r_{\parallel}^2 \rangle &= \sigma_{\parallel}^2 \neq 0, \\ v_{\perp}^{\text{short}} &\equiv v_{\alpha}^{\text{short}}, \quad \alpha = 4, \dots, 9 \Rightarrow \langle v_{\alpha}^{\text{short}} \rangle = 0, \\ \langle v_{\alpha}^{\text{short}} v_{\beta}^{\text{short}} \rangle &= \delta_{\alpha\beta} \sigma_{\text{short}}^2 \neq 0, \quad \alpha, \beta = 4, \dots, 9, \\ v_{\perp}^{\text{long}} &\equiv v_{\alpha}^{\text{long}}, \quad \alpha = 4, \dots, 9 \Rightarrow \langle v_{\alpha}^{\text{long}} \rangle = 0, \\ \langle v_{\alpha}^{\text{long}} v_{\beta}^{\text{long}} \rangle &= \delta_{\alpha\beta} \sigma_{\text{long}}^2 \neq 0, \quad \alpha, \beta = 4, \dots, 9, \\ \langle v_{\alpha}^{\text{short,long}} v_{\beta}^{\text{short,long}} \rangle &= 0. \end{aligned} \quad (3)$$

Our stochastic-foam model has vanishing correlators of odd powers of  $v_i$  of the recoil velocity, but nontrivial correlators of even powers. Above,  $\langle \dots \rangle$  indicates averaging over both quantum fluctuations and the ensemble of D-particles in the foam;  $A = 1, 2, 3$  denote the longitudinal dimensions of the D3-braneworld. The  $\delta v_A$ ,  $A = 1, 2, 3$  represent the recoil velocity components of the D-particle during scattering. In the absence of any interaction with matter strings there is still a velocity  $v_i$  expressing the quantum fluctuations of individual D-particles.

We consider the process of capture by a quantum-fluctuating (heavy, nonrelativistic) D-particle of an open string representing a neutral flavored excitation, such as a neutrino (cf. lower figure in Fig. 1). By momentum conservation during this scattering event one has

$$\begin{aligned} \langle (p_1 + p_2)_A \rangle &= \frac{M_s}{g_s} \langle \delta v_A \rangle = 0, \\ \frac{M_s}{2g_s} \langle (v_A)_1^2 \rangle + E_1 &= \frac{M_s}{2g_s} \langle (v_A + \delta v_A)_1^2 \rangle + E_2 + \delta E, \\ E_I &= \sqrt{p_A^2 + m_I^2}, \quad I = 1, 2. \end{aligned} \quad (4)$$

We have assumed that *kinematical* constraints can be estimated by treating the open-string states as relativistic free fields of typical mass  $m_I$ , and the D-particles as heavy (nonrelativistic) particles, with masses  $M_s/g_s \gg m_I$ , where  $M_s$  is the string scale, and  $g_s < 1$  is the (weak)




FIG. 1. Upper: schematic representation of a generic D-particle space-time foam model, in which matter particles are treated as open strings propagating on a D3-brane, and the higher-dimensional bulk space-time is punctured by D-particle defects. Lower: details of the process whereby an open-string state propagating on the D3-brane is captured by a D-particle defect, which then recoils. This process involves an intermediate composite state that persists for a period  $\delta t \sim \alpha' E$ , where  $E$  is the energy of the incident string state, which distorts the surrounding space-time during the scattering, leading to an effective refractive index but not birefringence. The components of the recoil velocity perpendicular to the D3 braneworld lead to vacuum energy contributions and thus target-space supersymmetry-breaking.

string coupling.  $\delta E$  denotes *brane energy fluctuations* in the transverse direction to the brane, which in view of (1) and (2) (with  $v_{D0} = v_{\perp}^{\text{short,long}}$ ), are due both to long and short-distance D-particles. In our dilute-foam approximation we may take

$$\delta E = \delta E_{\text{short}}(\sigma'_{\text{short}}(p)) + \delta E_{\text{long}}(\sigma'_{\text{long}}), \quad (5)$$

where according to our previous discussion only the short-distance stochastic-foam fluctuations may depend on the (average) momentum  $p$  of the propagating matter state. In view of the  $v_{\perp}^2$  dependence of the respective fluctuations (cf. (1) and (2)) both types of energy fluctuations are proportional to the respective velocity fluctuations  $\sigma'_{\perp}$ .

Averaging over foam fluctuations, and Eq. (4) implies that the (average) energy  $E_2$  of a string excitation, corresponding to a state of mass  $m_1$ , propagating in the foam, will be modified from its free value  $E_1 = \sqrt{p^2 + m_1}$  by:

$$E_2 = \sqrt{p^2 + m_1} - \frac{g_s}{2M_s} \sigma_{\parallel}^2 p^2 + \delta E_{\text{short}}(\sigma'_{\text{short}}(p)) + \delta E_{\text{long}}(\sigma'_{\text{long}}). \quad (6)$$

Since the signature of the quantum energy fluctuations is *not fixed*, all kinds of situations for the foam dressed masses are allowed in such processes. For instance, in case of flavored states, where in the mass-eigenstate spectrum, both masses are allowed, i.e., one may have  $m_1 > m_2$ , when  $\delta E < 0$  is predominantly due to short-range contributions (2), while  $m_1 > m_2$  in case  $\delta E$  is dominated by the long-range contributions. On the other hand, when the massive state is not flavored, then necessarily  $m_1 = m_2$  which is either achieved by the equality  $-\frac{g_s}{2M_s} \sigma_{\parallel}^2 p^2 + \delta E_{\text{short}}(\sigma'_{\text{short}}(p)) + \delta E_{\text{long}}(\sigma'_{\text{long}}) = 0$ , or by means of appropriately modified dispersion relations, and hence a vacuum refractive index proportional to the foam fluctuations. Notice that these selection rules are allowed, as they express a link between the transverse foam fluctuations to the longitudinal ones, which are free parameters of the model. The net  $\delta E$  can be both positive and negative, which would be compatible with oscillations.

A simple illustrative analysis can be effected by using a piecewise continuous distribution  $P(\eta)$  of D-particles in the (transverse or bulk direction)  $\eta$  parametrized as

$$P(\eta) = \begin{cases} \frac{\eta^3}{L^4}, & \text{when } 0 < \eta < a, \\ \frac{b}{L}, & \text{when } a < \eta < L - a, \\ -\frac{(\eta-L)^3}{L^4}, & L - a < \eta < L. \end{cases}$$

(We have suppressed a normalization factor  $N$  which is given by  $N = (b + \frac{a^4}{2L^4} - \frac{2ab}{L})^{-1}$ .) The total energy  $\mathcal{E}$  exerted on the D-brane by the D-particles  $\mathcal{E} \propto \int_0^L d\eta p(\eta) \tilde{\varepsilon}(\eta)$ , with  $\tilde{\varepsilon}(\eta)$  denoting the linear bulk energy density at position  $\eta$ , satisfies (upon using the structures (1) and (2) for  $\tilde{\varepsilon}(\eta)$  appropriately):

$$\mathcal{E} \propto - \left[ av_L \left( b + \frac{a^4}{5L^4} \right) + \frac{\alpha'^2 a}{L^4} v_S \right] + \frac{v_L}{2} \left( bL + \frac{a^4}{4L^3} \right), \quad (7)$$

where

$$v_L = \frac{(v_{\perp}^{\text{long}})^2}{8\pi}, \quad v_S = \frac{\pi(v_{\perp}^{\text{short}})^2}{12}. \quad (8)$$

The vanishing of  $\mathcal{E}$  leads to the condition

$$b \sim \frac{\alpha'^2 a}{L^4} \frac{v_S}{v_L \left( \frac{L}{2} - a \right)} \quad (9)$$

provided  $a \ll \alpha'^{(1/2)}$ , and so  $P(\eta)$  displays a substringy structure (consistent with the validity of the limiting forms of the potential energy that we have used). Now the coupling of D-particles to the matter on the brane can lead to a deformation of  $P(\eta)$  on the brane which can lead to  $\mathcal{E}$  having either sign.

This would be expected to be a generic effect given such a  $P(\eta)$ , a demonstration of which can be seen in the context of a simple toy model: a D-particle is pointlike from the point of view of observers on the brane. The scale of the strongest interactions is substringy and so, in comparison, the size of the bulk dimension is very large. The D-particle, as far as matter excitations of the string are concerned, has no quantum numbers. In order to gain some intuition about changes to  $P(\eta)$  due to D-particle brane interactions we will consider a very simple field theory where we represent the D-particle by a massive scalar field  $\phi$  and the matter excitation by a harmonic oscillator on the boundary. Generically the model can be represented as an action:

$$S = \frac{1}{2} \int_{\Omega} d^{d+1} \eta \sqrt{-G} (G^{MN} \phi_{,M} \phi_{,N} + m^2 \phi^2) - \frac{1}{2} \int_{\partial\Omega} d^d \eta \sqrt{-g} [g^{\mu\nu} q_{,\mu} q_{,\nu} + \mu^2 q^2 + 2\beta\phi q + \lambda\phi^2], \quad (10)$$

where  $\Omega$  is the bulk,  $\partial\Omega$  is the brane (which is the boundary of the bulk),  $G^{MN}$  is the bulk metric, and  $g^{\mu\nu}$  is the brane metric; the analysis is simpler for  $d = 0$  and suffices to illustrate the deformation of the distribution of  $P(\eta)$  owing to bulk-brane interactions. The quantum theory is determined by the classical modes. For  $d = 0$ , bulk  $-\infty < \eta < 0$ , brane position  $\eta = 0$ , conjugate momentum  $\pi(\eta, t)$ , the action  $S$  leads to a Hamiltonian [11]

$$H = \int_{-\infty}^0 \left( \frac{1}{2} \pi(\eta, t)^2 + \frac{1}{2} (\partial_{\eta} \phi(\eta, t))^2 + \frac{1}{2} m^2 \phi(\eta, t)^2 \right) d\eta + \frac{1}{2} \lambda \phi(0, t)^2 + \beta \phi(0, t) q(t) + \frac{1}{2} \mu^2 q(t)^2 + \frac{1}{2} p(t)^2. \quad (11)$$

The equation of motion is

$$\partial_t^2 \phi(\eta, t) = \partial_{\eta}^2 \phi(\eta, t) - m^2 \phi(\eta, t) - \delta(\eta) (\partial_{\eta} \phi(0, t) + \lambda \phi(0, t) + \beta q(t)), \quad (12)$$

which is equivalent to the solution of the massive Klein-Gordon equation with a boundary condition

$$\partial_x \phi(0, t) = -\lambda \phi(0, t) - \beta q(t). \quad (13)$$

The oscillator satisfies the equation of motion

$$\partial_t^2 q(t) = -\beta \phi(0, t) - \mu^2 q(t). \quad (14)$$

The mode solutions corresponding to (12) and (14) are

$$\begin{aligned} \phi(\eta, t) = & \int_0^{\infty} (\rho(\rho^2 - \mu^2 + m^2) \cos(\rho \eta) \\ & - (\lambda(\rho^2 - \mu^2 + m^2) + \beta^2) \sin(\rho \eta)) \\ & \times (a(\rho) \cos(\omega_{\rho} t) + b(\rho) \sin(\omega_{\rho} t)) d\rho \end{aligned} \quad (15)$$

and

$$q(t) = \int_0^\infty \beta \rho (a(\rho) \cos(\omega_\rho t) + b(\rho) \sin(\omega_\rho t)) d\rho, \quad (16)$$

where  $\omega_\rho = \sqrt{m^2 + \rho^2}$  and  $a(\rho)$  and  $b(\rho)$  are real. The equations of motion are linear and so the integral over  $\rho$  represents a linear superposition. These are solutions which propagate in the bulk. Square integrable states bound to the brane also exist by considering specific pure imaginary  $\rho$ . The asymptotic behavior as  $\eta \rightarrow -\infty$  has to be nondivergent and this selects the details of the coefficients in the solution. Once the solutions are substituted in  $H$  we find a description in terms of harmonic oscillators associated with the classical modes. Clearly from the mode structure for real  $\rho$  (or bulk modes) we have an oscillatory behavior with  $\eta$  and so this will lead to a modulation in  $P(\eta)$  as a result of (the analogue of) D-particle recoil and hence this will allow both positive and negative  $\delta E$ . The bound states do not participate in this modulation. A more detailed model involving two harmonic oscillators (with distinct frequencies) on the brane will be presented elsewhere but the qualitative features that we have presented here should not change.

In the next section a string theoretic picture of flavor changes owing to non-BPS (Bogomolny-Prasad-Sommerfield) (compactified) D-particles will be considered. We shall argue in favor of the positivity of the energy density of the *flavor vacuum* [3] and, hence, its appropriateness to represent the physical ground state of the D-foam in the presence of flavored matter.

### B. D-foam-induced fermion condensates and flavor-vacuum stability

In the above formalism, the presence of mixed-sign fluctuations may lead to negative contributions to the vacuum energy, implying an *instability* of the standard *mass-eigenstate vacuum* in the case of flavored particles.<sup>4</sup> The qualitative description of the role of D-particles in producing an instability in the mass vacuum can be put on a more explicit footing. In order to do this we will need to have a more detailed picture of stringy matter and how the standard model of particle physics (and beyond) emerge. This

<sup>4</sup>We also note in passing, that in the case of antiparticles, viewed as “negative energy” solutions of the free Dirac equation,  $\bar{E}_{<0} = -\sqrt{p^2 + m_1^2}$ , Eq. (6) implies  $E_2 = -\bar{E}_{2,>0} = -\sqrt{p^2 + m_1^2} - \frac{g_s}{2M_s} \sigma_\parallel^2 p^2 + \delta E_{\text{short}}(\sigma'_{\text{short}}(p)) + \delta E_{\text{long}}(\sigma'_{\text{long}})$ , where, on adopting Feynman’s interpretation, that antiparticles have positive energy but move “backwards” in time, we have denoted the positive energy of the flavored antiparticle after interaction with the foam by  $\bar{E}_{2,>0}$ . Solving for  $\bar{E}_{2,>0}$ , we thus obtain different modified dispersion relations between particle and antiparticle, due to the *relative sign difference* of the foam corrections, and thus induced *CPT* violation. The difference in energy between particles and antiparticles, implied by the foam contributions, may lead to neutrino/antineutrino oscillations, which however we shall not explore here.

represents a vast area known as string phenomenology where a multitude of detailed models involving stacks of D-branes and orientifolds interacting with stringy matter have been constructed. In D-foam models certain generic four-fermion interactions are induced among fermionic matter which should arise in a large class of models. The interactions are of a four-fermion vector-current-current form which can lead to a mass-matrix type mixing in the case of models where right-handed neutrinos exist. This type of four-fermion interaction describes the result of the splitting of open strings when interacting with the D-particle defects. Such interactions arise in D-foam models, based on type IA [12] or type IIB strings [13], as a consequence of intermediate open-string exchanges, stretched between D-particles and the braneworlds; these represent quantum fluctuations of the recoiling D-particle during its interaction with the propagating matter open-string states. Some of these exchanged states are U(1) gauge particles, where U(1) are Abelian gauge groups that are abundant in string models, not necessarily representing electromagnetism. As a result of such Abelian gauge particle exchanges, there will be induced, among others, attractive four-fermion interactions in a low-energy effective field-theory framework [14]. For chiral fermions,  $\psi$ ,  $\psi'$  for instance, such interactions (obtained by considering the corresponding tree-level (Veneziano-like) four-point string scattering amplitudes) assume the generic form [13,14]

$$\mathcal{L}_{4\text{-eff}} = \frac{\xi}{M_s^2} V^c \sum_{i,j} \sum_{a,b=L,R} \mathcal{G}_{ab} \bar{\psi}_i^a \gamma^\mu \psi_{ia} \bar{\psi}_j^b \gamma_\mu \psi'_{jb}, \quad (17)$$

where  $V^c$  denotes the compactification volume element for four space-time dimensions, in units of the string length  $\sqrt{\alpha'}$ ; the indices  $i, j$  refer to fermion species, including flavor,  $\mathcal{G}_{ab}$  are numerical coefficients depending on the (square of the) couplings of the particular interactions, and are of order  $O(1)$  as far as the string scale  $M_s$  is concerned, and  $L, R$  denote the appropriate chirality of the spinors  $\psi_i, \psi'_j$ . The constant  $\xi$  depends on the type of D-foam considered. The combination

$$\kappa = 8\pi \frac{V^c}{M_s^2} \quad (18)$$

defines the gravitational coupling of the four-dimensional effective theory.

For type IA string D-foam [7,12], the D-particle defects, being pointlike and electrically neutral, interact only with electrically neutral particles, since charge conservation prevents the splitting (cf. Figure 1) of charged open-string states when interacting with the defect. In this case, the fermions  $\psi$  represent neutrinos of the standard model, assuming these to be the dominant electrically neutral fermions in the low-energy world today. Thus,  $\xi = 1$  for such neutral excitations, while  $\xi = 0$  for charged fermions. On the other hand, as explained in [13] and mentioned

above, for D-foam models based on type IIB strings, the D-“particles” are only effectively pointlike, being represented by D3 brane defects wrapped up along three cycles appropriately, which then intersect the D7 branes of the model (representing after compactification to three spatial dimensions our braneworld). In such models, there are nontrivial charged particle interactions with the D-particles, of the type considered in Fig. 1, which however are suppressed as compared to the neutral particle interactions by a factor

$$\xi = \frac{(1.55)^4}{M_s^4 V_{A3} R'}, \quad (19)$$

where  $M_s$  is the string scale,  $V_{A3}$  denotes the volume of a cell on the D7 brane (playing the rôle of a brane Universe after compactification of four of its dimensions) containing a D3 particle in the foam, and  $R'$  is the fourth dimension of the D7 brane, transverse to the D3-brane dimensions. The corresponding four-fermion interactions assume the form (17) with  $\xi$  given by (19).

Let us consider flavored neutrinos which are chiral, where

$$\gamma_5 \psi_i = \pm \psi_i, \quad (20)$$

and the  $\pm$  sign depends on whether the fermion is left- or right-handed. Indeed, in models such as the standard model we encounter only one type of Chiral fermion, say left-handed, satisfying (20), with no right-handed neutrinos. Next, let us assume that there are *right-handed flavored fermions* (e.g., neutrinos) interacting with the extra U(1) stringy symmetries, which are different from the Abelian gauge group of the standard model interactions. In this sense, the presence of right-handed fermions is consistent with their nature as being neutral under the standard model group.<sup>5</sup> With the above in mind, we observe that, upon a Fierz rearrangement, the four-fermion interactions (17), involving both left- and right-handed flavored fermions, includes terms :

$$\begin{aligned} \mathcal{L}_{4\text{-eff}} \ni & \frac{\xi \mathcal{G} V^c}{M_s^2} \sum_{i,j} \left[ -(\bar{\psi}_i \psi_j)(\bar{\psi}_j \psi_i) + (\bar{\psi}_i \gamma_5 \psi_j)(\bar{\psi}_j \gamma_5 \psi_i) \right. \\ & + \frac{1}{2}(\bar{\psi}_i \gamma^\mu \psi_j)(\bar{\psi}_j \gamma_\mu \psi_i) \\ & \left. + \frac{1}{2}(\bar{\psi}_i \gamma_5 \gamma^\mu \psi_j)(\bar{\psi}_j \gamma_5 \gamma_\mu \psi_i) \right], \quad (21) \end{aligned}$$

where for brevity and concreteness we have assumed one type of non-standard-model interaction among neutrinos due to the intermediate string states, with  $\mathcal{G} > 0$  (which is proportional to the square of the pertinent coupling), and the spinors have both left- and right-handed components, i.e.,

<sup>5</sup>The assumptions on having right-handed neutrinos in models beyond the standard model is not uncommon, see for instance [15].

$$\psi_i = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}_i$$

$i = 1, \dots, N$  where  $N$  is the number of flavors. We can obtain *attractive* interactions in the scalar channel of the form

$$\mathcal{L}_{4\text{-eff}} \ni -\frac{\xi \mathcal{G} V^c}{M_s^2} \sum_{i,j} \langle (\bar{\psi}_i \psi_j) \rangle \langle (\bar{\psi}_j \psi_i) \rangle. \quad (22)$$

The condensates would lead to dynamical mass-matrix contributions:

$$m_{ij} = \frac{\xi \mathcal{G} V^c}{M_s^2} \langle \bar{\psi}_i \psi_j \rangle, \quad i, j = 1, \dots, N. \quad (23)$$

From the off-diagonal in flavor space parts of the condensate one would thus obtain *mixing*, while the diagonal parts would lead to *dynamically* generated *fermion-mass* contributions, which can lead to chiral symmetry-breaking.<sup>6</sup> We stress once again that such scalar condensates characterize only theories where right-handed fermions do exist, because otherwise the scalar four-fermion terms arising from the Fierz rearrangement (21) are cancelled by the pseudoscalar ones.

An important remark is in order here. The linearization of the attractive four-fermion interactions in the scalar channel by means of the nonpropagating Hubbard-Stratonovich condensate field  $\Delta_{ij}$ , i.e.,

$$\mathcal{L}_{4\text{-eff}} \ni -\sum_{i,j} \left( \frac{M_s^2}{4V^c |\xi| \mathcal{G}} \Delta_{ij} \Delta^{ij} + \Delta_{ij} (\bar{\psi}_j \psi_i) \right) \quad (24)$$

would lead, upon formation of the condensate, to *negative* contributions to the *vacuum energy* ( $\mathcal{O}(\langle \Delta_{ij} \rangle^2)$  terms above). However, in the D-foam situation, there are also the *positive* energy contributions arising from the (temporal components of the) repulsive vector four-fermion condensates in (21), which are nonzero as a result of the flavor vacuum that exists due to the induced mixing. Indeed, a nontrivial vacuum condensate of the time component of the current operator ( $\langle \sum_i \bar{\psi}_i \gamma^0 \psi_i \rangle = \langle \sum_i \psi_i^\dagger \psi_i \rangle$ ) is equivalent to a nonzero particle-species number, which is known to characterize the *flavor vacuum* [1–3] but *not* the mass-eigenstate one. These contributions could overcome the negative vacuum energy contributions on the braneworlds, arising from the interactions of the bulk D-particles with

<sup>6</sup>In the context of our D-brane Universes, the vacuum energy contributions due to the bulk D-particles (cf. (1) and (2)) lead to (anti)de Sitter space-times. Chiral symmetry-breaking in such nontrivial space-time backgrounds has been discussed in the context of early Universe studies in [16,17], with the conclusion that the dynamical formation of chiral condensates, due to four-fermion interactions, exhibits better UltraViolet (UV) behavior than in the corresponding flat space-time case [18]. We postpone a detailed discussion of such issues to a forthcoming publication.

the brane Universe (cf. (2)), leading to the *stability* of the flavor vacuum.

Incidentally, the fermion-number condensates *violate* not only Lorentz but also *CPT* symmetry. In the presence of such fermion-number condensates, the effective four-Fermi interactions (21), when linearized by means of appropriate Hubbard-Stratonovich fields, lead to a low-energy effective Lagrangian containing *CPT* and Lorentz-violating terms of the type appearing in the parametrization of Ref. [19]. This induced violation of Lorentz and *CPT* invariance is in agreement with the generic properties of the D-foam discussed previously.

### C. On the Scale of flavor-vacuum-condensate-induced supersymmetry-breaking

Before closing this section a final remark is in order about the nature of symmetries that may be broken by the aforementioned flavor-vacuum condensates in D-foam models and the associated order of magnitude of the pertinent symmetry-breaking scales. In addition to the chiral symmetries, that at any rate do not necessitate the introduction of a flavor vacuum, it is also supersymmetry that is broken in a novel nonperturbative way by the introduction of the flavor vacuum. In fact, the four-fermion interactions (17), due to intermediate-string-state exchanges, arise consistently with target-space supersymmetries of the respective string theories [14], so formally the respective Lagrangians can be made supersymmetric, in the sense of their invariance under supersymmetry transformations (local supersymmetry (supergravity) actually, in view of the models' coupling to gravity).

However, the formation of Lorentz-violating fermion-flavor-vacuum condensates will induce a novel breaking of global supersymmetry, which we shall discuss in the next section (any extension to supergravity is left for future works). It becomes obvious from the above discussion that the order of the induced supersymmetry-breaking is then dictated by the order of the respective condensates. In our stochastic-foam situations, the natural four-dimensional mass scale is  $1/\sqrt{\kappa}$  (18), which from a four-dimensional point of view may be identified with the Planck scale, and for large radii compactification is different from the string scale  $M_s$ , which can be as low as a few TeV in the modern version of string-theory. However, as discussed in [1,2], the induced condensates in D-foam models are calculated by subtracting appropriately the Minkowski vacuum contributions in the absence of D-particles. This implies [1,2] that the condensates are *proportional* to the average of stochastic fluctuations of the recoil velocities  $\sigma_{\text{short,long}}^2 \equiv \sigma'^2$  in (3), assuming for simplicity a common scale of D-foam recoil velocity fluctuations. Intuitively one can understand this dependence as follows: the flavor vacuum is Lorentz-violating, and in D-foam models the only source of local Lorentz violation are the stochastic fluctuations of the recoil velocities of the

D-particles (3). Thus, it is natural that the flavour-vacuum condensate is proportional to the latter. This makes the effective supersymmetry-breaking mass scale in D-foam models, due to flavor vacuum, of order

$$M^2 \sim \frac{1}{\kappa} \sigma'^2. \quad (25)$$

The parameter  $\sigma'^2$  is an arbitrary parameter of the stringy foam model, although naturalness criteria may lead us to assume that it is more or less of order one (but this assumption is non binding).

To recapitulate, the above analysis implies that the open-string states, representing flavored particles, when interacting with the D-foam, may *induce flavor mixing*, of the order of the foam fluctuations. The vacuum on the brane-world has broken supersymmetry, due to the relative (fluctuation) motion between the braneworld and the nearby D-particles, as well as *broken Lorentz invariance*, due to the fluctuating recoil velocities of the D-particle space-time defects. These features imply that the appropriate Fock space vacuum for such configurations cannot be the ordinary Lorentz invariant, supersymmetric vacuum. We have argued in [2] that the physically correct vacuum in this case is the *flavor vacuum* proposed by Blasone and Vitiello [3] to describe generically theories with mixing, which is *orthogonal* to the mass-eigenstate vacuum. Moreover, the fact that there is *particle creation* in our string case and in the flavor vacuum [3], as mentioned above, constitutes an additional consistency check of the identification of the latter with the physical vacuum in our D-foam model. Indeed, because of this property, the Lorentz-violating flavor vacuum leads to Lorentz- and *CPT*-violating vector condensates (temporal components) that contribute positively to the vacuum energy of the braneworld, cancelling negative anti-de Sitter type contributions on the brane vacuum energy (cf. (2)), and therefore restoring stability of the ground state. It was also argued in [2] that the particle-creation feature of the flavor vacuum leads to a dynamical appearance of a high-momentum cutoff, which is set by the mass scale of the flavored particles. The argument was based on the maximal amount of particle-creation below the cutoff.

In the following section, therefore, we proceed to study the properties of supersymmetric fluids, using an interesting toy example, that of the Wess-Zumino supersymmetric theory. In the above context, this QFT should be viewed as an effective theory on our braneworld punctured by the flavor-change-inducing D-particle foam. We shall assume simple mixing in the mass-matrix, which from the above microscopic point of view may be viewed as the result of four-fermion interactions in the presence of right-handed neutrinos. We shall first formulate the theory, and then demonstrate some generic but quite nontrivial properties, with important differences between the boson and fermion excitations, thereby implying further nontrivial and novel manifestations of supersymmetry “breaking”. As we shall

see, in supersymmetric theories both flavored bosons and fermions form condensates in the flavor vacuum, but the total vacuum energy turns out to be positive, and moreover, the equations of state between bosons and fermions turn out to be different. These features constitute a novel, and purely nonperturbative, way of breaking supersymmetry. In what follows we shall demonstrate these properties in free theories, reserving an extension to interacting field theory in a future publication [8].

### III. THE FLAVORED WESS-ZUMINO MODEL

The WZ model is historically a prototype where the basic features of a supersymmetric theory may be formulated in a concrete and quite instructive way. In our work, motivated by the considerations of the previous section, we shall be dealing with extensions of the simplest version of the free WZ model, in which the fields carry nontrivial flavor quantum number and exhibit mixing. In the next subsection, the model and its particle content, is described in some detail. We should again stress that the flat-space supersymmetric WZ model is viewed here as the simplest prototype of an effective low-energy theory, describing the dynamics of excitations on a spatially flat braneworld in the model of the previous section. The initial supersymmetry among the fields is inherited from the underlying supersymmetries of the string/brane theory; the SUSY-breaking is due to the relative brane motion and will be incorporated, in this context, by viewing the ‘‘flavor’’ vacuum as the physical vacuum, describing correctly the associated mixing. It is understood that a more sophisticated local supersymmetry (supergravity) effective Lagrangian should be used in realistic constructions, but this will not affect qualitatively the important features of the flavor vacuum revealed by the study of the toy WZ model.

#### A. Supersymmetric model

The Lagrangian of the model reads [20,21]

$$\mathcal{L} = \mathcal{L}_\psi + \mathcal{L}_S + \mathcal{L}_P \quad (26)$$

with

$$\begin{aligned} \mathcal{L}_\Psi &= \bar{\Psi}_f (i\not{\partial} - M_f) \Psi_f^T, \\ \mathcal{L}_S &= \partial_\mu \mathcal{S}_f \partial^\mu \mathcal{S}_f^T - \mathcal{S}_f M_b \mathcal{S}_f^T, \\ \mathcal{L}_P &= \partial_\mu \mathcal{P}_f \partial^\mu \mathcal{P}_f^T - \mathcal{P}_f M_b \mathcal{P}_f^T, \end{aligned} \quad (27)$$

where

$$\Psi_f \equiv (\psi_A, \psi_B), \quad \mathcal{S}_f \equiv (S_A, S_B), \quad \mathcal{P}_f \equiv (P_A, P_B),$$

$$M_f \equiv \begin{pmatrix} m_A & m_{AB} \\ m_{AB} & m_B \end{pmatrix},$$

$$M_b \equiv \begin{pmatrix} m_A^2 & m_{AB}^2 \\ m_{AB}^2 & m_B^2 \end{pmatrix}.$$

The fields  $\psi_\iota$ ,  $S_\iota$  and  $P_\iota$  (with  $\iota = A, B$ ) are, respectively, a Majorana spinor, a scalar and a pseudoscalar.<sup>7</sup> For our purposes, they are considered to be fields of definite flavor,  $A$  or  $B$ . The cross terms of the Lagrangian (26), i.e., terms involving products of fields of different flavors, disappear when one expresses the model in terms of new fields, obtained by appropriate rotations of  $\Psi_f$ ,  $\mathcal{S}_f$  and  $\mathcal{P}_f$

$$\begin{aligned} \mathcal{L}_\Psi &= \bar{\Psi}_m (i\not{\partial} - M_1) \Psi_m^T, \\ \mathcal{L}_S &= \partial_\mu \mathcal{S}_m \partial^\mu \mathcal{S}_m^T - \mathcal{S}_m M_2 \mathcal{S}_m^T, \\ \mathcal{L}_P &= \partial_\mu \mathcal{P}_m \partial^\mu \mathcal{P}_m^T - \mathcal{P}_m M_2 \mathcal{P}_m^T, \end{aligned} \quad (28)$$

with

$$\begin{aligned} \Psi_m &\equiv (\psi_1, \psi_2) \equiv \Psi_f \Omega^T, & \mathcal{S}_m &\equiv (S_1, S_2) = \mathcal{S}_f \Omega^T, \\ \mathcal{P}_m &\equiv (P_1, P_2) = \mathcal{P}_f \Omega^T, & M_1 &\equiv \begin{pmatrix} m_1 & 0 \\ 0 & m_2 \end{pmatrix}, \\ M_2 &\equiv \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix}, & \text{and } \Omega &\equiv \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}, \end{aligned}$$

where  $m_1$ ,  $m_2$  and  $\theta$  are suitable functions of  $m_A$ ,  $m_B$  and  $m_{AB}$ . The new Lagrangian, written as (28), is just the sum of free Lagrangians for the fields  $\psi_i$ ,  $S_i$  and  $P_i$ , all with the same mass  $m_i$  ( $i = 1, 2$ ). Therefore we can decompose these fields as

$$\begin{aligned} \psi_i(x) &= \sum_{r=1,2} \int \frac{d^3k}{(2\pi)^{3/2}} [a_i^r(\vec{k}) u_i^r(\vec{k}) e^{-i\omega_i(k)t} \\ &\quad + v_i^r(-\vec{k}) a_i^{r\dagger}(-\vec{k}) e^{i\omega_i(k)t}] e^{i\vec{k}\cdot\vec{x}}, \end{aligned} \quad (29)$$

$$\begin{aligned} S_i(x) &= \int \frac{d^3k}{(2\pi)^{3/2}} \frac{1}{\sqrt{2\omega_i(k)}} \\ &\quad \times [b_i(\vec{k}) e^{-i\omega_i(k)t} + b_i^\dagger(-\vec{k}) e^{i\omega_i(k)t}] e^{i\vec{k}\cdot\vec{x}}, \end{aligned} \quad (30)$$

$$\begin{aligned} P_i(x) &= \int \frac{d^3k}{(2\pi)^{3/2}} \frac{1}{\sqrt{2\omega_i(k)}} \\ &\quad \times [c_i(\vec{k}) e^{-i\omega_i(k)t} + c_i^\dagger(-\vec{k}) e^{i\omega_i(k)t}] e^{i\vec{k}\cdot\vec{x}}, \end{aligned} \quad (31)$$

with  $\{a_i^r(\vec{k}), a_j^{s\dagger}(\vec{q})\} = \delta_{rs} [b_i(\vec{k}), b_j^\dagger(\vec{q})] = \delta_{rs} [c_i(\vec{k}), c_j^\dagger(\vec{q})] = \delta_{rs} \delta_{ij} \delta^3(\vec{k} - \vec{q})$  and  $\omega_i \equiv \sqrt{\vec{k}^2 + m_i^2}$  with  $i = 1, 2$ . From now on, we will refer to these fields as the fields *in the mass representation*, to be contrasted with the ones *in the flavor representation*, which are characterized by definite flavor quantum number.

<sup>7</sup>The pseudoscalar nature of  $P$  plays a role in the interactive WZ case, which we do not analyze here; in the current work, it can be regarded just as another scalar field. Nevertheless, to distinguish it from  $S$ , we will keep the nomenclature *pseudoscalar*.



### B. Flavor-mixing formalism

Following the BV formalism [3], we introduce the operator  $G_\theta(t)$ , connecting fields between the flavor and mass representations as follows:

$$\begin{aligned}\phi_A(x) &= G_\theta^\dagger(t)\phi_1(x)G_\theta(t), \\ \phi_B(x) &= G_\theta^\dagger(t)\phi_2(x)G_\theta(t),\end{aligned}\quad (32)$$

with  $\phi = \psi, S, P$ . Using Baker-Campbell-Hausdorff (BCH) formula,  $e^Y X e^{-Y} = X + [X, Y] + \frac{1}{2!}[X, [X, Y]] + \frac{1}{3!}[X, [X, [X, Y]]] + \dots$ , and the (anti-)commutation relations between fields, it can be shown that

$$G_\theta(t) = e^\theta \int d\vec{x}(X_{12}(x) - X_{21}(x)), \quad (33)$$

with

$$X_{12}(x) \equiv \frac{1}{2}\psi_1^\dagger(x)\psi_2(x) + i\dot{S}_2(x)S_1(x) + i\dot{P}_2(x)P_1(x). \quad (34)$$

The *flavor vacuum* is defined as<sup>8</sup>

$$|0\rangle_f \equiv G_\theta^\dagger(t)|0\rangle, \quad (35)$$

where  $|0\rangle$  is usual vacuum defined by  $a_i^r(\vec{k})|0\rangle = b_i(\vec{k})|0\rangle = c_i(\vec{k})|0\rangle = 0$  ( $i = 1, 2$ ).

Using the operator  $G_\theta(t)$  it is also possible to define new operators

$$\begin{aligned}a_A^r(\vec{k}, t) &\equiv G_\theta^\dagger(t)a_1^r(\vec{k})G_\theta(t), \\ a_B^r(\vec{k}, t) &\equiv G_\theta^\dagger(t)a_2^r(\vec{k})G_\theta(t), \\ b_A(\vec{k}, t) &\equiv G_\theta^\dagger(t)b_1(\vec{k})G_\theta(t), \\ b_B(\vec{k}, t) &\equiv G_\theta^\dagger(t)b_2(\vec{k})G_\theta(t), \\ c_A(\vec{k}, t) &\equiv G_\theta^\dagger(t)c_1(\vec{k})G_\theta(t), \\ c_B(\vec{k}, t) &\equiv G_\theta^\dagger(t)c_2(\vec{k})G_\theta(t).\end{aligned}\quad (36)$$

Using again BCH formula, one can show that

$$\begin{aligned}a_A^r(\vec{k}, t) &= \cos\theta a_1^r(\vec{k}) - \sin\theta \sum_s (W^{rs}(\vec{k}, t)a_2^s(\vec{k}) \\ &\quad + Y^{rs}(\vec{k}, t)a_2^{s\dagger}(-\vec{k})), \\ a_B^r(\vec{k}, t) &= \cos\theta a_2^r(\vec{k}) + \sin\theta \sum_s (W^{sr*}(\vec{k}, t)a_1^s(\vec{k}) \\ &\quad + Y^{sr}(-\vec{k}, t)a_1^{s\dagger}(-\vec{k})),\end{aligned}\quad (37)$$

<sup>8</sup>Although the *flavor-vacuum* state depends on time (and therefore it would be more appropriate to write  $|0(t)\rangle_f$ ); nevertheless, for the sake of simplicity, we will omit from now on the  $t$  dependence in the notation. Remarkably, as we shall see below, the important features of this state, of interest to us here, turn out to be *time independent*.

$$\begin{aligned}b_A(\vec{k}, t) &= \cos\theta b_1(\vec{k}) + \sin\theta(U^*(k, t)b_2(\vec{k}) + V(k, t), \\ b_2^\dagger(-\vec{k})b_B(\vec{k}, t) &= \cos\theta b_2(\vec{k}) - \sin\theta(U(k, t)b_1(\vec{k}) \\ &\quad - V(k, t)b_1^\dagger(-\vec{k})),\end{aligned}\quad (38)$$

$$\begin{aligned}c_A(\vec{k}, t) &= \cos\theta c_1(\vec{k}) + \sin\theta(U^*(k, t)c_2(\vec{k}) \\ &\quad + V(k, t)c_2^\dagger(-\vec{k})), \\ c_B(\vec{k}, t) &= \cos\theta c_2(\vec{k}) - \sin\theta(U(k, t)c_1(\vec{k}) \\ &\quad - V(k, t)c_1^\dagger(-\vec{k})),\end{aligned}\quad (39)$$

where

$$\begin{aligned}W^{rs}(\vec{k}, t) &\equiv \frac{1}{2}(u_2^{r\dagger}(\vec{k})u_1^s(\vec{k}) + v_2^{s\dagger}(\vec{k})v_1^r(\vec{k}))e^{i(\omega_1 - \omega_2)t}, \\ Y^{rs}(\vec{k}, t) &\equiv \frac{1}{2}(u_2^{r\dagger}(\vec{k})v_1^s(-\vec{k}) + u_1^{s\dagger}(-\vec{k})v_2^r(\vec{k}))e^{i(\omega_1 + \omega_2)t},\end{aligned}\quad (40)$$

with  $u_i^r(\vec{k})$  and  $v_i^r(\vec{k})$  the spinors in (29)

$$\begin{aligned}U(k, t) &\equiv \frac{1}{2}\left(\sqrt{\frac{\omega_1}{\omega_2}} + \sqrt{\frac{\omega_2}{\omega_1}}\right)e^{-i(\omega_1 - \omega_2)t}, \\ V(k, t) &\equiv \frac{1}{2}\left(\sqrt{\frac{\omega_1}{\omega_2}} - \sqrt{\frac{\omega_2}{\omega_1}}\right)e^{i(\omega_1 + \omega_2)t}.\end{aligned}\quad (41)$$

These operators satisfy the (anti-)commutation relations

$$\begin{aligned}\{a_\kappa^r(\vec{q}, t), a_\kappa^{s\dagger}(\vec{p}, t)\} &= \delta_{rs}[b_\iota(\vec{q}, t), b_\kappa^\dagger(\vec{p}, t)] \\ &= \delta_{rs}[c_\iota(\vec{q}, t), c_\kappa^\dagger(\vec{p}, t)] \\ &= \delta_{rs}\delta_{\iota\kappa}\delta^3(\vec{q} - \vec{p})\end{aligned}\quad (42)$$

(all others being zero and  $\iota, \kappa = A, B$ ) and it is also true that  $a_i^r(\vec{k})|0\rangle_f = b_\iota(\vec{k})|0\rangle_f = c_\iota(\vec{k})|0\rangle_f = 0$ . Therefore we are allowed to build a new Fock space out of these ingredients (operators+*flavor vacuum*), in the usual way. This is actually a *new* Fock space, and not just a different representation of the old one constructed by means of  $a_1^\dagger, a_2^\dagger$  and  $|0\rangle$ , since all the new states are found orthogonal to the old ones in the thermodynamic limit (see [3] for an exhaustive discussion of flavor states, and *vacua* within the BV formalism). This physical inequivalence leads to interesting consequences. As we have said previously, one of those is the nontrivial violation of Lorentz symmetry, manifested through the nonzero vacuum expectation values of the stress-energy tensor, and also of SUSY in a particular way, as we now proceed to discuss.

### IV. FLAVOR-VACUUM-INDUCED SUSY-BREAKING

In order to study the properties of the *flavor vacuum*, we will calculate the *flavor vacuum* expectation value of the stress-energy tensor, namely,

$${}_f\langle 0|T_{\mu\nu}(x)|0\rangle_f. \quad (43)$$

The stress-energy tensor for the theory (28) is written as

$$T_{\mu\nu}(x) = \sum_{i=1,2} (2\partial_{(\mu}S_i(x)\partial_{\nu)}S_i(x) + 2\partial_{(\mu}P_i(x)\partial_{\nu)}P_i(x) + i\bar{\psi}_i(x)\gamma_{(\mu}\partial_{\nu)}\psi_i(x)) - \eta_{\mu\nu}\mathcal{L}. \quad (44)$$

In case of a perfect relativistic fluid, as the *flavor-vacuum* condensate turns out to be, the nonvanishing elements of this matrix, which are actually the ones on the diagonal, have a specific physical interpretation:  $T_{00}$  represents the energy density of the fluid and  $T_{ii}$  the pressure in the three spatial directions. In particular, we are interested in the equation of state, given by

$$w = \frac{{}_f\langle 0|T_{ii}(x)|0\rangle_f}{{}_f\langle 0|T_{00}(x)|0\rangle_f}. \quad (45)$$

An important remark is in order. A straightforward computation of  ${}_f\langle 0|T_{\mu\nu}(x)|0\rangle_f$  leads to a divergent result. In previous works, when the fermionic and the bosonic cases were treated separately, a *finite* result was achieved in two steps: first, a normal ordering with respect to the usual vacuum was defined

$${}_f\langle 0|\mathcal{O}:|0\rangle_f \equiv {}_f\langle 0|\mathcal{O}|0\rangle_f - \langle 0|\mathcal{O}|0\rangle, \quad (46)$$

then, a cutoff in the momentum space was introduced. In the context of supersymmetry, the first step, the normal ordering with respect to the usual vacuum, is not needed, since

$$\langle 0|T_{\mu\nu}(x)|0\rangle = 0. \quad (47)$$

A formal cutoff remains as a regulator of the theory, a physical interpretation for which can be provided by a microscopic theory (such as the brane model of [1,2], described briefly in Sec. II above). The BV formalism is used to provide an effective (Lorentz-violating) field theory in the low-energy limit: the cutoff appears dynamically in the theory, and is interpreted as the high energy scale, below which the low-energy effective BV field theory is valid.

### A. Outline of calculational steps

In order to demonstrate the novel SUSY-breaking induced by the flavor vacuum and describe the collective behavior (equation of state) of the relativistic fluids under consideration, we need first to evaluate the quantity (43), where the expectation value is taken with respect to the *flavor-vacuum* state. To this end, we first simplify the quantum operator structure of the various quantities involved, and then we deal with the remaining functions over the momenta.

Specifically, the stress-energy tensor (44) is first expressed in terms of the fields  $\psi_i(x)$ ,  $S_i(x)$  and  $P_i(x)$ , in the *mass representation*. On expressing the fields in terms

of the mass-eigenstate creation and annihilation (*mass ladder*) operators (cf. (29)–(31)), we can write

$$T_{\mu\nu}(x) = \sum_{i=1,2} \sum_{r,s} \int d\vec{p}d\vec{q} [L_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) a_i^{r\dagger}(\vec{p}) a_i^s(\vec{q}) + M_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) a_i^{r\dagger}(\vec{p}) a_i^{s\dagger}(\vec{q}) + N_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) a_i^r(\vec{p}) a_i^s(\vec{q}) + K_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) a_i^r(\vec{p}) a_i^{s\dagger}(\vec{q})], \quad (48)$$

with  $r, s = -1, 0, 1, 2$ , defining  $a_i^0(\vec{k}) \equiv b_i(\vec{k})$  and  $a_i^{-1}(\vec{k}) \equiv c_i(\vec{k})$ , with  $L, M, N, K$  suitable functions of the momenta. In (48), the bra  ${}_f\langle 0|$  and ket  $|0\rangle_f$  act on the *mass ladder operators*

$$\begin{aligned} &{}_f\langle 0|T_{\mu\nu}(x)|0\rangle_f \\ &= \sum_{i=1,2} \sum_{r,s} \int d\vec{p}d\vec{q} [L_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) {}_f\langle 0|a_i^{r\dagger}(\vec{p}) a_i^s(\vec{q})|0\rangle_f \\ &+ M_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) {}_f\langle 0|a_i^{r\dagger}(\vec{p}) a_i^{s\dagger}(\vec{q})|0\rangle_f \\ &+ N_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) {}_f\langle 0|a_i^r(\vec{p}) a_i^s(\vec{q})|0\rangle_f \\ &+ K_{\mu\nu}^{rs}(\vec{p}, \vec{q}, m_i) {}_f\langle 0|a_i^r(\vec{p}) a_i^{s\dagger}(\vec{q})|0\rangle_f]. \end{aligned} \quad (49)$$

In order to simplify the expectation values of pairs of *mass operators*, we use the formulas

$$\begin{aligned} a_1^r(\vec{k}) &= \cos\theta a_A^r(\vec{k}, t) + \sin\theta \sum_s (W^{rs}(\vec{k}, t) a_B^s(\vec{k}, t) + Y^{rs}(\vec{k}, t) a_B^{s\dagger}(-\vec{k}, t)) \\ a_2^r(\vec{k}) &= \cos\theta a_B^r(\vec{k}, t) - \sin\theta \sum_s (W^{sr*}(\vec{k}, t) a_A^s(\vec{k}, t) + Y^{sr}(-\vec{k}, t) a_A^{s\dagger}(-\vec{k}, t)), \end{aligned} \quad (50)$$

$$\begin{aligned} b_1(\vec{k}) &= \cos\theta b_A(\vec{k}, t) - \sin\theta (U^*(k, t) b_B(\vec{k}, t) + V(k, t) a_B^\dagger(-\vec{k}, t)) \\ b_2(\vec{k}) &= \cos\theta b_B(\vec{k}, t) + \sin\theta (U(k, t) b_A(\vec{k}, t) - V(k, t) a_A^\dagger(-\vec{k}, t)), \end{aligned} \quad (51)$$

$$\begin{aligned} c_1(\vec{k}) &= \cos\theta c_A(\vec{k}, t) - \sin\theta (U^*(k, t) c_B(\vec{k}, t) + V(k, t) a_B^\dagger(-\vec{k}, t)) \\ c_2(\vec{k}) &= \cos\theta c_B(\vec{k}, t) + \sin\theta (U(k, t) c_A(\vec{k}, t) - V(k, t) a_A^\dagger(-\vec{k}, t)), \end{aligned} \quad (52)$$

which can be obtained from (37) ( $W, Y, U, V$  being defined in (40) and (41)), the (anti-)commutation relations (42), as well as the property of the *flavor vacuum*

$$a_i^r(\vec{k}, t)|0\rangle_f = b_i(\vec{k}, t)|0\rangle_f = c_i(\vec{k}, t)|0\rangle_f = 0. \quad (53)$$

Being left with functions of the momenta, the last step is to simplify and evaluate the integrals over  $\vec{p}$  and  $\vec{q}$ .

We next proceed, therefore, to give the energy density and pressure of the fluid, obtained by following the above procedure, and discuss the associated novel SUSY-breaking mechanism induced by the flavor condensates in the BV formalism.

### B. Energy density and pressure

After straightforward but quite involved calculations, one finds that the flavored vacuum expectation value of the off-diagonal elements of the stress-energy tensor vanish,  ${}_f\langle 0|T_{\mu\nu}|0\rangle_f = 0$ ,  $\mu \neq \nu$ , as appropriate for a relativistic fluid (in a momentarily comoving rest frame). On the other hand, the energy density  $e$  and pressure  $p$  of the flavor vacuum for the theory (26) are determined to be:

$$\begin{aligned} e &= {}_f\langle 0|T_{00}(x)|0\rangle_f \\ &= \sin^2\theta \frac{(m_1 - m_2)^2}{2\pi^2} \int_0^\Lambda dk k^2 \left( \frac{1}{\omega_1(k)} + \frac{1}{\omega_2(k)} \right) \\ &= \sin^2\theta \frac{(m_1 - m_2)^2}{2\pi^2} f(\Lambda), \end{aligned} \tag{54}$$

$$\begin{aligned} p &= {}_f\langle 0|T_{ii}(x)|0\rangle_f \\ &= -\sin^2\theta \frac{m_1^2 - m_2^2}{2\pi^2} \int_0^\Lambda dk k^2 \left( \frac{1}{\omega_1(k)} - \frac{1}{\omega_2(k)} \right) \\ &= \sin^2\theta \frac{m_1^2 - m_2^2}{2\pi^2} g(\Lambda). \end{aligned} \tag{55}$$

The integrals in (54) and (55) can be evaluated analytically. This yields for the functions  $f(\Lambda)$  and  $g(\Lambda)$

$$\begin{aligned} f(\Lambda) &= \frac{1}{2} \left( \Lambda \omega_1(\Lambda) + \Lambda \omega_2(\Lambda) - m_1^2 \log \left[ \frac{\Lambda + \omega_1(\Lambda)}{m_1} \right] \right. \\ &\quad \left. - m_2^2 \log \left[ \frac{\Lambda + \omega_2(\Lambda)}{m_2} \right] \right) \\ g(\Lambda) &= \frac{1}{2} \left( \Lambda \omega_1(\Lambda) - \Lambda \omega_2(\Lambda) - m_1^2 \log \left[ \frac{\Lambda + \omega_1(\Lambda)}{m_1} \right] \right. \\ &\quad \left. + m_2^2 \log \left[ \frac{\Lambda + \omega_2(\Lambda)}{m_2} \right] \right), \end{aligned} \tag{56}$$

whereby  $\Lambda$  we denote the cutoff in momentum space. The behavior of the energy density and pressure as functions of the cutoff is shown in Figs. 2 and 3.

As we can see, the energy is *positive*. This is not surprising, since we know that in a supersymmetric field-theory model the state with the lowest energy is unique, and corresponds to the usual vacuum  $|0\rangle$ , for which  $\langle 0|T_{00}|0\rangle = 0$ , all other states having positive energies. The difference in energy of the “flavor vacuum” from the mass-eigenstate one is due to the condensation of massive flavored particle states [3,5]. The choice of this vacuum leads therefore to a *breaking of supersymmetry*, in a novel way to be explored further below.

On the other hand, the pressure is *negative*, leading to an equation of state





FIG. 2 (color online). The energy density as a function of the cutoff  $\Lambda$ , with the values:  $m_1 = 1$ ,  $m_2 = 0.9$ , in arbitrary units (sufficient for our purpose here, which is to demonstrate the dependence on  $\Lambda$  only) is plotted. The two plots correspond to two different cutoff regimes: the diagram on the left, shows the behavior of  $e(\Lambda)$  for small  $\Lambda$ . It approaches zero as  $e(\Lambda) \sim \Lambda^3$  for  $\Lambda \rightarrow 0$ ; the plot on the right depicts the behavior of the energy density for large  $\Lambda$ , where we have  $e(\Lambda) \sim \Lambda^2$ . As a typical scale of the theory, with respect to which we define the low and high cutoff scale, we take  $m_1 \approx m_2 = 1$ .




FIG. 3 (color online). The pressure as a function of the cutoff  $\Lambda$ , with the values  $\sin\theta = 2\pi^2$ ,  $m_1 = 1$ ,  $m_2 = 0.9$  is plotted. Again, on the left we have the behavior of  $p(\Lambda)$  for small  $\Lambda$  (compared with  $m_2 \approx m_1 = 1$ ), where  $p(\Lambda) \sim \Lambda^3$ , and on the right for large  $\Lambda$ , where  $p(\Lambda) \sim \log(\Lambda)$ .

$$w = \frac{(m_1 + m_2) g(\Lambda)}{(m_1 - m_2) f(\Lambda)}. \quad (57)$$

As we can see from Fig. 4, for small values of the cutoff (compared with  $m_1 \approx m_2$ ), the equation of state approaches  $-1$ . This is the region in which our *flavor vacuum* might give a contribution to the dark energy, in the form of an approximate cosmological constant. For greater values of the cutoff, the equation of state goes to zero as  $-\log(\Lambda)\Lambda^{-2}M^2$ , with  $M \equiv (m_1 + m_2)^2/2$ , and the fluid approaches that of a relativistic dust.<sup>9</sup>

### C. Different equation of state for bosons and fermions—strong and novel SUSY-breaking

In order to understand the above results and compare them with our previous work [1,2], it is useful to disentangle the various contributions to  $e$  and  $p$  coming from the boson and Majorana fermion fields.

Since the Lagrangian (26) is the sum of six free Lagrangians (two scalars, two pseudoscalars and two

<sup>9</sup>The reader should recall at this stage that an accelerated expanding Universe is obtained when  $w < -1/3$ , for the fluid that fills it in; the flavored WZ fluid respects this property, for sufficiently low values of the cutoff  $\Lambda$ . However, in the current work, we did not perform a self-consistent analysis of the flavor vacuum in such curved space-times, incorporating particle production, as done in [1,2]. Thus, the conclusions one can reach from the above flat space-time analysis is that this type of fluids can potentially find applications to cosmology, once the above step is performed.

Majorana spinors), the stress-energy tensor can be written as the sum of three different terms

$$T_{\mu\nu}(x) = T_{\mu\nu}^S(x) + T_{\mu\nu}^P(x) + T_{\mu\nu}^\psi(x), \quad (58)$$

where the various superscripts denote scalar ( $S$ ), pseudo-scalar ( $P$ ) and fermion ( $\psi$ ) contributions, respectively. The *flavor vacuum* expectation values  ${}_f\langle 0|T_{\mu\nu}^S(x)|0\rangle_f$ ,  ${}_f\langle 0|T_{\mu\nu}^P(x)|0\rangle_f$  and  ${}_f\langle 0|T_{\mu\nu}^\psi(x)|0\rangle_f$  are nonvanishing only for the diagonal elements.

As far as the energy density,  $e$ , is concerned, we have

$$\begin{aligned} {}_f\langle 0|\sum_{i=1,2} T_{00}^{S_i}(x)|0\rangle_f &= {}_f\langle 0|\sum_{i=1,2} T_{00}^{P_i}(x)|0\rangle_f \\ &= \int dk \frac{k^2}{2\pi^2} (\omega_1(k) + \omega_2(k)) \\ &\quad \times \left[ 1 + \sin^2\theta \frac{(\omega_1(k) - \omega_2(k))^2}{2\omega_1(k)\omega_2(k)} \right], \end{aligned} \quad (59)$$

$$\begin{aligned} {}_f\langle 0|\sum_{i=1,2} T_{00}^{\psi_i}(x)|0\rangle_f &= \int dk \frac{k^2}{\pi^2} (\omega_1(k) + \omega_2(k)) \\ &\quad \times \left[ -1 + \sin^2\theta \left( \frac{\omega_1(k)\omega_2(k) - m_1 m_2 - k^2}{\omega_1(k)\omega_2(k)} \right) \right]. \end{aligned} \quad (60)$$

On the other hand, the respective contributions to the pressure are




FIG. 4 (color online). The equation of state  $w$  as a function of the cutoff is plotted; The plot on the left corresponds to  $m_1 = 1$ ,  $m_2 = 0.9$ . In the plot on the right we depict a set of three curves, each of them corresponding to different values of the mass-difference parameter  $\delta$ :  $m_2 \equiv m + \delta$  and  $m_1 \equiv m$ , with  $m = 1$  and  $\delta = 0.5, 0.8, 0.99$ .

$$\begin{aligned}
 {}_f\langle 0 | \sum_{i=1,2} T_{ii}^{S_i}(x) | 0 \rangle_f &= {}_f\langle 0 | \sum_{i=1,2} T_{ii}^{P_i}(x) | 0 \rangle_f \\
 &= \int dk \frac{k^2}{2\pi^2} \left[ \frac{k^2}{3} \left( \frac{1}{\omega_1(k)} + \frac{1}{\omega_2(k)} \right) \right. \\
 &\quad \left. - \sin^2 \theta \frac{m_1^2 - m_2^2}{2} \left( \frac{1}{\omega_1(k)} - \frac{1}{\omega_2(k)} \right) \right], \quad (61)
 \end{aligned}$$

$${}_f\langle 0 | \sum_{i=1,2} T_{ii}^{\psi_i}(x) | 0 \rangle_f = - \int dk \frac{k^4}{3\pi^2} \left( \frac{1}{\omega_1(k)} + \frac{1}{\omega_2(k)} \right). \quad (62)$$

As we anticipated in the beginning of this section, such a theory needs normal ordering with respect to the usual vacuum (46). Now let us look at the individual equations of state:

$$w_b = \frac{{}_f\langle 0 | \sum_{i=1,2} T_{ii}^{S_i}(x) : | 0 \rangle_f}{{}_f\langle 0 | \sum_{i=1,2} T_{00}^{S_i}(x) : | 0 \rangle_f} = \frac{{}_f\langle 0 | \sum_{i=1,2} T_{ii}^{P_i}(x) : | 0 \rangle_f}{{}_f\langle 0 | \sum_{i=1,2} T_{00}^{P_i}(x) : | 0 \rangle_f} = -1, \quad (63)$$

and

$$w_f = \frac{{}_f\langle 0 | \sum_{i=1,2} T_{ii}^{\psi_i}(x) : | 0 \rangle_f}{{}_f\langle 0 | \sum_{i=1,2} T_{00}^{\psi_i}(x) : | 0 \rangle_f} = 0. \quad (64)$$

This means that the bosonic contribution to the vacuum condensate by itself would lead to a *pure cosmological*

*constant type* equation of state [1,2]. This behavior of the fluid is mitigated by the contribution of the Majorana fermionic fluid, that has a *vanishing* pressure.

When just low energies (induced by a low cutoff scale) are taken into account, the bosonic contribution drives the equation of state towards  $-1$ , but as the cutoff increases, the fermionic contribution gets stronger and stronger, leading  $w$  asymptotically to zero value, thus making the fluid behaving as dust in such high energy regimes. The different behavior between fermions and bosons indicates, in addition to the nonzero vacuum energy, a novel strong type of *SUSY-breaking* in the model, purely nonperturbative, induced by the various particle excitations of the flavor vacuum.

The choice of the cutoff is therefore crucial in determining the behavior of the fluid [5]. This implies the necessity for constructing models in which the cutoff scale is determined dynamically by the microscopic theory, as, for instance, the D-brane model of [1,2], a feature that cannot be captured by the simplified free WZ model considered here.

However, we should stress that the cutoff dependence of the condensate is a consequence of the fact that the computations have been done here in flat space-time, and thus some regularization is in order (the reader should compare, for instance, the cutoff dependence of the effective running coupling of the nonrenormalizable four-fermion interaction in the chiral-symmetry-breaking model of Ref. [18], which enters the expression for the respective condensates. When the analysis of chiral-symmetry-breaking is

performed in curved, and, in particular, de Sitter space-time [17], as appropriate for our D-foam framework, any potential ultraviolet (UV) infinities may be absorbed in the cosmological constant and Planck scale of the de Sitter space-time, as demonstrated explicitly in the related analysis of Refs. [16,17]. In our context one should expect a similar behavior, since the flavor-vacuum condensates are also induced in our D-foam model from appropriate (non-renormalizable) four-fermion interactions in the effective low-energy theory (17), which become better behaved in the UV in our (anti) de Sitter backgrounds. Hence the presence of a flavor-vacuum condensate is not an artefact of the UV cutoff.

## V. DISCUSSION

In this work we have examined a novel type of SUSY-breaking in a flavored supersymmetric WZ smodel, induced by mixing in the BV formalism. The strong breaking of SUSY is characterized not only by a nonvanishing flavor-vacuum energy but also by different equations of state between bosonic and fermionic particles. We have motivated the choice of the flavor vacuum over the mass-eigenstate vacuum by an explicit, and by *no means* generic, example of a space-time quantum foam in string theory. The vacuum of such a configuration broke target-space supersymmetry and Lorentz symmetry, as a result of the quantum fluctuations of massive space-time defects. As such, the ordinary supersymmetric and Lorentz invariant vacuum is not appropriate for the description of the underlying physics, and the *flavor vacuum* is a *necessity*. Despite the difference in the setup from those in the models previously studied [1,2], SUSY-breaking is a feature also in this context. Indeed, compared with [1,2], a flat universe instead of a FRW universe is assumed in the present work. Moreover, no MSW effect has been introduced. As a result, BV formalism by itself provides a mechanism for SUSY-breaking.

Such a mechanism is not a *spontaneous* SUSY-breaking in the usual sense: no potential is introduced and formally, if the model is not considered as an effective theory, a state with zero energy (denoted by  $|0\rangle$ ), exists. However, such a state does not represent the *physical* vacuum; in the sense of [3] only flavored particles can be produced/annihilated, if one does not insist on energy conservation at an interaction vertex (taking into account the time-energy uncertainty relation). From our D-particle-foam view point, energy is of course conserved overall in a process where a flavored particle is produced by the “vacuum.” The missing energy from a low-energy field-theory point of view [3], amounting to the difference in energy between the normal mass-eigenstate and flavor vacua, is accounted for in the string-theory picture by the recoil of the D-particle

defects in the foam and the induced condensation of particle-antiparticle flavored particles. The latter amount to stretched string states between the recoiling D-particle and the D3-braneworld in Fig. 1. Such *backreaction* effects of quantum flavored matter onto the surrounding space-time involve stringy/brany degrees of freedom that go beyond the local field-theory formalism.

The strong SUSY-breaking that characterizes such a state is a really new feature here. It would be interesting to discuss other, phenomenologically more realistic models, involving Dirac fermions and, in general, interacting fields [8]. This would allow us to check whether the Majorana fermion fluid zero pressure we have found here is a specific feature of the free WZ model or is valid in general and study the patterns of flavor-induced symmetry-breaking through the associated equations of state. One is tempted to interpret the zero total pressure of the fermionic fluid as a result of an extra contribution, as compared with the bosonic case, due to the Pauli exclusion principle, the so called degeneracy pressure. The latter is a positive contribution to the total pressure of a fermion fluid which is due to the extra force one has to exert as a consequence of the fact that two identical fermions cannot occupy the same quantum state at the same time. The force provided by this pressure sets a limit on the extent to which matter can be squeezed together without collapsing into, e.g., a neutron star or black hole. When two fermions (of the same flavor) are squeezed too close together, the exclusion principle requires them to have opposite spins in order to occupy the same energy level. To add another fermion of this flavor to a given volume (as required by the formation of a condensate) requires raising the fermion’s energy level, and this requirement for energy to compress the material appears as a (positive) pressure. In our cosmological situation, it may be that this positive contribution to the pressure cancels out the cosmological negative pressure of the fermionic fluid, leading to the dustlike behavior we find here. On the other hand, the bosons are not characterized by such degeneracy pressure contributions, and therefore their total (negative) pressure leads to an equation of state resembling that of a cosmological constant [1,2]. These issues, together with the extension of the formalism to incorporate interacting field theories, are currently under investigation.

## ACKNOWLEDGMENTS

The work of N. E. M. and S. S. is partially supported by the European Union through the Marie Curie Research and Training Network *UniverseNet* (MRTN-2006-035863), and W. T. acknowledges support from King’s College London.

- [1] N. E. Mavromatos and S. Sarkar, *New J. Phys.* **10**, 073009 (2008).
- [2] N. E. Mavromatos, S. Sarkar, and W. Tarantino, *Phys. Rev. D* **80**, 084046 (2009).
- [3] M. Blasone and G. Vitiello, *Ann. Phys. (N.Y.)* **244**, 283 (1995); **249**, 363(E) (1996).
- [4] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, England, 1982).
- [5] A. Capolupo, S. Capozziello, and G. Vitiello, *Phys. Lett. A* **363**, 53 (2007); **373**, 601 (2009), and references therein.
- [6] A. Capolupo, M. Di Mauro, and A. Iorio, [arXiv:1009.5041](https://arxiv.org/abs/1009.5041).
- [7] J. R. Ellis, N. E. Mavromatos, and M. Westmuckett, *Phys. Rev. D* **70**, 044036 (2004); **71**, 106006 (2005).
- [8] N. E. Mavromatos, Sarben Sarkar, and W. Tarantino (unpublished).
- [9] See for instance: J. Polchinski, *String Theory* (Cambridge University Press, Cambridge, England, 1998), Vol. 2.
- [10] N. E. Mavromatos and S. Sarkar, *Phys. Rev. D* **72**, 065016 (2005).
- [11] A. George, *J. Phys. A* **38**, 7399 (2005).
- [12] J. R. Ellis, N. E. Mavromatos, and D. V. Nanopoulos, *Phys. Lett. B* **665**, 412 (2008).
- [13] T. Li, N. E. Mavromatos, D. V. Nanopoulos, and D. Xie, *Phys. Lett. B* **679**, 407 (2009).
- [14] I. Antoniadis, K. Benakli, and A. Laugier, *J. High Energy Phys.* **05** (2001) 044; D. Lust, O. Schlotterer, S. Stieberger, and T. R. Taylor, *Nucl. Phys.* **B828**, 139 (2010); D. Lust, S. Stieberger, and T. R. Taylor, *Nucl. Phys.* **B808**, 1 (2009).
- [15] There are many models employing right-handed neutrinos, ranging from purely phenomenological ones to microscopic models obtained from extensions of the standard model and string theory. These models contain four-neutrino interactions that may lead to condensates, with effects in the standard model physics and/or cosmology. For a very partial (and by no means complete) list of recent works, see G. Barenboim and J. Rasero, *J. High Energy Phys.* **03** (2011) 097, where cosmological consequences of a phenomenologically introduced four-right-handed-neutrino interactions have been explored. A. Aparici, A. Santamaria, and J. Wudka, *J. Phys. G* **37**, 075012 (2010); F. del Aguila, J. de Blas, R. Szafron, J. Wudka, and M. Zralek, *Phys. Lett. B* **683**, 282 (2010); R. Tatar, Y. Tsuchiya, and T. Watari, *Nucl. Phys.* **B823**, 1 (2009); M. Endo and T. Shindou, *J. High Energy Phys.* **09** (2009) 037; S. Antusch, J. Kersten, M. Lindner, and M. Ratz, *Nucl. Phys.* **B658**, 203 (2003).
- [16] P. Candelas and D. J. Raine, *Phys. Rev. D* **12**, 965 (1975).
- [17] F. Giacosa, R. Hofmann, and M. Neubert, *J. High Energy Phys.* **02** (2008) 077; S. Alexander, T. Biswas, and G. Calcagni, *Phys. Rev. D* **81**, 043511 (2010); **81**, 069902 (2010).
- [18] Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961); **124**, 246 (1961).
- [19] V. A. Kostelecky and M. Mewes, *Phys. Rev. D* **70**, 076002 (2004); See also: S. H. S. Alexander, [arXiv:0911.5156](https://arxiv.org/abs/0911.5156).
- [20] J. M. Figueroa-O'Farrill, [arXiv:hep-th/0109172](https://arxiv.org/abs/hep-th/0109172).
- [21] M. F. Sohnius, *Phys. Rep.* **128**, 39 (1985).