

Non-Abelian statistics versus the Witten anomaly

John McGreevy and Brian Swingle

Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

(Received 27 June 2011; published 15 September 2011)

This paper is motivated by prospects for non-Abelian statistics of deconfined particlelike objects in $3 + 1$ dimensions, realized as solitons with localized Majorana zero modes. To this end, we study the fermionic collective coordinates of magnetic monopoles in $3 + 1$ -dimensional spontaneously-broken $SU(2)$ gauge theories with various spectra of fermions. We argue that a single Majorana zero mode of the monopole is not compatible with cancellation of the Witten $SU(2)$ anomaly. We also compare this approach with other attempts to realize deconfined non-Abelian objects in $3 + 1$ dimensions.

DOI: [10.1103/PhysRevD.84.065019](https://doi.org/10.1103/PhysRevD.84.065019)

PACS numbers: 14.80.Hv, 03.65.Ge, 05.30.Pr, 73.20.-r

I. INTRODUCTION

Point particles in $3 + 1$ dimensions cannot have non-Abelian statistics because of the triviality of the topology of their configuration space [1]. However, a particlelike object with extra structure can have a configuration space with more interesting homotopy. Inspired by ideas from topological insulators [2], Teo and Kane [3] recently made a specific proposal in this direction. The objects in question are hedgehogs of a 3-component order parameter, coupled to fermionic excitations that are gapped in the presence of a nonzero order parameter. Freedman *et al.* [4] show that these objects exhibit what they call *projective ribbon statistics*; the data needed to specify a configuration include the preimage under the order parameter map of the north pole and a nearby point.

The hedgehog defects support real fermionic zero modes, and multiple hedgehogs are associated with a non-local Hilbert space generated by the zero mode operators. Motions of the hedgehogs implement unitary transformations in the nonlocal Hilbert space, a concept familiar from topological quantum computing in $2 + 1$ dimensions [5]. Because exchanging identical particles leads to a nontrivial unitary transformation of the quantum state instead of merely a phase, we say such objects have non-Abelian statistics. The presence of Majorana zero modes and the nontrivial configuration space are both crucial to this story.

Freedman *et al.* also point out the following problematic feature of the model of Teo and Kane: if the order parameter field has a nonzero stiffness, a single hedgehog is not a finite-energy configuration. Configurations with zero net hedgehog number can have finite energy, but there will be a confining force between the hedgehogs due to gradient energy in the order parameter field. This energy cost will scale at least linearly with the separation between the hedgehogs. The cost may be even higher in the absence of full $SU(2)$ symmetry for the order parameter (and such symmetry is unlikely given that the order parameter involves both superconducting and particle-number-conserving terms). This makes it difficult to imagine adiabatically moving these solitons around each other.

Putting aside the issues with using this proposal as a platform for quantum computing (note further that braiding of Majoranas does not provide a set of universal gates), we would like to confront the conceptual question of whether it is possible in principle to deconfine such non-Abelian particles in $3 + 1$ dimensions. We are also interested more generally in what happens to Majorana zero modes when the relevant order parameter field begins to fluctuate. If we were able to deconfine non-Abelian particles in $3 + 1$ dimensions, there would be profound practical and conceptual implications.

One suggestion for removing the confining energy follows the analogous step in the study of vortices in two dimensions: gauge the rotation symmetry in the order parameter space. If all directions are gauge equivalent, there need not be a confining energy between the hedgehogs, which in the resulting gauge theory are 't Hooft-Polyakov monopoles [6,7]. (There will be a magnetic Coulomb force between the monopoles, but this falls off with their separation.) But $3 + 1$ dimensional $SU(2)$ gauge theory with the requisite fermion content, namely, a single Weyl doublet (i.e. eight Majorana fields), suffers from the Witten $SU(2)$ anomaly [8] (as [4] also observe). One implication of this is that the gauge field partition sum vanishes identically. Another pathology resulting from the anomaly is a violation of fermion parity by the gauge dynamics. Specifically, an instanton creates a single fermion in violation of fermion parity. The addition of an adjoint Higgs field (relative to the discussion of [8]) does not change the structure of the fermion determinant which is responsible for the fatal factor of -1 (which it acquires under the gauge transformations which represent the nontrivial element of $\pi_4(SU(2))$), as we argue below in Sec. II A.

We will construct below a microscopically-consistent theory which, in a range of energy scales, looks like this Witten-anomalous $SU(2)$ gauge theory in the Higgs phase with a single Weyl doublet. The spectrum of fermionic particles with $E < M_W$ is identical to that of the theory described above; at these energies, the Witten anomaly is cancelled by a certain Wess-Zumino-Witten term made from the adjoint scalar and the gauge field. (This situation

is similar, but not identical, to models discussed by d’Hoker and Farhi [9,10].) However, this term is ill-defined when the order parameter vanishes, as it does in the core of the monopole, and we must provide a UV completion to address the question of whether the monopole has a Majorana zero mode. The simplest UV completion of this model involves adding in another Weyl fermion doublet.

Before proceeding with an analysis of the $SU(2)$ gauge theory, we pause to consider an alternative possible route to deconfine the localized objects hosting Majorana modes. Instead of gauging the $SU(2)$ symmetry that is spontaneously broken by the order parameter field in the Teo-Kane model, we can consider disordering the broken phase into a liquidlike phase without any broken symmetry [4,11]. Importantly, we must achieve this disordering without proliferating the monopole defects that hosted Majorana modes, otherwise we will trivially lose the localized Majorana mode. The simplest-to-describe disordered phase has a description in terms of an emergent $U(1)$ gauge field, and the hedgehog defects, assuming they have finite energy, become magnetic monopoles in the $U(1)$ gauge theory: we are again led to a description in terms of magnetic charges in an Abelian gauge theory. (We describe other possibilities for disordered phases in the last section.) Now the important question is do the Majorana modes survive the disordering process, and if so, are the magnetic monopoles in this theory deconfined (only interacting via a long range Coulomb interaction) particles carrying Majorana zero modes? Again the question of the survival of the Majorana modes requires short-distance information about the theory. Later we will return to this question for the disordered state, arguing on general grounds that this particular scenario is unlikely.

To clarify, our desiderata for deconfined non-Abelian excitations in $3 + 1$ dimensions are as follows. First, we will discuss the desired form of the regulated theory at high energies and then the form of the theory at low energies. From the point of view of condensed matter physics, we would most like to have a microscopic lattice model involving only spinlike or electronlike degrees of freedom that enters a phase where there are deconfined particles. We do not accept as a valid realization a model that contains Majorana fermion degrees of freedom in a microscopic lattice model. We make this requirement because we do not want to put the Majoranas in “by hand.” However, we would permit a Majorana based lattice model provided we could reinterpret it as an intermediate scale description arising from a truly microscopic model of electrons, likely in the presence of superconductivity (a bipartite lattice is a sufficient condition). From the point of view of high energy physics, we would like to have an anomaly free gauge theory coupled to fermions and scalars that has a nonperturbative regularization of some type, be it lattice gauge theory or string theory. In the high energy way of thinking, we do not require the absence of gauge fields in the micro-

scopic description, for example, we would accept an asymptotically free gauge theory interacting with Dirac fermions.

In the low energy theory, we have two general interests. First, any putative non-Abelian particlelike excitations should have a clearly defined configuration space. We should have a clear understanding of the nonlocality inherent in this configuration space that permits otherwise pointlike objects to have interesting statistics. Second, it must be possible to perform motions of the non-Abelian excitations without high energy cost, without dramatically exciting other degrees of freedom, without violating causality or unitarity, and without producing decoherence in the space of “protected” states. For example, decoherence due to unscreened gauge fields limits our ability to superpose states with macroscopically different charge configurations. We emphasize especially the issue of the low energy configuration space. This space must be rich enough to support representations of its fundamental group that are nonlocal, as with non-Abelian anyon representations of the braid group in $2 + 1$. The symmetric group is known to be insufficient for this purpose [12,13], and indeed as a finite group its image in any unitary group must be quite limited.

In this paper we study the possibility of non-Abelian particlelike excitations in a $3 + 1$ dimensional field theory. In particular, we explore the apparent conflict between a single Majorana zero mode of the ’t Hooft-Polyakov monopole (we will refer to such an object as a “Majorana monopole”) and microscopic consistency of the $SU(2)$ gauge theory.

The outline of the paper is as follows. In the next section we generalize the classic analysis of Jackiw and Rebbi [14] to construct the zero mode solution of the Dirac equation in the Witten-anomalous theory described above. In Sec. III, we discuss the cancellation of the Witten anomaly and its effects on the zero mode structure of the monopole. In Sec. IV, we discuss an instructive example in $4 + 1$ dimensions. In Sec. V, we provide general arguments for obstructions to Majorana monopoles in $3 + 1$ dimensions following the desiderata described above.

Related work appears in [15], which studies an interesting fermion dimer model whose low energy physics includes Majorana monopoles interacting with gauge fields as well as gapless fermionic degrees of freedom. Some features of three-dimensional non-Abelian particles appear to be realized in their model, but we emphasize that their conclusions do not contradict our own; our analysis suggests that the gapless fermions are essential. Reference [15] also studies a five-dimensional model similar to the one discussed in Sec. IV.

II. MAJORANA MONOPOLES IN AN ANOMALOUS THEORY

Consider an $SU(2)$ gauge theory in $3 + 1$ dimensions with a scalar field Φ in the adjoint representation; we will

suppose that the action for Φ is such that in the ground state it breaks SU(2) down to U(1). Include also a single SU(2) doublet of Weyl fermions, χ ; altogether there are $2^3 = 8$ real fermion degrees of freedom. This is half as many fermion degrees of freedom as considered by Jackiw and Rebbi in their 3 + 1-d discussion [14] and the same number as considered by Witten [8]. As we demonstrate next, this theory suffers from the Witten anomaly—if we try to quantize the gauge field, we get nonsense. Specifically, the partition function vanishes and expectation values of gauge-invariant observables are undefined. For the discussion in this section the bosonic fields Φ, A will therefore be treated as background fields.

Consider the fermion Lagrange density

$$L_{\text{fermions}} = \chi^\dagger i \bar{\sigma}^\mu D_\mu \chi - \frac{1}{2} \lambda \chi^T i \sigma^2 i \tau^2 \vec{\tau} \cdot \vec{\Phi} \chi + \text{H.c.} \quad (2.1)$$

Here $\chi_{\alpha a}$ is a (left-handed) Weyl doublet of SU(2): $\alpha = 1, 2$ is a spin index, $a = 1, 2$ is a gauge index. $\bar{\sigma}^\mu = (1, -\vec{\sigma})^\mu$. The covariant derivative is defined as $(D_\mu \chi)_{\alpha a} = \partial_\mu \chi_{\alpha a} - i g A_{\mu ab} \chi_{\alpha b}$, where $A_{\mu ab}$ is the SU(2) gauge field. λ is a complex coupling constant. Note that because χ is in a pseudoreal representation of both the SU(2) gauge group and the Lorentz group, the object $\chi^C \equiv \chi^T i \sigma^2 i \tau^2$ transforms in the conjugate representation of both groups. There is no nonvanishing, gauge-invariant and Lorentz-invariant mass term (not involving the Higgs field Φ) with this field content. We will comment in Sec. III D on the effect of Lorentz-breaking terms of the form $\chi^\dagger \chi$.

This theory has two independent mass scales: the mass of the W bosons, $m_W = g v$ (v is the vev of the adjoint Higgs field, g is the SU(2) gauge coupling at the scale m_W), and the mass of the fermion λv .

A. Persistence of Witten anomaly

The addition of the adjoint scalar Φ and its coupling to the fermion doublet does not modify the anomalous transformation law of the fermion determinant. That this is the case can be seen by embedding the theory in an SU(3) gauge theory with a perturbative gauge anomaly as in [16–18]. The relevant theory has an SU(3) adjoint scalar $\tilde{\Phi}$, an SU(3) triplet of Weyl fermions $\tilde{\chi}$, and an SU(3) triplet of scalars Y , with the coupling

$$L_{\text{SU}(3)} \supset \tilde{\chi}_a Y_b \epsilon_{abc} \tilde{\Phi}_{cd} \tilde{\chi}_d, \quad (2.2)$$

where $a = 1, 2, 3$ is a triplet index. Condensing the scalar triplet $\langle Y \rangle = \lambda$ breaks the SU(3) down to SU(2), and the coupling (2.2) reduces to the desired coupling between the Weyl fermions charged under the unbroken SU(2) and the adjoint scalar in (2.1). The form of the perturbative SU(3) anomaly is unaffected by the addition of scalars and so the calculation of the variation of the fermion measure by integrating the SU(3) anomaly [16–18] is unmodified compared to the theory without scalar fields.

B. The Majorana zero mode

The Dirac equation which results from varying (2.1) is

$$0 = \delta_{\tilde{\chi}} S_{\text{fermion}} = -i \bar{\sigma}^\mu D_\mu \chi + \lambda^\dagger i \sigma^2 \Phi \cdot \tau i \tau^2 \chi^*. \quad (2.3)$$

We consider this Dirac equation in the background of the 't Hooft-Polyakov monopole solution,

$$A_0^B = 0; \quad A_i^B = \epsilon_{ijB} \hat{r}^j A(r); \quad \Phi^B = \hat{r}^B \phi(r) \quad (2.4)$$

($B = 1, 2, 3$ is an adjoint index) with

$$A(r) \stackrel{r \rightarrow \infty}{\approx} 1/r, \quad \phi(r) \stackrel{r \rightarrow \infty}{\approx} v. \quad (2.5)$$

A zero-energy solution of (2.3) is of the form $\chi_{\alpha a} = i \tau_{\alpha a}^2 g(r)$ (where α is the spin index and a is the SU(2) index). This is the same ansatz as in equation A4 of [14]. With this substitution, the zero mode equation reduces to

$$(\partial_i + 2 \hat{r}_i A) g + i \lambda \phi \hat{r}_i g^* = 0. \quad (2.6)$$

By rephasing the χ field, we can assume WLOG that λ is real and positive. The solution for g is then

$$g(r) = c e^{-\pi i/4} e^{-\int^r (\lambda \phi - 2A)}, \quad (2.7)$$

where c is a *real* constant. We emphasize that the phase of the normalizable solution is determined by normalizability of the solution at large r .

Quantizing this fermionic collective coordinate gives a Majorana fermion acting on the monopole Hilbert space, which is represented by a unique state. This leads inevitably to non-Abelian statistics for the monopoles, in the same manner as expected for vortices in $p + ip$ superconductors or the Pfaffian quantum Hall state [5, 19–22]. Briefly, two widely-separated monopoles will have two Majorana zero modes, which can be combined into $c = \frac{\gamma_1 + i \gamma_2}{2}$, with $\{c, c^\dagger\} = 1$; this algebra must be represented by a two-state system. Interchanging the monopoles adiabatically implements the operator

$$U_{1 \rightleftharpoons 2} = \exp\left(\frac{\pi}{4} \gamma_1 \gamma_2\right) = \exp\left(i \frac{\pi}{4} (1 - 2c^\dagger c)\right).$$

With two pairs of monopoles we could perform operations which do not commute with each other.

We note that the coupling of χ to the gauge field does not play a crucial role in generating this zero mode; since [by (2.5)] the dominant term in the exponent of (2.7) at large r comes from the scalar profile, the gauge field can be set to zero without interfering with the zero mode. The existence of the zero mode solution without the gauge field essentially follows from the analysis of [3].

If only this were a consistent quantum system. We describe one pathology of this system. Recall that in the case of a Dirac fermion there is a complex fermion zero mode in the ungauged theory. Once the SU(2) symmetry is gauged, the low energy gauge group is U(1) and hedgehog configurations become magnetic monopoles. Now what

happens to the two states living on a hedgehog in the ungauged theory? In fact [14], they become bosonic, having charge $\pm 1/2$ under the unbroken $U(1)$ due to the low energy $U(1)$ theta term of π . To see this, assume that the charge $-1/2$ state is bosonic, then when we add a fermion in the zero mode of charge 1 we reach a state of charge $1/2$ which would appear to differ in spin by $1/2$ from the bosonic state. But we have forgotten the gauge field which adds extra angular momentum. Indeed, a unit charge orbiting a minimal monopole leads to a gauge field configuration with angular momentum given by a half integer. This extra half integer angular momentum when combined with the bare half integer angular momentum of the fermion leads again to a bosonic state. In fact, we can check from the structure of the zero mode that the position and spin of the fermion are correlated so that no matter where the fermion is measured its spin will always compensate the \hat{r} angular momentum contribution coming from the field.

Now the puzzle: in the case of a single Weyl fermion, we found that the complex fermion was replaced by a real zero mode, but what should happen when we turn on the gauge field? Heuristically, we should obtain half of the pair of states with charges $\pm 1/2$. Let \mathcal{O}_{+1} be an operator that moves us from the $-1/2$ to the $1/2$ charge state so that \mathcal{O}_{+1} carries charge 1. By analogy with the definition of the Majorana fermion, an apparently interesting combination to consider is $\mathcal{O}_{+1} + \mathcal{O}_{+1}^\dagger$, but this operator creates states that decohere in the presence of the fluctuating $U(1)$ field; we can identify no candidate for the pointer states into which they should decohere. Is the Witten anomaly to blame? The simplest resolution of the Witten anomaly, namely, adding a second identical Weyl doublet, removes the spectre of decoherence by adding an extra real zero mode in the monopole core allowing for complex solutions, as we will see next.

To summarize, we found an $SU(2)$ gauge theory where magnetic monopoles of an unbroken $U(1)$ gauge field appear to carry Majorana zero modes. However, this theory suffers from the Witten anomaly rendering all gauge-invariant observables ill-defined. Related pathologies include a violation of fermion number by instantons and decoherent $U(1)$ charge superpositions. In what follows, we try to cure the Witten anomaly while preserving the zero mode structure of the monopole.

III. CANCELLING THE WITTEN ANOMALY

It is possible to cancel the Witten anomaly by adding to the action a certain functional of the adjoint scalar. To see that this is the case, consider integrating out a Weyl fermion χ_2 coupled to the scalar field as above:

$$\ln \int D\chi_2 \exp(iS_{\text{ferm}}[\chi_2]) = \Gamma[\Phi, A] + \text{nonuniversal stuff.} \quad (3.1)$$

The functional Γ defined by this equation is well behaved because of the gap in the fermion spectrum. The both-hand side of Eq. (3.1) must shift by $\pi \pmod{2\pi}$ under an $SU(2)$ gauge transformation representing the nontrivial class of $\pi_4(SU(2))$. The fact that the nonuniversal, short-distance stuff on the right-hand side (rhs) does not accomplish this shift follows because it is not sensitive to the topology of spacetime.

It is difficult to give an explicit expression for the functional Γ . Naively, the WZW term for $SU(2)$ vanishes identically. However, our term is not quite the usual WZW term since it arises from a Pfaffian rather than a determinant, i.e. it is invariant under only real-linear basis changes. A similar situation with different fermion representations arises in [9,10], where the effective action contains terms taking the form of the gauge variation of our functional Γ . In Sec. II A, we have determined the anomalous transformation of Γ by embedding into a theory with a perturbative anomaly; this trick does not immediately determine the form of Γ itself. It would be useful to find an explicit expression for this functional.

One thing about Γ , however, is certain: it is ill-defined when the order parameter Φ is not invertible. A simple argument for this is that only when Φ is invertible are the fermion degrees of freedom gapped. Therefore, in any field configuration where Φ vanishes, such as the core of the magnetic monopole, a model where the Witten anomaly is cancelled by the variation of $\Gamma[\Phi]$ requires a UV completion.

The simplest way to do this is obviously to integrate in the second Weyl doublet χ_2 by which we proved the existence of Γ ; we study this possibility next. Are there other ways? In the final section, we will argue that the answer is “no”.

A. Generic couplings in the two-Weyl-doublet theory

Consider the fermion lagrangian density

$$L_{2\text{fermions}} = \chi^{I\dagger} i\bar{\sigma}^\mu D_\mu \chi_I - \lambda^{IJ} \chi_I^T i\sigma^2 i\tau^2 \vec{\tau} \cdot \vec{\Phi} \chi_J + \text{H.c.} - m^{IJ} \chi_I^T i\sigma^2 i\tau^2 \chi_J + \text{H.c.} \quad (3.2)$$

Here $\chi_{I\alpha a}$ are a pair of (left-handed) Weyl doublets of $SU(2)$: $I = 1, 2$ is a flavor index, $\alpha = 1, 2$ is a spin index, $a = 1, 2$ is a gauge index. Altogether there are now $2^3 = 8$ complex fermion degrees of freedom. This is the same set of fermion degrees of freedom considered by Jackiw and Rebbi and twice as many as considered by Witten.

We now comment on symmetries of this action and simplifications that can be made by field redefinitions of the fermions. The Yukawa coupling term is more explicitly written as

$$\lambda^{IJ} \chi_I^T i\sigma^2 i\tau^2 \vec{\tau} \cdot \vec{\Phi} \chi_J + \text{H.c.} = \lambda^{IJ} \chi_I^T i\sigma^2 i\tau^2 \vec{\tau} \cdot \vec{\Phi} \chi_J + \lambda^{\dagger IJ} \chi_I^\dagger \vec{\tau} \cdot \vec{\Phi} i\sigma^2 i\tau^2 \chi_J^*.$$

The matrix λ is symmetric, $\lambda^{IJ} = \lambda^{JI}$ by Fermi statistics. A general complex symmetric matrix is not diagonalizable but rather has different right and left eigenvalues. A complex symmetric matrix has a singular value decomposition (SVD) (called Takagi decomposition) of the form

$$\lambda = WdW^T, \quad (3.3)$$

where d is diagonal with real, positive entries

$$d = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix},$$

and W is unitary.

Rephasing the fermion fields by a unitary rotation $U = (U^{-1})^\dagger$

$$\chi_I \rightarrow U_I^J \chi_J \quad (3.4)$$

changes the coupling matrix λ by

$$\lambda \rightarrow U\lambda U^T = UWdW^T U^T. \quad (3.5)$$

Choosing $U = W^{-1}$ gives $\lambda = d$.

By Fermi statistics, the Dirac mass matrix m_{IJ} is anti-symmetric, $m_{IJ} = m\epsilon_{IJ}$. The effect of the rephasing (3.7) on the Dirac mass is therefore $m_{IJ} \rightarrow m_{IJ} \det U$. Having fixed our freedom to rephase the fermions, the phase of the Dirac mass m will be significant. Global symmetries can constrain the phase of m . In particular, with a Hermitian mass matrix, $m = m^\dagger$, the model preserves a CP symmetry which acts by $\chi \mapsto i\sigma^2 i\tau^2 \chi^*$.

When λ is purely off-diagonal

$$\lambda = \begin{pmatrix} 0 & \lambda_0 \\ \lambda_0 & 0 \end{pmatrix}$$

the system admits an extra $U(1)$ symmetry under which

$$\chi_1 \mapsto e^{i\theta} \chi_1, \quad \chi_2 \mapsto e^{-i\theta} \chi_2. \quad (3.6)$$

When the Dirac mass vanishes, the resulting model is identical to the model studied in [14]. To see this, construct from the two left-handed Weyl doublets a single Dirac fermion

$$\Psi \equiv \begin{pmatrix} \chi_1 \\ \chi_2^* i\tau^2 i\sigma^2 \end{pmatrix}. \quad (3.7)$$

Then the action (3.2), with λ off diagonal and $\lambda_0 \equiv \lambda_0^R + i\lambda_0^I$, is

$$L_{2\text{fermions}} = \bar{\Psi} i\not{D}\Psi - \bar{\Psi}(\lambda_0^R + i\lambda_0^I \gamma^5) \vec{\tau} \cdot \vec{\Phi} \Psi + m \bar{\Psi} \Psi. \quad (3.8)$$

Returning to the SVD form of the action, this is equivalent to the case where the diagonal entries are equal $\lambda_1 = \lambda_2$. In this basis, the $U(1)$ symmetry acts as the $SO(2)$ rotation

$$\chi_1 + i\chi_2 \mapsto e^{i\theta} (\chi_1 + i\chi_2). \quad (3.9)$$

The general two-Weyl-doublet theory now has three mass scales: the mass of the W bosons, $m_W = gv$, and the masses of the two Weyl fermions $\lambda_{1,2}v$. In the regime

$$\lambda_1 v \ll m_W \ll \lambda_2 v \quad (3.10)$$

we have a large window of energies in which the bulk spectrum is that of the Witten-anomalous theory studied above.

We note that the theory with two Weyl doublets admits a Lorentz-violating (but gauge-invariant and rotation-invariant) mass term of the form

$$L_{\text{NR}} = M^{IJ} \chi_I^\dagger \chi_J. \quad (3.11)$$

We will comment below in Sec. III D on its effects on the zero mode structure.

B. FZMs in the two-Weyl-doublet theory

The Dirac equation is now

$$\begin{aligned} 0 &= \delta_{\bar{\chi}_I} S_{\text{fermion}} \\ &= -i\vec{\sigma} \cdot D_\mu \chi_I + \lambda_{IJ}^\dagger i\sigma^2 \Phi \cdot \tau i\tau^2 \chi_J^* + m_{IJ}^\dagger i\sigma^2 i\tau^2 \chi_J. \end{aligned} \quad (3.12)$$

When the Dirac mass $m = 0$, in the basis where λ is diagonal, the zero mode equations for $\chi_{1,2}$ decouple, and each is of the form of (2.7). There are then two real solutions:

$$\chi_{I\alpha a}(r) = i\tau_{\alpha a}^2 g_I, \quad g_I = c_I e^{-\pi i/4} e^{-\int^r (\lambda_i \phi - 2A)}. \quad (3.13)$$

C. Effect of the Dirac mass

With a nonzero Dirac mass, the zero mode equations for $\chi_{1,2}$ are coupled. A nonzero Dirac mass requires any putative zero mode solution to include also a triplet component, i.e. to have the more general form

$$\chi_{\alpha a I} = i\tau_{\alpha a}^2 g_I + i(\tau^2 \tau^i)_{\alpha a} g_I^i. \quad (3.14)$$

The zero-energy Dirac equation is

$$0 = i\vec{\sigma} \cdot \vec{D}\chi - \lambda \vec{\tau} \cdot \vec{\Phi} i\sigma^2 i\tau^2 \chi^* + m i\sigma^2 i\tau^2 \chi^*. \quad (3.15)$$

Here we have assumed $m^\dagger = +m$, and more specifically

$$\lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \quad m = i \begin{pmatrix} 0 & \mu \\ -\mu & 0 \end{pmatrix} \quad (3.16)$$

with $\lambda_{1,2}, \mu$ real and positive. The reality of μ (which implies that the Dirac mass matrix is Hermitian) is not fully general; we return to this point anon.

Following [14], let

$$\chi_{\alpha a I} = \mathcal{M}_{abI} i\tau_{ba}^2 = (\delta_{ab} g_I + \sigma^i_{ab} g_I^i) \epsilon_{ba}. \quad (3.17)$$

This decomposition incorporates the breaking of $SU(2)_{\text{gauge}} \times SU(2)_{\text{spin}}$ and decomposes $\chi \in (2, 2)$ into irreps of the unbroken $SU(2)$. It reduces the Dirac equation to the two equations:

$$\begin{aligned}
0 &= i\vec{\nabla}g - 2i\hat{r}Ag - 3\vec{g} \times \hat{r}A + \lambda^\dagger g^* \phi \hat{r} - m^\dagger \vec{g}^* \\
&\quad + i\vec{\nabla} \times \vec{g} + \lambda \phi \hat{r} \times \vec{g}^* \\
0 &= i\vec{\nabla} \cdot \vec{g} + 2i\vec{g} \cdot \hat{r}A - \lambda^\dagger \vec{g}^* \cdot \hat{r}\phi + m^\dagger g^*. \quad (3.18)
\end{aligned}$$

The last term in (3.18) forces us to include a nonzero \vec{g} when $m \neq 0$. The equations for g, \vec{g} are (not too surprisingly) similar to [14] equation A7a, b, with extra terms coming from the Dirac mass.

We make an ansatz of the form $\vec{g} = \hat{r}g_r(r)$. This eliminates the curl terms in the Dirac equation, leaving

$$\begin{aligned}
0 &= i\vec{\nabla}g - 2i\hat{r}Ag - \lambda^\dagger g^* \phi \hat{r} - m^\dagger \vec{g}^* \\
0 &= i\vec{\nabla} \cdot \vec{g} + 2iA\vec{g} \cdot \hat{r} + \lambda^\dagger \vec{g}^* \cdot \hat{r}\phi + m^\dagger g^*. \quad (3.19)
\end{aligned}$$

We choose the phases of g, g_r so that $ig = g^*$

$$g \equiv \alpha h, \quad g_r \equiv \alpha^{-1} h_r, \quad \alpha \equiv e^{-\pi i/4}. \quad (3.20)$$

The Dirac equation becomes

$$0 = \vec{\nabla}h - 2\hat{r}Ah + \lambda^\dagger h \phi \hat{r} + \mu \epsilon \vec{h} \quad (3.21)$$

$$0 = \vec{\nabla} \cdot \vec{h} - 2A\vec{h} + \lambda^\dagger \vec{h} \cdot \hat{r}\phi - \mu \epsilon h. \quad (3.22)$$

In (3.21) and (3.22), all complex phases are explicit. With the assumption (3.16), we have $\lambda^\dagger = \lambda$ is diagonal. The ϵ symbol acts on the IJ flavor indices and is the only thing which does.

The particular solution of (3.22) for \vec{h} given the source h is

$$h_r = +\mu \epsilon r^{-2} e^{-\tilde{H}} \int^r s^2 e^{\tilde{H}} h(s), \quad (3.23)$$

where $\tilde{H} \equiv \int^r (\tilde{\lambda}\phi - 2A)$ and $\tilde{\lambda}$ has the property that

$$\tilde{\lambda}m = m\lambda, \quad (3.24)$$

which in turn requires

$$\tilde{\lambda} = \begin{pmatrix} \lambda_2 & 0 \\ 0 & \lambda_1 \end{pmatrix}. \quad (3.25)$$

Plugging the solution (3.23) into (3.21) (and remembering that $\epsilon^2 = -1$) gives

$$\partial_r h + (\lambda\phi - 2A)h = +\mu^2 r^{-2} e^{\tilde{H}} \int^r ds s^2 e^{-\tilde{H}} h. \quad (3.26)$$

Substituting $h = e^{-H}\gamma$ with $H \equiv \int^r (\lambda\phi - 2A)$ gives

$$r^2 \partial_r \gamma = \mu^2 e^{H-\tilde{H}} \int^r ds s^2 e^{\tilde{H}-H} \gamma. \quad (3.27)$$

Differentiating (3.27) (and thereby introducing an extra integration constant) gives the linear second-order ODE for γ :

$$r^{-2} \partial_r e^{\tilde{H}-H} r^2 \partial_r \gamma = \mu^2 e^{\tilde{H}-H} \gamma. \quad (3.28)$$

We know the asymptotic behavior of the solutions at large and small r . At small r , $H, \tilde{H} \rightarrow 0$, and the Eq. (3.28) reduces to the Helmholtz equation

$$\nabla^2 \gamma \equiv r^{-2} \partial_r r^2 \partial_r \gamma = \mu^2 \gamma, \quad (3.29)$$

whose solutions are

$$\gamma \stackrel{r \rightarrow 0}{\approx} c^{(-)} \frac{e^{-\mu r}}{r} + c^{(+)} \frac{e^{+\mu r}}{r}. \quad (3.30)$$

The combination of these solutions which also solves the integro-differential Eq. (3.27) in the small- r regime has $c^{(+)} = -c^{(-)} \equiv -c$:

$$\gamma \stackrel{r \rightarrow 0}{\approx} \frac{c}{r} (e^{-\mu r} - e^{+\mu r}). \quad (3.31)$$

Note that only the combination $\tilde{H} - H = \lambda_T \mathbb{1} \int^r \phi$ (where $\mathbb{1}_{IJ} \equiv \delta_{IJ}$, $\lambda_T = \lambda_1 - \lambda_2$) enters this equation. We emphasize that there is one such solution for each value of the flavor index $I = 1, 2$ belled by a real integration constant c_I :

$$\gamma_I \stackrel{r \rightarrow 0}{\approx} \frac{c_I}{r} (e^{-\mu r} - e^{+\mu r}). \quad (3.32)$$

In the special case where the eigenvalues of λ are degenerate, $\lambda_T = 0$, the equation for γ is exactly the Helmholtz equation. In this case, the solution (3.32) is exact.

At large r , $\phi(r) \approx r^{-\infty} v$ and $A(r) \approx r^{-\infty} \frac{a_0}{r}$. Therefore

$$H \stackrel{r \rightarrow \infty}{\approx} \begin{pmatrix} \lambda_1 v r & 0 \\ 0 & \lambda_2 v r \end{pmatrix}, \quad \tilde{H} - H \stackrel{r \rightarrow \infty}{\approx} \lambda_T v r \mathbb{1}. \quad (3.33)$$

To discuss the normalizability of the solutions at large r , we distinguish various parameter regimes.

- (i) If $\mu = 0$, both solutions γ_I in (3.32) are normalizable for all $\lambda_{1,2}$. Varying the signs or phases of $\lambda_{1,2}$ is innocuous; it merely changes the overall phase of the zero mode solution and can be absorbed in a field redefinition.
- (ii) For small μ ,

$$\mu < \frac{1}{2} |\lambda_1 - \lambda_2| v, \quad (3.34)$$

both zero modes are still normalizable.

- (iii) Since the zero mode wave functions involve products of exponentials of the form $e^{\mu r} e^{-\lambda v r}$, one might have thought (pantingly) that one zero mode would become non-normalizable, e.g. for μ in between the two Yukawa-induced fermion masses

$$\lambda_1 v < \mu < \lambda_2 v. \quad (3.35)$$

This hope is not realized—there is no change in the normalizability of the modes at $\mu = \lambda_1 v$.

- (iv) For μ larger than the geometric mean of the fermion masses,

$$\sqrt{\lambda_1 \lambda_2} v < \mu, \quad (3.36)$$

both modes are non-normalizable. There is no value of the parameters for which an odd number of Majorana modes are normalizable.

It is interesting to note that we are free to tune the effective sizes of the two real zero modes independently of each other. By adjusting λ_1 and λ_2 we can produce a shell-like configuration of zero modes. More precisely, by making one of the fermion masses very heavy, we can arrange [in the parameter regime (3.35)] for only one zero mode to have a sizable wave function until very close to the monopole core, as shown in Fig. 1. Whether this separation of scales could in principle allow for interesting physical effects is not clear to us.

Note that the variation of the bulk fermion spectrum with μ corroborates the understanding of the normalizability properties of the zero modes presented above. The product of the bulk fermion masses is the determinant of the fermion mass matrix

$$M = v\lambda \otimes \tau^3 + m \otimes \mathbb{1} = \begin{pmatrix} \lambda_1 v & i\mu & 0 & 0 \\ -i\mu & \lambda_2 v & 0 & 0 \\ 0 & 0 & -\lambda_1 v & i\mu \\ 0 & 0 & -i\mu & -\lambda_2 v \end{pmatrix} \quad (3.37)$$

which is

$$\det M = (\lambda_1 \lambda_2 v^2 - \mu^2)^2. \quad (3.38)$$

Comparing (3.38) to the condition for normalizability of the zero modes on the monopole, (3.36), we see that precisely when the zero modes become marginally normalizable, there is a massless fermion in the bulk. For μ above the critical value, the zero modes leak out of the monopole core and join the bulk states.

If m is not Hermitian, any rephasing analogous to (3.20) produces overconstraining equations: the solutions are forced to have nonzero energy. As we discussed above, a CP symmetry can enforce Hermiticity of m .

D. Nonrelativistic mass

The nonrelativistic mass $m\chi^\dagger\chi$ appears in the Dirac equation in precisely the same way as the energy. In fact, this term is nothing but a chemical potential for the chiral symmetry, and thus it clearly breaks Lorentz invariance while preserving rotational symmetry. As the full chiral symmetry is anomalous, this term produces a finite density of fermions carrying a nonconserved charge. This symmetry is also explicitly broken by the scalar coupling, and so even without the anomaly the chiral symmetry is broken as in a superconductor. As the fermion spectrum remains fully gapped in the presence of the scalar coupling, we expect that the nonrelativistic mass does not seriously affect the zero-mode spectrum. This must be true in the ungauged




FIG. 1 (color online). Top: The profile of the zero mode solution for $\mu = 0$. Bottom: One of the profiles in the parameter range (3.35), exhibiting the ringlike structure.

theory of a single Weyl doublet coupled to a scalar field in the adjoint, as such a theory has only a single Majorana mode on a hedgehog that cannot pair and disappear.

IV. FIVE-DIMENSIONAL REALIZATION

Consider $SU(2)$ Yang-Mills theory in $4 + 1$ dimensions with a five-dimensional Dirac fermion in the doublet representation, and an adjoint scalar in its condensed phase. Identify the fourth spatial dimension $y \simeq y + 2\pi R$. Consider a kink-antikink configuration of the five-dimensional Dirac mass $M(y)$ of the fermion, with the kink and antikink on opposite sides of the circle, that is

$$M = +m, y \in (0, \pi R), \quad M = -m, y \in (\pi R, 2\pi R). \quad (4.1)$$

The kink and antikink each support a four-dimensional massless Weyl fermion (for a useful review, see [23]). We can arrange for the four-dimensional coupling to the scalar field that we have been considering by using the fact that spinor representations in five-dimensions are also pseudoreal (the Lorentz group is equivalent to a symplectic group which has a real invariant form). Using the four-dimensional chiral basis, a five-dimensional spinor may be

written $\Psi = [\psi_L, \psi_R]^T$. The combination $\psi_L^T i\sigma^2 \psi_L + \psi_R^T i\sigma^2 \psi_R$ is manifestly invariant under four-dimensional Lorentz transformations. The extra four transformations in the five-dimensional Lorentz group, generated by $[\gamma^4, \gamma^\mu]$, act infinitesimally like $\delta\psi_L = \epsilon\sigma^\mu \psi_R$ and $\delta\psi_R = -\epsilon\bar{\sigma}^\mu \psi_L$. The invariance under five-dimensional Lorentz transformations then follows from the identity $(\sigma^\mu)^T i\sigma^2 + i\sigma^2 \bar{\sigma}^\mu = 0$. (That is, the symplectic invariant of $SO(4, 1)$ is $\Omega \equiv \gamma^1 \gamma^3$.) The full coupling is then

$$\psi_L^T i\sigma^2 i\tau^2 \Phi \psi_L + \psi_R^T i\sigma^2 \tau^2 \Phi \psi_R = \Psi^T i\tau^2 \Phi \Omega \Psi.$$

We would like to view this model in analogy with lattice realizations of a single 2 + 1-dimensional Dirac fermion on the boundary of a 3 + 1-dimensional lattice. The extra dimension allows one to evade the lattice doubling no-go theorems [24–26]. The Witten anomaly seems to be cancelled by inflow from the bulk. The precise meaning of the previous sentence could be clarified given an explicit expression for the WZW functional $\Gamma[A, \Phi]$.

This model is unsatisfactory in at least three ways. First, its five-dimensional nature may make it hard to realize in the laboratory. Secondly, five-dimensional Yang-Mills theory is not asymptotically free and must be completed at short distances somehow (string theory gives interesting ways to do this, e.g. [27]; this model can also be latticized). Thirdly, if we allow the profile of the mass to fluctuate, the kink and antikink can annihilate each other. Nevertheless, the model is instructive.

The model has many mass scales: the W-boson mass M_W , the Kaluza-Klein scale R^{-1} , the Dirac mass m , the inverse thickness of the kink, and an extreme UV cutoff above which the gauge theory succumbs to higher-energy physics. The last two we suppose to be inaccessibly high.

At energies $E \ll 1/R$, this model reduces to the two-doublet theory studied in the previous section.

Having added an extra spatial dimension, monopoles (whose topological charge is characterized by a nontrivial element of $\pi_2(S^2)$) now become stringlike objects which we refer to as “monopole strings”. Consider a monopole string winding around the fifth dimension at some point in three-dimensional space, $\vec{r} = 0$. From a four-dimensional point of view, this appears to be a magnetic monopole. This follows from the fact that the monopole string current $J_M^{\mu\nu}$ sources the five-dimensional U(1) field strength via $dF = J_M$. Where the vanishing loci of the order parameter Φ and the five-dimensional Dirac mass intersect, the five-dimensional Dirac equation will support localized Majorana zero modes.

This model demonstrates that the two Majorana modes need not pair up. Here their wave functions are separated in the extra dimension. In the regime $m \gg R^{-1}$, their overlap is exponentially small.

To illustrate the physics of this five-dimensional construction, we consider a configuration of four monopole strings each parallel to the compact direction and wrapping

once around it. Each of the $i = 1, \dots, 4$ monopole strings intersects each of the $a = 1, 2$ domain walls once for a total of $4 * 2 = 8$ Majorana modes that we label γ_{ia} . These operators satisfy the algebra $\{\gamma_{ia}, \gamma_{jb}\} = 2\delta_{ij}\delta_{ab}$ and are real $\gamma_{ia}^+ = \gamma_{ia}$. From the point of view of four-dimensional physics, a natural basis for this space of states comes by forming complex fermions $c_i = \frac{\gamma_{i1} + i\gamma_{i2}}{2}$ made from Majoranas at the same point in the noncompact directions. Using these fermion operators we can build a space of 2^4 states which further subdivides into an eight-dimensional subspace of even fermion parity and an eight-dimensional subspace of odd fermion parity.

Three important questions must now be answered. First, what states can be produced by creation of such a system from the vacuum state (or any other state without such a configuration of monopole strings)? Second, what operations on the monopole strings can be carried out without large energy cost? Third, what decoherence-free superpositions are possible?

The first question has two immediate answers. The simplest local (from the four-dimensional point of view) vacuumlike state is the state annihilated by all the c_i defined above. The Majorana modes γ_{i1} and γ_{i2} can be viewed as the ends of a “quantum wire” as in [28], and it is quite natural from the four-dimensional point of view to pair up these Majoranas. The second immediate answer comes from thinking about the creation process by which such a monopole string configuration could be formed. For example, we could take monopoles 1 and 4 to have magnetic charge 1 and monopoles 2 and 3 to have magnetic charge -1 . Then we could pair create 1, 2 and 3, 4 from the vacuum state. With this process in mind, and remembering that the Majorana wave function overlap in the compact direction can be made exponentially small, a natural initial state would be that state annihilated by complex fermions formed from Majoranas on neighboring monopoles (independently for each domain wall). This state also has even fermion parity but is not equal to the state annihilated by all the c_i .

As for low energy operations, we must at least require no macroscopic stretching of the monopole strings beyond that required to have the monopole wrap the compact direction. If \mathcal{T} is the monopole tension, then the mass of the monopole string is $2\pi\mathcal{T}R$. In order to perform operations on the zero mode Hilbert space, we would like to entertain motions of the monopole strings. However, we must move an entire monopole string at once in order to avoid a large energy cost associated with stretching the monopole string. This always means exchanging pairs (coming from the two domain walls) of Majorana modes. For example, consider exchanging monopole strings 2 and 3 in the configuration above. This implements the operator

$$U_{2\rightleftharpoons 3} = \exp\left(\frac{\pi}{4}(\gamma_{21}\gamma_{31} + \gamma_{22}\gamma_{32})\right),$$

but this operation can be reexpressed in terms of the c_i as

$$U_{2\rightleftharpoons 3} = \exp\left(\frac{\pi}{2}(c_2 c_3^+ + c_2^+ c_3)\right).$$

This operator acts trivially on states with $c_2^+ c_2 = c_3^+ c_3$ and exchanges pairs of states with $c_2^+ c_2 = 1 - c_3^+ c_3 \text{ mod } 2$. In other words, it simply moves around local fermions from the four-dimensional perspective. Note also that we have not included the dynamics of the gauge field during this exchange process. We note in passing that there is interesting physics associated with the dynamics of the gauge field, particularly the role of instantons; for example, an instanton localized along a line in five-dimensional spacetime describes a conduit via which fermions tunnel from one wall to the other. Since the four-dimensional local basis effectively stacks the two Weyl fermions on top of each other, the physics should be qualitatively similar to the case of a single Dirac fermion in four-dimensions discussed above. In particular, once the gauge field motions are included, we find that the states built from the c_i operators are actually all bosonic because of the extra angular momentum coming from the gauge field.

Finally, what about decoherence-free superpositions? The four-dimensional local basis c_i seems naively decoherence free, but another regime is possible where the smallest scale is R^{-1} . In this regime, the Abelian gauge field resulting from the Higgsing of $SU(2)$ looks five dimensional and may even decohere the fermions in the four-dimensional local basis generated by the c_i . However, in this case we are faced with the question: decohere to what? There seems to be no local basis once the gauge field is allowed to fluctuate in the 5th dimension. There is also no superconductivity to justify forming decoherence-free superpositions of different charge states. In fact, there is an even simpler configuration that can cause concern. Consider a single monopole string forming a closed loop which does not wrap the extra dimension but still punctures one of the domain walls twice. Now this configuration may cost a lot of energy and be unstable, but assuming we could hold the monopole string in place, we appear to have two Majorana modes on a single domain wall but again with no obvious local basis to decohere into. We are again faced with the question: decohere to what?

To resolve these issues, we need to bring in a thus-far neglected piece of the puzzle. In four-dimensions the $SU(2)$ monopole has a collective coordinate, a rotor degree of freedom corresponding to the unbroken $U(1)$ charge. The excitations of the rotor generate the familiar dyon states of the monopole. In the five-dimensional model we have a new complication: instead of a single quantum mechanical rotor, we are faced with a rotor degree of freedom for each point on the monopole string. Thus the monopole string supports a finite-size realization of the $1+1$ dimensional XY model, a $c=1$ conformal field theory. These gapless degrees of freedom can significantly affect the physics. Charge will be dynamically screened by

the gapless rotor degrees of freedom living in the monopole string core. In the parameter regime where the compact radius is large, we have five-dimensional $U(1)$ gauge theory, and the only configurations of the Majorana zero modes that remain decoherence free are those connected by a single rotor string, and they are still linearly confined by the monopole string tension. Thus for strings wrapping the compact direction the decoherence-free subspace is always the four-dimensional local basis and we recover the low energy physics of the Dirac fermion coupled to a scalar in four-dimensions as we must. We can also consider monopole strings as above that intercept only one domain wall, but here the Majorana zero modes are bound by the monopole string stretching between them, the same string that screens their gauge charge.

V. CONCLUSIONS AND GENERAL ARGUMENTS

Despite a promising attempt, we have not found a consistent field theory with Majorana monopoles that are not linearly confined. We would like to argue that this conclusion is general and will do so from a variety of points of view.

A. Monopole configuration space

If we had found a consistent gauge theory with unpaired Majorana operators on the cores of monopoles, we would have been in serious trouble. Indeed, the fundamental group of the bare N -monopole configuration space is precisely S_N , and we know that this group has no interesting nonlocal representations [1,12,13,29].¹ The existence of extended magnetic field lines does not help since the static magnetic field configuration is completely specified by the positions of the monopoles via the magnetic Gauss law. One might have hoped that the Dirac string, which is the remnant of the ribbon that proved so essential in the ungauged theory [4], could play a similar role here. However, this string is unphysical as its position can be moved using gauge transformations. For example, in lattice $U(1)$ gauge theory the Dirac string is completely meaningless and undefined. Thus the only remaining possibility is the existence of some subtle topological information encoded in the existence of the Dirac string (but not its precise position) in certain UV completions of $U(1)$ gauge theory. We can find no such data and although we do not prove it cannot be found, we regard this possibility as quite remote. The main point is simply that the configuration of monopoles in a Coulomb phase is insufficient to support non-Abelian particles. One would have to add extra data beyond the monopole positions in any model that realized non-Abelian particlelike excitations.

¹Deligne's theorem [29] proves that replacing the braid group by the symmetric group gives "local" theories called $\text{Rep}(G, \mu)$. We have bosons ($\mu = 1$) or fermions ($\mu = -1$) with a local "internal" symmetry G .

B. Callias index and anomaly

Here we make a precise connection between the Majorana number mod two and the Witten anomaly. Roughly, we can relate the Witten anomaly to the chiral anomaly mod two; in turn we can relate the chiral anomaly mod two to the Majorana number of the monopole.

In a theory with a Witten anomaly, a chiral rotation by π is an element of the gauge group [30,31], i.e. $(-1)^F$ acts in precisely the same way as a gauge rotation $e^{i\pi\tau_3}$ for some gauge generator τ_3 . One way to think about this statement is that there are no gauge-neutral excitations which carry unit fermion number; this means that the fermion number and the gauge charge are the same mod two. The chiral anomaly mod two is therefore in fact a gauge anomaly [30,31]. In the Witten-anomalous theory, the chiral anomaly—i.e. the fact that an instanton violates the chiral charge by one unit (destroys a rh fermion *or* creates a left-hand (lh) fermion)—means that the instanton must also violate the gauge symmetry (despite the fact that there is no local gauge anomaly).²

The chiral anomaly mod two in turn is related by (a generalization of) Callias' index theorem [34] to the number of real fermion zero modes of the monopole. The result proved by Callias counts the index of a complex-linear Dirac operator; this is the number of *complex* zero modes weighted by some version of chirality. Because of the coupling to the Higgs field Φ , our Dirac operator is only real-linear, and we wish to count its real zero modes (in a monopole background), mod two. This kind of zero mode counting has been considered in [35], and they concluded that the Chern number indeed counts the Majorana number mod two. Thus it seems that within the setup of microscopic fermions coupled to an SU(2) gauge field and an adjoint scalar, the existence of an unpaired Majorana zero mode in the ungauged theory is unavoidably related to the presence of the Witten anomaly in the gauge theory.

C. More arguments from low energy

More generally, we could ask if deconfinement is possible via the disordering route mentioned in the introduction. This scenario has at least two problems. First, as we

²We note in passing that the fermion-number violation by instantons seems to be a *symptom* of the Witten anomaly, rather than an equivalent statement. We say this for the following reason. Recently [32,33] it has been argued that it is possible to modify gauge theories by restricting the instanton sum, for example, to even instanton numbers. This would make the partition function π periodic in θ solve the obstruction given in Eqs. (18, 19) of [31]. However, applying the path-integral method for accomplishing this modification given in [33] to a Witten-anomalous theory does not change the fact that the fermion determinant faithfully represents $\pi_4(\text{SU}(2))$ and therefore does not prevent the gauge field path integral from vanishing.

argued above, the configuration space of monopoles is too trivial to support non-Abelian particles. It appears we must gauge away the ribbon data or disorder it away. Second, given the unbroken global SU(2) symmetry in the disordered phase, the quantum numbers of local excitations should be consistent with the unbroken symmetry. It is hard to see how we can build a sensible real zero mode without doing violence to the SU(2) group structure.

This question can be addressed in more detail using the slave particle techniques which have been developed for the study of spin liquids (i.e. disordered ground states of quantum spin systems). In the disordering scheme described in the introduction, we write the order parameter in terms of bosons z_α as $n^i = z^+ \tau^i z$. In a fractionalized phase with unbroken SU(2) symmetry the doublet fermions will be screened by z and become SU(2)-neutral; however, the resulting SU(2)-singlet fermions will carry an internal U(1) gauge charge. As before, for any hope of success we must disorder the n field without condensing hedgehogs. Assuming we can do this, hedgehogs will become monopoles of the emergent U(1) gauge field. It appears difficult to form the necessary decoherence-free superpositions of fermions charged under the internal U(1) to produce Majorana zero modes on the monopole cores. We also still have the problem of the monopole configuration space. Thus we argue that such a phase is either impossible or the number of Majorana zero modes on the monopole changes across the phase transition.

It is possible for the U(1) gauge symmetry to be found in a Higgs phase; in this case the Majorana solitons are monopoles in a superconductor which again are linearly confined by magnetic flux tubes, and it is perfectly consistent to have localized states of indefinite charge.

We can consider other possibilities where there is no U(1) gauge symmetry at any energy scale. For example, one could try to decompose the order parameter as $n^i = b^T i\tau^2 \tau^i b$ with b a two component complex doublet of bosons. Now the disordered phase will only have an emergent Z_2 gauge field, but the original order parameter has an extra U(1) symmetry associated with $b \rightarrow e^{i\theta} b$ (whereas the SU(2) transformation is $b \rightarrow e^{i\theta\tau^3} b$). In other words, n must be complex. Even if we break this symmetry in the Hamiltonian we can still unwind hedgehog configurations using the extra scalar degrees of freedom. This is to be expected since the hedgehog would have turned into a localized object in the Z_2 gauge theory, but there is no local object in such a theory in 3 + 1 dimensions (the vortex from 2 + 1 is now a vortex line in 3 + 1).

The possibility remains that a 3 + 1-dimensional *lattice* model exists with deconfined Majorana monopoles, i.e. that the continuum limit (our starting point) fails to capture some crucial element. Certain kinds of lattice models that begin with Majorana fermions may, not surprisingly, more easily produce Majorana excitations. If these models cannot be realized with a “proper” regularization involving

only complex fermions coupled to superconductivity, then we are tempted to regard them as too artificial. We can easily design a network of Kitaev quantum wires in three spatial dimensions that reproduce the topological aspects of the Teo-Kane model; however, there is no $SU(2)$ symmetry (it is reduced to a discrete subgroup), and the confinement is still linear. Without the full $SU(2)$ symmetry we cannot gauge the model. Furthermore, there can be no four-dimensional lattice realization of the Teo-Kane model with full $SU(2)$ symmetry since such a lattice model, when attached to the surface of the five-dimensional model above, would produce a trivial surface. Put differently, if such a lattice model did exist it could be trivially gauged, and we would face the Witten-anomalous gauge theory again.

We started from a desire to produce deconfined non-Abelian particlelike excitations in $3 + 1$ dimensions. Specifically, we were interested in localized objects displaying what could be called Majorana statistics. The perhaps simplest route to deconfinement led to an anomalous gauge theory. In attempting to cure the anomaly, we found repeatedly that deconfinement requires the number of Majorana zero modes to be even, giving ordinary statistics. We have made many attempts: high energy fermionic matter, extra dimensions, disordered phases exhibiting emergent gauge fields, but none led to deconfined non-Abelian particles. This is all completely consistent

with general expectations about the nature of particle excitations in three-dimensional space. We conclude with a few comments for future work. We always find linear confinement, but this may not be the most general situation. For example, we can argue that gauging only a subgroup of the full $SU(2)$ symmetry still leaves linear confinement intact. So how strongly bound must such non-Abelian particles be in general? Finally, there remains the prospect that with the right low energy data, deconfined non-Abelian particles would be possible. Although we have ruled out many promising paths to this goal, it would be very exciting to see such a possibility realized elsewhere.

ACKNOWLEDGMENTS

J. M. thanks Frank Wilczek for sharing ideas on a model similar to the gauged Teo-Kane model in January 2009. We would like to thank Andrea Allais, Jae Hoon Lee, Chetan Nayak, Zhenghan Wang, Mike Mulligan, T. Senthil for useful discussions and comments. We thank T. Senthil for collaboration in the initial stages of this work. Three-dimensional graphic from the Logo Company [36]. The work of J. M. is supported in part by funds provided by the U. S. Department of Energy (D. O. E.) under cooperative research agreement DE-FG0205ER41360 and in part by the Alfred P. Sloan Foundation.

-
- [1] J. M. Leinaas and J. Myrheim, *Nuovo Cimento Soc. Ital. Fis. B* **37**, 1 (1977).
 - [2] L. Fu and C. L. Kane, *Phys. Rev. Lett.* **100**, 096407 (2008).
 - [3] J. C. Y. Teo and C. L. Kane, *Phys. Rev. Lett.* **104**, 046401 (2010).
 - [4] M. Freedman *et al.*, [arXiv:1005.0583](https://arxiv.org/abs/1005.0583).
 - [5] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, *Rev. Mod. Phys.* **80**, 1083 (2008).
 - [6] G. 't Hooft, *Nucl. Phys. B* **79**, 276 (1974).
 - [7] A. M. Polyakov, *JETP Lett.* **20**, 194 (1974).
 - [8] E. Witten, *Phys. Lett. B* **117**, 324 (1982).
 - [9] E. D'Hoker and E. Farhi, *Phys. Lett. B* **134**, 86 (1984).
 - [10] E. D'Hoker and E. Farhi, *Nucl. Phys. B* **248**, 59 (1984).
 - [11] T. Senthil, discussions therewith (2010).
 - [12] S. Doplicher, R. Haag, and J. E. Roberts, *Commun. Math. Phys.* **23**, 199 (1971).
 - [13] S. Doplicher, R. Haag, and J. E. Roberts, *Commun. Math. Phys.* **35**, 49 (1974).
 - [14] R. Jackiw and C. Rebbi, *Phys. Rev. D* **13**, 3398 (1976).
 - [15] M. Freedman, C. Nayak, M. Hastings, and X.-L. Qi, [arXiv:1107.2731](https://arxiv.org/abs/1107.2731).
 - [16] E. Witten, *Nucl. Phys. B* **223**, 433 (1983).
 - [17] S. Elitzur and V. P. Nair, *Nucl. Phys. B* **243**, 205 (1984).
 - [18] F. R. Klinkhamer, *Phys. Lett. B* **256**, 41 (1991).
 - [19] C. Nayak and F. Wilczek, *Nucl. Phys. B* **479**, 529 (1996).
 - [20] N. Read and D. Green, *Phys. Rev. B* **61**, 10267 (2000).
 - [21] D. A. Ivanov, *Phys. Rev. Lett.* **86**, 268 (2001).
 - [22] M. Sato, *Phys. Lett. B* **575**, 126 (2003).
 - [23] D. B. Kaplan, [arXiv:0912.2560](https://arxiv.org/abs/0912.2560).
 - [24] H. B. Nielsen and M. Ninomiya, *Nucl. Phys. B* **185**, 20 (1981).
 - [25] H. B. Nielsen and M. Ninomiya, *Nucl. Phys. B* **193**, 173 (1981).
 - [26] H. B. Nielsen and M. Ninomiya, *Phys. Lett. B* **105**, 219 (1981).
 - [27] N. Seiberg, *Phys. Lett. B* **388**, 753 (1996).
 - [28] A. Y. Kitaev, *Phys. Usp.* **44**, 131 (2001).
 - [29] P. Deligne, *Moscow Math. J.* **2**, 227 (2002).
 - [30] J. Goldstone (1984).
 - [31] S. P. de Alwis, *Phys. Rev. D* **32**, 2837 (1985).
 - [32] S. Hellerman, A. Henriques, T. Pantev, E. Sharpe, and M. Ando, *Adv. Theor. Math. Phys.* **11**, 751 (2007).
 - [33] N. Seiberg, *J. High Energy Phys.* **07** (2010) 070.
 - [34] C. Callias, *Commun. Math. Phys.* **62**, 213 (1978).
 - [35] L. Santos, Y. Nishida, C. Chamon, and C. Mudry, *Phys. Rev. B* **83**, 104522 (2011).
 - [36] (2011), http://thelogocompany.net/3d_logos.htm.