

# Symmetrical parametrizations of the lepton mixing matrix

W. Rodejohann<sup>1,\*</sup> and J. W. F. Valle<sup>2,†</sup><sup>1</sup>Max-Planck-Institut für Kernphysik, Postfach 103980, 69029 Heidelberg, Germany<sup>2</sup>AHEP Group, Institut de Física Corpuscular—C.S.I.C./Universitat de València, Edificio Institutos de Paterna, Apt 22085, E-46071 Valencia, Spain

(Received 14 September 2011; published 21 October 2011)

Advantages of the original symmetrical form of the parametrization of the lepton mixing matrix are discussed. It provides a conceptually more transparent description of neutrino oscillations and lepton number violating processes like neutrinoless double beta decay, clarifying the significance of Dirac and Majorana phases. It is also ideal for parametrizing scenarios with light sterile neutrinos.

DOI: 10.1103/PhysRevD.84.073011

PACS numbers: 14.60.Lm, 14.60.Pq

## I. INTRODUCTION

Since the historic discovery of neutrino oscillations, massive neutrinos currently provide the most direct and testable evidence for physics beyond the standard model (SM) of particle physics. At low energies, nine (seven) parameters must be determined depending on whether neutrinos are Majorana (Dirac) particles [1].<sup>1</sup> Here we tacitly assume the former, more general, and theoretically preferred case. Parametrizing the lepton mixing matrix [2,3] in a convenient and intuitive manner is very helpful for data handling and interpretation of the physics, such as neutrino oscillation searches in upcoming long-baseline experiments [4,5] or searches for neutrinoless double beta decay [6,7]. While the former are sensitive to the Dirac phase, Majorana phases [1,8–11] are crucial to describe the latter.

The Particle Data Group (PDG) has adopted a parametrization of the lepton mixing matrix in which it is a product of three consecutive rotations multiplied with a

diagonal phase matrix  $P$  containing the Majorana phases [12]. The mixing matrix can be written as

$$U = R_{23}(\theta_{23}; 0)R_{13}(\theta_{13}; \delta)R_{12}(\theta_{12}; 0)P, \quad (1)$$

where  $R_{ij}(\theta; \varphi)$  is a rotation around the  $ij$  axis, e.g.

$$R_{13}(\theta_{13}; \delta) = \begin{pmatrix} \cos\theta_{13} & 0 & \sin\theta_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -\sin\theta_{13}e^{i\delta} & 0 & \cos\theta_{13} \end{pmatrix}. \quad (2)$$

The position of the Dirac phase  $\delta$  is the convention chosen by the PDG. The two Majorana phases, denoted here  $\alpha$  and  $\beta$ , are usually put inside  $P$ , to the right of the mixing matrix:  $P = \text{diag}(e^{i\alpha}, e^{i\beta}, 1)$ . However, there is no consensus notation yet in what concerns the parametrization of these phases, neither for their names (e.g.  $\phi_1$  and  $\phi_2$  or  $\varphi_1$  and  $\varphi_2$  or  $\sigma$  and  $\rho$ , sometimes with a minus sign, sometimes divided by two, etc.), nor for their position within the matrix ( $P = \text{diag}(1, e^{i\alpha}, e^{i\beta})$ , or  $P = \text{diag}(e^{i\alpha}, e^{i\beta}, 1)$ , etc.). The mixing matrix Eq. (1) is explicitly given by

$$U \equiv \tilde{U}P = \begin{pmatrix} c_{12}c_{13}e^{i\alpha} & s_{12}c_{13}e^{i\beta} & s_{13}e^{-i\delta} \\ -(s_{12}c_{23} + c_{12}s_{23}s_{13}e^{i\delta})e^{i\alpha} & (c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta})e^{i\beta} & s_{23}c_{13} \\ (s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta})e^{i\alpha} & -(c_{12}s_{23} + s_{12}c_{23}s_{13}e^{i\delta})e^{i\beta} & c_{23}c_{13} \end{pmatrix}. \quad (3)$$

The elements  $|U_{e1}|$  and  $|U_{\mu 3}|$  are known with good accuracy, thus the two “large” angles  $\theta_{12}$  and  $\theta_{23}$  are well determined by solar and atmospheric neutrino oscillation data. There is also recent evidence for a nonzero value of the “small” element  $|U_{e3}|$  from the T2K Collaboration [13] and the global neutrino oscillation data sample [14,15]. While there are eight equivalent ways to parametrize the mixing matrix [16], the above order of rotation is useful in the sense that experimentally the straightforwardly measur-

able elements  $|U_{e1}|$ ,  $|U_{e3}|$ , and  $|U_{\mu 3}|$  allow to directly extract the angles in the three-neutrino lepton mixing matrix.<sup>2</sup> In contrast, the other equivalent parametrizations do not share this property.

## II. A MORE CONVENIENT PARAMETRIZATION OF THE MIXING MATRIX

The above form  $U$  is nothing but a rewriting of the “symmetrical” form  $K$  proposed in Ref. [1] (apart from factor ordering, which was left unspecified in the original paper). Here we would like to argue in favor of

<sup>2</sup>See Ref. [17] for a recent application of the other possible parametrizations and [18] for a rare use of the symmetrical parametrization we will study in this work.

\*werner.rodejohann@mpi-hd.mpg.de

†valle@ific.uv.es

<sup>1</sup>Many more parameters exist if neutrino masses arise from type-I seesaw; the parametrization in Ref. [1] covers all seesaw cases.

the conceptual advantages of the original symmetrical presentation of the lepton mixing matrix [1]. For the case of three neutrinos it is given as

$$K = \omega_{23}(\theta_{23}; \phi_{23})\omega_{13}(\theta_{13}; \phi_{13})\omega_{12}(\theta_{12}; \phi_{12}), \quad (4)$$

where each of the  $\omega$ 's is effectively  $2 \times 2$ , characterized by an angle and a  $CP$  phase, e.g.

$$K = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13}e^{-i\phi_{12}} & s_{13}e^{-i\phi_{13}} \\ -s_{12}c_{23}e^{i\phi_{12}} - c_{12}s_{13}s_{23}e^{-i(\phi_{23}-\phi_{13})} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{-i(\phi_{12}+\phi_{23}-\phi_{13})} & c_{13}s_{23}e^{-i\phi_{23}} \\ s_{12}s_{23}e^{i(\phi_{12}+\phi_{23})} - c_{12}s_{13}c_{23}e^{i\phi_{13}} & -c_{12}s_{23}e^{i\phi_{23}} - s_{12}s_{13}c_{23}e^{-i(\phi_{12}-\phi_{13})} & c_{13}c_{23} \end{pmatrix}. \quad (5)$$

Here all three  $CP$  violating phases are physical [9]:  $\phi_{12}$ ,  $\phi_{23}$ , and  $\phi_{13}$ . Even though the parametrization is fully ‘‘symmetric’’ there is a basic difference between Dirac and Majorana phases. In order to understand this let us use the identity [1]

$$P^{-1}KP = \omega_{23}(\theta_{23}; \phi_{23} - \beta)\omega_{13}(\theta_{13}; \phi_{13} - \alpha) \times \omega_{12}(\theta_{12}; \phi_{12} + \beta - \alpha), \quad (6)$$

which allows us, up to unphysical phases, to identify the Dirac phase as

$$\delta \leftrightarrow \phi_{13} - \phi_{12} - \phi_{23}. \quad (7)$$

This formula relates the Dirac phase, which denotes the phase responsible for  $CP$  violation in neutrino oscillations in the PDG parametrization, with the equivalent phase in the symmetrical representation. Note that it obeys field rephasing invariance, as it should. Moreover, in contrast to the PDG description, in the symmetrical form  $CP$  violation in neutrino oscillations is immediately recognized as a three-generation phenomenon involving the phases of all three generations,<sup>3</sup> an important conceptual advantage.

$$J_{CP} = \Im\{U_{e1}^*U_{\mu 3}^*U_{e3}U_{\mu 1}\} = \begin{cases} \frac{1}{8}\sin 2\theta_{12}\sin 2\theta_{23}\sin 2\theta_{13}\cos\theta_{13}\sin\delta & \text{(PDG),} \\ \frac{1}{8}\sin 2\theta_{12}\sin 2\theta_{23}\sin 2\theta_{13}\cos\theta_{13}\sin(\phi_{13} - \phi_{12} - \phi_{23}) & \text{(symmetrical),} \end{cases}$$

and shows the same result as Eq. (7).

### A. Application to lepton number violating phenomena

As is well known, there are two conceptually different kinds of  $CP$  violating phenomena [1]. In the language of the PDG parametrization, one is associated to the ‘‘Dirac phase’’  $\delta$  and is the exact analogue to the  $CP$  phase in the quark mixing matrix, responsible for the area of the Cabibbo-Kobayashi-Maskawa unitarity triangle, while

<sup>3</sup>In this sense the Dirac phase has an intrinsic geometric meaning, like the curl of two vectors or the area of a triangle.

$$\omega_{13} = \begin{pmatrix} c_{13} & 0 & e^{-i\phi_{13}}s_{13} \\ 0 & 1 & 0 \\ -e^{i\phi_{13}}s_{13} & 0 & c_{13} \end{pmatrix}.$$

Explicitly, the symmetrical parametrization of the lepton mixing matrix,  $K$  can be written as

Recall that the neutrino oscillation probability for a  $\nu_\alpha \rightarrow \nu_\beta$  flavor transition is given by

$$P(\nu_\alpha \rightarrow \nu_\beta) = \left| \sum_j U_{\alpha j}^* U_{\beta j} e^{-i\frac{m_j^2}{2E}L} \right|^2 = \delta_{\alpha\beta} - 4 \sum_{i>j} \Re\{U_{\alpha i}^* U_{\alpha j} U_{\beta i} U_{\beta j}^*\} \sin^2\left(\frac{\Delta m_{ij}^2}{4E}L\right) + 2 \sum_{i>j} \Im\{U_{\alpha i}^* U_{\alpha j} U_{\beta i} U_{\beta j}^*\} \sin\left(\frac{\Delta m_{ij}^2}{2E}L\right),$$

where  $E$  is the neutrino energy,  $L$  is the distance traveled by the neutrino, and  $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$  ( $m_i$  being positive mass eigenvalues) are the neutrino mass-squared differences. Here  $\Re$  and  $\Im$  denote real and imaginary parts. For three families there is only one independent imaginary part  $\Im\{U_{\alpha i}^* U_{\alpha j} U_{\beta i} U_{\beta j}^*\}$ , which is responsible for  $CP$  violation in neutrino oscillations. Comparing this invariant with the PDG and symmetrical parametrizations gives

the other one is associated to the two ‘‘Majorana phases’’  $\alpha$  and  $\beta$ , which do not show up in neutrino oscillations [8–11] but do affect lepton number violating amplitudes.

In what follows we will briefly discuss the role of Majorana phases in determining the rates characterizing neutrinoless double beta decay and neutrino-anti-neutrino oscillations [9].

A suitable parametrization of Majorana phases plays a *very important* role in interpreting the effective mass parameter characterizing the amplitude for neutrinoless double beta decay (see Ref. [7] for a recent review). Its explicit form reads

$$\langle m \rangle = \left| \sum_j U_{ej}^2 m_j \right| = \begin{cases} |c_{12}^2 c_{13}^2 m_1 e^{2i\alpha} + s_{12}^2 c_{13}^2 m_2 e^{2i\beta} + s_{13}^2 m_3 e^{2i\delta}| & \text{(PDG),} \\ |c_{12}^2 c_{13}^2 m_1 + s_{12}^2 c_{13}^2 m_2 e^{2i\phi_{12}} + s_{13}^2 m_3 e^{2i\phi_{13}}| & \text{(symmetrical).} \end{cases} \quad (8)$$

Only the two Majorana phases should appear in  $\langle m \rangle$  [9]. However, this is not at all clear in the PDG presentation. In contrast, the symmetrical parametrization provides a manifestly transparent description in which only the two Majorana phases appear in  $\langle m \rangle$ , as it should. Currently nuclear matrix element uncertainties prevent the extraction of Majorana phases from neutrinoless double beta decay. However, should these be circumvented and should the determination of the Majorana phases become an issue, then the symmetrical parametrization will surely be preferred over the PDG one.

It has long been known that the lepton mixing matrix characterizing the charged current interaction of Majorana neutrinos in gauge theories may have complex entries that conserve  $CP$  [19]. These special  $CP$  conserving situations are associated with Wolfenstein's  $CP$  signs [20], when neutrino mass states are  $CP$  eigenstates with  $CP$  parity  $\eta_{CP} = \pm i$ . There are four possible inequivalent sign configurations in the sum in Eq. (8):  $(+++)$ ,  $(+--)$ ,  $(+-+)$ , and  $(++-)$ . These are in correspondence to special values of the Majorana phases, namely,  $\phi_{12} = \phi_{13} = 0$ ,  $\phi_{12} = \phi_{13} = \pi/2$ ,  $\phi_{13} = \pi/2$  with  $\phi_{12} = 0$ , and  $\phi_{13} = 0$  with  $\phi_{12} = \pi/2$ , respectively. Majorana phases would also show up in processes analogous to neutrinoless double beta decay, such as decays like  $K^+ \rightarrow \pi^- \mu^+ \mu^+$ , whose amplitude would be proportional to  $\sum U_{\mu i}^2 m_i$ , and have extremely low branching ratios, see Ref. [7] and references therein.

Let us now comment on neutrino-antineutrino oscillations. A Gedanken experiment looking for antineutrinos  $\bar{\nu}_\beta$  in a beam of neutrinos  $\nu_\alpha$  has been suggested in Ref. [9] in order to clarify the physical nature of Majorana phases at the two-generation level. In the three-generation case the probability for such a process is given as

$$\begin{aligned} P(\nu_\alpha \rightarrow \bar{\nu}_\beta) &= \frac{1}{E^2} \left| \sum_j U_{\alpha j} U_{\beta j} m_j e^{-iE_j t} \right|^2 \\ &= \frac{1}{E^2} \left| \sum_{i,j} U_{\alpha j} U_{\beta j} U_{\alpha i}^* U_{\beta i}^* m_i m_j e^{-i(E_j - E_i)t} \right|. \end{aligned} \quad (9)$$

leading to complicated transition probability expressions, which will not be explicitly given here. The least unrealistic channel is represented by  $\nu_e$  to  $\bar{\nu}_e$  transitions, because the transition probabilities go with the ratio of mass over energy squared, and electron neutrinos can be produced with much lower energy than the other flavors. For three families,  $P(\nu_e \rightarrow \bar{\nu}_e)$  depends only on the Majorana

phases  $\phi_{12}$  and  $\phi_{13}$  for the symmetrical parametrization, whereas the PDG case leads to a dependence on all three phases. Note the analogy to the effective mass discussed above.

## B. Application to seesaw and sterile neutrinos

We now turn to the lepton mixing matrix characterizing models containing gauge singlets such as seesaw models. Their most general form was presented within the symmetrical parametrization in Ref. [1], covering seesaw schemes of all types, type I, type II, and type III. Since it applies to an arbitrary number  $m$  of nondoublet leptons (singlets in type I and II, triplets in type III) this parametrization also covers low-scale in addition to high-scale seesaw schemes. To a good approximation, in the standard high-scale seesaw case neutrino oscillations are well described by the simplest unitary form of the lepton mixing matrix used above in Eqs. (1) or (4). In contrast, for the low-scale seesaw schemes [21–24] neutrino oscillations involve only a unitarity-violating truncation of the full mixing matrix, see, for example, Refs. [25,26]. In both cases one has an “effective” neutrino oscillation description with  $m = 0$ , i.e. the extra neutral states are too heavy to take part in the oscillation phenomena. Because these possibilities have already been widely discussed in the literature, here we will focus on the alternative possibility that singlets are light enough to participate in oscillations, in the simple case of  $m = 1$ , i.e. one “sterile” neutrino plus three active ones. This possibility has recently regained attention [27]. In terms of the mixing matrix, a useful order of rotation is 34-24-14-23-13-12. We have, therefore,

$$\begin{aligned} U &= \omega_{34}(\theta_{34}; 0) \omega_{24}(\theta_{24}, \delta_{24}) \omega_{14}(\theta_{14}; \delta_{14}) \omega_{23}(\theta_{23}; 0) \\ &\quad \times \omega_{13}(\theta_{13}; \delta_{13}) \omega_{12}(\theta_{12}; 0) P, \end{aligned} \quad (10)$$

with  $P = \text{diag}(e^{i\alpha}, e^{i\beta}, e^{i\gamma})$  in the sense of the PDG description, or in the symmetrical form:

$$\begin{aligned} K &= \omega_{34}(\theta_{34}; \phi_{34}) \omega_{24}(\theta_{24}, \phi_{24}) \omega_{14}(\theta_{14}; \phi_{14}) \\ &\quad \times \omega_{23}(\theta_{23}; \phi_{23}) \omega_{13}(\theta_{13}; \phi_{13}) \omega_{12}(\theta_{12}; \phi_{12}). \end{aligned} \quad (11)$$

First note that the number of rotations for  $3 + 1$  neutrino types (six) is exactly the number of phases (three Dirac and three Majorana), a characteristic feature of the symmetrical parametrization [1].

Consider first the effective mass characterizing neutrinoless double beta decay in this case, which is given as

$$\langle m \rangle = \begin{cases} |c_{12}^2 c_{13}^2 c_{14}^2 m_1 e^{2i\alpha} + s_{12}^2 c_{13}^2 c_{14}^2 m_2 e^{2i\beta} + s_{13}^2 c_{14}^2 m_3 e^{2i(\gamma-\delta_{13})} + s_{14}^2 m_4 e^{-2i\delta_{14}}| & \text{(PDG),} \\ |c_{12}^2 c_{13}^2 c_{14}^2 m_1 + s_{12}^2 c_{13}^2 c_{14}^2 m_2 e^{2i\phi_{12}} + s_{13}^2 c_{14}^2 m_3 e^{2i\phi_{13}} + s_{14}^2 m_4 e^{2i\phi_{14}}| & \text{(symmetrical).} \end{cases}$$

Obviously the symmetrical parametrization has advantages over the one of PDG. Indeed, again, the  $CP$  conserving cases corresponding to different  $CP$  signs are obtained by choosing the three Majorana phases  $\phi_{12,13,14}$  to be  $\pi/2$  or zero.

Regarding oscillations, the three Dirac phases in the PDG description are  $\delta_{13}$ ,  $\delta_{14}$ , and  $\delta_{24}$ . While there are nine [28] independent  $J_{\alpha\beta}^{ij} = \Im\{U_{\alpha i}^* U_{\beta j}^* U_{\alpha j} U_{\beta i}\}$ , only three independent  $CP$  asymmetries  $P(\nu_\alpha \rightarrow \nu_\beta) - P(\nu_\beta \rightarrow \nu_\alpha)$  exist. In the symmetrical parametrization, the relevant independent phase combinations appearing in oscillation probabilities are

$$\begin{aligned} I_{123} &= \phi_{12} + \phi_{23} - \phi_{13}, \\ I_{124} &= \phi_{12} + \phi_{24} - \phi_{14}, \\ I_{134} &= \phi_{13} + \phi_{34} - \phi_{14}. \end{aligned} \quad (12)$$

Each of the phase combinations  $I_{ijk}$ , with  $i < j < k$ , is seen to span three generations, as necessary for the existence of  $CP$  violation in neutrino oscillations. Note that there is a fourth possible combination,  $I_{234}$ . However, since

$$I_{123} + I_{134} - I_{124} = I_{234} \quad (13)$$

holds, this fourth invariant is not independent. This condition actually implies that  $I_{ijk}$  is a 2-cocycle [29]. This is true for an arbitrary number  $N_s$  of additional sterile neutrinos: noting that the  $\frac{1}{2}N_s(N_s - 1)$  rotations between sterile neutrinos are unphysical, it is easy to see that for  $N$  massive neutrinos, including  $N_s = N - 3$  sterile neutrinos, there are  $N - 1 = N_s + 2$  Majorana phases and  $2N - 5 = 2N_s + 1$  Dirac phases. Each massless neutrino results in one Majorana phase less. The total number of  $3(N - 2) = 3(N_s + 1)$  phases corresponds exactly to the number of physical rotations, which is  $\frac{1}{2}N(N - 1) - \frac{1}{2}N_s(N_s - 1) = 3(N - 2)$ . The symmetrical parametrization is therefore tailor-made also for concisely describing the phenomenology of sterile neutrinos.

It has been argued that current neutrino data might imply in fact that two sterile neutrinos are present [30]. A possible order of the nine physical rotations is

$$\begin{aligned} K &= \omega_{25}(\theta_{25}; \phi_{25})\omega_{34}(\theta_{34}; \phi_{34})\omega_{35}(\theta_{35}; \phi_{35}) \\ &\times \omega_{24}(\theta_{24}; \phi_{24})\omega_{23}(\theta_{23}; \phi_{23})\omega_{15}(\theta_{15}; \phi_{15}) \\ &\times \omega_{14}(\theta_{14}; \phi_{14})\omega_{13}(\theta_{13}; \phi_{13})\omega_{12}(\theta_{12}; \phi_{12}). \end{aligned} \quad (14)$$

Again, the effective mass characterizing neutrinoless double beta decay automatically looks straightforward:

$$\begin{aligned} \langle m \rangle &= |c_{12}^2 c_{13}^2 c_{14}^2 c_{15}^2 m_1 + s_{12}^2 c_{13}^2 c_{14}^2 c_{15}^2 e^{2i\phi_{12}} \\ &\quad + s_{13}^2 c_{14}^2 c_{15}^2 m_3 e^{2i\phi_{13}} + s_{14}^2 c_{15}^2 m_4 e^{2i\phi_{14}} \\ &\quad + s_{15}^2 m_5 e^{2i\phi_{15}}|, \end{aligned} \quad (15)$$

involving just the four physical Majorana phases. Regarding  $CP$  violation in oscillations, there are

$$\binom{5}{3} - 3 = 10 - 3 = 7$$

different  $I_{ijk}$  combinations with  $i < j < k$  and  $i, j \in \{1, 2, 3, 4, 5\}$ , where the subtraction of 3 stems from the cases that have both 4 and 5 in  $ijk$ :

$$I_{123}, I_{124}, I_{125}, I_{134}, I_{135}, I_{234}, I_{235}. \quad (16)$$

There also exist

$$\binom{5}{4} - 3 = 5 - 3 = 2$$

“sumrules” in analogy to Eq. (13), namely,

$$I_{123} + I_{134} - I_{124} = I_{234} \quad \text{and} \quad I_{123} + I_{135} - I_{125} = I_{235}. \quad (17)$$

Hence, at the end there are five physical Dirac  $CP$  violating phase combinations affecting neutrino oscillation probabilities; for instance one could choose

$$I_{123} = \phi_{12} + \phi_{23} - \phi_{13}, \quad (18)$$

$$I_{124} = \phi_{12} + \phi_{24} - \phi_{14}, \quad (19)$$

$$I_{134} = \phi_{13} + \phi_{34} - \phi_{14}, \quad (20)$$

$$I_{125} = \phi_{12} + \phi_{25} - \phi_{15}, \quad (21)$$

$$I_{135} = \phi_{13} + \phi_{35} - \phi_{15}. \quad (22)$$

The phase relevant for  $CP$  violation in the short-baseline oscillations  $\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu$  sector is  $\Im\{U_{e4}^* U_{\mu 5}^* U_{\mu 4} U_{e5}\} \propto \sin(\phi_{14} - \phi_{15} - \phi_{24} + \phi_{25}) = \sin(I_{125} - I_{124})$ .

The generalization to  $N_s$  sterile neutrinos is now clear: In more mathematical language, adopted from Ref. [29], one may define the operator  $\delta$  (with  $\delta^2 = 0$ ) such that [cf. Eq. (13)]

$$\delta I_{1234} \equiv F_{1234}^{(4)} = I_{123} + I_{134} - I_{124} - I_{234} = 0. \quad (23)$$

For general  $i, j, k, l$  with  $i < j < k < l$  one has  $\delta I_{ijkl} = 0$ . For five active generations one has ten  $I_{ijk}$  and five  $F_{ijkl}^{(4)} = 0$ , one of which can be expressed by the other four, for instance,

$$F_{2345}^{(4)} = F_{1235}^{(4)} + F_{1345}^{(4)} - F_{1234}^{(4)} - F_{1245}^{(4)} \quad \text{or} \quad \delta F_{12345} = 0. \quad (24)$$

Therefore, the standard result of  $10 - (5 - 1) = 6$  Dirac phases is obtained. If all generations were active, then the number of independent phase combinations would be

$$\binom{N}{3} - \left[ \binom{N}{4} - \binom{N}{5} \right] = \binom{N}{0} - \left[ \binom{N}{1} - \binom{N}{2} \right] \\ = \frac{1}{2}(N-1)(N-2). \quad (25)$$

One simply subtracts the number of  $\frac{1}{2}N_s(N_s - 1) = \frac{1}{2}(N-3)(N-4)$  unphysical rotations from this result to obtain the already quoted  $2N - 5 = 2N_s + 1$  Dirac phases. The first binomial in Eq. (25) is the number of  $ijk$  combinations with  $i < j < k$ , the second binomial is the number of sumrules between them, and the third binomial describes the linear relations existing between the sumrules. For instance, five active generations have

$$\binom{5}{3} = 10$$

different  $I_{ijk}$  combinations with  $i < j < k$  and  $i, j \in \{1, 2, 3, 4, 5\}$ . A number of

$$\binom{5}{4} = 5$$

sumrules exist, with

$$\binom{5}{5} = 1$$

linear relation between them.

### III. FINAL REMARKS

In summary, the issue of a proper parametrization scheme for the lepton mixing matrix will become relevant as experiments reach sensitivity to  $CP$  violation, either of Dirac or Majorana type—even more so if present indications for sterile neutrinos are confirmed in upcoming experiments. While the form given in the PDG is just a rewriting of the symmetrical form proposed in Ref. [1], we have advocated here the conceptual advantages of the original symmetrical presentation for the description of neutrino oscillations and, especially, of lepton number violating processes like neutrinoless double beta decay.

### ACKNOWLEDGMENTS

This work was supported by the ERC under the Starting Grant MANITOP, by the Deutsche Forschungsgemeinschaft in the Transregio 27 “Neutrinos and beyond—weakly interacting particles in physics, astrophysics and cosmology” (W.R.) and by the Spanish MICINN under grants FPA2008-00319/FPA and MULTIDARK Consolider CSD2009-00064, by Prometeo/2009/091, and by the EU grant UNILHC PITN-GA-2009-237920 (J.V.). We wish to thank Morimitsu Tanimoto and the organizers of the Mini Workshop on Neutrinos at the IPMU, where this paper was initiated. W.R. would like to thank Jisuke Kubo and the Yukawa Institute for Theoretical Physics at Kyoto University for hospitality during the YITP workshop “Summer Institute 2011,” where part of this paper was written.

- 
- [1] J. Schechter and J. W. F. Valle, *Phys. Rev. D* **22**, 2227 (1980).  
 [2] B. Pontecorvo, *Zh. Eksp. Teor. Fiz.* **33**, 549 (1957).  
 [3] Z. Maki, M. Nakagawa, and S. Sakata, *Prog. Theor. Phys.* **28**, 870 (1962).  
 [4] A. Bandyopadhyay *et al.* (ISS Physics Working Group), *Rep. Prog. Phys.* **72**, 106201 (2009).  
 [5] H. Nunokawa, S. J. Parke, and J. W. F. Valle, *Prog. Part. Nucl. Phys.* **60**, 338 (2008).  
 [6] F. T. Avignone, S. R. Elliott, and J. Engel, *Rev. Mod. Phys.* **80**, 481 (2008).  
 [7] W. Rodejohann, *Int. J. Mod. Phys. E* **20**, 1833 (2011).  
 [8] S. M. Bilenky, J. Hosek, and S. T. Petcov, *Phys. Lett.* **94B**, 495 (1980).  
 [9] J. Schechter and J. W. F. Valle, *Phys. Rev. D* **23**, 1666 (1981).  
 [10] M. Doi, T. Kotani, H. Nishiura, K. Okuda, and E. Takasugi, *Phys. Lett.* **102B**, 323 (1981).  
 [11] A. de Gouvea, B. Kayser, and R. N. Mohapatra, *Phys. Rev. D* **67**, 053004 (2003).  
 [12] K. Nakamura *et al.*, *J. Phys. G* **37**, 075021 (2010).  
 [13] K. Abe *et al.* (T2K Collaboration), *Phys. Rev. Lett.* **107**, 041801 (2011).  
 [14] T. Schwetz, M. Tortola, and J. W. F. Valle, *New J. Phys.* **13**, 063004 (2011); **13**, 109401 (2011).  
 [15] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo, and A. M. Rotunno, *Phys. Rev. D* **84**, 053007 (2011).  
 [16] H. Fritzsch and Z.-z. Xing, *Phys. Lett. B* **517**, 363 (2001).  
 [17] W. Rodejohann, H. Zhang, and S. Zhou, arXiv:1107.3970.  
 [18] Y. H. Ahn, H.-Y. Cheng, and S. Oh, arXiv:1107.4549.  
 [19] J. Schechter and J. W. F. Valle, *Phys. Rev. D* **24**, 1883 (1981); **25**, 283(E) (1982).  
 [20] L. Wolfenstein, *Phys. Lett.* **107B**, 77 (1981).  
 [21] R. N. Mohapatra and J. W. F. Valle, *Phys. Rev. D* **34**, 1642 (1986).



- [22] M. C. Gonzalez-Garcia and J. W. F. Valle, *Phys. Lett. B* **216**, 360 (1989).
- [23] E. Akhmedov *et al.*, *Phys. Lett. B* **368**, 270 (1996); *Phys. Rev. D* **53**, 2752 (1996).
- [24] M. Malinsky, J. C. Romao, and J. W. F. Valle, *Phys. Rev. Lett.* **95**, 161801 (2005).
- [25] D. Forero, S. Morisi, M. Tortola, and J. W. F. Valle, [arXiv:1107.6009](https://arxiv.org/abs/1107.6009).
- [26] H. Hettmansperger, M. Lindner, and W. Rodejohann, *J. High Energy Phys.* 04 (2011) 123.
- [27] G. Mention *et al.*, *Phys. Rev. D* **83**, 073006 (2011).
- [28] W.-l. Guo and Z.-z. Xing, *Phys. Rev. D* **65**, 073020 (2002).
- [29] M. Gronau, R. Johnson, and J. Schechter, *Phys. Rev. D* **32**, 3062 (1985).
- [30] M. Sorel, J. Conrad, and M. Shaevitz, *Phys. Rev. D* **70**, 073004 (2004).