Finsler geometric extension of Einstein gravity

Christian Pfeifer* and Mattias N. R. Wohlfarth[†]

Institut für Theoretische Physik und Zentrum für Mathematische Physik, Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany (Received 10 January 2012; published 8 March 2012)

(Received to January 2012, published 8 Match 2012

We construct gravitational dynamics for Finsler spacetimes in terms of an action integral on the unit tangent bundle. These spacetimes are generalizations of Lorentzian metric manifolds which satisfy necessary causality properties. A coupling procedure for matter fields to Finsler gravity completes our new theory that consistently becomes equivalent to Einstein gravity in the limit of metric geometry. We provide a precise geometric definition of observers and their measurements and show that the transformations, by means of which different observers communicate, form a groupoid that generalizes the usual Lorentz group. Moreover, we discuss the implementation of Finsler spacetime symmetries. We use our results to analyze a particular spacetime model that leads to Finsler geometric refinements of the linearized Schwarzschild solution.

DOI: 10.1103/PhysRevD.85.064009

PACS numbers: 04.50.Kd, 02.40.-k, 04.20.Cv

I. INTRODUCTION

The weak equivalence principle states that the trajectories of small test bodies, neither affected by gravitational tidal forces nor by forces other than gravity, are independent of their internal structure and composition [1]. Experimentally, this principle is confirmed with extremely high precision [2]; in gravity theory, it has been implemented already by Newton who postulated that gravitational mass should equal inertial mass, and then by Einstein who formulated the motion of test bodies in terms of geodesics on Lorentzian spacetime. These trajectories extremize the Lorentzian length integral which is interpreted physically as proper time. The fundamental geometric object entering this construction is the Lorentzian spacetime metric. This observation led Einstein to the development of a gravity theory that determines the metric and so provides a dynamical background geometry for point particles, observers and physical fields.

The essential point in Einstein's implementation of the weak equivalence principle by a clock postulate is the use of geometric concepts, not the particular choice of metric geometry. Indeed, more general geometries can be used which automatically realize this principle. Here, we consider Finsler geometry [3,4] which generalizes metric geometry by providing a very general length functional for curves $\tau \mapsto \gamma(\tau)$ on a manifold M,

$$S[\gamma] = \int d\tau F(\gamma(\tau), \dot{\gamma}(\tau)). \tag{1}$$

The Finsler function F maps the points of the curve and the attached four-velocities into real numbers and is homogeneous of degree 1 in its second argument to ensure the

reparametrization invariance of $S[\gamma]$. The usual length measure associated to a Lorentzian metric *g* is obtained for the special case $F(\gamma, \dot{\gamma}) = \sqrt{|g_{ab}(\gamma)\dot{\gamma}^a\dot{\gamma}^b|}$. Physically, we interpret the integral above as a generalized clock postulate and point particle action.

On this basis, we will develop a consistent gravity theory which determines the Finsler function dynamically. Our construction builds on the precise definition of physical Finsler spacetimes in Ref. [5], where we established a minimal set of requirements on the function F so that it can describe a geometric spacetime background suitable for physics. In particular, Finsler spacetimes provide a well-defined notion of causality and the possibility to formulate field-theory actions. A generalization of gravity based on Finsler geometry has the potential to explain various issues that are not naturally explained by Einstein gravity. Indeed, it has been argued that Finsler geometry in principle can address the rotational curves of galaxies and the acceleration of the Universe without introducing dark matter [6] or dark energy [7] and that it admits sufficiently complex causal structures which allow a consistent geometric explanation of superluminal neutrino propagation [8]. Here, we will find that Finsler gravity may also explain the fly-by anomaly [9] in the Solar System.

Our presentation is structured as follows. We will begin in Sec. II with a brief review of our definition of Finsler spacetimes and of the mathematical tools and geometric objects needed. Moreover, we will define field theory action integrals on Finsler spacetimes. Equipped with these concepts, we will show in Sec. III how to model observers on Finsler spacetimes and discuss how they perform measurements. We will prove that two different observers are related by a transformation composed out of a certain parallel transport and a Lorentz transformation. The set of these transformations has the algebraic structure of a groupoid that reduces to the usual Lorentz group in the metric geometry limit. In Sec. IV, we will present our

^{*}christian.pfeifer@desy.de

[†]mattias.wohlfarth@desy.de

CHRISTIAN PFEIFER AND MATTIAS N.R. WOHLFARTH

new theory of Finsler gravity, including a matter coupling principle, which geometrically extends Einstein gravity without introducing new fundamental scales. We will derive the Finsler gravity field equation by variation and prove that it reduces consistently to the Einstein field equations in the metric limit. Symmetries of Finsler spacetimes will be introduced in Sec. V. The maximally symmetric solution of vacuum Finsler gravity turns out to be standard Minkowski spacetime. Section VI considers a spherically symmetric perturbation around this vacuum which is found to be a refinement of the Schwarzschild solution which shows how Finsler gravity could resolve the fly-by anomaly. We conclude in Sec. VII. Appendix A presents technical details of our derivations.

II. FINSLER SPACETIME GEOMETRY

The central idea behind the generalization of Einstein gravity presented in this article is the description of spacetime and its dynamics by Finsler geometry instead of metric geometry. In this section, we will review the basic geometric concepts available on Finsler spacetimes [5]. These were introduced as a generalization of Lorentzian metric spacetimes; they allow full control of the null geometry and a clean definition of causality, which is essential for the description of light and observers. In particular, we will describe nonlinear connections and curvature; moreover, we will explain how to obtain welldefined field theory actions.

A. Basic concepts

The definition of Finsler spacetimes involves the eightdimensional tangent bundle *TM* which is the union of all tangent spaces to the underlying four-dimensional event manifold *M*. Thus, any point $P \in TM$ is a tangent vector to *M* at some point $p \in M$; there is a natural projection $\pi: TM \to M, P \mapsto p$. It is convenient to use the so-called induced coordinates on the tangent bundle which are constructed as follows. Let (*x*) be coordinates on some open neighborhood $U \subset M$ of $p = \pi(P)$. With respect to these, we can express $P = y^a \frac{\partial}{\partial x^a}|_{x(p)}$; the induced coordinates of *P* are then (*x*(*p*), *y*). The corresponding induced coordinate basis of *TTM* will be denoted by $\{\partial_a = \frac{\partial}{\partial x^a}, \bar{\partial}_a = \frac{\partial}{\partial y^a}\}$, and that of its dual T^*TM by $\{dx^a, dy^a\}$.

Definition 1. A Finsler spacetime (M, L, F) is a fourdimensional, connected, Hausdorff, paracompact, smooth manifold M equipped with a continuous function $L: TM \rightarrow \mathbb{R}$ on the tangent bundle, which has the following properties:

- (i) *L* is smooth on the tangent bundle without the zero section *TM* \ {0};
- (ii) *L* is positively homogeneous of real degree $n \ge 2$ with respect to the fiber coordinates of *TM*,

$$L(x, \lambda y) = \lambda^n L(x, y) \quad \forall \ \lambda > 0; \tag{2}$$

(iii) L is reversible in the sense

$$|L(x, -y)| = |L(x, y)|;$$
(3)

(iv) the Hessian g_{ab}^{L} of L with respect to the fiber coordinates is nondegenerate on $TM \setminus A$, where Ahas measure zero and does not contain the null set $\{(x, y) \in TM | L(x, y) = 0\},\$

$$g_{ab}^{L}(x, y) = \frac{1}{2}\bar{\partial}_{a}\bar{\partial}_{b}L; \qquad (4)$$

(v) the unit timelike condition holds, i.e., for all $x \in M$, the set

$$\Omega_{x} = \left\{ y \in T_{x}M || L(x, y)| = 1, \\ g_{ab}^{L}(x, y) \text{ has signature } (\epsilon, -\epsilon, -\epsilon, -\epsilon), \\ \epsilon = \frac{|L(x, y)|}{L(x, y)} \right\}$$
(5)

contains a nonempty closed connected component $S_x \subset \Omega_x \subset T_x M$.

The Finsler function associated to *L* is $F(x, y) = |L(x, y)|^{1/n}$ and the Finsler metric $g_{ab}^F = \frac{1}{2}\bar{\partial}_a\bar{\partial}_bF^2$.

Lorentzian metric spacetimes (M, g) arise from definition 1 in case $L(x, y) = g_{ab}(x)y^ay^b$; then $g^L = \epsilon g^F = g$, and S_x is the set of unit g-timelike vectors. For general function L, the relation between the inverse Finsler metric g^F and the inverse Lagrange metric g^L is given by

$$g^{Fab} = \frac{nL}{2|L|^{2/n}} \left(g^{Lab} + \frac{2(n-2)}{n(n-1)L} y^a y^b \right).$$
(6)

The definition of Finsler spacetimes guarantees a welldefined causal structure by the existence of timelike vectors that form an open convex cone with null boundary in every local tangent space, see Fig. 1. It provides full control of the geometry along the null directions where L(x, y) = 0: we could show that the geometric concepts of connections and curvature that will be discussed below can be extended to this set, which is not possible in standard textbook formulations of Finsler spaces. Furthermore, by constructing a theory of electrodynamics on Finsler spacetimes, we proved that light propagates along null directions. For further details, see Ref. [5].

The geometry of Finsler spacetimes is formulated by extending the standard language of Finsler geometry, see e.g. Refs. [3,4]; this is a special geometry on the tangent bundle based on the Finsler function *F* that appears in the clock postulate. The basic geometric object deduced from the Finsler function is the Cartan nonlinear connection. Any nonlinear connection on *TM* is equivalent to a unique decomposition of all tangent spaces to *TM* as $T_PTM =$ $H_PTM \oplus V_PTM$, see Fig. 2. While the vertical bundle *VTM* is canonically spanned by { $\bar{\partial}_a$ }, the horizontal bundle



FIG. 1 (color online). Geometric structures implemented in every tangent space T_xM by definition 1 of Finsler spacetimes. The solid lines show the guaranteed cone of timelike vectors with the shells of unit timelike vectors and null boundary. The dotted lines indicate a potentially more complex null structure.

HTM is spanned by $\{\delta_a = \partial_a - N^b{}_a \bar{\partial}_b\}$, where the $N^a{}_b(x, y)$ are the coefficients of the nonlinear connection. Then, the dual bundle $(VTM)^*$ is spanned by $\{\delta y^a = dy^a + N^a{}_b dx^b\}$, and $(HTM)^*$ by $\{dx^a\}$. For the Cartan nonlinear connection, the connection coefficients are determined by the fundamental functions *F* or *L* as

$$N^{a}{}_{b} = \frac{1}{4}\bar{\partial}_{b} [g^{Faq}(y^{p}\partial_{p}\bar{\partial}_{q}F^{2} - \partial_{q}F^{2})]$$

$$= \frac{1}{4}\bar{\partial}_{b} [g^{Laq}(y^{p}\partial_{p}\bar{\partial}_{q}L - \partial_{q}L)].$$
(7)

The equality of the two expressions for $N^a{}_b$ follows from the proof of theorem 2 in Ref. [5], which makes particular use of the Euler theorem $y^a \bar{\partial}_a f(x, y) = mf(x, y)$ for *m*-homogeneous functions. Note that the right-hand side is also valid on the null structure where *F* is not even differentiable.

Under tangent bundle changes of coordinates induced from coordinate transformations $(x) \rightarrow (\tilde{x}(x))$ of the manifold, the nonlinear connection coefficients $N^a{}_b$ transform in such a way that the horizontal/vertical basis $\{\delta_a, \bar{\partial}_a\}$ of *TTM* and the dual basis $\{dx^a, \delta y^a\}$ transform as vectors or one-forms over the manifold,



FIG. 2 (color online). Tangent bundle geometry: decomposition of $T_{(x,y)}TM$ into horizontal and vertical parts.

$$(\delta_a, \bar{\delta}_a) \to \left(\frac{\partial \tilde{x}^b}{\partial x^a} \tilde{\delta}_b, \frac{\partial \tilde{x}^b}{\partial x^a} \tilde{\delta}_b \right) ,$$

$$(dx^a, \delta y^a) \to \left(\frac{\partial x^a}{\partial \tilde{x}^b} d\tilde{x}^b, \frac{\partial x^a}{\partial \tilde{x}^b} \delta \tilde{y}^b \right) .$$

$$(8)$$

Tensors on *TM* that obey the standard tensor transformation law under manifold-induced coordinate changes are called distinguished, or, in short, "d-tensors." An example for a d-vector is the horizontal lift X^H of a vector X in T_xM to $H_{(x,y)}TM$,

- - h

$$X = X^a \partial_{a|x} \mapsto X^H = X^a \delta_{a|(x,y)}.$$
 (9)

The inverse map is given by the push forward along the projection, $\pi_*X^H = X$. The horizontal tangent spaces to *TM* can thus be identified with the tangent spaces to the manifold *M*; this will be important for the identification of field components of tensors over the tangent bundle with field components measured over the spacetime manifold.

A d-tensor of particular importance for our construction of gravitational dynamics for Finsler spacetimes is the curvature of the Cartan nonlinear connection which measures the integrability of the horizontal bundle *HTM*. It is defined as

$$R^{a}{}_{bc} = [\delta_{b}, \delta_{c}]^{a}$$

= $-y^{q} (\delta_{b} \Gamma^{\delta a}{}_{qc} - \delta_{c} \Gamma^{\delta a}{}_{qb} + \Gamma^{\delta a}{}_{pb} \Gamma^{\delta p}{}_{qc}$
 $- \Gamma^{\delta a}{}_{pc} \Gamma^{\delta p}{}_{qb}), \qquad (10)$

using as a short-hand notation the generalized Christoffel symbols

$$\Gamma^{\delta a}{}_{bc} = \frac{1}{2}g^{Faq}(\delta_b g^F_{qc} + \delta_c g^F_{qb} - \delta_q g^F_{bc})$$
$$= \frac{1}{2}g^{Laq}(\delta_b g^L_{qc} + \delta_c g^L_{qb} - \delta_q g^L_{bc}), \qquad (11)$$

which are related to the nonlinear connection coefficients via $N^a{}_b = \Gamma^{\delta a}{}_{bc} y^c$. The proof of the above equality is given in theorem 2 of Ref. [5].

In order to formulate covariant differential equations in the language of Finsler geometry, a linear covariant derivative ∇ that acts on tensor fields over *TM* is needed. This must be compatible with the horizontal/vertical structure so that the identification of horizontal tangent spaces with tangent spaces to the manifold stays intact under transports. In the literature exist four such covariant derivatives with slightly different properties. We will employ the Cartan linear covariant derivative defined by

$$\nabla_{\delta_{a}}\delta_{b} = \Gamma^{\delta q}{}_{ab}\delta_{q}, \qquad \nabla_{\delta_{a}}\bar{\partial}_{b} = \Gamma^{\delta q}{}_{ab}\bar{\partial}_{q},
\nabla_{\bar{\partial}_{a}}\delta_{b} = \frac{1}{2}g^{Fpq}\bar{\partial}_{p}g^{F}_{ab}\delta_{q}, \qquad \nabla_{\bar{\partial}_{a}}\bar{\partial}_{b} = \frac{1}{2}g^{Fpq}\bar{\partial}_{p}g^{F}_{ab}\bar{\partial}_{q}.$$
(12)

In later sections, we will often need the Cartan linear covariant derivative with respect to horizontal directions; we will then use the abbreviation $\nabla_a \equiv \nabla_{\delta_a}$. The action of ∇ on the following Sasaki-type metric on *TM*,

$$G = -g_{ab}^{F} dx^{a} \otimes dx^{b} - \frac{1}{F^{2}} g_{ab}^{F} \delta y^{a} \otimes \delta y^{b}, \qquad (13)$$

gives zero. Hence, this metric is covariantly constant with respect to the Cartan linear connection; it will play a crucial role as an integration measure on *TM*.

This completes our quick review of geometric objects needed in this article. For later use, observe that all the objects definable from F, i.e., the Finsler metric g^F , the Cartan nonlinear connection $N^a{}_b$, its curvature, and the Cartan linear connection, are invariant under the transformation $L \rightarrow L^k$. In the metric limit L(x, y) = $g_{ab}(x)y^ay^b$, they reduce to known constructions from metric geometry: the curvature essentially becomes the Riemann curvature tensor, $R^a{}_{bc}(x, y) = -R^a{}_{dbc}(x)y^d$; the generalized Christoffel symbols become the usual Christoffel symbols, $\Gamma^{\delta a}{}_{bc}(x, y) = \Gamma^a{}_{bc}(x)$; the nonlinear connection now is a linear connection, $N^a{}_b(x, y) =$ $\Gamma^a{}_{bc}(x)y^c$; the Cartan linear covariant derivative in horizontal directions becomes the Levi-Civita covariant derivative while it becomes trivial in vertical directions.

B. Action integrals

We saw that all geometric objects on Finsler spacetimes are homogeneous tensor fields on the tangent bundle. The same will be true for physical fields in our construction. The formulation of field theory actions needs the welldefined integration procedure developed in Ref. [5]. Since integrals over the tangent bundle over homogeneous functions diverge, we must consider integrals over the unit tangent bundle

$$\Sigma = \{ (x, y) \in TM | F(x, y) = 1 \}.$$
(14)

On this domain, we have a natural volume element constructed from the pullback of the Sasaki-type metric (13). Thus, actions arise as integrals of scalar functions f on TMrestricted to Σ as $f_{|\Sigma}$.

To perform such integrals explicitly, we change coordinates on *TM* from the induced $Z^A = (x^a, y^a)$ to the more convenient coordinates $\hat{Z}^A = (\hat{x}^a(x), u^\alpha(x, y), R(x, y))$, where $\alpha = 1...3$, the u^α are zero-homogeneous, and R = F(x, y). Now, Σ is defined by R = 1 and described by coordinates (\hat{x}, u) . The coordinate transformation matrices on *TM* are

$$\frac{\partial \hat{Z}^{A}}{\partial Z^{B}} = \begin{bmatrix} \delta^{a}_{b} & 0\\ \partial_{b} u^{\alpha} & \bar{\partial}_{b} u^{\alpha}\\ \partial_{b} |L|^{1/r} & \bar{\partial}_{b} |L|^{1/r} \end{bmatrix},$$
(15)
$$\frac{\partial Z^{A}}{\partial \hat{Z}^{B}} = \begin{bmatrix} \delta^{a}_{b} & 0 & 0\\ \hat{\partial}_{b} y^{a} & \partial_{u^{\beta}} y^{a} & \frac{y^{a}}{R} \end{bmatrix}$$

and satisfy the invertibility properties

$$\frac{\partial \hat{Z}^{A}}{\partial Z^{C}} \frac{\partial Z^{C}}{\partial \hat{Z}^{B}} = \begin{bmatrix} \delta^{a}_{b} & 0 \\ \partial_{b} u^{\alpha} + \bar{\partial}_{c} u^{\alpha} \hat{\partial}_{b} y^{c} & \bar{\partial}_{c} u^{\alpha} \partial_{u^{\beta}} y^{c} & \bar{\partial}_{c} u^{\alpha} \frac{y^{c}}{R} \\ \partial_{b} |L|^{1/r} + \bar{\partial}_{c} |L|^{1/r} \hat{\partial}_{b} y^{c} & \bar{\partial}_{c} |L|^{1/r} \partial_{u^{\beta}} y^{c} & 1 \end{bmatrix} = \begin{bmatrix} \delta^{a}_{b} & 0 \\ 0 & \delta^{\alpha}_{\beta} & 0 \\ 0 & 0 & 1 \end{bmatrix},$$
(16a)
$$\frac{\partial Z^{A}}{\partial \hat{Z}^{C}} \frac{\partial \hat{Z}^{C}}{\partial Z^{B}} = \begin{bmatrix} \delta^{a}_{b} & 0 \\ \hat{\partial}_{b} y^{a} + \partial_{b} u^{\gamma} \partial_{u^{\gamma}} y^{a} + \frac{y^{a}}{R} \partial_{b} |L|^{1/r} & \partial_{u^{\gamma}} y^{a} \bar{\partial}_{b} u^{\gamma} + \frac{y^{a}}{R} \bar{\partial}_{b} |L|^{1/r} \end{bmatrix} = \begin{bmatrix} \delta^{a}_{b} & 0 \\ 0 & \delta^{a}_{b} \end{bmatrix}.$$
(16b)

We now calculate the pullback of the Sasaki-type metric to Σ in order to determine the relevant volume form. First, we transform Eq. (13) to the new coordinates (\hat{x} , u, R), which yields

$$G = -g^{F}_{ab}d\hat{x}^{a} \otimes d\hat{x}^{b} - \frac{1}{R^{2}}h^{F}_{\alpha\beta}\delta u^{\alpha} \otimes \delta u^{\beta} - \frac{1}{R^{2}}dR \otimes dR$$
(17)

in terms of $h_{\alpha\beta}^F = g_{ab}^F \partial_{\alpha} y^a \partial_{\beta} y^b$ and $\delta u^{\alpha} = du^{\alpha} + (\bar{\partial}_b u^{\alpha} N^b{}_a - \partial_a u^{\alpha}) d\hat{x}^a$. Then the pullback to Σ with R = 1 becomes

$$G^* = -g^F_{ab|\Sigma} d\hat{x}^a \otimes d\hat{x}^b - h^F_{\alpha\beta|\Sigma} \delta u^\alpha \otimes \delta u^\beta.$$
(18)

Using the short-hand notation $g^F = |\det g^F_{ab}|$ and $h^F = |\det h^F_{\alpha\beta}|$, a well-defined integral over Σ of a homogeneous tangent bundle function f now reads

$$\int_{\Sigma} d^4 \hat{x} d^3 u \sqrt{g^F h^F}_{|\Sigma} f(x, y)_{|\Sigma}.$$
 (19)

For tangent bundle functions $A^a(x, y)$ that are homogeneous of degree *m*, the following formulae for integration by parts hold

$$\int_{\Sigma} d^4 \hat{x} d^3 u \sqrt{g^F h^F} |_{\Sigma} (\delta_a A^a)|_{\Sigma}$$

= $-\int_{\Sigma} d^4 \hat{x} d^3 u \sqrt{g^F h^F} |_{\Sigma} [(\Gamma^{\delta p}{}_{pa} + S_a) A^a]|_{\Sigma},$ (20a)

$$\int_{\Sigma} d^{4}\hat{x} d^{3}u \sqrt{g^{F} h^{F}} |_{\Sigma} (\bar{\partial}_{a} A^{a})|_{\Sigma}$$

$$= -\int_{\Sigma} d^{4}\hat{x} d^{3}u \sqrt{g^{F} h^{F}} |_{\Sigma} [(g^{Fpq} \bar{\partial}_{a} g^{F}_{pq} - (m+3)y^{p} g^{F}_{pa}) A^{a}]|_{\Sigma}, \qquad (20b)$$

where $S^{a}_{bc} = \Gamma^{\delta a}_{bc} - \bar{\partial}_{b}N^{a}_{b}$ and $S_{a} = S^{p}_{pa}$. These formulae can be proven with the help of the coordinate transformation relations (16).

The definitions and mathematical techniques presented in this section are the foundation of the following new developments. In the next section, we present how observers are modeled and perform measurements before we turn to the construction of a gravity theory for Finsler spacetimes.

III. OBSERVERS AND MEASUREMENTS

In order to study physics on general Finsler spacetimes, it is necessary to define a mathematical model of physical observers and to determine how they measure time, spatial distances, and physical fields. Guided by general relativity, freely falling observers move on trajectories that extremize the proper time integral; the variation of Eq. (1) here leads to Finsler geodesics. Moreover, a model of observers requires four tangent vectors that build an orthonormal frame; then measurable quantities are the components of physical fields with respect to this frame, evaluated at the observer's position. We explicitly calculate the illustrative example of an observer's measurement of the speed of light on which the results of Ref. [8] are based. To compare measurements of different observers, it is necessary to communicate the results obtained by one observer to another. This communication is realized by a certain class of transformations between different observers; we will show that these transformations have the algebraic structure of a groupoid that generalizes the usual Lorentz group in metric geometry.

A. Orthonormal observer frames

An observer moves along a spacetime curve $\tau \mapsto \gamma(\tau)$ in M with timelike tangents. The parametrization can be chosen so that $\dot{\gamma} \in S_{\gamma}$ is unit timelike; according to definition 1, we now have $|L(\gamma, \dot{\gamma})| = 1$, and the signature of $g_{ab}^{L}(\gamma, \dot{\gamma})$ is Lorentzian.¹ Then the clock postulate (1) tells us that $\dot{\gamma}$ must be interpreted as the local unit time direction of the observer. We may write the normalization condition in the form $g_{(\gamma,\dot{\gamma})}^{F}(e_{0}, e_{0}) = 1$ using the horizontal lift $e_{0} = \dot{\gamma}^{H}$ of $\dot{\gamma}$, see Eq. (9).

To identify the three-space seen by an observer, we will complete e_0 to a four-dimensional basis e_{μ} of $H_{(\gamma, \dot{\gamma})}TM$; as explained before, the projections of the e_{α} for $\alpha = 1...3$ by π_* into $T_{\gamma}M$ then are identified as the spatial tangent directions to the manifold. We determine the three horizontal vectors e_{α} by the condition $g_{(\gamma, \dot{\gamma})}^F(e_0, e_{\alpha}) = 0$. This construction is justified by the observation that a horizontal three-space is defined by a conormal horizontal one-form. The only linearly independent one-form available in terms of geometric data is the vertical form $dL = \bar{\partial}_a L \delta y^a$. This can be mapped globally to the horizontal one-form $d\tilde{L} = \bar{\partial}_a L dx^a$, which is proportional to the Cartan one-form known from Finsler geometry and is a Lagrangian analogue of the Poincaré one-form in Hamiltonian mechanics. The condition $d\tilde{L}_{(\gamma,\dot{\gamma})}(e_\alpha) = 0$ on e_α is equivalent to that stated above in terms of the Finsler metric. We remark that the e_α may depend less trivially on $\dot{\gamma} = \pi_* e_0$ than in Lorentzian geometry because of their defining equation $g_{(\gamma,\dot{\gamma})}^F(e_0, e_\alpha) = 0$.

The definition of unique unit directions in the threedimensional span $\langle e_{\alpha} \rangle$ requires orthonormalization. For this purpose, we use the Finsler metric to set $g_{(\gamma,\dot{\gamma})}^{F}(e_{\alpha}, e_{\beta}) \sim \delta_{\alpha\beta}$, assuming definite signature. The choice of the metric g^{F} for orthonormalization is preferred over that of g^{L} , since only g^{F} is invariant under $L \rightarrow L^{k}$ as are the geometrical objects on Finsler spacetimes, see Sec. II A. We will now prove a useful theorem on the relation between the signatures of the metrics g^{L} and g^{F} where both are defined; a corollary will then confirm our assumption of a definite signature of the Finsler metric on $\langle e_{\alpha} \rangle$.

Theorem 1— On the set $TM \setminus (A \cup \{L = 0\})$, the metric g^L is nondegenerate of signature $(-1_m, 1_p)$ for natural numbers m, p with m + p = 4. Then, the Finsler metric has the same signature, where L(x, y) > 0, and reversed signature $(-1_p, 1_m)$, where L(x, y) < 0.

The observer's time direction e_0 is in S_{γ} which is contained in $TM \setminus (A \cup \{L = 0\})$. This tells us that the metric $g_{(\gamma,\dot{\gamma})}^L$ has signature $(-1_3, 1_1)$ for $L(\gamma, \dot{\gamma}) > 0$ and $(-1_1, 1_3)$ for $L(\gamma, \dot{\gamma}) < 0$. We also know that $g_{(\gamma,\dot{\gamma})}^F(e_0, e_0) = 1$. Hence, we conclude from theorem 1:

Corollary— The Finsler metric $g_{(\gamma,\dot{\gamma})}^F$ evaluated at the tangent bundle position of an observer has Lorentzian signature $(-1_3, 1_1)$, and the unit spatial directions satisfy $g_{(\gamma,\dot{\gamma})}^F(e_\alpha, e_\beta) = -\delta_{\alpha\beta}$.

Proof of Theorem 1— By the definition of Finsler spacetimes, the metric g^L is nondegenerate on $TM \setminus A$, hence also on the smaller set excluding the null structure on which g^F is defined. Now, observe that if an inner product is given by a matrix $C_{ab} = A_{ab} + B_a B_b$ and A_{ab} has indefinite signature $(-1_m, 1_p)$, then the signature of C_{ab} is found to be $(-1_{m+1}, 1_{p-1})$ for $A^{-1ab}B_aB_b < -1$ and $(-1_m, 1_p)$ for $A^{-1ab}B_aB_b > -1$; for $A^{-1ab}B_aB_b = -1$, the result is the once-degenerate signature $(-1_{m-1}, 1_p, 0_1)$. This can be seen in a Sylvester normal form basis for A by using the remaining SO(m, p) freedom. We can apply this result to our situation by identifying

$$A = \frac{nL}{2|L|^{2/n}} (g^L)^{-1}, \qquad B = \left(\frac{n-2}{(n-1)|L|^{2/n}}\right)^{1/2} y \quad (21)$$

¹In the following, we often use the very intuitive notation $(\gamma, \dot{\gamma})$ for points of the tangent bundle $\dot{\gamma} \in T_{\gamma}M \subset TM$, which is analogous to the coordinate representation $(x(\gamma), y(\dot{\gamma}))$.

from Eq. (6). It follows that $A^{-1}(B, B) = n - 2 > -1$; then, the signature of g^F is the signature of A, i.e., that of g^L up to the sign of L.

We summarize our construction of observers' frames and measurements into a precise definition.

Definition 2— Let (M, L, F) be a Finsler spacetime. Physical observers along worldlines $\tau \mapsto \gamma(\tau)$ in M are described by a frame basis $\{e_{\mu}\}$ of $H_{(\gamma, \dot{\gamma})}TM$ which

- (i) has a timelike vector e_0 in the sense $\pi_* e_0 = \dot{\gamma} \in S_{\gamma}$; and
- (ii) is g^F -orthogonal, $g^F_{(\gamma,\dot{\gamma})}(e_{\mu}, e_{\nu}) = -\eta_{\mu\nu}$.

They measure the components of horizontal tensor fields over *TM* with respect to their frame at their tangent bundle position $(\gamma, \dot{\gamma})$.

The Minkowski metric $\eta_{\mu\nu}$ in this definition has signature $(-1_1, 1_3)$. We emphasize again that the frame $\{e_{\mu}\}$ in $H_{(\gamma, \hat{\gamma})}TM$ can be identified one-to-one with a frame $\{\pi_*e_{\mu}\}$ in $T_{\gamma}M$, or reversely by the horizontal lift. From Eq. (6), one can show that the observer frame has the nice property to diagonalize g^F and g^L simultaneously. We will now discuss the measurement procedure in more detail and present the example of how an observer measures the speed of light.

B. Measuring the speed of light

Definition 2 of the observer frame includes the statement that a physical observable is given by the components of a horizontal tensor field with respect to the observer's frame, evaluated at her position on the tangent bundle, i.e., at her position on the manifold and her four-velocity. The motivation for this is as follows. The geometry of Finsler spacetimes is formulated on the tangent bundle TM, and hence matter tensor fields coupling to this gravitational background must also be defined over TM. Not all such tensor fields can be interpreted as tensor fields from the perspective of the spacetime manifold M. This interpretation requires that the tensor fields be horizontal; then, they are multilinear maps built on the horizontal space H_PTM and its dual which are identified with the tangent space $T_{\pi(P)}M$ and its dual. Consider the example of a 2-form field Φ over TM; in the horizontal/vertical basis, this expands as

$$\Phi = \Phi_{1ab}(x, y)dx^a \wedge dx^b + 2\Phi_{2ab}(x, y)dx^a \wedge \delta y^b + \Phi_{3ab}(x, y)\delta y^a \wedge \delta y^b.$$
(22)

Only the purely horizontal part $\Phi_{1ab}(x, y)dx^a \wedge dx^b$ has a clear interpretation. Note that such horizontal tensor fields are automatically d-tensor fields and have the same number of components as a tensor field of same rank on M. The difference is that the components depend on the tangent bundle position. The measurement of a horizontal tensor field by an observer at the tangent bundle position $(\gamma, \dot{\gamma})$ clearly requires an observer frame of $H_{(\gamma,\dot{\gamma})}TM$ in order to read out the components.

We emphasize that the dependence of observables on the four-velocity of the observer is not surprising. Neither is it problematic as long as observers can communicate their results. In general relativity, observables are the components of tensor fields over M with respect to the observer's frame in $T_{\gamma}M$; they clearly depend on $\dot{\gamma}$ which induces the splitting of $T_{\gamma}M$ into time and space directions. On Finsler spacetimes, the dependence of observables on the observer's four-velocity is not only present in the time/space split of $H_{(\gamma,\dot{\gamma})}TM$, but also in the argument of the tensor field components.

As a simple example, we discuss the measurement of the spatial velocity of a point particle that moves on a worldline ρ with horizontal tangent $\dot{\rho}$. This can be expanded in the orthonormal frame of an observer as $\dot{\rho} = \dot{\rho}^0 e_0 + \dot{\rho}^a = \dot{\rho}^0 e_0 + \dot{\rho}^a e_{\alpha}$, where we recall that $e_0 = \dot{\gamma}$ is the observer's four-velocity. The time $\dot{\rho}^0$ passes while the particle moves in spatial direction $\dot{\rho}^{\alpha}$, so the spatial velocity \vec{v} and its square v^2 are

$$\vec{v} = \frac{\dot{\vec{\rho}}}{\dot{\rho}^{0}}, \qquad v^{2} = \frac{\delta_{\alpha\beta}\dot{\rho}^{\alpha}\dot{\rho}^{\alpha}}{(\dot{\rho}^{0})^{2}} = -\frac{g_{(\gamma,\dot{\gamma})}^{F}(\vec{\rho},\vec{\rho})}{g_{(\gamma,\dot{\gamma})}^{F}(\dot{\rho},\dot{\gamma})^{2}}.$$
 (23)

As a consequence of this formula, we may derive the speed of light seen by a given observer. As discussed in Ref. [5], light propagates on null worldlines ρ with $L(\rho, \dot{\rho}) = 0$, which is equivalent to $F(\rho, \dot{\rho})^2 = 0$. We can use this fact to replace the Finsler metric in the formula for the velocity above. Taylor-expanding $F(\rho, \dot{\rho}^0 e_0 + \dot{\rho})^2 = 0$ around $\dot{\rho} = 0$ yields

$$0 = (\dot{\rho}^{0})^{2} + g^{F}_{(\rho,\dot{\gamma})}(\dot{\vec{\rho}}, \dot{\vec{\rho}}) + \sum_{k=3}^{\infty} \frac{(\dot{\rho}^{0})^{2-k}}{k!} \bar{\partial}_{c_{1}} \dots \bar{\partial}_{c_{k}} F(\rho, \dot{\gamma})^{2} \dot{\vec{\rho}}^{c_{1}} \dots \dot{\vec{\rho}}^{c_{k}}.$$
 (24)

Evaluating this formula at the position of the observer $\rho = \gamma$ and dividing by $(\dot{\rho}^0)^2$, we immediately obtain an expression for the speed of light $c^2_{(\gamma,\dot{\gamma})}(\dot{\vec{\rho}})$, i.e., the speed of light traveling in spatial direction $\dot{\vec{\rho}}$ and measured by the observer $(\gamma, \dot{\gamma})$:

$$c_{(\gamma,\dot{\gamma})}^{2}(\dot{\vec{\rho}}) = 1 + \sum_{k=3}^{\infty} \frac{(\dot{\rho}^{0})^{-k}}{k!} \bar{\partial}_{c_{1}} \dots \bar{\partial}_{c_{k}} F(\gamma,\dot{\gamma})^{2} \dot{\vec{\rho}}^{c_{1}} \dots \dot{\vec{\rho}}^{c_{k}}.$$
(25)

The $\dot{\rho}^0$ are determined by solving the null condition $L(\gamma, \dot{\rho}^0 \dot{\gamma} + \dot{\vec{\rho}}) = 0$; on a generic Finsler spacetime, there can be more than one solution since the null structure can be very complicated. From Eq. (25), we see that the measured speed of light depends on the higher-thansecond-order derivatives of the squared Finsler function; these vanish in the metric limit where we thus reobtain $c^2_{(\gamma,\dot{\gamma})}(\dot{\vec{\rho}}) = 1$ independent of the observer and the spatial direction of the light ray. The formulae (23) and (24) enable us to compare experimental results on particle and

light velocities with predictions on specific Finsler spacetime models. In Ref. [8], we used this to study the possibility of superluminal particle propagation.

C. Generalized Lorentz transformations

We already stressed the importance that observers should be able to communicate their measurements. Consider two observers whose worldlines meet at a point $x \in M$. Since observers by definition 2 measure the components of horizontal tensor fields in their frame and at their tangent bundle position, we need to determine which transformation uniquely maps an observer frame $\{e_{\mu}\}$ in $H_{(x,y)}TM$ to a second observer frame $\{f_{\mu}\}$ in $H_{(x,z)}TM$. Their respective four-velocities, or time directions, y = $\pi_* e_0$ and $z = \pi_* f_0$ generically are different, so that the two observer frames are objects in tangent spaces to TM at different points. As a consequence, we will now demonstrate that the transformations between observers consist of two parts: the first is a transport of the frame $\{e_{\mu}\}$ from (x, y) to (x, z), and the second will turn out to be a Lorentz transformation.

Theorem 2— Consider two observer frames $\{e_{\mu}\}$ in $H_{(x,y)}TM$ and $\{f_{\mu}\}$ in $H_{(x,z)}TM$ on a Finsler spacetime (M, L, F). If z is in a sufficiently small neighborhood around $y \in T_x M$, then the following procedure defines a unique map $\{e_{\mu}\} \mapsto \{f_{\mu}\}$:

- (i) Let t → v(t) be a vertical autoparallel of the Cartan linear connection that connects v(0) = (x, y) to v(1) = (x, z); this satisfies π_{*} v = 0 and ∇_v v = 0. Determine a frame {ê_μ(v(t))} along v(t) by parallel transport ∇_v ê_μ = 0 with the initial condition ê_μ(v(0)) = e_μ.
- (ii) Find the unique Lorentz transformation Λ so that $f_{\mu} = \Lambda^{\nu}_{\mu} \hat{e}_{\nu}(v(1)).$

Proof We first show that the curve v required in (i) exists. The verticality condition $\pi_* \dot{v} = 0$ implies $\dot{v} = \dot{v}^a \bar{\partial}_a$; the definition of the Cartan linear connection (12) then tells us that $\nabla_{\dot{v}} \dot{v} = 0$ is equivalent to solving $\ddot{v}^a + \frac{1}{2}g^{Fap}\bar{\partial}_p g_{bc}^F \dot{v}^b \dot{v}^c = 0$. This has a unique solution connecting (x, y) to any point (x, z) in a sufficiently small neighborhood in $T_x M$. Now let $\{\hat{e}_\mu(v(t))\}$ be the parallelly transported vector fields $\nabla_{\dot{v}} \hat{e}_\mu = 0$ with $\hat{e}_\mu(v(0)) = e_\mu$. The properties of the Cartan linear connection ensure that the \hat{e}_μ are horizontal fields. Observe also that $\nabla_{\dot{v}}(g_v^F(\hat{e}_\mu, \hat{e}_v)) = 0$ along the curve v since g^F is covariantly constant under ∇ . It follows that

$$g_{\nu(t)}^{F}(\hat{e}_{\mu}(\nu(t)), \hat{e}_{\nu}(\nu(t))) = -\eta_{\mu\nu}$$
(26)

is independent of *t* and holds, in particular, at the final point of the transport v(1) = (x, z). Now $\{\hat{e}_{\mu}(v(1))\}$ and $\{f_{\mu}\}$ are orthonormal frames with respect to g^{F} in $H_{(x,z)}TM$; hence, they are related by a unique Lorentz transformation as stated in point *(ii)* of the theorem. \Box



FIG. 3 (color online). Transformation between two observer frames: the frame $\{e_{\mu}\}$ in $H_{(x,y)}TM$ is first parallely transported to $\{\hat{e}_{\mu}\}$ in $H_{(x,z)}TM$ and second, Lorentz transformed into the final frame $\{f_{\mu}\}$.

The procedure described in theorem 2 provides a map between the frames of two observers at the same point of the manifold $x \in M$, but with different four-velocities $y, z \in S_x \subset T_x M$; we display the two parts of this procedure as $\Lambda \circ P_{y \to z}$, i.e., as parallel transport followed by Lorentz transformation, which is illustrated in Fig. 3. The combined maps transform observers uniquely into one another as long as the autoparallel v connecting the vertically different points in *TM* exists and is unique. This is certainly the case if (x, y) and (x, z) are sufficiently close to each other. Whether the geometric structure of a specific, or maybe all, Finsler spacetimes is such that unique transformations between all observers exist requires further investigation.

In the observer transformations on generic Finsler spacetimes, there appears an additional ingredient that is not present on metric spacetimes. Before applying the Lorentz transformation to the frame, one has to perform a parallel transport in the vertical tangent space. In the metric limit, the vertical covariant derivative becomes trivial so that the parallelly transported frame does not change at all along the curve v. In this special case, the transformation of an observer thus reduces to $\Lambda \circ id_{y \to z}$, which is fully determined by a Lorentz transformation.

The observer transformations on Finsler spacetimes essentially have the algebraic structure of a groupoid that reduces to the Lorentz group in the metric limit. We first review the general definition of a groupoid and then show how this applies to our case.

Definition 3— A groupoid G consists of a set of objects G_0 and a set of arrows G_1 . Every arrow A is assigned a source e = s(A) and a target f = t(A) by the maps $s: G_1 \rightarrow G_0$ and $t: G_1 \rightarrow G_0$; one writes this as $A: e \rightarrow f$. For arrows A and B whose source and target match as t(A) = s(B) there exists an associative multiplication $G_1 \times G_1 \rightarrow G_1$, $(A, B) \mapsto BA$ with

$$s(BA) = s(A),$$
 $t(BA) = t(B),$ $C(BA) = (CB)A.$
(27)

A unit map $G_0 \to G_1$, $e \mapsto \mathbb{1}_e$ where $\mathbb{1}_e: e \to e$ exists so that

$$\mathbb{1}_{t(A)}A = A = A \mathbb{1}_{s(A)}.$$
(28)

For every arrow A exists an inverse arrow A^{-1} that satisfies

/ 15

$$s(A^{-1}) = t(A), t(A^{-1}) = s(A), (29)$$

$$A^{-1}A = \mathbb{1}_{s(A)}, AA^{-1} = \mathbb{1}_{t(A)}.$$

Groupoids are generalizations of groups. These can be expressed as groupoids with a single object in G_0 ; then, the arrows correspond to group elements, all of which can be multiplied since sources and targets always match. The multiplication is associative, and the identity element and inverse elements exist.

Consider $G_0 = S_x \subset T_x M$ as the set of unit timelike vectors which contains the different four-velocities of observers at the point $x \in M$. Let the arrows in G_1 be the set of all maps between two observer frames at x, which are defined by the procedure stated in theorem 2. In case the involved vertical autoparallels connect the four-velocities uniquely, the sets G_0 and G_1 define a groupoid: source and target of a map $A = \Lambda \circ P_{y \to z}$ between two frames $\{e_{\mu} \in$ $H_{(x,y)}TM$ and $\{f_{\mu} \in H_{(x,z)}TM\}$ are simply given by $s(A) = y \in S_x$ and $t(A) = z \in S_x$; the multiplication BA is defined by applying the procedure of theorem 2 to construct the map between s(A) and t(B), which gives the properties (27); we choose the unit map $\mathbb{1}_{v}$ that provides (28) as $\mathbb{1}_y = \mathbb{1} \circ id_{y \to y}$, i.e., as trivial parallel transport of the frame $\{e_{\mu} \in H_{(x,y)}TM\}$ with respect to the Cartan linear connection along the vertical autoparallel that stays at (x, y) followed by the identity Lorentz transformation. Finally, we define the inverse $A^{-1} = \Lambda^{-1} \circ P_{z \to v}$, where $P_{z \rightarrow y}$ denotes parallel transport backwards along the unique vertical autoparallel connecting (x, y) and (x, z), which is also used for $P_{y \to z}$; to check the properties (29), one simply shows that parallel transport of the frames and Lorentz transformation commute. Thus, we have shown the following result:

Theorem 3— On Finsler spacetimes (M, L, F), the transformations between observer frames $\{e_{\mu} \in H_{(x,y)}TM\}$ at $x \in M$ that are attached to points $(x, y) \in U_x \subset S_x$ define a groupoid G under the condition that any pair of points in U_x can be connected by a unique vertical autoparallel of the Cartan linear connection.

We already discussed that the transformations of observer frames reduce to the form $A = \Lambda \circ id_{v \to z}$ in the limit of metric geometry. Hence, the only information contained in the reduced groupoid \tilde{G} with $\tilde{G}_0 = S_x$ and $\tilde{G}_1 = \{\Lambda \circ id_{v \to z}\}$ is given by the Lorentz transformations. In mathematically precise language, this can be expressed as the equivalence of G to the Lorentz group seen as a

groupoid \mathcal{H} with a single object $H_0 = \{x\}$ and arrows $H_1 = \{\Lambda\}$. The functor $\varphi: \tilde{\mathcal{G}} \to \mathcal{H}$ establishing the equivalence can be defined by the projection $\varphi_0 =$ $\pi: \tilde{G}_0 \to H_0$ and by $\varphi_1: \tilde{G}_1 \to H_1$, $\Lambda \circ id_{y \to z} \Lambda$. Indeed, φ can be checked to be injective, full, and essentially surjective, and so it makes \tilde{G} and \mathcal{H} equivalent. See Ref. [10] for details on the required mathematical definitions.

IV. GRAVITATIONAL DYNAMICS

We have now reviewed the basic concept of Finsler spacetime geometry and laid the foundations for the interpretation of physics on these backgrounds. We have seen that well-defined observers exist which communicate with each other by means of groupoid transformations that generalize the Lorentz group in metric geometry. As emphasized in previous work [5,8], physical predictions in our generalized geometric framework require gravitational dynamics for the fundamental geometry function L to determine specific spacetime solutions. In this section, we for the first time present a Finsler gravity action along with a consistent minimal coupling principle between gravity and matter. We begin our presentation with the variation of the pure gravity action, before the coupling of Finsler gravity to matter is discussed in Sec. IV B; the full field equation is derived in Sec. IV C. Moreover, we prove that the Finsler gravity field equation becomes equivalent to the Einstein equations in the metric limit.

A. Action and vacuum equations

On Finsler spacetime, the simplest curvature scalar built from the nonlinear curvature tensor R^a_{bc} , which is relevant for the tidal acceleration of Finsler geodesics, is $\mathcal{R} =$ $R^{a}_{\ ab}y^{b}$. This contains the lowest number of derivatives on the fundamental function L without involving additional d-tensors besides the curvature, like $S^a_{\ bc}$ or $\bar{\partial}_a g^F_{\ bc}$. Recall that integrals are well-defined over the sevendimensional unit tangent bundle Σ with coordinates (\hat{x}, u) , as discussed in Sec. II B. These two facts directly lead to our Finsler gravity action

$$S_G[L] = \int d^4 \hat{x} d^3 u [\sqrt{g^F h^F} \mathcal{R}]_{|\Sigma}.$$
 (30)

The gravitational field equation in vacuum now is obtained by variation with respect to L. To perform this variation for an *m*-homogeneous function f(x, y) on TM restricted to Σ , it is useful to realize that

$$\delta(f_{|\Sigma}) = (\delta f)_{|\Sigma} - \frac{m}{n} f_{|\Sigma} \frac{\delta L}{L}, \qquad (31)$$

where *n* is the homogeneity of *L*. With the help of this formula and the results for integration by parts in Eq. (20), we can derive the vacuum field equations in three steps. The first uses the variation formula above with $f(x, y) = \sqrt{g^F h^F} R^a_{\ ab} y^b$ and m = 5, which yields

$$\delta S_G[L] = \int d^4 \hat{x} d^3 u \bigg[\delta(\sqrt{g^F h^F} \mathcal{R}) - \frac{5}{n} \sqrt{g^F h^F} \mathcal{R} \frac{\delta L}{L} \bigg]_{|\Sigma}.$$
(32)

The second step is the variation of the volume element which leads to

$$\delta S_G[L] = \int d^4 \hat{x} d^3 u \sqrt{g^F h^F} \sum_{|\Sigma|} \left[\left(g^{Fpq} \delta g^F_{pq} - \frac{6}{n} \frac{\delta L}{L} \right) \mathcal{R} + y^b \delta R^a{}_{ab} \right]_{|\Sigma'},$$
(33)

while in the third step, we use the following identities:

$$\int d^{4}\hat{x}d^{3}u[\sqrt{g^{F}h^{F}}g^{Fpq}\delta g^{F}_{pq}\mathcal{R}]_{|\Sigma}$$
$$=\int d^{4}\hat{x}d^{3}u\left[\sqrt{g^{F}h^{F}}g^{Fab}\bar{\partial}_{a}\bar{\partial}_{b}\mathcal{R}\frac{\delta L}{nL}\right]_{|\Sigma},\qquad(34a)$$

$$\int d^{4}\hat{x}d^{3}u[\sqrt{g^{F}h^{F}}y^{b}\delta R^{a}{}_{ab}]_{|\Sigma}$$

$$= \int d^{4}\hat{x}d^{3}u\left[\sqrt{g^{F}h^{F}}2g^{Fab}(\nabla_{a}S_{b} + S_{a}S_{b} + \bar{\partial}_{a}\nabla S_{b})\frac{\delta L}{nL}\right]_{|\Sigma},$$
(34b)

to arrive at the final form of the variation of the Finsler gravity action (30):

$$\delta S_G[L] = \int d^4 \hat{x} d^3 u \sqrt{g^F h^F} \sum_{|\Sigma|} [g^{Fab} \bar{\partial}_a \bar{\partial}_b \mathcal{R} - 6\mathcal{R} + 2g^{Fab} (\nabla_a S_b + S_a S_b + \bar{\partial}_a \nabla S_b)]_{|\Sigma|} \frac{\delta L}{nL}.$$
(35)

For further details of this variation, we refer the reader to Appendix A 1. Now, we can read off the vacuum field equation on Σ as

$$[g^{Fab}\bar{\partial}_{a}\bar{\partial}_{b}\mathcal{R} - 6\mathcal{R} + 2g^{Fab}(\nabla_{a}S_{b} + S_{a}S_{b} + \bar{\partial}_{a}\nabla S_{b})]_{|\Sigma} = 0.$$
(36)

Observe that all terms in the bracket are zerohomogeneous on TM, except the second term \mathcal{R} that has homogeneity two. Since $(\mathcal{R})_{|\Sigma} = (\mathcal{R}/F^2)_{|\Sigma}$, we can replace the second term by \mathcal{R}/F^2 , which is now also zerohomogeneous. Hence, the equation can be lifted to TM in the form

$$g^{Fab}\bar{\partial}_a\bar{\partial}_b\mathcal{R} - \frac{6}{F^2}\mathcal{R} + 2g^{Fab}(\nabla_a S_b + S_a S_b + \bar{\partial}_a \nabla S_b) = 0.$$
(37)

It seems as if this equation could be invalid on $\{L = 0\} = \{F = 0\}$, where *F* is not differentiable so that the Finsler metric g^F does not exist. However, this is not the case: the equation is valid also on the null structure. To see this, one expresses g^F through g^L with the help of formula (6) and multiplies by F^2 . The resulting equation is well-defined wherever g^L is nondegenerate and, in particular, on the null structure. Note that Eq. (37) is invariant under the transformation $L \rightarrow L^k$, which will be a guiding principle for matter coupling below.

In the metric limit $L = g_{ab}(x)y^ay^b$, the tensors in the Finsler gravity equation reduce as $\mathcal{R} = -y^ay^bR_{ab}$ and $S_a = 0$, where R_{ab} is the Ricci tensor of the metric g. Accordingly, the field equation becomes

$$2R + \frac{6}{F^2} R_{ab} y^a y^b = 0, (38)$$

which is equivalent to the Einstein vacuum equations $R_{ab} = 0$ by differentiating twice with respect to y. We conclude that a family of solutions of the Finsler gravity vacuum equation (37) is induced by solutions $g_{ab}(x)$ of the vacuum Einstein equations via the fundamental functions $L_k = (g_{ab}(x)y^ay^b)^k$. In Sec. VI, we will present a solution of the Finsler gravity vacuum equation beyond metric geometry.

B. Consistent matter coupling

Above, we have achieved a consistent generalization of vacuum Einstein gravity from metric spacetimes to Finsler spacetimes. Next, we will show that this generalization can be completed by the coupling of matter fields. For this purpose, we will discuss a minimal coupling principle that generates consistent matter field actions on Finsler spacetimes from their well-known counterparts on metric spacetimes. In the discussion, we restrict our attention to *p*-form fields; spinor fields have to be investigated further. In Sec. IV C, we will deduce the complete gravity equations with energy-momentum source term.

Consider an action $S_m[g, \phi]$ for a physical *p*-form field ϕ on a Lorentzian spacetime (M, g),

$$\tilde{S}_m[g,\phi] = \int_M d^4x \sqrt{g} \mathcal{L}(g,\phi,\mathrm{d}\phi).$$
(39)

The corresponding matter action on Finsler spacetime is obtained by lifting \tilde{S}_m to the tangent bundle *TM* equipped with the Sasaki-type metric G(x, y) defined in Eq. (13) in the following way:

 (i) consider the Lagrangian density L(...) of the standard theory on M as a contraction prescription that forms a scalar function from various tensorial objects;

- (ii) replace the Lorentzian metric g(x) in $\mathcal{L}(...)$ by the Sasaki-type metric G(x, y);
- (iii) replace the *p*-form field $\phi(x)$ on *M* by a zerohomogeneous *p*-form field² $\Phi(x, y)$ on *TM*;
- (iv) introduce Lagrange multipliers λ for all not purely horizontal components of Φ ;
- (v) finally, integrate over the unit tangent bundle Σ with the volume form given by the pullback G^* of the Sasaki-type metric.

The result of this procedure is the Finsler spacetime field theory action

$$S_m[L, \Phi, \lambda] = \int_{\Sigma} d^4 \hat{x} d^3 u [\sqrt{g^F h^F} (\mathcal{L}(G, \Phi, d\Phi) + \lambda (1 - P^H) \Phi)]_{|\Sigma}.$$
(40)

The projection P^H projects to the purely horizontal part of the *p*-form Φ ; in the example of the general two-form on *TM* displayed in Eq. (22), we have

$$P^H \Phi = \Phi_{1ab} dx^a \wedge dx^b. \tag{41}$$

The Lagrange multiplier guarantees that the on-shell degrees of freedom of Φ are precisely those with a clear physical interpretation as fields along the manifold M, as discussed in Sec. III B. The minimal coupling principle for matter to Finsler spacetime presented above is a slightly refined version of that in Ref. [5], where we discussed electrodynamics on Finsler spacetime. The only modification here is in the definition of the Sasaki-type metric G(x, y) which is now defined in terms of the Finsler metric g^F instead of g^L . This change ensures that the resulting matter action $S_m[L, \Phi, \lambda]$ is invariant under $L \rightarrow L^k$ in the same way as the pure gravity action. Nevertheless, the results obtained for electrodynamics on Finsler spacetime in Ref. [5] are unchanged by the refined coupling principle presented here.

The matter field equations obtained by extremizing the action with respect to the *p*-form field Φ and the Lagrange multiplier λ can be studied most easily if expressed in components with respect to the horizontal/vertical basis. The calculation is performed in detail in Appendix A 2. We display the results with the convention that barred indices denote vertical components, unbarred indices now denote horizontal components, and capital indices both horizontal and vertical components. Variation with respect to the Lagrange multiplier yields the constraints

$$\Phi_{\bar{a}_1...\bar{a}_i a_{i+1}...a_p} = 0, \quad \forall \ i = 1...p.$$
(42)

Variation for the purely horizontal components of Φ gives

$$\frac{\partial \mathcal{L}}{\partial \Phi_{a_1\dots a_p}} - (p+1)(\nabla_q + S_q) \frac{\partial \mathcal{L}}{\partial (\mathrm{d}\Phi_{qa_1\dots a_p})} - (\bar{\partial}_{\bar{q}} + g^{Fmn} \bar{\partial}_{\bar{q}} g^F_{mn} - 4g^F_{\bar{q}q} y^q) \frac{\partial \mathcal{L}}{\partial (\mathrm{d}\Phi_{\bar{q}a_1\dots a_p})} = 0,$$
(43)

which determines the evolution of the physical field components, while variation with respect to the remaining components produces

$$\lambda^{\bar{a}_{1}A_{2}...A_{p}} = -\frac{\partial \mathcal{L}}{\partial \Phi_{\bar{a}_{1}A_{2}...A_{p}}} + (p+1)(\nabla_{q} + S_{q})\frac{\partial \mathcal{L}}{\partial(d\Phi_{q\bar{a}_{1}A_{2}...A_{p}})} + \frac{p(p+1)}{2}\frac{\partial \mathcal{L}}{\partial(d\Phi_{PQA_{2}...A_{p}})}\gamma^{\bar{a}_{1}}_{PQ} + (\bar{\partial}_{\bar{q}} + g^{Fmn}\bar{\partial}_{\bar{q}}g^{F}_{mn} - 4g^{F}_{\bar{q}q}y^{q})\frac{\partial \mathcal{L}}{\partial(d\Phi_{\bar{q}\bar{a}_{1}A_{2}...A_{p}})},$$

$$(44)$$

which fixes the components of the Lagrange multiplier. The $\gamma_{PQ}^{\bar{a}}$ are the commutator coefficients of the horizontal/vertical basis.

Our coupling principle is consistent with the metric limit, i.e., the equations of motion obtained from the Finsler spacetime action, reduce to the equations of motion on Lorentzian spacetime in the case $L = g_{ab}(x)y^ay^b$ and $\Phi_{A_1...A_p}(x, y) = \phi_{A_1...A_p}(x)$. Then, we have the geometric identity $S_a = 0$; moreover

$$d\Phi_{a_1\dots a_{p+1}} = (p+1)\partial_{[a_1}\phi_{a_2\dots a_{p+1}]},\tag{45}$$

using the constraints (42) and the fact that the horizontal derivative acts as a partial derivative on the *y*-independent *p*-form components. Finally,

$$\frac{\partial \mathcal{L}}{\partial (\mathrm{d}\Phi_{\bar{q}a_1\dots a_p})} = 0 \tag{46}$$

because, as a consequence of our coupling principle where the Sasaki-type metric is block-diagonal in the horizontal/ vertical basis, the vertical index of $d\Phi_{\bar{q}a_1...a_p}$ must appear in $\mathcal{L}(G, \Phi, d\Phi)$ contracted via g^F into either a vertical derivative or into components of Φ with at least one vertical index. In the metric limit, vertical derivatives give zero, while the constraints (42) guarantee that all components of Φ with at least one vertical index vanish. Combining these observations shows that Eq. (43) reduces to

$$\frac{\partial \mathcal{L}}{\partial \Phi_{a_1\dots a_p}} - (p+1)\nabla_q \frac{\partial \mathcal{L}}{\partial (\mathrm{d}\Phi_{qa_1\dots a_p})} = 0, \qquad (47)$$

where ∇ now operates in the same way as the Levi-Civita connection of the metric *g*. Again, as a consequence of our minimal coupling principle with the block-diagonal form

- 0

²A two-form field Φ as in Eq. (22), for example, is zerohomogeneous if and only if its components Φ_1 , Φ_2 and Φ_3 have the homogeneities 0, -1, and -2, respectively; these are cancelled by the homogeneity of δy .

of the Sasaki-type metric in the horizontal/vertical basis, we can conclude in the metric limit that

$$\frac{\partial \mathcal{L}(G, \Phi, d\Phi)}{\partial \Phi_{a_1...a_p}} = \frac{\partial \mathcal{L}(g, \phi, d\phi)}{\partial \phi_{a_1...a_p}},$$

$$\frac{\partial \mathcal{L}(G, \Phi, d\Phi)}{\partial (d\Phi_{qa_1...a_p})} = \frac{\partial \mathcal{L}(g, \phi, d\phi)}{\partial (d\phi_{qa_1...a_p})}$$
(48)

so that Eq. (47) becomes equivalent to the standard *p*-form field equation of motion on metric spacetime.

Our minimal coupling procedure for matter fields to Finsler spacetime can be applied immediately for instance to the scalar field, as done in Ref. [8]. Note that it can be easily extended to the case of interacting form fields of any degree with metric spacetime action

$$\tilde{S}_m[g,\phi_1,\phi_2,\ldots] = \int_M d^4x \sqrt{g} \mathcal{L}(g,\phi_1,\mathrm{d}\phi_1,\phi_2,\mathrm{d}\phi_2,\ldots).$$
(49)

The minimal coupling procedure then leads to the action

$$S_{m}[L, \Phi_{1}, \lambda_{1}, \Phi_{2}, \lambda_{2}, \ldots]$$

$$= \int_{\Sigma} d^{4}\hat{x} d^{3}u \left[\sqrt{g^{F} h^{F}} (\mathcal{L}(G, \Phi_{1}, d\Phi_{1}, \Phi_{2}, d\Phi_{2}, \ldots) + \sum_{I} \lambda_{I} (1 - P^{H}) \Phi_{I}) \right]_{|\Sigma}.$$
(50)

The equations of motion for each field ϕ_I have the same form as in the single field case, and the metric limit leads to the standard field equations by arguments that proceed in a completely analogous way as before.

In the standard formulation of electrodynamics, the action is a functional $\tilde{S}_m[g, A, dA]$ of a one-form potential A, but the classical physical field is F = dA. Our minimal coupling principle to obtain an action on Finsler space-times cannot be applied immediately to this situation: the problem is that the Lagrange multiplier then only kills the vertical components of the lift of A, but does not guarantee that the lift of F is purely horizontal. This problem is solved in Ref. [5] by starting from an equivalent interacting action of the form $\tilde{S}_m[g, A, dA, F, dF]$, which provides the complete set of Maxwell equations F = dA and $d \star_g F = 0$ by variation. Now, the minimal coupling principle entails that both the lifted fields A and F are purely horizontal and can be interpreted physically.

C. Gravity field equations and metric limit

We are now in the position to study the interplay between the matter actions S_m introduced in Eq. (40) and the pure Finsler gravity action S_G in Eq. (30). Their sum provides a complete description of gravity and classical matter fields on Finsler spacetimes:

$$S[L, \Phi, \lambda] = \kappa^{-1} S_G[L] + S_m[L, \Phi, \lambda]$$

= $\kappa^{-1} \int d^4 \hat{x} d^3 u [\sqrt{g^F h^F} \mathcal{R}]_{|\Sigma}$
+ $\int d^4 \hat{x} d^3 u [\sqrt{g^F h^F} \mathcal{L}(G, \Phi, d\Phi)]_{|\Sigma}$
+ $\int d^4 x d^3 u [\sqrt{g^F h^F} \lambda (1 - P^H) \Phi]_{|\Sigma}.$ (51)

As usual, the matter field equations following from this are the same as for the pure matter action. The gravitational field equations are obtained by variation with respect to the fundamental geometry function L. The variation of S_m with respect to L is

$$\delta S_m = \int d^4 \hat{x} d^3 u \left(\frac{\delta S_m}{\delta L} \delta L \right)_{|\Sigma}$$

=
$$\int d^4 \hat{x} d^3 u \left(\sqrt{g^F h^F} \frac{nL}{\sqrt{g^F h^F}} \frac{\delta S_m}{\delta L} \right)_{|\Sigma} \frac{\delta L}{nL}$$
(52)

and leads us to the definition of the energy-momentum scalar $T_{|\Sigma}$ on the unit tangent bundle as

$$T_{|\Sigma} = \left(\frac{nL}{\sqrt{g^F h^F}} \frac{\delta S_m}{\delta L}\right)_{|\Sigma}.$$
 (53)

With this definition, the complete gravitational field equations on Finsler spacetime including energy-momentum sources formally become

$$[g^{Fab}\bar{\partial}_{a}\bar{\partial}_{b}\mathcal{R} - \frac{6}{F^{2}}\mathcal{R} + 2g^{Fab}(\nabla_{a}S_{b} + S_{a}S_{b} + \bar{\partial}_{a}\nabla S_{b})]_{|\Sigma}$$

= $-\kappa T_{|\Sigma}.$ (54)

As in the vacuum case with $T_{|\Sigma} = 0$, these equations can be lifted to *TM*. The terms in the bracket on the left-hand side are all zero-homogeneous and can be lifted trivially. The terms in *T* without the restriction on the right-hand side in principle can result from variation with different homogeneities; to lift these, one simply multiplies each term by the appropriate power of *F* in order to make it zerohomogeneous. This is the same procedure applied in Sec. IVA to the gravity side.

The gravitational constant κ will now be determined so that the gravitational field equation on Finsler spacetimes becomes equivalent to the Einstein equations in the metric limit. Variation with respect to *L* of the concrete form of the matter action in Eq. (51) and performing the metric limit, i.e., $g_{ab}^F(x, y) = -g_{ab}(x)$ for observers and $\Phi_{A_1...A_p}(x, y) = \phi_{A_1...A_p}(x)$, the gravity equation (54) becomes

$$2g^{ab}R_{ab} + 6\frac{R_{ab}y^{a}y^{b}}{|g_{pq}y^{p}y^{q}|} = -\kappa \left(4\mathcal{L} - 4g_{ab}\frac{\partial\mathcal{L}}{\partial g_{ab}} - 24\frac{y_{a}y_{b}}{|g_{pq}y^{p}y^{q}|}\frac{\partial\mathcal{L}}{\partial g_{ab}}\right).$$
(55)

The detailed calculation of this result is involved and can be found in Appendix A 2. Introducing the standard energy-momentum tensor of *p*-form fields on Lorentzian metric spacetimes $\tilde{T}^{ab} = g^{ab} \mathcal{L} + 2 \frac{\partial \mathcal{L}}{\partial g_{ab}}$ and its trace $\tilde{T} = \tilde{T}^{ab}g_{ab} = 4\mathcal{L} + 2g_{ab} \frac{\partial \mathcal{L}}{\partial g_{ab}}$, we can rewrite the equation above as

$$2R - 6\frac{R_{ab}y^{a}y^{b}}{g_{pq}y^{p}y^{q}} = -\kappa \Big(-2\tilde{T} + 12\frac{\tilde{T}^{ab}y_{a}y_{b}}{g_{pq}y^{p}y^{q}}\Big), \quad (56)$$

if evaluated at *g*-timelike observer four-velocities *y*. Now, we take a second derivative with respect to *y*, contract with g^{-1} , reinsert the result, and conclude

$$\left(R_{ab} - \frac{1}{2}g_{ab}R\right)y^a y^b = 2\kappa \tilde{T}_{ab}y^a y^b.$$
(57)

Since there is no y-dependence beyond the explicit one, a second derivative with respect to y yields the Einstein equations, if we choose the gravitational constant $\kappa = \frac{4\pi G}{c^4}$.

The gravity equation on Finsler spacetime including the coupling to matter therefore is

$$-g^{Fab}\bar{\partial}_{a}\bar{\partial}_{b}\mathcal{R} + \frac{6}{F^{2}}\mathcal{R} - 2g^{Fab}(\nabla_{a}S_{b} + S_{a}S_{b} + \bar{\partial}_{a}\nabla S_{b})$$
$$= \frac{4\pi G}{c^{4}}T,$$
(58)

with zero-homogeneous source function T on TM. Observe that this field equation including the matter part is invariant under $L \rightarrow L^k$ by construction of the coupling principle, as the vacuum equation is. This leads to the interesting conclusion that every solution $g_{ab}(x)$ of the Einstein equations induces a family L_k of solutions of the Finsler gravity solution with $L_k = (g_{ab}(x)y^ay^b)^k$.

In order to find further solutions of this highly complicated partial-differential equation, we will study symmetries of Finsler spacetimes in the next section. Then, we present a solution of the linearized Finsler gravity equation in Sec. VI which turns out to be a geometric refinement of the linearized Schwarzschild solution of general relativity.

V. FINSLER SPACETIME SYMMETRIES

In the previous section, we deduced the Finsler gravity field equation including matter sources and showed that it is consistent with the Einstein equations in the metric geometry limit. Our new field equation is a highly complex differential equation; in order to simplify the task of finding analytic solutions, we wish to consider symmetric spacetimes. We begin this section by defining symmetries of Finsler spacetimes and show how this concept is a generalization of the symmetries of Lorentzian spacetimes. We explicitly present the general structure of the fundamental geometry function L for the spherically, cosmologically, and maximally symmetric case. In the next section, we will then use our results about symmetric Finsler spacetimes to solve the linearized Finsler gravity equation.

A. Definition

On a symmetric Lorentzian manifold (M, g), the metric is invariant under certain diffeomorphisms. Similarly, we wish to define symmetries of a Finsler spacetime (M, L, F) as an invariance of the fundamental geometry function L. Consider a diffeomorphism generated by the vector field $X = \xi^a(x)\partial_a$; this acts as a coordinate change on local coordinates on M as $(x^a) \rightarrow (x^a + \xi^a)$ and on the induced coordinates on the tangent bundle TM as $(x^a, y^a) \rightarrow (x^a + \xi^a, y^a + y^q \partial_q \xi^a)$. Hence, the diffeomophism on M induces a diffeomorphism on TM that is generated by the vector field $X^C = \xi^a \partial_a + y^q \partial \xi^a \bar{\partial}_a$, called the complete lift of X. The idea of implementing symmetries via complete lifts in a Finsler geometry setting appears already in Ref. [11]. Here, we want to make this concept precise for Finsler spacetimes:

Definition 4— A symmetry of a Finsler spacetime (M, L, F) is a diffeomorphism generated by a vector field Y over the tangent bundle TM so that Y(L) = 0 and Y is the complete lift X^C of a vector field X over M. A Finsler spacetime is called symmetric if it possesses at least one symmetry.

The following theorem summarizes important properties of Finsler spacetime symmetries. The symmetry generators form a Lie algebra with the commutator of vector fields on TM, and they are isomorphic to a Lie algebra of vector fields on M which becomes the usual symmetry algebra of Lorentzian manifolds in the metric geometry limit. This not only shows that definition 4 of symmetry is consistent with that of Lorentzian spacetimes, but also that the usual Killing vectors, e.g., those for spherical symmetry, can be used to study symmetries of Finsler spacetimes.

Theorem 4— Let S be the set of symmetry-generating vector fields of a Finsler spacetime.

- (i) (S, [·, ·]) is a Lie subalgebra of the set of vector fields over *TM*;
- (ii) (S, [·, ·]) is isomorphic to the Lie subalgebra (π_∗(S), [·, ·]) of the set of vector fields over M;
- (iii) in the metric geometry limit, $(\pi_*(S), [\cdot, \cdot])$ becomes the symmetry algebra of the emerging Lorentzian spacetime.

Proof-

- (i) Let $Y \in S$; then Y(L) = 0 and $(\pi_*Y)^C Y = 0$. Both properties are linear, so that S is a vector subspace of the Lie algebra of all vector fields on TM. It remains to be proven that the commutator of two elements $Y_1, Y_2 \in S$ closes in S. It is clear that $[Y_1, Y_2](L) = 0$; to show that $(\pi_*[Y_1, Y_2])^C = [Y_1, Y_2]$, one uses that $Y_i = X_i^C$ for some vector fields X_i on M and that $[X_1^C, X_2^C] = [X_1, X_2]^C$ (see Ref. [12]).
- (ii) The inverse for π_* on $\pi_*(S)$ is given by the complete lift; hence, S and $\pi_*(S)$ are isomorphic as vector spaces. The Lie algebra structure is preserved in both directions because of $[X_1^C, X_2^C] = [X_1, X_2]^C$, and hence also $\pi_*[Y_1, Y_2] = [\pi_*Y_1, \pi_*Y_2]$.

FINSLER GEOMETRIC EXTENSION OF EINSTEIN GRAVITY

(iii) For $Y = X^C \in S$, we have $\xi^a \partial_a L + y^q \partial_q \xi^a \bar{\partial}_a L = 0$. In the metric geometry limit $L(x, y) = g_{ab}(x)y^a y^b$, and, hence, $y^p y^q (\xi^a \partial_a g_{pq} + g_{ap} \partial_q \xi^a + g_{aq} \partial_p \xi^a) = y^p y^q \mathcal{L}_X g_{pq}(x) = 0$. Since the Lie-derivative of the metric *g* does not depend on the fiber coordinates of the tangent bundle, we conclude $\mathcal{L}_X g_{pq}(x) = 0$. This is the condition that defines *X* as the symmetry generator of a metric spacetime. \Box

We now wish to study the implications of spherical, cosmological, and maximal symmetry for the fundamental function L of a Finsler spacetime.

PHYSICAL REVIEW D 85, 064009 (2012)

B. Spherical symmetry

Consider a Finsler spacetime (M, L, F) and coordinates $(t, r, \theta, \phi, y^t, y^r, y^{\theta}, y^{\phi})$ on its tangent bundle. Spherical symmetry is defined by the following three vector fields, which generate spatial rotations and form the algebra $\mathfrak{so}(3)$,

$$X_{4} = \sin\phi \partial_{\theta} + \cot\theta \cos\phi \partial_{\phi},$$

$$X_{5} = -\cos\phi \partial_{\theta} + \cot\theta \sin\phi \partial_{\phi},$$

$$X_{6} = \partial_{\phi}.$$
(59)

Their complete lifts are obtained via the procedure described in the previous section:

$$K_4^C = \sin\phi\partial_\theta + \cot\theta\cos\phi\partial_\phi + y^\phi\cos\phi\bar{\partial}_\theta - \left(y^\theta\frac{\cos\phi}{\sin^2\theta} + y^\phi\cot\theta\sin\phi\right)\bar{\partial}_\phi,\tag{60a}$$

$$X_5^C = -\cos\phi\partial_\theta + \cot\theta\sin\phi\partial_\phi + y^\phi\sin\phi\bar{\partial}_\theta - \left(y^\theta\frac{\sin\phi}{\sin^2\theta} - y^\phi\cot\theta\cos\phi\right)\bar{\partial}_\phi,\tag{60b}$$

$$X_6^C = \partial_{\phi}. \tag{60c}$$

Applying the symmetry condition $X_6^C(L) = 0$ implies $\partial_{\phi}L = 0$, while using $X_4^C(L) = 0$ and $X_5^C(L) = 0$ to deduce $(\sin\phi X_4^C - \cos\phi X_5^C)(L) = 0$ and $(\cos\phi X_4^C + \sin\phi X_5^C)(L) = 0$ yields

$$\partial_{\theta}L = y^{\phi} \cot\theta \bar{\partial}_{\phi}L, \qquad y^{\phi} \sin^2\theta \bar{\partial}_{\theta}L = y^{\theta} \bar{\partial}_{\phi}L. \tag{61}$$

In order to analyze the implications of these equations on L, we introduce new coordinates

$$u(\theta) = \theta, \qquad v(y^{\theta}) = y^{\theta}, \qquad w(\theta, y^{\theta}, y^{\phi})^2 = (y^{\theta})^2 + \sin^2\theta(y^{\phi})^2, \tag{62}$$

while keeping (t, y^t, r, y^r, ϕ) . The associated transformation of the derivatives

$$\partial_t = \partial_t, \qquad \partial_r = \partial_r, \qquad \partial_\theta = \frac{w^2 - v^2}{w} \cot u \partial_w + \partial_u, \qquad \partial_\phi = \partial_\phi,$$
 (63a)

$$\bar{\partial}_t = \bar{\partial}_t, \qquad \bar{\partial}_r = \bar{\partial}_r, \qquad \bar{\partial}_\theta = \frac{v}{w} \partial_w + \partial_v, \qquad \bar{\partial}_\phi = \sin u \frac{\sqrt{(w^2 - v^2)}}{w} \partial_w,$$
(63b)

makes the equations (61) equivalent to the simple constraints $\partial_u L = 0$ and $\partial_v L = 0$.

Hence, we conclude from the analysis of the symmetry conditions $X_i^C(L) = 0$ that the most general spherically symmetric Finsler spacetime is described by a fundamental function which is *n*-homogeneous in (y^t, y^r, w) and of the form

$$L(t, r, \theta, \phi, y^t, y^r, y^\theta, y^\phi) = L(t, r, y^t, y^r, w(\theta, y^\theta, y^\phi)),$$
(64)

where $w(\theta, y^{\theta}, y^{\phi})$ is defined in Eq. (62).

C. Cosmological and maximal symmetry

After our discussion of the spherically symmetric case in full detail above, we will now present the results of a similar analysis first for cosmologically and second for maximally symmetric Finsler spacetimes.

Cosmological symmetry describes an isotropic and homogeneous spacetime. This is a much more symmetric situation than in the spherically symmetric scenario and is implemented by requiring the following six vector fields to be symmetry generators, see Ref. [13]:

$$X_1 = \chi \sin\theta \cos\phi \partial_r + \frac{\chi}{r} \cos\theta \cos\phi \partial_\theta - \frac{\chi}{r} \frac{\sin\phi}{\sin\theta} \partial_\phi,$$
(65a)

$$X_2 = \chi \sin\theta \sin\phi \partial_r + \frac{\chi}{r} \cos\theta \sin\phi \partial_\theta + \frac{\chi}{r} \frac{\cos\phi}{\sin\theta} \partial_\phi,$$
(65b)

$$X_3 = \chi \cos\theta \partial_r - \frac{\chi}{r} \sin\theta \partial_\theta, \tag{65c}$$

$$X_4 = \sin\phi \partial_\theta + \cot\theta \cos\phi \partial_\phi, \qquad X_5 = -\cos\phi \partial_\theta + \cot\theta \sin\phi \partial_\phi, \qquad X_6 = \partial_\phi, \tag{65d}$$

where we write $\chi = \sqrt{1 - kr^2}$ and k is constant. The complete lifts of these vector fields are listed in Appendix A 3. Applying the symmetry conditions $X_i^C(L) = 0$ to the fundamental function L and introducing the new coordinates

$$q(r) = r, \quad s(y^r) = y^r, \quad u(\theta) = \theta, \quad v(y^\theta) = y^\theta,$$
(66a)

$$w_C(r,\theta,y^r,y^\theta,y^\phi)^2 = \frac{(y^r)^2}{1-kr^2} + r^2((y^\theta)^2 + \sin^2\theta(y^\phi)^2),$$
(66b)

while keeping (t, y^t) , yields the following result: the cosmological fundamental function L is *n*-homogeneous in (y^t, w_c) and has the form

$$L(t, r, \theta, \phi, y^t, y^r, y^\theta, y^\phi) = L(t, y^t, w_C(r, \theta, y^r, y^\theta, y^\phi)).$$
(67)

The constant k only appears in the expression for the coordinate w_C . The value of w_C can be understood as the metric length measure on a three-dimensional manifold of constant curvature k. The same metric appears in the spatial part of the standard Robertson-Walker metric.

For the study of maximally symmetric Finsler spacetimes, we use some notation from Ref. [14], where such spacetimes are constructed from embeddings into a fivedimensional manifold. Symmetry vectors generating maximal symmetry are given by

$$X_{\alpha} = C(x)\alpha^{c}\partial_{c}, \qquad X_{\Omega} = \Omega_{b}^{a}x^{b}\partial_{a}, \qquad (68)$$

with $C(x) = \sqrt{1 - KC_{pq}x^p x^q}$, constant *K*, and constant 4×4 matrices C_{ab} and $\Omega^a{}_b$. There are four linearly independent vector fields X_α and six X_Ω by requiring the condition $\Omega^q{}_bC_{qa} = -\Omega^q{}_aC_{qb}$; their complete lifts are

$$X_{\alpha}^{C} = C(x)\alpha^{c}\partial_{c} - y^{b}\frac{KC_{bm}x^{m}}{C(x)}\alpha^{c}\bar{\partial}_{c},$$

$$X_{\Omega}^{C} = \Omega^{a}{}_{b}x^{b}\partial_{a} + y^{b}\Omega^{a}{}_{b}\bar{\partial}_{a}.$$
(69)

Evaluating the symmetry conditions $X_{\alpha}^{C}(L) = 0$ and $X_{\Omega}^{C}(L) = 0$ on the fundamental function and introducing new coordinates

$$u^{a}(x) = x^{a},$$

$$v^{\gamma}(y) = y^{\gamma},$$

$$w_{M}(x, y)^{2} = C_{ab}y^{a}y^{b} + \frac{K}{C(x)^{2}}C_{ap}x^{p}y^{a}C_{bq}x^{q}y^{b}$$

$$= g_{ab}(x)y^{a}y^{b},$$
(70)

where γ runs over any three indices in {0, 1, 2, 3}, yields the following result: the maximally symmetric fundamental function *L* is *n*-homogeneous in w_M and of the form

$$L(x, y) = L(w_M(x, y)) = Aw_M(x, y)^n.$$
 (71)

The final equality is obtained from Euler's theorem for homogeneous functions.

Observe that the maximally symmetric fundamental function always describes a metric geometry, see Eq. (70). Hence, all maximally symmetric Finsler spacetimes are Lorentzian spacetimes, and the gravity equation (58) is equivalent to Einstein's equations. Thus, we can immediately conclude that the only maximally symmetric, source-free vacuum solution of our Finsler gravity equation is the Minkowski-metric-induced fundamental function $L = \eta_{ab}y^ay^b$ and its powers. In the expression for w_M above, this corresponds to $C_{ab} = \eta_{ab}$ and K = 0. Maximally symmetric spacetimes with $K \neq 0$ can only be obtained as solutions of the Finsler gravity equation by adding a cosmological constant term, similarly as in general relativity.

VI. LOWEST-ORDER EFFECTS IN THE SOLAR SYSTEM

In this section, we will study Finsler spacetimes that describe mild deviations from Lorentzian geometry. In this situation, the complicated Finsler gravity field equation allows a simplified treatment. After a general discussion of the linearized field equation, we will employ what we learned about spacetime symmetries to present a spherically symmetric solution. This particular model turns out to be a refinement of the linearized Schwarzschild solution of general relativity, and we will argue that it should be capable of modelling unexplained effects in the Solar System like the fly-by anomaly.

A. Finsler modifications of Lorentzian geometry

Recall that the fundamental functions $L = L_0$ and $L = (L_0)^k$ define the same geometry, and that this is respected by the Finsler gravity field equation. Hence, the following class of fundamental functions gives us good control over deviations from Lorentzian metric geometry:

$$L = (g_{ab}(x)y^a y^b)^k + h(x, y) = G(x, y)^k + h(x, y).$$
(72)

Here, h(x, y) is a 2*k*-homogeneous function that causes the Finsler modifications of the Lorentzian metric spacetime (M, g). The abbreviation G(x, y) should not be confused with the Sasaki-type metric on *TM*.

Recall the Finsler gravity vacuum field equation from Eq. (37):

$$g^{Fab}\bar{\partial}_a\bar{\partial}_b\mathcal{R} - \frac{6}{F^2}\mathcal{R} + 2g^{Fab}(\nabla_a S_b + S_a S_b + \bar{\partial}_a \nabla S_b) = 0.$$
(73)

We will now expand this equation to linear order in the modification h(x, y), where $G(x, y) \neq 0$. In the following calculations, we suppress all higher-order terms. We

introduce $l = \frac{G^{1-k}}{k}h$ and $l_{ab} = \frac{1}{2}\bar{\partial}_a\bar{\partial}_b l$ to expand the Finsler function and Finsler metric as

$$F^{2} \simeq \frac{G}{|G|}(G+l), \qquad g_{ab}^{F} \simeq \frac{G}{|G|}(g_{ab}+l_{ab}).$$
 (74)

The coefficients of the nonlinear connection are calculated from Eq. (7):

$$N^{a}_{\ b} \simeq y^{m} \Gamma^{a}_{\ bm} - \frac{1}{2} g_{cq} \Gamma^{c}_{\ mn} y^{m} y^{n} \bar{\partial}_{b} l^{aq} - l^{aq} g_{cq} \Gamma^{c}_{\ bm} y^{m} + \frac{1}{2} g^{aq} (\partial_{b} l_{qm} + \partial_{m} l_{qb} - \partial_{q} l_{bm}) y^{m}.$$
(75)

Here, the $\Gamma^a{}_{bc}$ are the Christoffel symbols of the metric g, and in the following, ∇ acts as the Levi-Civita connection. Note that the zeroth-order term, for $l \rightarrow 0$, is the metric linear connection. The curvature and the tensor S can be expressed with the help of the short-hand notation

$$T^{a}_{\ bc} = \frac{1}{2}g^{aq}(\nabla_{b}l_{qc} + \nabla_{c}l_{qb} - \nabla_{q}l_{bc})$$
(76)

as

$$\mathcal{R} = R^{a}{}_{ab}y^{b} \simeq -y^{a}y^{b}R_{ab}[g] - \nabla_{a}(y^{b}y^{c}T^{a}{}_{bc}) + \nabla(y^{c}T^{a}{}_{ac}),$$
(77a)

$$S_a = \Gamma^{\delta p}{}_{pa} - \bar{\partial}_p N^p{}_a \simeq -y^q \bar{\partial}_a T^p{}_{pq}.$$
(77b)

The zeroth-order term in \mathcal{R} is determined by the Ricci tensor of g, while $S_a \rightarrow 0$. Collecting all terms in the gravitational field, Eq. (73) finally yields

$$0 \simeq -2 \frac{G}{|G|} R[g] + \frac{6}{G} y^a y^b R_{ab}[g] + \left[2 \frac{G}{|G|} l^{ab} R_{ab}[g] \right] + \frac{6G - 2l}{G^2} y^a y^b R_{ab}[g] + g^{ab} \bar{\partial}_a \bar{\partial}_b (-\nabla_a (y^b y^c T^a{}_{bc}) + \nabla (y^c T^a{}_{ac})) - \frac{6}{G} (-\nabla_a (y^b y^c T^a{}_{bc}) + \nabla (y^c T^a{}_{ac})) - 2g^{ab} (\nabla_a y^q \bar{\partial}_b T^p{}_{pq} + \bar{\partial}_a \nabla y^q \bar{\partial}_b T^p{}_{pq}) \right].$$
(78)

The zeroth-order contribution in the first line is equivalent to the Einstein vacuum equations, as discussed in Sec. IVA. The first-order terms in square brackets determine the Finsler modification of the unperturbed metric background solution. The details of how to rewrite the different terms of this equation in terms of the perturbation h in the fundamental function L, see Eq. (72), instead of l, can be found in Appendix A 4.

B. Refinements to the linearized Schwarzschild solution

We will now use our results on symmetries and on the linearization of vacuum Finsler gravity around metric spacetimes to derive a particular model that refines the linearized Schwarzschild solution and can be used to study Solar System physics.

Recall from Sec. V B that the dependence of the general spherically symmetric fundamental function in tangent bundle coordinates induced by (t, r, θ, ϕ) is restricted to $L(t, r, y^t, y^r, w(\theta, y^{\theta}, y^{\phi}))$, where $w^2 = (y^{\theta})^2 + \sin^2\theta(y^{\phi})^2$. We wish to study such a spherically symmetric fundamental function that describes a Finsler modification of Lorentzian geometry. For simplicity, we consider a bimetric four-homogeneous Finsler spacetime that perturbs the maximally symmetric vacuum solution of Finsler gravity which is given by Minkowski spacetime. We assume $L = (\eta_{ab}y^ay^b)^2 + \eta_{ab}y^ay^bh_{cd}y^cy^d = (\eta_{ab}y^ay^b)(\eta_{cd} + h_{cd})y^cy^d$ with $h_{ab} = \text{diag}(a(r), b(r), c(r)r^2, c(r)r^2\sin^2\theta)$. This ansatz has the explicit form

$$L(r, y^{t}, y^{r}, w) = (-y^{t^{2}} + y^{r^{2}} + r^{2}w^{2})([-1 + a(r)]y^{t^{2}} + [1 + b(r)]y^{r^{2}} + [1 + c(r)]r^{2}w^{2}).$$
 (79)

Observe that the function c(r) cannot be transformed away by defining a new radial coordinate. Although this could remove c(r) from the metric in the right-hand bracket, such a coordinate change would generate extra terms in the metric appearing in the left-hand bracket. Therefore, the existence of the function c(r) as a physical degree of freedom is a Finsler geometric effect that appears as a consequence of the bimetric spacetime structure assumed here.

We will now solve the linearized Finsler gravity equation (78) for a(r), b(r), and c(r) with the ansatz (79). Sorting the equation with respect to powers in y^t , y^r , and w gives rise to three equations that have to be satisfied:

$$-2a' - ra'' = 0,$$

$$ra'' + 2b' - 4c' - 2rc'' = 0,$$
 (80)

$$ra'' + 2b + rb' - 2c - 4rc' - r^2c'' = 0.$$

The solution of these equations is

ra

$$a(r) = -\frac{A_1}{r} + A_2,$$

$$b(r) = -\frac{A_1}{r} + \frac{A_3}{r^2},$$

$$c(r) = \frac{A_4}{r} - \frac{A_3}{r^2}.$$

(81)

We will now study the properties of this specific firstorder Finsler spacetime solution and compare it to the linearized Schwarzschild spacetime. We use the linearized expression for the nonlinear connection coefficients in Eq. (75) to analyze the Finsler geodesic equation that is derived by extremizing the proper time integral (1). For a curve with coordinates $x(\tau)$, this has the form $\ddot{x}^a + N^a{}_b(x, \dot{x})\dot{x}^b = 0$. As usual in spherical symmetry, setting $\theta = \frac{\pi}{2}$ solves one of the four component equations immediately; the remaining equations are CHRISTIAN PFEIFER AND MATTIAS N.R. WOHLFARTH

$$0 = \ddot{r} - \frac{1}{2} \frac{A_1}{r^2} \dot{t} \dot{r}, \qquad (82a)$$

$$0 = \ddot{r} - \frac{1}{4} \frac{A_1}{r^2} \dot{t}^2 + \frac{1}{4} \left(\frac{A_1}{r^2} - 2 \frac{A_3}{r^3} \right) \dot{r}^2 + \left(-r - \frac{A_1}{2} - \frac{A_4}{4} + \frac{1}{2} \frac{A_3}{r} \right) \dot{\phi}^2, \qquad (82b)$$

$$0 = \ddot{\phi} + \frac{2}{r} \left(1 - \frac{1}{4} \frac{A_4}{r} + \frac{1}{2} \frac{A_3}{r^2} \right) \dot{\phi} \dot{r}.$$
 (82c)

From these equations, we find two constants of motion

$$E = i \left(1 + \frac{1}{2} \frac{A_1}{r} \right), \qquad \ell = r^2 \left(1 + \frac{1}{2} \frac{A_4}{r} - \frac{1}{2} \frac{A_3}{r^2} \right) \dot{\phi}.$$
(83)

These can be used to deduce the orbit equation from the affine normalization condition that $F(x, \dot{x}) = 1$ along the Finsler geodesic; we employ (74) and write σ for the sign of $\eta_{ab}\dot{x}^a\dot{x}^b = -\dot{t}^2 + \dot{r}^2 + r^2\dot{\phi}^2$ to obtain

$$\frac{1}{2}\dot{r}^{2} = \frac{E^{2}}{2}\left(1 - \frac{A_{2}}{2}\right) + \frac{1}{2}\sigma\left(1 + \frac{A_{1}}{2r}\right) - \frac{\ell^{2}}{2r^{2}}\left(1 + \frac{A_{1}}{2r} - \frac{A_{4}}{2r}\right) + \frac{A_{3}}{4r^{2}}(\sigma - E^{2}).$$
 (84)

The geodesic equations, the constants of motion, and the orbit equation are well-suited to compare the bimetric linearized Finsler solution with the linearized Schwarzschild solution. To see the differences to this solution of Einstein gravity, we first note that A_2 can be absorbed into a redefinition of E, and hence can be assumed to be zero. Second, we introduce the Schwarzschild radius r_0 to redefine $A_1 = -2r_0(1 + a_1)$, $A_3 = 2\ell^2 a_3/(E^2 - \sigma)$ and $A_4 = 2r_0a_4$ in terms of dimensionless small constants a_1 , a_3 , and a_4 . Then, the orbit equation becomes

$$\frac{1}{2}\dot{r}^2 = \frac{E^2}{2} + \frac{\sigma}{2} - \frac{\sigma r_0}{2r}(1+a_1) - \frac{\ell^2}{2r^2}(1+a_3) + \frac{r_0\ell^2}{2r^3}(1+a_1+a_4).$$
(85)

In the special case $a_1 = a_3 = a_4 = 0$, this is precisely the orbit equation in the linearized Schwarzschild geometry, see Ref. [15]; the same limit also applies to the geodesic equations and the constants of motion.

The Finsler geometric refinements of the metric Schwarzschild geometry are encoded in the constants a_1 , a_3 , and a_4 . These can in principle be fitted to data from Solar System experiments. Indeed, there are certain observations that cannot be fully explained by the Schwarzschild solution [9], for instance, the fly-by anomaly: for several spacecrafts, it has been reported that swing-by maneuvers lead to a small unexplained velocity increase. This corresponds to a change in the shape of the orbit of the spacecraft. Such a change can in principle be modeled by Finsler refinements; the perturbations a_1 , a_3 , and a_4 certainly



FIG. 4. Numerical fly-by solutions of the geodesic equations for linearized Schwarzschild geometry (dashed line) and the bimetric Finsler refinement (solid line) with $a_1 = 0$, $a_3 \approx 0.156$, and $a_4 = 0.1$. The mass is centered at the origin and has Schwarzschild radius $r_0 = 0.1$. The initial conditions are r(0) = 0.5, $\dot{r}(0) = 0.02$, $\phi(0) = 0$, $\dot{\phi}(0) = 1.1$, and t(0) = 0 for both curves, and $\dot{t}(0)$ is calculated from the respective unit normalization condition $F(x, \dot{x}) = 1$.

provide possibilities to alter the wideness of the swing-by orbit as compared to that expected from Einstein gravity. This can be confirmed by simple numerical calculations, see Fig. 4.

We have seen that Finsler geometries exist that are extremely close to metric geometries. Our specific example of a spherically symmetric bimetric perturbation around Minkowski spacetime could be reinterpreted as a geometry close to the linearized Schwarzschild solution of Einstein gravity. The more complex causal structure, however, leads to additional constants that modify the geodesic equations and, in particular, the shape of test particle orbits. This could be a means to explain the fly-by anomaly in the Solar System. We emphasize that this consequence already at first-order perturbation theory gives a glimpse on the potential of Finsler gravity.

VII. DISCUSSION

Finsler geometry is fundamentally based on the reparamtrization invariant length integral (1). In physics, this integral can be used as a very general clock postulate on the one hand, and on the other as an action for massive point particles which automatically guarantees the very precisely tested weak equivalence principle. In previous work [5], we formulated a set of minimal requirements for the application of Finsler geometry to the description of spacetime. This led us to definition 1 of Finsler spacetimes (M, L, F), which have sufficient structure to provide notions of causality and are mathematically controlled generalizations of Lorentzian geometry.

In this article, we constructed an action for Finsler gravity from first principles. Our theory of Finsler gravity (51) fully includes the description of matter fields which

are coupled to Finsler spacetime by a lifting principle that generates the appropriate action from the standard Lagrangian on Lorentzian spacetime. We derived the gravitational field equation by variation with respect to the fundamental geometry function L and could show that it consistently becomes equivalent to the Einstein field equations in the metric geometry limit. Hence, Einstein gravity can be seen as special case of our gravity theory based on Finsler geometry. We presented the geometric definition 2 of observers on Finsler spacetimes along with a clear interpretation of how they measure physical fields. By theorem 2, we were able to characterize the class of transformations that relates two different observers at the same point of the spacetime manifold; these transformations have the algebraic structure of a groupoid as proven in theorem 3. Any observer transformation can be understood as the composition of a usual Lorentz transformation and an identification of the two observers' four-velocities by a geometrically well-defined parallel transport. For the limiting case of metric geometries, this parallel transport trivializes so that the transformation groupoid becomes equivalent to the standard Lorentz group. In this sense, Finsler spacetimes generalize Lorentz invariance instead of violating it.

As a further formal development, we presented definition 4 of symmetries of Finsler spacetimes and theorem 4 that shows some of their basic properties. We applied this notion to determine the most general fundamental geometry functions consistent with spherical, cosmological, or maximal symmetry. Maximally symmetric Finsler spacetimes are, in fact, maximally symmetric metric spacetimes, and the only maximally symmetric source-free vacuum solution of Finsler gravity is Minkowski spacetime. As a concrete application of the results of this article, we studied a simple spherically symmetric bimetric perturbation around this flat metric vacuum. We found a first-order solution of the Finsler gravity equation, which is a refinement of the linearized Schwarzschild solution of Einstein gravity. With a special choice of the parameters in our solution, the resulting geodesics are identical to those of linearized Schwarzschild spacetime. This demonstrates that weak-field gravitational experiments may not be sufficient to distinguish Finsler spacetimes from Lorentzian metric spacetimes. But we saw that the full set of parameters in our model solution could be capable to resolve the fly-by anomaly in the Solar System.

This is very promising and creates a strong motivation for more intensive studies of our new theory of gravity. It is natural to ask whether the additional degrees of freedom of Finsler spacetime solutions as compared to metric spacetimes may lead to new insights on the dark matter distributions in galaxies or dark energy in the Universe. These could be effects of a fundamentally more complex spacetime geometry instead of being particle physics phenomena. It will be possible to study these questions once solutions of Finsler gravity, especially for spherical symmetry and cosmology, become available. For cosmology, it will also be necessary to study perfect fluid sources for Finsler spacetimes.

We saw in Ref. [8] that Finsler spacetimes can provide a geometric explanation of the OPERA measurements of superluminal neutrinos [16]. The velocity difference between the neutrinos and the speed of light recognized as the boundary velocity of observers depends not only on the energy and mass of the neutrino, but also on the underlying spacetime geometry. So also in this context, solutions for spherical symmetry and cosmology are needed in order to understand the size of the effect for the different observed neutrino sources.

Further important topics for future research are the coupling of spinor fields, the analysis of field theories on the generalized causal structure of Finsler spacetimes, and their quantization.

ACKNOWLEDGMENTS

C. P. and M. N. R. W. thank Niklas Hübel, Claudio Dappiaggi, Andreas Degner, Klaus Fredenhagen, Manuel Hohmann, Matthias Lange, Falk Lindner, Gunnar Preuss and Felix Tennie for inspiring discussions. They acknowledge full financial support from the German Research Foundation DFG under Grant No. WO 1447/1-1.

APPENDIX A: TECHNICAL DETAILS

This appendix presents technical details for several derivations in the main text. In particular, we show how to perform the variation of the Finsler gravity and matter field actions on the unit tangent bundle; for completeness, we state the complete lifts of the cosmological symmetry generators; and we display some additional material on the linearized Finsler gravity equations.

1. Variation of the gravity action

The Finsler gravitational field equation presented in Sec. IVA can be deduced from our new Finsler gravity action as follows. Before we consider the variation of the matter part with respect to the fundamental geometry function L in the next section, we here take a look at the pure gravity action (30):

$$S_G[L] = \int d^4 \hat{x} d^3 u [\sqrt{g^F h^F} R^a{}_{ab} y^b]_{|\Sigma}.$$
(A1)

The integrand is homogeneous of degree five; to obtain the first intermediate step (32) of the variation, we use the facts that for f(x, y) homogeneous of degree k holds $f(x, y)|_{\Sigma} = \frac{f(x,y)}{F(x,y)^k}$ and that $\delta_L(f(x, y)|_{\Sigma}) = (\delta_L f(x, y))|_{\Sigma} - \frac{k}{n} f(x, y)|_{\Sigma} \frac{\delta L}{L}$.

The second step (33) is obtained by using the coordinate transformation formulae (16) and the fact that $\delta(\partial_{\alpha} y^{a}) = -y^{a} \partial_{\alpha} \left(\frac{\delta L}{nL}\right)$ to calculate

CHRISTIAN PFEIFER AND MATTIAS N.R. WOHLFARTH

$$h^{F\alpha\beta}\delta h^{F}{}_{\alpha\beta} = (g^{Fab}\bar{\partial}_{a}u^{\alpha}\bar{\partial}_{b}u^{\beta})(\partial_{\alpha}y^{c}\partial_{\beta}y^{d}\delta g^{F}{}_{cd} + 2\partial_{\alpha}\delta y^{c}\partial_{\beta}y^{d}g^{F}{}_{cd})$$
$$= g^{Fab}\delta g^{F}{}_{ab} - \frac{2}{n}\frac{\delta L}{L}, \qquad (A2)$$

which in turn is used to deduce

$$\delta(\sqrt{g^{F}h^{F}}R^{a}{}_{ab}y^{b}) = \sqrt{g^{F}h^{F}} \left(\left[g^{Fab} \delta g^{F}{}_{ab} - \frac{1}{n} \frac{\delta L}{L} \right] \times R^{a}{}_{ab}y^{b} + \delta R^{a}{}_{ab}y^{b} \right).$$
(A3)

The formulae (34) used in the third step of the variation are basically obtained by means of integration by parts (20). For a function f(x, y) that is *k*-homogeneous in *y*, the following holds:

$$\int d^{4}\hat{x}d^{3}u(\sqrt{g^{F}h^{F}}g^{Fab}\delta g^{F}_{ab}f)|_{\Sigma}$$

$$= \int d^{4}\hat{x}d^{3}u\left(\sqrt{g^{F}h^{F}}(f(k+4)(2-k) + F^{2}g^{Fab}\bar{\partial}_{a}\bar{\partial}_{b}f)\frac{\delta L}{nL}\right)_{|\Sigma};$$
(A4)

choosing $f = R^a{}_{ab}y^b$, which has k = 2, proves formula (34a). To show Eq. (34b), we first write $S^a{}_{bc} = -y^q \bar{\partial}_b \Gamma^{\delta a}{}_{ac}$ and use

$$\delta R^a{}_{bc} = -2y^d \nabla_{[b} \delta \Gamma^{\delta a}{}_{c]d} + 2y^p S^a{}_{q[b} \Gamma^{\delta q}{}_{c]p}$$
(A5)

to equate

$$\begin{split} \delta R^{a}{}_{ab}y^{b} &= -2y^{b}y^{q} (\nabla_{[a} \delta \Gamma^{\delta a}{}_{b]q} - \frac{1}{2}S_{c} \delta \Gamma^{\delta c}{}_{bq}) \\ &= -\nabla_{a} (y^{b}y^{q} \delta \Gamma^{\delta a}{}_{bq}) + y^{b} \nabla_{b} (y^{q} \delta \Gamma^{\delta a}{}_{aq}) \\ &+ S_{c} \delta \Gamma^{\delta c}{}_{bq} y^{b} y^{q}. \end{split}$$
(A6)

The integration by parts formulae (20) and

$$y^{b}y^{q}\delta\Gamma^{\delta a}{}_{bq} = \frac{1}{2}g^{Lap}(y^{b}\nabla_{b}\bar{\partial}_{p}\delta L - \nabla_{p}\delta L)$$
$$= \frac{|L|^{2/n}}{nL}g^{Fab}(y^{b}\nabla_{b}\bar{\partial}_{p}\delta L - \nabla_{p}\delta L)$$
$$+ \frac{(2-n)}{nL}y^{a}y^{b}\nabla_{b}\delta L \qquad (A7)$$

then yield the desired equation

$$\int d^{4}\hat{x}d^{3}u(\sqrt{g^{F}h^{F}}\delta R^{a}{}_{ab}y^{b})|_{\Sigma}$$

$$= \int d^{4}\hat{x}d^{3}u(\sqrt{g^{F}h^{F}}2S_{c}\delta\Gamma^{\delta c}{}_{bq}y^{b}y^{q})|_{\Sigma}$$

$$= \int d^{4}\hat{x}d^{3}u\left(\sqrt{g^{F}h^{F}}2F^{2}g^{F}_{ab}(\nabla_{a}S_{b}+S_{a}S_{b}+\delta_{a}\nabla S_{b})\frac{\delta L}{nL}\right)|_{\Sigma}.$$
(A8)

Combining these three steps as we did in Sec. IVA finally produces the Finsler gravity vacuum field equation (37).

2. Variation of the matter action

In Sec. IV B, we presented a coupling principle of matter fields to Finsler gravity. The crucial steps of the derivation of the constraints (42), equations of motion (43) and (44), and of the metric limit of the complete gravity equation (54), including the matter source terms, shall be presented here. Recall the matter action for a *p*-form field $\Phi(x, y)$ on Finsler spacetime arises from a lift of the standard *p*-form action on Lorentzian spacetime as

$$S_m[L, \Phi, \lambda] = \int_{\Sigma} d^4x d^3u [\sqrt{g^F h^F} (\mathcal{L}(G, \Phi, d\Phi) + \lambda(1 - P^H)\Phi)]_{|\Sigma}.$$
 (A9)

In order to perform the variation, we consider all objects in the horizontal/vertical basis of *TTM* where *G* is diagonal, see Eq. (13). In the following, the M, N, \ldots label both horizontal and vertical indices, \bar{a}, \bar{b}, \ldots label vertical indices, and $a, b \ldots$ label horizontal indices. Then,

$$\mathcal{L}(G, \Phi, d\Phi) + \lambda(1 - P^{H})\Phi$$

$$= \mathcal{L}(G_{MN}, \Phi_{M_{1}\dots M_{p}}, d\Phi_{AM_{1}\dots M_{p}}) + \lambda^{\bar{a}_{1}M_{2}\dots M_{p}}\Phi_{\bar{a}_{1}M_{2}\dots M_{p}},$$
(A10)

and the variation of this Lagrangian can now be written as follows:

$$\delta(\mathcal{L} + \lambda(1 - P^{H})\Phi)$$

$$= \frac{\partial \mathcal{L}}{\partial G_{MN}} \delta G_{MN} + \frac{\partial \mathcal{L}}{\partial \Phi_{M_{1}...M_{p}}} \delta \Phi_{M_{1}M_{p}}$$

$$+ \frac{\partial \mathcal{L}}{\partial (d\Phi_{NM_{1}...M_{p}})} \delta (d\Phi_{NM_{1}M_{p}})$$

$$+ \lambda^{\bar{a}_{1}M_{2}...M_{p}} \delta \Phi_{\bar{a}_{1}M_{2}...M_{p}} + \delta \lambda^{\bar{a}_{1}M_{2}...M_{p}} \Phi_{\bar{a}_{1}M_{2}...M_{p}}.$$
(A11)

We can immediately read off the variation with respect to the Lagrange multiplier components which produces Eq. (42). Hence, the Lagrange multiplier λ sets to zero all components of Φ with at least one vertical index, so that only purely horizontal components remain on-shell.

The expansion of $d\Phi$ in components with respect to the horizontal/vertical basis yields

$$d \Phi_{NM_1...M_p} = (p+1)D_{[N} \Phi_{M_1...M_p]} - \frac{p(p+1)}{2} \gamma^Q_{[NM_1} \Phi_{|Q|M_2...M_p]}, \quad (A12)$$

where we write $D_M = \delta^a_M \delta_a + \delta^{\bar{a}}_M \bar{\partial}_a$, and γ^Q_{MN} denote the commutator coefficients of the horizontal/vertical basis. Their only nonvanishing components are given by $\gamma^{\bar{a}}_{bc} = [\delta_b, \delta_c]^{\bar{a}} = R^{\bar{a}}_{bc}$ and $\gamma^{\bar{a}}_{\bar{b}c} = [\bar{\partial}_b, \delta_c]^{\bar{a}} = \bar{\partial}_b N^{\bar{a}}_c$. One now uses the integration by parts formulae (20) to obtain the variation of the matter action with respect to Φ ; this produces the equations of motion (43) and (44).

Finally, the source term for the gravity field equation is obtained by variation of the matter action S_m in Eq. (A9) with respect to the fundamental geometry function L. This not only includes the variation (A11) but also that of the volume element which can be read off from Eq. (A3). We will now show that the metric limit of Finsler gravity plus matter is consistent; this can be done on-shell where we may use the Lagrange multiplier constraints to set all explicitly appearing $\Phi_{\bar{a}_1M_2...M_k}$ to zero. Then the variation of S_m with respect to L becomes

$$\delta S_m[L, \Phi] = \int_{\Sigma} d^4 x d^3 u \sqrt{g^F h^F} \Big[(g^{Fab} \bar{\partial}_a \bar{\partial}_b \pounds + 4\pounds) \frac{\delta L}{nL} \\ + \frac{\partial \pounds}{\partial G_{MN}} \delta G_{MN} + \frac{\partial \pounds}{\partial (d\Phi_{NM_1...M_k})} \\ \times \delta (d\Phi_{NM_1M_k}) \Big]_{|\Sigma} \\ = \int_{\Sigma} d^4 x d^3 u \sqrt{g^F h^F} \Big[(g^{Fab} \bar{\partial}_a \bar{\partial}_b \pounds + 4\pounds) \frac{\delta L}{nL} \\ + \frac{\partial \pounds}{\partial g^F_{ab}} \delta g^F_{ab} + \frac{\partial \pounds}{\partial g^F_{\bar{a}\bar{b}}} \delta \Big(\frac{g^F_{\bar{a}\bar{b}}}{F^2} \Big) \\ + \frac{\partial \pounds}{\partial (d\Phi_{ba_1...a_k})} \delta (\delta_{[b} \Phi_{a_1...a_k]}) \Big]_{|\Sigma}.$$
(A13)

In order to determine the energy-momentum scalar $T_{|\Sigma}$ defined in (53) on a generic Finsler spacetime, one has to calculate all terms in the expression above carefully. However, in the metric geometry limit, the last two terms vanish. Indeed, $\frac{\partial f}{\partial g_{ab}^T}$ is always composed from terms with vertical indices that must be either of the type $\bar{\partial}\Phi$ or contain components of Φ with at least one vertical index; the last term is proportional to $\delta N \bar{\partial} \Phi$; in the metric limit $\bar{\partial}\Phi$ vanishes; and the vertical index components of Φ are zero on-shell. Therefore, the remaining terms that are relevant in the metric limit are

$$\delta S_m[L, \Phi] \to \int_{\Sigma} d^4 x d^3 u \sqrt{g^F h^F} {}_{|\Sigma} \left[(g^{Fab} \bar{\partial}_a \bar{\partial}_b \mathcal{L} + 4\mathcal{L}) \right] \\ \times \frac{\delta L}{nL} + \frac{\partial \mathcal{L}}{\partial g^F_{ab}} \delta g^F_{ab} \right]_{|\Sigma}.$$
(A14)

The rewriting $\delta g^F_{ab} = \frac{1}{2} \bar{\partial}_a \bar{\partial}_b \delta F^2$ and subsequent integration by parts yields

$$\int_{\Sigma} d^{4}x d^{3}u \sqrt{g^{F} h^{F}}_{|\Sigma} \left[\frac{\partial \mathcal{L}}{\partial g^{F}_{ab}} \delta g^{F}_{ab} \right]_{|\Sigma}$$

$$= \int_{\Sigma} d^{4}x d^{3}u \sqrt{g^{F} h^{F}}_{|\Sigma} \left[-\bar{\partial}_{c} K^{c} + (-g^{Fij}\bar{\partial}_{c} g^{F}_{ij} + 4g^{F}_{ic} y^{i}) K^{c} \right]_{|\Sigma} \frac{\delta L}{nL}, \quad (A15)$$

with

$$K^{c} = \left(-g^{Fij}\bar{\partial}_{d}g^{F}_{ij} + \frac{4}{F^{2}}g^{F}_{id}y^{i}\right)\frac{\partial \mathcal{L}}{\partial g^{F}_{cd}} - \bar{\partial}_{d}\frac{\partial \mathcal{L}}{\partial g^{F}_{cd}}.$$
 (A16)

Applying the metric limit now means to consider $L(x, y) = g_{ab}(x)y^a y^b$ with the consequence that $g_{ab}^F(x, y) = -g_{ab}(x)$ for timelike y. The expression for K^c reduces to $K^c \rightarrow \frac{4}{F^2}g_{id}y^i\frac{\partial f}{\partial g_{cd}}$ and $\bar{\partial}_c K^c \rightarrow (\frac{8}{F^4}g_{id}y^ig_{jc}y^j + \frac{4}{F^2}g_{cd})\frac{\partial f}{\partial g_{cd}}$. Collecting all terms in the variation of the matter action in the metric geometry limit finally yields

$$\delta S_m[L, \Phi] \rightarrow \int_{\Sigma} d^4 x d^3 u \sqrt{g^F h^F}_{|\Sigma} \times \left[4\mathcal{L} - 4g_{cd} \frac{\partial \mathcal{L}}{\partial g_{cd}} - 24y^c y^d \frac{\partial \mathcal{L}}{\partial g_{cd}} \right]_{|\Sigma} \times \frac{\delta L}{nL}, \tag{A17}$$

from which we can read off the expression for the source term $T_{|\Sigma}$,

$$T_{|\Sigma} \rightarrow \left(4\mathcal{L} - 4g_{cd}\frac{\partial \mathcal{L}}{\partial g_{cd}} - 24y^{c}y^{d}\frac{\partial \mathcal{L}}{\partial g_{cd}}\right)_{|\Sigma}.$$
 (A18)

The lift of this expression to *TM* requires making all terms zero-homogeneous by multiplication with the appropriate powers of F(x, y), which here means multiplication of the third term by $F(x, y)^{-2}$. The result confirms Eq. (55) that was used to prove the consistency of Finsler gravity with Einstein gravity in the metric geometry limit.

3. Complete lifts of cosmological symmetry generators

We deduced the most general fundamental geometry function L for Finsler spacetimes with cosmological symmetries in Sec. V. The derivation requires the complete lifts of the symmetry-generating vector fields (65) which we display here explicitly:

$$X_{1}^{C} = \chi \left(\sin\theta \cos\phi \partial_{r} + \frac{\chi}{r} \cos\theta \cos\phi \partial_{\theta} - \frac{\chi}{r} \frac{\sin\phi}{\sin\theta} \partial_{\phi} \right) + (y^{r}\chi' \sin\theta \cos\phi + y^{\theta}\xi \cos\theta \cos\phi - y^{\phi}\xi \sin\theta \sin\phi) \bar{\partial}_{r} + \left(y^{r} \left(\frac{\chi}{r} \right)' \cos\theta \cos\phi - y^{\theta} \frac{\chi}{r} \sin\theta \cos\phi - y^{\phi} \frac{\chi}{r} \cos\theta \sin\phi \right) \bar{\partial}_{\theta} + \left(-y^{r} \left(\frac{\chi}{r} \right)' \frac{\sin\phi}{\sin\theta} + y^{\theta} \frac{\chi}{r} \frac{\sin\phi}{\sin^{2}\theta} \cos\theta - y^{\phi} \frac{\chi}{r} \frac{\cos\phi}{\sin\theta} \right) \bar{\partial}_{\phi},$$
(A19)

$$X_{2}^{C} = \chi \sin\theta \sin\phi \partial_{r} + \frac{\chi}{r} \cos\theta \sin\phi \partial_{\theta} + \frac{\chi}{r} \frac{\cos\phi}{\sin\theta} \partial_{\phi} + (y^{r}\chi'\sin\theta \sin\phi + y^{\theta}\xi \cos\theta \sin\phi + y^{\phi}\xi \sin\theta \cos\phi)\bar{\partial}_{r} + \left(y^{r}\left(\frac{\chi}{r}\right)'\cos\theta \cos\phi - y^{\theta}\frac{\chi}{r}\sin\theta \sin\phi + y^{\phi}\frac{\chi}{r}\cos\theta \cos\phi\right)\bar{\partial}_{\theta} + \left(y^{r}\left(\frac{\chi}{r}\right)'\frac{\cos\phi}{\sin\theta} - y^{\theta}\frac{\chi}{r}\frac{\cos\phi}{\sin^{2}\theta}\cos\theta - y^{\phi}\frac{\chi}{r}\frac{\sin\phi}{\sin\theta}\right)\bar{\partial}_{\phi},$$
(A20)

$$X_3^C = \chi \cos\theta \partial_r - \frac{\chi}{r} \sin\theta \partial_\theta + (y^r \chi' \cos\theta - y^\theta \chi \sin\theta) \bar{\partial}_r - \left(y^r \left(\frac{\chi}{r}\right)' \sin\theta + y^\theta \frac{\chi}{r} \cos\theta\right) \bar{\partial}_\theta.$$
(A21)

The complete lifts X_4^C , X_5^C , and X_6^C are stated in Eqs. (60). In the formulae above, we use the abbreviation $\chi = \sqrt{1 - kr^2}$, and primes denote differentiation with respect to the coordinate *r*.

4. Linearization identities

In order to study the Finsler gravitational field equation perturbatively, we have considered a class of Finsler spacetimes that are mild deviations from metric geometry in Sec. VIA. Here, we list for completeness how to rewrite the appearing geometric objects in terms of the perturbation h instead of the variable l used in the main text.

First, we rewrite various derivatives acting on *l* in terms of derivatives acting on *h*:

$$\nabla_a \nabla_b l = \frac{G^{1-k}}{k} \nabla_a \nabla_b h, \tag{A22a}$$

$$\bar{\partial}_a l = \frac{G^{1-k}}{k} \bar{\partial}_a h + \frac{(1-k)G^{-k}}{k} 2g_{ai} y^i h,$$
(A22b)

$$\nabla_a \nabla \bar{\partial}_q l = \frac{G^{1-k}}{k} \nabla_a \nabla \bar{\partial}_q h + \frac{2(1-k)G^{-k}}{k} g_{qi} y^i \nabla_a \nabla h, \tag{A22c}$$

$$\bar{\partial}_a \bar{\partial}_b l = \frac{G^{1-k}}{k} \bar{\partial}_a \bar{\partial}_b h + \frac{2(1-k)G^{-k}}{k} \left(g_{bi} y^i \bar{\partial}_a h + g_{ai} y^i \bar{\partial}_b h + \left(g_{ab} - \frac{2k}{G} g_{ai} y^i g_{bj} y^j \right) h \right), \tag{A22d}$$

$$\nabla \nabla \bar{\partial}_a \bar{\partial}_b l = \frac{G^{1-k}}{k} \nabla \nabla \bar{\partial}_a \bar{\partial}_b h + \frac{2(1-k)G^{-k}}{k} \nabla \nabla \left(g_{bi} y^i \bar{\partial}_a h + g_{ai} y^i \bar{\partial}_b h + \left(g_{ab} - \frac{2k}{G} g_{ai} y^i g_{bj} y^j \right) h \right), \quad (A22e)$$

$$g^{ab}l_{ab} = \frac{1}{2}g^{ab}\bar{\partial}_a\bar{\partial}_b l = \frac{G^{1-k}}{2k}g^{ab}\bar{\partial}_a\bar{\partial}_b h + \frac{2(1-k)(k+2)}{k}G^{-k}h,$$
 (A22f)

$$\nabla_q g^{ab} l_{ab} = \frac{1}{2} g^{ab} \nabla_q \bar{\partial}_a \bar{\partial}_b l = \frac{G^{1-k}}{2k} \nabla_q g^{ab} \bar{\partial}_a \bar{\partial}_b h + \frac{2(1-k)(k+2)}{k} G^{-k} \nabla_q h.$$
(A22g)

These identities can now be employed to determine the curvature scalar $\mathcal{R} = R^a_{\ ab} y^b$ of the Cartan nonlinear connection, which we use as the basic ingredient in our construction of Finsler gravity:

$$R^{a}{}_{ab}y^{b} = -y^{a}y^{b}R_{ab}[g(x)] - \frac{G^{1-k}}{2k}g^{ag}\left(\nabla_{a}\nabla\bar{\partial}_{q}h - \nabla_{a}\nabla_{q}h - \frac{1}{2}\nabla\nabla\bar{\partial}_{a}\bar{\partial}_{q}h\right) - \frac{2(1-k)G^{-k}}{2k}g^{aq}$$

$$\times \left(g_{qi}y^{i}\nabla_{a}\nabla h - \frac{1}{2}\nabla\nabla\left(g_{qi}y^{i}\bar{\partial}_{a}h + g_{ai}y^{i}\bar{\partial}_{q}h + \left(g_{aq} - \frac{2k}{G}g_{ai}y^{i}g_{qj}y^{j}\right)h\right)\right)$$

$$= -y^{a}y^{b}R_{ab}[g(x)] - \frac{G^{1-k}}{2k}g^{ag}(\nabla_{a}\nabla\bar{\partial}_{q}h - \nabla_{a}\nabla_{q}h - \frac{1}{2}\nabla\nabla\bar{\partial}_{a}\bar{\partial}_{q}h)$$

$$+ \frac{(1-k)(1+k)G^{-k}}{k}\nabla\nabla h.$$
(A23)

Finally, we display how to rewrite the d-tensor S and various derivatives acting on it; using the notation tr $h = g^{ab} \bar{\partial}_a \bar{\partial}_b h$, we find:

FINSLER GEOMETRIC EXTENSION OF EINSTEIN GRAVITY

PHYSICAL REVIEW D 85, 064009 (2012)

$$S_{p} = -\frac{G^{1-k}}{4k} y^{q} \bar{\partial}_{p} \nabla_{q}(\operatorname{tr}h) - \frac{(1-k)(k+2)}{k} G^{-k} y^{q} \bar{\partial}_{p} \nabla_{q}h - \frac{(1-k)G^{-k}}{2k} g_{pi} y^{i} \nabla(\operatorname{tr}h) + 2(1-k)(k+2)G^{-(1+k)} g_{pi} y^{i} \nabla h,$$
(A24a)

$$\nabla_{a}S_{b} = -\frac{G^{1-k}}{4k}y^{q}\nabla_{a}\bar{\partial}_{b}\nabla_{q}\mathrm{tr}h - \frac{(1-k)(k+2)}{k}G^{-k}y^{q}\nabla_{a}\bar{\partial}_{b}\nabla_{q}h - \frac{(1-k)G^{-k}}{2k}g_{bi}y^{i}\nabla_{a}\nabla(\mathrm{tr}h) + 2(1-k)(k+2)G^{-(1+k)}g_{bi}y^{i}\nabla_{a}\nabla h, \qquad (A24b)$$

$$g^{ab}\nabla_{a}S_{b} = -\frac{G^{1-k}}{4k}y^{q}g^{ab}\nabla_{a}\bar{\partial}_{b}\nabla_{q}(\mathrm{tr}h) - \frac{(1-k)(k+2)}{k}G^{-k}y^{q}g^{ab}\nabla_{a}\bar{\partial}_{b}\nabla_{q}h - \frac{(1-k)G^{-k}}{2k}\nabla\nabla(\mathrm{tr}h) + 2(1-k)(k+2)G^{-(1+k)}\nabla\nabla h,$$
(A24c)

$$g^{ab}y^{p}\bar{\partial}_{a}\nabla_{p}S_{b} = -\frac{(1-k)(2+k)}{k}G^{-k}(y^{q}y^{p}g^{ab}\bar{\partial}_{a}\nabla_{p}\bar{\partial}_{b}\nabla_{q}h + g^{ab}\nabla\bar{\partial}_{a}\nabla_{b}h)$$

$$-\frac{G^{1-k}}{4k}(y^{q}y^{p}g^{ab}\bar{\partial}_{a}\nabla_{p}\bar{\partial}_{b}\nabla_{q}(\mathrm{tr}h) + g^{ab}\nabla\bar{\partial}_{a}\nabla_{b}(\mathrm{tr}h))$$

$$+ 2(1-k)(2+k)(2k+3)G^{-(k+1)}\nabla\nabla h - \frac{(1-k)(2k+1)}{2k}G^{-k}\nabla\nabla(\mathrm{tr}h).$$
(A24d)

- [1] C. M. Will, Living Rev. Relativity 9, 3 (2005).
- [2] S. Baessler, B.R. Heckel, E.G. Adelberger, J.H. Gundlach, U. Schmidt, and H.E. Swanson, Phys. Rev. Lett. 83, 3585 (1999).
- [3] D. Bao, S.S. Chern, and Z. Shen, *An Introduction to Riemann-Finsler Geometry* (Springer, New York, 2000).
- [4] I. Bucataru and R. Miron, *Finsler-Lagrange Geometry* (Editura Academiei Romane, Bucharest, Romania, 2007).
- [5] C. Pfeifer and M. N. R. Wohlfarth, Phys. Rev. D 84, 044039 (2011).
- [6] Z. Chang and X. Li, Phys. Lett. B 668, 453 (2008).
- [7] Z. Chang and X. Li, Phys. Lett. B 676, 173 (2009).
- [8] C. Pfeifer and M. N. R. Wohlfarth, arXiv:1109.6005.

- [9] C. Lammerzahl, O. Preuss, and H. Dittus, arXiv:gr-qc/ 0604052.
- [10] J.C. Baez, A.E. Hoffnung, and C.D. Walker, arXiv:0812.4864.
- [11] X. Li and Z. Chang, arXiv:1010.2020.
- [12] K. Yano and S. Ishihara, *Tangent and Cotangent Bundles* (Marcel Dekker, New York 1973).
- [13] J. Garecki, Gen. Relativ. Gravit. 27, 55 (1995).
- [14] S. Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972).
- [15] S. M. Carroll, *Spacetime and geometry* (Addison Wesley, San Francisco, 2004).
- [16] T. Adam et al. (OPERA Collaboration), arXiv:1109.4897.