

$N = 1$ non-Abelian tensor multiplet in four dimensions

Hitoshi Nishino* and Subhash Rajpoot†

Department of Physics & Astronomy, California State University, 1250 Bellflower Boulevard, Long Beach, California 90840
 (Received 21 November 2011; revised manuscript received 15 February 2012; published 18 May 2012)

We carry out the $N = 1$ supersymmetrization of a physical non-Abelian tensor with nontrivial consistent couplings in four dimensions. Our system has three multiplets: (i) The usual non-Abelian vector multiplet (A_μ^I, λ^I) , (ii) A non-Abelian tensor multiplet (TM) $(B_{\mu\nu}^I, \chi^I, \varphi^I)$, and (iii) A compensator vector multiplet (CVM) (C_μ^I, ρ^I) . All of these multiplets are in the adjoint representation of a non-Abelian group G . Unlike topological theory, all of our fields are propagating with kinetic terms. The C_μ^I -field plays the role of a Stueckelberg compensator absorbed into the longitudinal component of $B_{\mu\nu}^I$. We give not only the component Lagrangian, but also a corresponding superspace reformulation, reconfirming the total consistency of the system. The adjoint representation of the TM and CVM is further generalized to an arbitrary real representation of general $SO(N)$ gauge group. We also couple the globally $N = 1$ supersymmetric system to supergravity, as an additional nontrivial confirmation.

DOI: 10.1103/PhysRevD.85.105017

PACS numbers: 12.60.Jv, 11.15.-q, 11.30.Pb

I. INTRODUCTION

Recently, the long-standing problem with non-Abelian tensors [1] has been solved by de Wit, Samtleben, and Nicolai [2,3]. The original motivation in Ref. [2] was to generalize the tensor and vector field interactions in manifestly $E_{6(+6)}$ -covariant formulation of five-dimensional (5D) maximal supergravity by gauging non-Abelian subgroups. In Ref. [3], this work was further related to M-theory [4] by confirming the representation assignments under the duality group of the gauge charges. The underlying hierarchies of these tensor and vector gauge fields are presented with the consistency of general gaugings.

The hierarchy in Refs. [2,3] has been further applied to the conformal supergravity in 6D [5]. In Ref. [5], the ‘‘minimal tensor hierarchy’’ as a special case of the more general hierarchy in Refs. [2,3] has been discussed. This hierarchy consists of A_μ^r and 2-form gauge potentials $B_{\mu\nu}^I$, with two labels, r and I . Also introduced is the 3-form gauge potential $C_{\mu\nu\rho}^r$ with the index r , which is dual to r of A_μ^r . The field strengths of vector and 2-form gauge potentials are defined by [5]

$$\mathcal{F}_{\mu\nu}^r \equiv 2\partial_{[\mu}A_{\nu]}^r + h_I^r B_{\mu\nu}^I, \quad (1.1a)$$

$$\begin{aligned} \mathcal{H}_{\mu\nu\rho}^I &\equiv 3D_{[\mu}B_{\nu\rho]}^I + 6d_{rs}^I A_{[\mu}^r \partial_{\nu}A_{\rho]}^s \\ &\quad - 2f_{pq}^s d_{rs}^I A_{[\mu}^r A_{\nu}^p A_{\rho]}^q + g^{Ir} C_{\mu\nu\rho}^r. \end{aligned} \quad (1.1b)$$

The prescription for tensor-vector system, which we will be based upon, is described with Eq. (3.22) in Ref. [5]. To be more specific, we consider in the present paper the product of two identical gauge groups $G \times G$ [6], whose adjoint indices are, respectively, r, s, \dots and r', s', \dots . Accordingly, we use the coefficients

*hnishino@csulb.edu
 †rajpoot@csulb.edu

$$f_{rs}^t = f_{rs}^t, \quad f_{rs'}^t = -f_{s'r}^t = +\frac{1}{2}f_{rs'}^t, \quad (1.2a)$$

$$d_{rs'}^t = d_{s'r}^t = -\frac{1}{2}f_{rs'}^t, \quad h_s^r = \delta_s^r, \quad (1.2b)$$

where f_{rs}^t is the structure constant of a non-Abelian gauge group. We use the same field content arising by this prescription.

Since the outstanding paper Ref. [5] gives the extensive details of how to get our system from Refs. [2,3,6], there is nothing new to explain, except for our notational preparation. In our notation, the field strengths of the B - and C -fields are, respectively, G and H , defined by

$$G_{\mu\nu\rho}^I \equiv +3D_{[\mu}B_{\nu\rho]}^I - 3f^{IJK}C_{[\mu}^J F_{\nu\rho]}^K, \quad (1.3a)$$

$$H_{\mu\nu}^I \equiv +2D_{[\mu}C_{\nu]}^I + gB_{\mu\nu}^I. \quad (1.3b)$$

The gauge transformations for B , C , and A -fields are

$$\begin{aligned} \delta_\alpha(B_{\mu\nu}^I, C_\mu^I, A_\mu^I) &= (-f^{IJK}\alpha^J B_{\mu\nu}^K, \\ &\quad -f^{IJK}\alpha^J C_\mu^K, +D_\mu\alpha^I), \end{aligned} \quad (1.4a)$$

$$\delta_\beta(B_{\mu\nu}^I, C_\mu^I, A_\mu^I) = (+2D_{[\mu}\beta_{\nu]}^I, -g\beta_\mu^I, 0), \quad (1.4b)$$

$$\delta_\gamma(B_{\mu\nu}^I, C_\mu^I, A_\mu^I) = (-f^{IJK}F_{\mu\nu}^J \gamma^K, D_\mu\gamma^I, 0). \quad (1.4c)$$

As Eq. (1.3b) or Eq. (1.4b) shows, C_μ^I is a vectorial Stueckelberg field, absorbed into the longitudinal component of $B_{\mu\nu}^I$. Because of the general hierarchy [2,3], all field strengths are invariant:

$$\delta_\alpha(G_{\mu\nu\rho}^I, H_{\mu\nu}^I, F_{\mu\nu}^I) = -f^{IJK}\alpha^J(G_{\mu\nu\rho}^K, H_{\mu\nu}^K, F_{\mu\nu}^K), \quad (1.5a)$$

$$\delta_\beta(G_{\mu\nu\rho}^I, H_{\mu\nu}^I, F_{\mu\nu}^I) = 0,$$

$$\delta_\gamma(G_{\mu\nu\rho}^I, H_{\mu\nu}^I, F_{\mu\nu}^I) = 0. \quad (1.5b)$$

Since the hierarchy given in Refs. [2,3] guarantees the gauge invariance of all field strengths, the construction of purely bosonic Lagrangian is straightforward. Consider the action $I_1 \equiv \int d^4x g^2 \mathcal{L}_1^1$ with

$$\mathcal{L}_1 \equiv -\frac{1}{12}(G_{\mu\nu\rho}{}^I)^2 - \frac{1}{4}(H_{\mu\nu}{}^I)^2 - \frac{1}{4}(F_{\mu\nu}{}^I)^2. \quad (1.6)$$

The gauge invariances of all field strength also guarantee the consistency of the A -, B -, and C -field equations, such as the divergence $D_\nu(\delta\mathcal{L}_1/\delta B_{\mu\nu}{}^I) \doteq 0$.² Since we will do similar confirmation for supersymmetric system later, we skip the details for the purely bosonic system.

The purpose of our present paper is to supersymmetrize this system. The rest of our paper is organized as follows. In Sec. II, we give the component formulation of $N = 1$ TM. In Sec. III, we give the superspace reformulation of component result. In Sec. IV, we give the generalization to non-adjoint representation of $G = SO(N)$ case. In Sec. V, we give the supergravity coupling to non-Abelian TM, as supporting evidence for the consistency of the global case.

Sec. VI is for concluding remarks. The Appendix is devoted to purely bosonic systems of non-Abelian tensors with much simpler structures than have been presented in arbitrary space-time dimensions with arbitrary signature. An example of tensor-vector duality $G = F^*$ in $D = 2 + 4$ dimensions, and its dimensional reduction (DR) into the self-dual Yang-Mills $F = F^*$ in $D = 2 + 2$, is also presented.

II. COMPONENT FORMULATION OF $N = 1$ TM

The supersymmetrization of the purely bosonic system Eq. (1.6) is rather straightforward, except for a subtlety to be mentioned later. Our system has three multiplets: (i) A TM $(B_{\mu\nu}{}^I, \chi^I, \varphi^I)$, (ii) A CVM $(C_\mu{}^I, \rho^I)$, and (iii) A YM vector multiplet $(A_\mu{}^I, \lambda^I)$. Our total action $I \equiv \int d^4x g^2 \mathcal{L}$ has the Lagrangian

$$\begin{aligned} \mathcal{L} = & -\frac{1}{12}(G_{\mu\nu\rho}{}^I)^2 + \frac{1}{2}(\bar{\chi}^I \not{D} \chi^I) - \frac{1}{2}(D_\mu \varphi^I)^2 - \frac{1}{2}g^2(\varphi^I)^2 - g(\bar{\chi}^I \not{\rho}^I) - \frac{1}{4}(H_{\mu\nu}{}^I)^2 + \frac{1}{2}(\bar{\rho}^I \not{D} \rho^I) - \frac{1}{4}(F_{\mu\nu}{}^I)^2 \\ & + \frac{1}{2}(\bar{\lambda}^I \not{D} \lambda^I) - \frac{1}{2}g f^{IJK}(\bar{\lambda}^I \chi^J) \varphi^K + \frac{1}{2} f^{IJK}(\bar{\lambda}^I \gamma^\mu \rho^J) D_\mu \varphi^K + \frac{1}{12} f^{IJK}(\bar{\lambda}^I \gamma^{\mu\nu\rho} \rho^J) G_{\mu\nu\rho}{}^K \\ & + \frac{1}{4} f^{IJK}(\bar{\rho}^I \gamma^{\mu\nu} \chi^J) F_{\mu\nu}{}^K - \frac{1}{4} f^{IJK}(\bar{\lambda}^I \gamma^{\mu\nu} \chi^J) H_{\mu\nu}{}^K - \frac{1}{2} f^{IJK} F_{\mu\nu}{}^I H^{\mu\nu J} \varphi^K, \end{aligned} \quad (2.1)$$

up to quartic-order terms $\mathcal{O}(\phi^4)$.

It is clear that the scalar φ^I has its mass g , while there is a mixture between χ^I and ρ^I , again with the same mass g . As has been mentioned after Eq. (1.4), $C_\mu{}^I$ plays the role of Stueckelberg field [7], being absorbed into the longitudinal component of $B_{\mu\nu}{}^I$. Eventually, the kinetic term of the C -field becomes the mass term of $B_{\mu\nu}{}^I$. Accordingly, the DOFs for the massive TM fields are $B_{\mu\nu}{}^I$ (3); ρ^I and χ^I (4); and φ^I (1), up to the adjoint index I .

Our action I is invariant under global $N = 1$ supersymmetry

$$\begin{aligned} \delta_Q B_{\mu\nu}{}^I &= +(\bar{\epsilon} \gamma_{\mu\nu} \chi^I) - 2f^{IJK} C_{[\mu}{}^J (\delta_Q A_{\nu]}{}^K), \\ \delta_Q \chi^I &= +\frac{1}{6}(\gamma^{\mu\nu\rho} \epsilon) G_{\mu\nu\rho}{}^I - (\gamma^\mu \epsilon) D_\mu \varphi^I \end{aligned} \quad (2.2a)$$

$$+\frac{1}{2} f^{IJK} [+\epsilon(\bar{\lambda}^J \rho^K) - (\gamma_5 \gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_5 \gamma_\mu \rho^K) - (\gamma_5 \epsilon)(\bar{\lambda}^J \gamma_5 \rho^K)], \quad (2.2b)$$

$$\delta_Q \varphi^I = +(\bar{\epsilon} \chi^I), \quad (2.2c)$$

$$\delta_Q C_\mu{}^I = +(\bar{\epsilon} \gamma_\mu \rho^I) + f^{IJK}(\bar{\epsilon} \gamma_\mu \lambda^J) \varphi^K, \quad (2.2d)$$

$$\begin{aligned} \delta_Q \rho^I &= +\frac{1}{2}(\gamma^{\mu\nu} \epsilon) H_{\mu\nu}{}^I - g \epsilon \varphi^I - \frac{1}{2} f^{IJK}(\gamma^{\mu\nu} \epsilon) F_{\mu\nu}{}^J \varphi^K + \frac{1}{4} f^{IJK} [+\epsilon(\bar{\lambda}^J \chi^K) - (\gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_\mu \chi^K) \\ & + \frac{1}{2}(\gamma^{\mu\nu} \epsilon)(\bar{\lambda}^J \gamma_{\mu\nu} \chi^K) - (\gamma_5 \gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_5 \gamma_\mu \chi^K) - (\gamma_5 \epsilon)(\bar{\lambda}^J \gamma_5 \chi^K)], \end{aligned} \quad (2.2e)$$

$$\delta_Q A_\mu{}^I = +(\bar{\epsilon} \gamma_\mu \lambda^I), \quad (2.2f)$$

$$\delta_Q \lambda^I = +\frac{1}{2}(\gamma^{\mu\nu} \epsilon) F_{\mu\nu}{}^I + \frac{1}{2} f^{IJK}(\gamma_5 \epsilon)(\bar{\rho}^J \gamma_5 \chi^K), \quad (2.2g)$$

up to cubic terms $\mathcal{O}(\phi^3)$ in fields. The fermionic quadratic terms in Eqs. (2.2b), (2.2e), and (2.2g) are fixed in superspace formulation, as will be explained later. In the *conventional* dimensions with all the bosonic (or fermionic)

fields with 1 (or 3/2) mass dimensions,³ these terms lead to nonrenormalizability. For example, the left side of (2.2b) has dimension 3/2, while its right side for the $\epsilon(\bar{\lambda}^J \rho^K)$ term has $(-1/2) + (3/2) + (3/2) = 5/2$. In other words, there

¹The reason we need the factor g^2 in the action is due to the mass-dimension assignments of our fields.

²We use the symbol \doteq for a field equation to distinguish from an algebraic equation.

³Our bosonic (or fermionic) fields have dimensions 0 (or 1/2), in contrast to the conventional dimensions 1 (or 3/2).

is an implicit coupling constant ℓ with the dimension of length in front of fermionic quadratic terms. This feature is also related to the existence of Pauli terms which are non-renormalizable, already at a *globally* supersymmetric system. These features are similar to supergravity [8], even though our system so far has only *global* supersymmetry.

The usual non-Abelian gauge transformation δ_α and our tensorial gauge transformation δ_β and δ_γ -transformation are exactly the same as Eq. (1.4), while all the fermionic fields are transforming only under δ_α , as the B and C -fields do, so that there arises no problem with the δ_β and δ_γ -invariances of the field strengths as in Eq. (1.5). These immediately lead to the invariances of our action $\delta_\alpha I = 0$, $\delta_\beta I = 0$, and $\delta_\gamma I = 0$.

The Bianchi identities (BIDs) for our field strengths G , H , and F are:

$$D_{[\mu} G_{\nu\rho]}^I - \frac{3}{2} f^{IJK} F_{[\mu\nu}^J H_{\rho\sigma]}^K \equiv 0, \quad (2.3a)$$

$$D_{[\mu} H_{\nu\rho]}^I - \frac{1}{3} g G_{\mu\nu\rho}^I \equiv 0, \quad (2.3b)$$

$$D_{[\mu} F_{\nu\rho]}^I \equiv 0. \quad (2.3c)$$

$$+ \not{D} \lambda^I - \frac{1}{2} g f^{IJK} \chi^J \varphi^K + \frac{1}{2} f^{IJK} (\gamma^\mu \rho^J) D_\mu \varphi^K - \frac{1}{4} f^{IJK} (\gamma^{\mu\nu} \chi^J) H_{\mu\nu}^K + \frac{1}{12} f^{IJK} (\gamma^{\mu\nu\rho} \rho^J) G_{\mu\nu\rho}^K \doteq 0, \quad (2.5a)$$

$$+ \not{D} \chi^I - g \rho^I + \frac{1}{2} g f^{IJK} \lambda^J \varphi^K - \frac{1}{4} f^{IJK} (\gamma^{\mu\nu} \lambda^J) H_{\mu\nu}^K + \frac{1}{4} f^{IJK} (\gamma^{\mu\nu} \rho^J) F_{\mu\nu}^K \doteq 0, \quad (2.5b)$$

$$+ \not{D} \rho^I - g \chi^I + \frac{1}{2} f^{IJK} (\gamma^\mu \lambda^J) D_\mu \varphi^K - \frac{1}{12} f^{IJK} (\gamma^{\mu\nu\rho} \lambda^J) G_{\mu\nu\rho}^K + \frac{1}{4} f^{IJK} (\gamma^{\mu\nu} \chi^J) F_{\mu\nu}^K \doteq 0, \quad (2.5c)$$

$$+ D_\nu F_{\mu}{}^{\nu I} + g f^{IJK} \varphi^J D_\mu \varphi^K + \frac{1}{2} g f^{IJK} (\bar{\lambda}^J \gamma_\mu \lambda^K) + f^{IJK} H_{\mu\nu}^J D^\nu \varphi^K - \frac{1}{2} f^{IJK} G_{\mu\rho\sigma}^J H^{\rho\sigma K} + \frac{1}{2} f^{IJK} (\bar{\chi}^J D_\mu \rho^K) + \frac{1}{2} f^{IJK} (\bar{\rho}^J D_\mu \chi^K) \doteq 0, \quad (2.5d)$$

$$+ D_\rho G^{\mu\nu\rho I} - g H^{\mu\nu I} - \frac{1}{2} f^{IJK} D_\rho (\bar{\lambda}^J \gamma^{\mu\nu\rho} \rho^K) + g f^{IJK} F^{\mu\nu J} \varphi^K - \frac{1}{2} g f^{IJK} (\bar{\lambda}^J \gamma^{\mu\nu} \chi^K) \doteq 0, \quad (2.5e)$$

$$+ D_\mu^2 \varphi^I - g f^{IJK} (\bar{\lambda}^J \chi^K) - g^2 \varphi^I - \frac{1}{2} f^{IJK} F_{\mu\nu}^J H^{\mu\nu K} \doteq 0, \quad (2.5f)$$

$$+ D_\nu H^{\mu\nu I} - \frac{1}{2} f^{IJK} F_{\rho\sigma}^J G^{\mu\rho\sigma K} - \frac{1}{2} f^{IJK} (\bar{\chi}^J D^\mu \lambda^K) - \frac{1}{2} f^{IJK} (\bar{\lambda}^J D^\mu \chi^K) + \frac{1}{2} g f^{IJK} (\bar{\lambda}^J \gamma^\mu \rho^K) - f^{IJK} F^{\mu\nu J} D_\nu \varphi^K \doteq 0. \quad (2.5g)$$

In the derivation of these field equations, we have also used other field equations in order to simply their final expressions, as a conventional prescription.

In the previous computation, we do not attempt to fix the $\mathcal{O}(\phi^3)$ -terms in field equations, or equivalently the fermionic $\mathcal{O}(\phi^4)$ -terms in the Lagrangian. There are several remarks about these terms. First, our system is nonrenormalizable as supergravity theory [8], as has been mentioned after Eq. (2.2). Accordingly, the (fermion)²-terms in the fermionic transformations such as Eqs. (2.2b), (2.2e), and (2.2g) are accompanied by the implicit constant ℓ carrying the dimension of (length). In supergravity theory [8], this is the gravitational coupling κ . In our Lagrangian, all the quartic-fermion terms carry ℓ^2 , so that the

Relevantly, the nontrivial δ_Q -transformations of the field strengths are

$$\delta_Q G_{\mu\nu\rho}^I = +3(\bar{\epsilon} \gamma_{[\mu\nu} D_{\rho]} \lambda^I) + 3f^{IJK} (\delta_Q A_{[\mu}^J) H_{\nu\rho]}^K - 3f^{IJK} (\delta_Q C_{[\mu}^J) F_{\nu\rho]}^K, \quad (2.4a)$$

$$\delta_Q H_{\mu\nu}^I = -2(\bar{\epsilon} \gamma_{[\mu} D_{\nu]} \rho^I) + g(\bar{\epsilon} \gamma_{\mu\nu} \chi^I) + 2f^{IJK} D_{[\mu} [(\delta_Q A_{\nu]}^J) \varphi^K], \quad (2.4b)$$

$$\delta_Q F_{\mu\nu}^I = -2(\bar{\epsilon} \gamma_{[\mu} D_{\nu]} \lambda^I), \quad (2.4c)$$

reflecting the presence of Chern-Simon terms.

Note that our YMVM and CVM have *on-shell* DOF 2 + 2, while *off-shell* DOF is 3 + 4, because we have *not* added the D -auxiliary field. On the other hand, our TM is in the *off-shell* formulation because the total off-shell DOF is 4 + 4, because the off-shell DOF of each field are $[(4 - 1) \cdot (4 - 2)]/2 = 3$ for $B_{\mu\nu}$, 4 for χ , and 1 for φ .

The field equations for λ^I , χ^I , ρ^I , A_μ^I , $B_{\mu\nu}^I$, φ^I , and C_μ^I are, respectively, ⁴

Lagrangian has the mass dimension +4. Accordingly, a typical Noether term has the structure $\ell \Psi^2 \partial \Phi$, which produces the terms of the form $\ell^2 \epsilon \Psi^3 \partial \Phi$ via $\delta_Q \Psi \approx \ell \epsilon \Psi^2$. Here, Ψ (or Φ) is a general fermionic (or bosonic) fundamental field. These $\ell^2 \epsilon \Psi^3 \partial \Phi$ -terms are canceled by the variation of the fermionic quartic terms $\ell^2 \Psi^4$, via $\delta_Q \Psi \approx \epsilon \partial \Phi$. In other words, the structure of these cancellations associated with quartic-fermion terms is parallel to supergravity [8], since ℓ is analogous to κ .

⁴These equations are fixed up to $\mathcal{O}(\phi^3)$ -terms, due to the quartic fermion terms in the Lagrangian.

However, in our peculiar system, this cancellation mechanism may be *not* simply parallel to conventional supergravity [8]. For example, there may be $\ell^2\Psi^2\Phi\partial\Phi$ -type terms in the action, while $\ell^2\epsilon\Psi^2\Phi$ -type terms in the transformation rules may exist, because both of them yield $\ell^2\epsilon\Psi^3\partial\Phi$ -type terms, canceling each other in $\delta_Q I$. At the present time, we do not know if such terms arise, because the $\ell^2\epsilon\Psi^2\Phi$ -type terms in transformations are at $\mathcal{O}(\phi^3)$, while $\ell^2\Psi^2\Phi\partial\Phi$ -type terms in the action are at $\mathcal{O}(\phi^4)$. In fact, even in the superspace reconfirmation in the next section, we have fixed only the $\mathcal{O}(\phi^1)$ - and $\mathcal{O}(\phi^2)$ -terms in the transformation rules for fermions, such as Eqs. (3.2d)–(3.2f), but *not* cubic terms $\mathcal{O}(\phi^3)$. Our consistent principle in this paper is to fix only $\mathcal{O}(\phi^1)$ -, $\mathcal{O}(\phi^2)$ -, and $\mathcal{O}(\phi^3)$ -terms in the Lagrangian and $\mathcal{O}(\phi^1)$ - and $\mathcal{O}(\phi^2)$ -terms in all transformation rules, while fixing $\mathcal{O}(\phi^1)$ - and $\mathcal{O}(\phi^2)$ -terms in all field equations. However, we try to fix neither $\mathcal{O}(\phi^4)$ -terms in the Lagrangian nor $\mathcal{O}(\phi^3)$ -terms in all transformation rules, nor $\mathcal{O}(\phi^3)$ -terms in all field equations. We do *not* specify that each field meant by ϕ is fermionic or bosonic in this paper, either.

Second, as an additional difference from supergravity [8], the fermionic quartic terms do *not* contain any gravitino. This implies that we cannot use the conventional technique of “supercovariantizing” fermionic field equations. Because of this feature, as well as the abovementioned possible non-purely fermionic $\ell^2\Psi^2\Phi\partial\Phi$ -type terms, the quartic terms $\mathcal{O}(\phi^4)$ at $\mathcal{O}(\ell^2)$ will be more involved than conventional supergravity [8], which are tedious. For these reasons, we do not attempt to fix them in this paper.

Third, according to the past experience in supergravity theory [8], it is understood that the series in terms of κ in a

Lagrangian will stop at a finite order, such as the quartic-fermion terms at $\mathcal{O}(\kappa^2)$ [8]. However, at the present time, we do *not* know whether this is also the case with our globally supersymmetric system. This is because of the abovementioned differences of our system from supergravity [8], and therefore the analogy with supergravity might be not valid in our system.

Fourth, since we have already fixed the cubic terms in the Lagrangian, they seem sufficient for nontrivial and consistent couplings as a supersymmetric system.

III. SUPERSPACE REFORMULATION OF $N = 1$ TM

As a reconfirmation of the total consistency of our system, we reformulate our theory in terms of superspace language. Our basic superspace BIDs for the superfield strengths $F_{AB}{}^I$, $G_{ABC}{}^I$, and $H_{AB}{}^I$ are⁵

$$+\frac{1}{6}\nabla_{[A}G_{BCD)}{}^I - \frac{1}{4}T_{[AB]}{}^E G_{E|CD)} - \frac{1}{4}f^{IJK}F_{[AB}{}^J H_{CD)}{}^K \equiv 0, \quad (3.1a)$$

$$+\frac{1}{2}\nabla_{[A}H_{BC)}{}^I - \frac{1}{2}T_{[AB]}{}^D H_{D|C)}{}^I - gG_{ABC}{}^I \equiv 0, \quad (3.1b)$$

$$+\frac{1}{2}\nabla_{[A}F_{BC)}{}^I - \frac{1}{2}T_{[AB]}{}^D F_{D|C)}{}^I \equiv 0. \quad (3.1c)$$

These BIDs are the superspace generalizations of the component BIDs in Eq. (2.3), with the supertorsion terms added for local Lorentz indices, as usual in superspace.

Our basic superspace constraints at mass dimensions $0 \leq d \leq 1$ are

$$T_{\alpha\beta}{}^c = +2(\gamma^c)_{\alpha\beta}, \quad G_{\alpha\beta c}{}^I = +2(\gamma_c)_{\alpha\beta}\varphi^I, \quad (3.2a)$$

$$G_{abc}{}^I = -(\gamma_{bc}\chi^I)_\alpha, \quad H_{ab}{}^I = -(\gamma_b\rho^I)_\alpha - f^{IJK}(\gamma_b\lambda^J)_\alpha\varphi^K, \quad (3.2b)$$

$$F_{ab}{}^I = -(\gamma_b\lambda^I)_\alpha, \quad \nabla_\alpha\varphi^I = -\chi_\alpha^I, \quad (3.2c)$$

$$\nabla_\alpha\chi_\beta^I = -\frac{1}{6}(\gamma^{cde})_{\alpha\beta}G_{cde}{}^I - (\gamma^c)_{\alpha\beta}\nabla_c\varphi^I - \frac{1}{2}f^{IJK}[+C_{\alpha\beta}(\bar{\lambda}^J\rho^K) - (\gamma_5\gamma^c)_{\alpha\beta}(\bar{\lambda}^J\gamma_5\gamma_c\rho^K) - (\gamma_5)_{\alpha\beta}(\bar{\lambda}^J\gamma_5\rho^K)], \quad (3.2d)$$

$$\begin{aligned} \nabla_\alpha\rho_\beta^I &= +\frac{1}{2}(\gamma^{cd})_{\alpha\beta}H_{cd}{}^I + gC_{\alpha\beta}\varphi^I - \frac{1}{2}f^{IJK}(\gamma^{cd})_{\alpha\beta}F_{cd}{}^J\varphi^K - \frac{1}{4}f^{IJK}[+C_{\alpha\beta}(\bar{\lambda}^J\chi^K) + (\gamma^c)_{\alpha\beta}(\bar{\lambda}^J\gamma_c\chi^K) \\ &\quad - \frac{1}{2}(\gamma^{cd})_{\alpha\beta}(\bar{\lambda}^J\gamma_{cd}\chi^K) - (\gamma_5\gamma^c)_{\alpha\beta}(\bar{\lambda}^J\gamma_5\gamma_c\chi^K) - (\gamma_5)_{\alpha\beta}(\bar{\lambda}^J\gamma_5\chi^K), \end{aligned} \quad (3.2e)$$

$$\nabla_\alpha\lambda_\beta^I = +\frac{1}{2}(\gamma^{cd})_{\alpha\beta}F_{cd}{}^I - \frac{1}{2}(\gamma_5)_{\alpha\beta}f^{IJK}(\bar{\rho}^J\gamma_5\chi^K). \quad (3.2f)$$

All other components, such as $G_{\alpha\beta\gamma}{}^I$, $T_{\alpha\beta}{}^\gamma$, $T_{ab}{}^c$, $H_{\alpha\beta}{}^I$, etc., at $d \leq 1$ are zero. Note that (fermion)²-terms in Eqs. (3.2d) through (3.2f) have been determined in superspace by satisfying BIDs at $d = 1$. Note that these results are valid up to $\mathcal{O}(\phi^3)$ -terms, which we do not attempt to fix in this paper. However, all the $\mathcal{O}(\phi^2)$ -terms have been included, as was also mentioned at the end of the previous section.

⁵Only in this superspace section, we use the indices $A = (a, \alpha)$, $B = (b, \beta)$, \dots for superspace coordinates, where $a, b, \dots = 0, 1, 2, 3$ (or $\alpha, \beta, \dots = 1, 2, 3, 4$) are for bosonic (or fermionic) coordinates. In superspace, the (anti)symmetrization convention, e.g., $X_{[AB]} \equiv X_{AB} - (-1)^{AB}X_{BA}$ is different from our component notation.

There are also useful relationships obtained from $d = +3/2$ BIDs:

$$\begin{aligned} \nabla_\alpha G_{bcd} = & -\frac{1}{2}(\gamma_{[bc}\nabla_{d]}\chi^\alpha) - \frac{1}{2}f^{IJK}(\gamma_{[b|}\lambda^J)_\alpha H_{|cd]}^K \\ & + \frac{1}{2}f^{IJK}(\gamma_{[b|}\rho^J)_\alpha F_{|cd]}^K, \end{aligned} \quad (3.3a)$$

$$\begin{aligned} \nabla_\alpha H_{bc}^I = & +(\gamma_{[b}\nabla_{c]}\rho^I)_\alpha - g(\gamma_{bc}\chi^I)_\alpha \\ & - f^{IJK}\nabla_{[b}[(\gamma_{c]}\lambda^J)_\alpha \varphi^K], \end{aligned} \quad (3.3b)$$

$$\nabla_\alpha F_{bc}^I = +(\gamma_{[b}\nabla_{c]}\lambda^I)_\alpha, \quad (3.3c)$$

up to $\mathcal{O}(\phi^3)$ -terms. Note the existence of the $\mathcal{O}(\phi^2)$ -terms in Eqs. (3.3a) and (3.3b), reflecting the corresponding terms in the component results in Eqs. (2.4a) and (2.4b).

As usual, the satisfaction of all the BIDs in superspace by the constraints Eqs. (3.2) and (3.3) is straightforward to perform, from the dimension $d = 0$ to $d = 3/2$, as usual. In particular, the (Fermions)² terms in Eqs. (3.2d) through (3.2f) are the results of our superspace reformulation.

The fermionic λ and ρ -field Eqs. (2.5a) and (2.5c) are obtained as usual by computing $\{\nabla_\alpha, \nabla_\beta\}\lambda^{\beta I}$ and

$\{\nabla_\alpha, \nabla_\beta\}\rho^{\beta I}$, while the χ -field Eq. (2.5b) is shown to be consistent with the component Lagrangian. As has been mentioned, since the TM is *off-shell* multiplet, we *cannot* get the χ -field Eq. (2.5b) in superspace directly, but we can show that Eq. (2.5b) is consistent in superspace. The bosonic field Eqs. (2.5d)–(2.5g) are obtained by applying another fermionic derivative on the fermionic field Eqs. (2.5a)–(2.5c).

IV. GENERALIZATION TO NON-ADJOINT REPRESENTATIONS OF $G = SO(N)$

We have so far considered the case for the TM and CVM both carrying only the adjoint representation. We can generalize this result to other, more general representations, such as an arbitrary real representation of a $SO(N)$ -type gauge group.⁶

To be more specific, we consider the TM $(B_{\mu\nu}^i, \chi^i, \varphi^i)$ and the CVM (C_μ^i, ρ^i) , where the index i is for any real representation of a gauge group $G = SO(N)$. Let $(T^I)^{jk}$ be the generator of the group G . Then our action $I' \equiv \int d^4x g^2 \mathcal{L}'$ has the Lagrangian⁷

$$\begin{aligned} \mathcal{L}' = & -\frac{1}{12}(G_{\mu\nu\rho}^i)^2 + \frac{1}{2}(\bar{\chi}^i \not{D} \chi^i) - \frac{1}{2}(D_\mu \varphi^i)^2 - \frac{1}{2}g^2(\varphi^i)^2 - g(\bar{\rho}^i \chi^i) - \frac{1}{4}(H_{\mu\nu}^i)^2 + \frac{1}{2}(\bar{\rho}^i \not{D} \rho^i) - \frac{1}{4}(F_{\mu\nu}^I)^2 \\ & + \frac{1}{2}(\bar{\lambda}^I \not{D} \lambda^I) - \frac{1}{2}g(T^I)^{jk}(\bar{\lambda}^I \chi^j) \varphi^k + \frac{1}{2}(T^I)^{jk}(\bar{\lambda}^I \gamma^\mu \rho^j) D_\mu \varphi^k + \frac{1}{12}(T^I)^{jk}(\bar{\lambda}^I \gamma^{\mu\nu\rho} \rho^j) G_{\mu\nu\rho}^k \\ & + \frac{1}{4}(T^I)^{jk}(\bar{\rho}^j \gamma^{\mu\nu} \chi^k) F_{\mu\nu}^I - \frac{1}{4}(T^I)^{jk}(\bar{\lambda}^I \gamma^{\mu\nu} \chi^j) H_{\mu\nu}^k - \frac{1}{2}(T^I)^{jk} F_{\mu\nu}^I H^{\mu\nu j} \varphi^k \end{aligned} \quad (4.1)$$

up to quartic terms $\mathcal{O}(\phi^4)$. Our action I' is invariant under global $N = 1$ supersymmetry

$$\delta_Q B_{\mu\nu}^i = +(\bar{\epsilon} \gamma_{\mu\nu} \chi^i) - 2(T^J)^{ik} C_{[\mu|}^k (\delta_Q A_{|\nu]}^J), \quad (4.2a)$$

$$\delta_Q \chi^i = +\frac{1}{6}(\gamma^{\mu\nu\rho} \epsilon) G_{\mu\nu\rho}^i - (\gamma^\mu \epsilon) D_\mu \varphi^i - \frac{1}{2}(T^J)^{ik} [+\epsilon(\bar{\lambda}^J \chi^k) - (\gamma_5 \gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_5 \gamma_\mu \chi^k) - (\gamma_5 \epsilon)(\bar{\lambda}^J \gamma_5 \chi^k)], \quad (4.2b)$$

$$\delta_Q \varphi^i = +(\bar{\epsilon} \chi^i), \quad (4.2c)$$

$$\delta_Q C_\mu^i = +(\bar{\epsilon} \gamma_\mu \rho^i) - (T^J)^{ik} (\bar{\epsilon} \gamma_\mu \lambda^J) \varphi^k, \quad (4.2d)$$

$$\begin{aligned} \delta_Q \rho^i = & +\frac{1}{2}(\gamma^{\mu\nu} \epsilon) H_{\mu\nu}^i - g \epsilon \varphi^i + \frac{1}{2}(T^J)^{ik} (\gamma^{\mu\nu} \epsilon) F_{\mu\nu}^J \varphi^k - \frac{1}{4}(T^J)^{ik} [+\epsilon(\bar{\lambda}^J \chi^k) - (\gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_\mu \chi^k) \\ & + \frac{1}{2}(\gamma^{\mu\nu} \epsilon)(\bar{\lambda}^J \gamma_{\mu\nu} \chi^k) - (\gamma_5 \gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_5 \gamma_\mu \chi^k) - (\gamma_5 \epsilon)(\bar{\lambda}^J \gamma_5 \chi^k)], \end{aligned} \quad (4.2e)$$

$$\delta_Q A_\mu^I = +(\bar{\epsilon} \gamma_\mu \lambda^I), \quad (4.2f)$$

$$\delta_Q \lambda^I = +\frac{1}{2}(\gamma^{\mu\nu} \epsilon) F_{\mu\nu}^I - \frac{1}{2}(T^J)^{jk} (\gamma_5 \epsilon) (\bar{\rho}^j \gamma_5 \chi^k). \quad (4.2g)$$

The essential point is that all the cubic-order terms contain one component field A_μ^I or λ^I with the index I , and the remaining two component fields out of either TM or CVM carry the indices j and k . So the cancellation structure is parallel to the adjoint-representation case, e.g., with the structure constant f^{IJK} replaced by the matrix $-(T^J)^{ik}$ in $D_\mu \chi^I = \partial_\mu \chi^I + g f^{IJK} A_\mu^J \chi^K \longrightarrow D_\mu \chi^I = \partial_\mu \chi^I - g (T^J)^{ik} A_\mu^J \chi^k$. Accordingly, the Stueckelberg mechanism [7]

⁶We can also consider the complex representation for $SU(N)$ -type gauge groups.

⁷Since the metric for the gauge group $G = SO(N)$ is positive definite, we do *not* distinguish the upper or lower indices for $i, j, \dots = 1, 2, \dots, \dim R$, where R is a real representation of G .

works in a parallel fashion, because C_μ^i is absorbed into the longitudinal component of $B_{\mu\nu}^i$, both in the same representation **R**.

V. COUPLING TO $N = 1$ SUPERGRAVITY

Once we have established the $N = 1$ global system of non-Abelian TM with nontrivial and consistent interactions, the next natural step is to make $N = 1$ supersymmetry *local*, coupling to $N = 1$ supergravity.

This coupling is rather straightforward because most of the basic structure is parallel to the usual matter coupling to supergravity, except for certain couplings to be mentioned later. Our result for the Lagrangian $\tilde{\mathcal{L}}$ of our action is $\tilde{I} \equiv \int d^4x g^2 \tilde{\mathcal{L}}$:

$$\begin{aligned}
e^{-1} \tilde{\mathcal{L}} = & -\frac{1}{4}R(\omega) - [\bar{\psi}_\mu \gamma^{\mu\nu\rho} D_\nu(\omega) \psi_\rho] - \frac{1}{12}(G_{\mu\nu\rho})^2 + \frac{1}{2}[\bar{\chi}^I \not{D}(\omega) \chi^I] - \frac{1}{2}(D_\mu \varphi^I)^2 - \frac{1}{4}(F_{\mu\nu}^I)^2 + \frac{1}{2}[\bar{\lambda}^I \not{D} \lambda^I] \\
& - \frac{1}{4}(H_{\mu\nu}^I)^2 + \frac{1}{2}[\bar{\rho}^I \not{D}(\omega) \rho^I] - g(\bar{\chi}^I \rho^I) - \frac{1}{2}g^2(\varphi^I)^2 - \frac{1}{2}g f^{IJK}(\bar{\lambda}^I \chi^J) \varphi^K - \frac{1}{4}f^{IJK}(\bar{\lambda}^I \gamma^{\mu\nu} \chi^J) H_{\mu\nu}^K \\
& + \frac{1}{12}f^{IJK}(\bar{\lambda}^I \gamma^{\mu\nu\rho} \rho^J) G_{\mu\nu\rho}^K + \frac{1}{4}f^{IJK}(\bar{\rho}^I \gamma^\mu \chi^J) F_{\mu\nu}^K - \frac{1}{2}f^{IJK} F_{\mu\nu}^I H^{\mu\nu J} \varphi^K + \frac{1}{2}f^{IJK}(\bar{\lambda}^I \gamma^{\mu\nu} \rho^J) D_\mu \varphi^K \\
& + (\bar{\psi}_\mu \gamma^\nu \gamma^\mu \chi^I) D_\nu \varphi^I + \frac{1}{6}(\bar{\psi}_\mu \gamma^{\rho\sigma\tau} \gamma^\mu \chi^I) G_{\rho\sigma\tau}^I - \frac{1}{2}(\bar{\psi}_\mu \gamma^{\rho\sigma} \gamma^\mu \lambda^I) F_{\rho\sigma}^I - \frac{1}{2}(\bar{\psi}_\mu \gamma^{\rho\sigma} \gamma^\mu \rho^I) H_{\rho\sigma}^I - g(\bar{\psi}_\mu \gamma^\mu \rho^I) \varphi^I,
\end{aligned} \tag{5.1}$$

up to $\mathcal{O}(\phi^4)$ terms.

Our action \tilde{I} is now invariant under local $N = 1$ supersymmetry

$$\delta_Q e_\mu^m = -2(\bar{\epsilon} \gamma^m \psi_\mu), \tag{5.2a}$$

$$\delta_Q \psi_\mu = +D_\mu(\hat{\omega})\epsilon - \frac{1}{6}(\gamma_\mu^{\rho\sigma\tau} \epsilon) \hat{G}_{\rho\sigma\tau}^I \varphi^I, \tag{5.2b}$$

$$\delta_Q B_{\mu\nu}^I = +(\bar{\epsilon} \gamma_{\mu\nu} \chi^I) - 2f^{IJK} C_{[\mu}^J (\delta_Q A_{\nu]}^K) - 4(\bar{\epsilon} \gamma_{[\mu} \psi_{\nu]}) \varphi^I, \tag{5.2c}$$

$$\delta_Q \chi^I = +\frac{1}{6}(\gamma^{\mu\nu\rho} \epsilon) \hat{G}_{\mu\nu\rho}^I - (\gamma^\mu \epsilon) \hat{D}_\mu \varphi^I + \frac{1}{2}f^{IJK} [+\epsilon(\bar{\lambda}^J \rho^K) - (\gamma_5 \gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_5 \gamma_\mu \rho^K) - (\gamma_5 \epsilon)(\bar{\lambda}^J \gamma_5 \rho^K)], \tag{5.2d}$$

$$\delta_Q \varphi^I = +(\bar{\epsilon} \chi^I), \tag{5.2e}$$

$$\delta_Q C_\mu^I = +(\bar{\epsilon} \gamma_\mu \rho^I) + f^{IJK}(\bar{\epsilon} \gamma_\mu \lambda^J) \varphi^K, \tag{5.2f}$$

$$\begin{aligned}
\delta_Q \rho^I = & +\frac{1}{2}(\gamma^{\mu\nu} \epsilon) \hat{H}_{\mu\nu}^I - g\epsilon \varphi^I - \frac{1}{2}f^{IJK}(\gamma^{\mu\nu} \epsilon) \hat{F}_{\mu\nu}^J \varphi^K + \frac{1}{4}f^{IJK} [+\epsilon(\bar{\lambda}^J \chi^K) - (\gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_\mu \chi^K) \\
& + \frac{1}{2}(\gamma^{\mu\nu} \epsilon)(\bar{\lambda}^J \gamma_{\mu\nu} \chi^K) - (\gamma_5 \gamma^\mu \epsilon)(\bar{\lambda}^J \gamma_5 \gamma_\mu \chi^K) - (\gamma_5 \epsilon)(\bar{\lambda}^J \gamma_5 \chi^K)],
\end{aligned} \tag{5.2g}$$

$$\delta_Q A_\mu^I = +(\bar{\epsilon} \gamma_\mu \lambda^I), \tag{5.2h}$$

$$\delta_Q \lambda^I = +\frac{1}{2}(\gamma^{\mu\nu} \epsilon) \hat{F}_{\mu\nu}^I + \frac{1}{2}f^{IJK}(\gamma_5 \epsilon)(\bar{\rho}^J \gamma_5 \chi^K), \tag{5.2i}$$

up to $\mathcal{O}(\phi^3)$ terms. The supercovariant field strengths are defined as usual in supergravity [8] by

$$\hat{F}_{\mu\nu}^I \equiv +2\partial_{[\mu} A_{\nu]}^I + g f^{IJK} A_\mu^J A_\nu^K - 2(\bar{\psi}_{[\mu} \gamma_{\nu]} \lambda^I) = F_{\mu\nu}^I - 2(\bar{\psi}_{[\mu} \gamma_{\nu]} \lambda^I), \tag{5.3a}$$

$$\begin{aligned}
\hat{G}_{\mu\nu\rho}^I \equiv & +3D_{[\mu} B_{\nu\rho]}^I - 3f^{IJK} C_{[\mu}^J F_{\nu\rho]}^K - 3(\bar{\psi}_{[\mu} \gamma_{\nu\rho]} \chi^I) + 6(\bar{\psi}_{[\mu} \gamma_{\nu]} \gamma_{\rho]} \psi_{|\rho|}) \varphi^I \\
& + G_{\mu\nu\rho}^I - 3(\bar{\psi}_{[\mu} \gamma_{\nu\rho]} \chi^I) + 6(\bar{\psi}_{[\mu} \gamma_{\nu]} \gamma_{\rho]} \psi_{|\rho|}) \varphi^I,
\end{aligned} \tag{5.3b}$$

$$\hat{H}_{\mu\nu}^I \equiv +2D_{[\mu} C_{\nu]}^I + g B_{\mu\nu}^I - 2(\bar{\psi}_{[\mu} \gamma_{\nu]} \rho^I) = H_{\mu\nu}^I - 2(\bar{\psi}_{[\mu} \gamma_{\nu]} \rho^I), \tag{5.3c}$$

$$\hat{D}_\mu \varphi^I \equiv +D_\mu \varphi^I - (\bar{\psi}_\mu \chi^I). \tag{5.3d}$$

Certain remarks are in order. First, the last term in Eq. (5.1) of the type $g(\bar{\psi}\gamma\rho)\varphi$ is related to the φ -linear term in $\delta_Q\rho$ in Eq. (5.2g). Second, the $\delta_Q B_{\mu\nu}$ contains the $(\bar{\epsilon}\gamma\psi)\varphi$ -term. This is consistent with $G_{\alpha\beta c}{}^I = +2(\gamma_c)_{\alpha\beta}\varphi^I$ in (3.2a) in superspace. Third, for the $g\psi\rho\chi$ -terms, we need nontrivial Fierz rearrangement. To be more specific, there are three contributions to this sector: (i) $g(\bar{\psi}\gamma\rho)\varphi$, (ii) $ge(\bar{\chi}\rho)$, and (iii) $(\bar{\psi}\gamma\gamma\rho)H$ -terms. This rearrangement is highly nontrivial, showing the consistency of our total system.

As the couplings to supergravity in Eq. (5.1) show, our original *globally* supersymmetric system shares certain features with supergravity, such as fermionic bilinear terms. Such terms are common in supergravity [8] but *not* in conventional global supersymmetry. Our original *global* system already possessed the feature of *local* $N = 1$ supersymmetry. As was mentioned following Eq. (2.2), the conventional dimensional analysis tells that such terms imply nonrenormalizability. In other words, our *globally* supersymmetric system already had a hidden gravitational constant κ providing negative mass dimension. In a sense, this feature resembles σ -models with nonrenormalizable couplings, sharing certain features with gravity interactions.

VI. CONCLUDING REMARKS

In this paper, we have carried out the $N = 1$ supersymmetrization in 4D of a non-Abelian tensor with consistent couplings, as a special case [6] of the minimal tensor hierarchy discussed in Ref. [5], which is further a special case of more general hierarchy in Refs. [2,3]. We have given both the component and superspace formulations of our system, providing the nontrivial consistency of our system. Our CVM ($C_{\mu}{}^I, \rho^I$) plays the role of a Stueckelberg [7] compensator multiplet, being absorbed into the TM ($B_{\mu\nu}{}^I, \chi^I, \varphi^I$), making the latter massive.

We have also generalized the adjoint-representation case to the general real representation for $G = SO(N)$. The action invariance works in a fashion parallel to the former. We foresee no obstruction against generalizing these results further to the complex representation of, e.g., $G = SU(N)$ group. Finally, we have also coupled the global $N = 1$ system to $N = 1$ supergravity up to quartic terms. This has provided a nontrivial confirmation for the total consistency of the non-Abelian TM.

It has been known that certain problem exists in the quantization of the Stueckelberg model [7] for non-Abelian gauge groups [9]. The common problem is that the longitudinal components of the gauge field do not decouple from the physical Hilbert space, upsetting the renormalizability and unitarity of the system [9]. For this issue, we clarify our standpoints as follows: First of all, our theory is *not* renormalizable from the outset due to Pauli couplings. Our theory makes stronger sense when couplings to supergravity are also taken into account, as we have done in Sec. V. Moreover, there are certain theories in

4D, such as nonlinear sigma models, which are not renormalizable but are not excluded from the outset. So we do not go into the renormalizability issue in this paper. Second, thanks to $N = 1$ supersymmetry, our system has a good chance to have a better quantum behavior compared with nonsupersymmetric systems.

As shown in the Appendix, the purely bosonic part of our system can be generalized to arbitrary space-time dimensions with arbitrary signatures. The key ingredient is the tensor $B_{\mu_1\cdots\mu_{p+1}}{}^I$ and a Stueckelberg-type [7] compensator $C_{\mu_1\cdots\mu_p}{}^I$.

The potential importance of the result in this paper is $N = 1$ supersymmetry that has better quantum behavior compared with nonsupersymmetric cases. We have presented a new *supersymmetric* physical system with Stueckelberg mechanism that solves both the problem with non-Abelian tensor and the problem with extra vector fields in the non-singlet representation of a non-Abelian gauge group.

ACKNOWLEDGMENTS

This work is supported in part by Department of Energy grant DE-FG02-10ER41693.

APPENDIX: HIGHER-DIMENSIONAL APPLICATION OF PURELY BOSONIC SYSTEM

In this appendix, we generalize the purely bosonic part of our system in 4D into arbitrary space-time dimensions with arbitrary signatures. We also apply it to the case of tensor-vector duality in 6D, and perform a DR to 4D. Our field content is $(A_{\mu}{}^I, B_{[n-1]}{}^I, C_{[n-2]}{}^I)$.⁸

We generalize the definitions of field strengths Eqs. (1.3a) and (1.3b) to arbitrary space-time dimension D as

$$G_{\mu_1\cdots\mu_n}{}^I \equiv +nD_{[\mu_1}B_{\mu_2\cdots\mu_n]}{}^I - \frac{n(n-1)}{2} \times f^{IJK}C_{[\mu_1\cdots\mu_{n-2}}{}^J F_{\mu_{n-1}\mu_n]}{}^K, \quad (A1a)$$

$$H_{\mu_1\cdots\mu_{n-1}}{}^I \equiv +(n-1)D_{[\mu_1}C_{\mu_2\cdots\mu_{n-1}]}{}^I + gB_{\mu_1\cdots\mu_{n-1}}{}^I. \quad (A1b)$$

The YM field strength F is the same as in Eq. (5.3a). The BIDs for these field strengths are

$$D_{[\mu}F_{\nu\rho]}{}^I \equiv 0, \quad (A2a)$$

$$D_{[\mu_1}G_{\mu_2\cdots\mu_{n+1}]}{}^I \equiv +\frac{n}{2}f^{IJK}F_{[\mu_1\mu_2]}{}^J H_{[\mu_3\cdots\mu_{n+1}]}{}^K, \quad (A2b)$$

$$D_{[\mu_1}H_{\mu_2\cdots\mu_n]}{}^I \equiv +\frac{1}{n}gG_{\mu_1\cdots\mu_n}{}^I. \quad (A2c)$$

The α -, β -, and γ -transformations for $A_{\mu}{}^I, B_{[n-1]}{}^I$, and $C_{[n-2]}{}^I$ are the generalizations of our 4D case:

⁸We use symbols like $[n]$ for totally antisymmetric indices $\mu_1\mu_2\cdots\mu_n$ in order to save space.

$$\delta_\alpha(A_{\mu}^I, B_{[n-1]}^I, C_{[n-2]}^I) = (D_\mu \alpha^I, -g f^{IJK} \alpha^J B_{[n-1]}^K, -g f^{IJK} \alpha^J C_{[n-2]}^K), \quad (\text{A3a})$$

$$\delta_\alpha(F_{\mu\nu}^I, G_{[n]}^I, H_{[n-1]}^I) = -g f^{IJK} \alpha^J (F_{\mu\nu}^K, G_{[n]}^K, H_{[n-1]}^K), \quad (\text{A3b})$$

$$\delta_\beta B_{\mu_1 \dots \mu_{n-1}}^I = +(n-1) D_{[\mu_1} \beta_{\mu_2 \dots \mu_{n-1}]}^I, \quad \delta_\beta A_\mu^I = 0, \quad (\text{A3c})$$

$$\delta_\beta C_{\mu_1 \dots \mu_{n-2}}^I = -g \beta_{\mu_1 \dots \mu_{n-2}}^I, \quad (\text{A3d})$$

$$\delta_\beta (F_{\mu\nu}^I, G_{[n-1]}^I, H_{[n-2]}^I) = 0, \quad (\text{A3e})$$

$$\delta_\gamma C_{\mu_1 \dots \mu_{n-2}}^I = +(n-2) D_{[\mu_1} \gamma_{\mu_2 \dots \mu_{n-2}]}^I, \quad \delta_\gamma A_\mu^I = 0, \quad (\text{A3f})$$

$$\delta_\gamma B_{\mu_1 \dots \mu_{n-1}}^I = + \frac{(n-1)(n-2)}{2} f^{IJK} \gamma_{[\mu_1 \dots \mu_{n-3}]}^J F_{[\mu_{n-2} \mu_{n-1}]}^K, \quad (\text{A3g})$$

$$\delta_\gamma (F_{\mu\nu}^I, G_{[n-1]}^I, H_{[n-2]}^I) = 0. \quad (\text{A3h})$$

Eq. (A3d) shows that the C -field is a Stueckelberg field absorbed into the longitudinal components of the B -field.

A typical action $I'' \equiv \int d^D x g^2 \mathcal{L}''$ is given by the Lagrangian

$$\mathcal{L}'' = -\frac{1}{2(n!)} (G_{[n]}^I)^2 - \frac{1}{2 \cdot (n-1)!} (H_{[n-1]}^I)^2 - \frac{1}{4} (F_{\mu\nu}^I)^2, \quad (\text{A4})$$

yielding the B - and C -field equations

$$\frac{\delta \mathcal{L}}{\delta B_{[n-1]}^I} = \frac{1}{(n-1)!} (D_\mu G^{\mu[n-1]I} - g H^{[n-1]I}) \doteq 0, \quad (\text{A5a})$$

$$\begin{aligned} \frac{\delta \mathcal{L}}{\delta C_{[n-2]}^I} &= \frac{1}{(n-2)!} (D_\nu H^{\nu[n-2]I} \\ &+ \frac{1}{2} f^{IJK} F_{\rho\sigma}^J G^{[n-2]\rho\sigma K}) \doteq 0. \end{aligned} \quad (\text{A5b})$$

As in the 4D case, it is straightforward to show the consistency

$$0 \stackrel{?}{=} D_\mu \left(\frac{\delta \mathcal{L}}{\delta B_{\mu[n-2]}^I} \right) \equiv -\frac{I}{n-I} g \left(\frac{\delta \mathcal{L}}{\delta C_{[n-2]}^I} \right) \doteq 0, \quad (\text{A6a})$$

$$\begin{aligned} 0 \stackrel{?}{=} D_\mu \left(\frac{\delta \mathcal{L}}{\delta C_{\mu[n-3]}^I} \right) &\equiv + \frac{n-I}{2} f^{IJK} F_{\rho\sigma}^J \left(\frac{\delta \mathcal{L}}{\delta B_{[n-3]\rho\sigma}^K} \right) \\ &\doteq 0 \quad (\text{Q.E.D.}). \end{aligned} \quad (\text{A6b})$$

We next apply our result to 6D with the signature $(-, -, +, +, +, +)$, and consider the duality condition

$$\begin{aligned} F_{\mu\nu}^I &\stackrel{*}{=} + \frac{1}{24} \epsilon_{\mu\nu}{}^{\rho\sigma\tau\lambda} G_{\rho\sigma\tau\lambda}^I, \\ G_{\mu\nu\rho\sigma}^I &\stackrel{*}{=} + \frac{1}{2} \epsilon_{\mu\nu\rho\sigma}{}^{\tau\lambda} F_{\tau\lambda}^I. \end{aligned} \quad (\text{A7})$$

This duality looks similar to Eq. (3.6) in Ref. [5], but the existence of the physical scalar field ϕ^I in the latter makes the fundamental difference.

We have first to confirm the consistency of Eq. (A7) with the G and H -BIDs. First, the rotation of the second equation in Eq. (A7) gives

$$\begin{aligned} 0 &\stackrel{?}{=} + \epsilon^{\mu\nu\rho\sigma\tau\lambda} D_\nu (G_{\rho\sigma\tau\lambda}^I - \frac{1}{2} \epsilon_{\rho\sigma\tau\lambda}{}^{\omega\psi} F_{\omega\psi}^I) \\ &\equiv + \epsilon^{\mu\nu\rho\sigma\tau\lambda} (2 f^{IJK} F_{\nu\rho}^J H_{\sigma\tau\lambda}^K) - 24 D_\nu F^{\mu\nu I} \\ &= -24 (D_\nu F^{\mu\nu I} - \frac{1}{12} \epsilon^{\mu\nu\rho\sigma\tau\lambda} f^{IJK} F_{\nu\rho}^J H_{\sigma\tau\lambda}^K). \end{aligned} \quad (\text{A8})$$

In the second identity in Eq. (A8), we have used the G -BID Eq. (A2b). The first term in the last line is the kinetic term of A_μ^I , so that its last term is its source term. Second, in order to see if Eq. (A8) has consistent solutions, we can confirm the conservation of the source term by applying D_μ on Eq. (A8) based on H -BID Eqs. (A2c) and (A7), but we skip the details here.

We next show that the usual self-duality relationship in $D = 2 + 2$

$$F_{\mu\nu}^I \stackrel{*}{=} + \frac{1}{2} \epsilon_{\mu\nu}{}^{\rho\sigma} F_{\rho\sigma}^I \quad (\text{A9})$$

is embedded into Eq. (A7). To this end, we use hat symbols both on fields and indices in 6D and no hats on 4D quantities from now on. We also use $\hat{\mu}, \hat{\nu}, \dots = 1, 2, 3, 4, 5, 6$ and $\mu, \nu, \dots = 1, 2, 3, 4$, while $\alpha, \beta, \dots = 5, 6$. Our basic ansätze for the DR are

$$\hat{G}_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}}^I \stackrel{*}{=} + \hat{F}_{[\hat{\mu}\hat{\nu}}^I \hat{P}_{\hat{\rho}\hat{\sigma}]}, \quad \hat{P}_{\hat{\mu}\hat{\nu}} \equiv + \hat{\delta}_{\hat{\mu}} \hat{X}_{\hat{\nu}} - \hat{\delta}_{\hat{\nu}} \hat{X}_{\hat{\mu}}, \quad \hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}^I \stackrel{*}{=} + \frac{1}{2} g \hat{F}_{[\hat{\mu}\hat{\nu}}^I \hat{X}_{\hat{\rho}]}, \quad (\text{A10a})$$

$$\hat{P}_{\hat{\mu}\hat{\nu}} = \epsilon_{\alpha\beta} \quad (\text{for } \hat{\mu} = \alpha, \hat{\nu} = \beta), \quad \hat{F}_{\hat{\mu}\hat{\nu}}^I = \hat{F}_{\mu\nu}^I = F_{\mu\nu}^I \quad (\text{for } \hat{\mu} = \mu, \hat{\nu} = \nu), \quad (\text{A10b})$$

$$\hat{\epsilon}_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}\hat{\tau}\hat{\lambda}} = \hat{\epsilon}_{\mu\nu\rho\sigma\alpha\beta} = \epsilon_{\mu\nu\rho\sigma} \epsilon^{\alpha\beta} \quad (\text{for } [\hat{\mu}\hat{\nu}\hat{\rho}\hat{\sigma}\hat{\tau}\hat{\lambda}] = [\mu\nu\rho\sigma\alpha\beta]). \quad (\text{A10c})$$

Other components, such as $\hat{P}_{\mu\beta}$, are all zero. We can confirm that Eqs. (A10) are consistent with the BIDs Eqs. (A2b) and (A2c). It is easy to show that the $[\alpha\beta]$ and $[\mu\alpha]$ components of the first equation in Eq. (A7) are satisfied, while the $[\mu\nu]$

component gives directly the 4D self-duality Eq. (A9). Thus the 4D self-duality $F = {}^* \tilde{F}$ is indeed embedded in the 6D duality Eq. (A7).

We next generalize the 6D result to the $D = 2m + 2$ with the signature $(-, -, \overbrace{+, \dots, +}^{2m})$. The duality condition Eq. (A7) is generalized to

$$\hat{F}_{\hat{\mu} \hat{\nu}}^I = + \frac{1}{(2m)!} \hat{\epsilon}_{\hat{\mu} \hat{\nu} \hat{\rho}_1 \dots \hat{\rho}_{2m}} \hat{G}_{\hat{\rho}_1 \dots \hat{\rho}_{2m}}^I, \quad \hat{G}_{\hat{\rho}_1 \dots \hat{\rho}_{2m}}^I = + \frac{1}{2} \hat{\epsilon}_{\hat{\rho}_1 \dots \hat{\rho}_{2m}}^{\hat{\mu} \hat{\nu}} \hat{F}_{\hat{\mu} \hat{\nu}}^I. \quad (\text{A11})$$

As in the 6D case, we can first confirm the consistency with BIDs. We can next confirm the current conservation, whose details are skipped here.

The previous ansätze for 6D case in Eq. (A10) are generalized to

$$\hat{G}_{\hat{\mu}_1 \dots \hat{\mu}_{2m}}^I = + c \hat{F}_{[\hat{\mu}_1 \hat{\mu}_2]}^I \hat{P}_{|\hat{\mu}_3 \hat{\mu}_4|}^{(1)} \dots \hat{P}_{|\hat{\mu}_{2m-1} \hat{\mu}_{2m}|}^{(m-1)}, \quad \hat{P}_{\hat{\mu} \hat{\nu}}^{(k)} \equiv \hat{\partial}_{\hat{\mu}} \hat{X}_{\hat{\nu}}^{(k)} - \hat{\partial}_{\hat{\nu}} \hat{X}_{\hat{\mu}}^{(k)}, \quad (\text{A12a})$$

$$\hat{H}_{\hat{\mu}_1 \dots \hat{\mu}_{2m-1}}^I = + \frac{1}{m} c g \hat{F}_{[\hat{\mu}_1 \hat{\mu}_2]}^I \hat{P}_{|\hat{\mu}_3 \hat{\mu}_4|}^{(1)} \dots \hat{P}_{|\hat{\mu}_{2m-3} \hat{\mu}_{2m-2}|}^{(m-2)} \hat{X}_{|\hat{\mu}_{2m-1}|}, \quad (\text{A12b})$$

$$\hat{P}_{\hat{\mu} \hat{\nu}}^{(k)} = \hat{P}_{2k+3, 2k+4}^{(k)} = -\hat{P}_{2k+4, 2k+3}^{(k)} = \epsilon_{2k+3, 2k+4}^{(k)} = -\epsilon_{2k+4, 2k+3}^{(k)} = +1 \quad (\text{for } \hat{\mu} = 2k + 3, \hat{\nu} = 2k + 4; k = 1, \dots, m - 1), \quad (\text{A12c})$$

$$\hat{F}_{\hat{\mu} \hat{\nu}}^I = F_{\mu\nu}^I \quad (\text{for } \hat{\mu} = \mu, \hat{\nu} = \nu), \quad (\text{A12d})$$

$$\hat{\epsilon}_{\hat{\mu}_1 \dots \hat{\mu}_{2m+2}} = \epsilon^{\mu\nu\rho\sigma} \epsilon^{\alpha_1 \dots \alpha_{2m-2}} = \epsilon^{\mu\nu\rho\sigma} \epsilon_{(1)}^{[\alpha_1 \alpha_2]} \dots \epsilon_{(m-1)}^{[\alpha_{2m-3} \alpha_{2m-2}]} \quad (\text{for } [\hat{\mu}_1 \dots \hat{\mu}_{2m+2}] = [\mu\nu\rho\sigma\alpha_1 \dots \alpha_{2m-2}]). \quad (\text{A12e})$$

where c is a constant to be fixed later.

As before, we can also confirm the G - and H -BIDs for Eq. (A11). The constant c in Eq. (A12a) is fixed by getting the 4D self-duality in the $[\mu\nu]$ component of the first equation in Eq. (A11):

$$F_{\mu\nu}^I = + \frac{1}{(2m)!} \hat{\epsilon}_{\mu\nu \hat{\rho}_1 \dots \hat{\rho}_{2m}} \hat{G}_{\hat{\rho}_1 \dots \hat{\rho}_{2m}}^I = + \frac{\binom{2m}{2}}{(2m)!} \hat{\epsilon}_{\mu\nu}^{\rho\sigma\alpha_1 \dots \alpha_{2m-2}} \hat{G}_{\rho\sigma\alpha_1 \dots \alpha_{2m-2}}^I = + \frac{1}{2} c \left[\frac{1}{(m-1)! \cdot (2m-3)!!} \right]^2 \epsilon_{\mu\nu}^{\rho\sigma} F_{\rho\sigma}^I. \quad (\text{A13})$$

For this to agree with $F = {}^* \tilde{F}$, we get $c = [(m-1)! \cdot (2m-3)!!]^2$. The remaining components $[\alpha\beta]$ and $[\mu\alpha]$ are trivially satisfied.

The previous mechanism for $D = 2m + 2$ is further generalized to $D = 2m + 1$ with the signature

$(-, -, \overbrace{+, \dots, +}^{2m-1})$, with the duality condition

$$\hat{F}_{\hat{\mu} \hat{\nu}}^I = + \frac{1}{(2m-1)!} \hat{\epsilon}_{\hat{\mu} \hat{\nu} \hat{\rho}_1 \dots \hat{\rho}_{2m-1}} \hat{G}_{\hat{\rho}_1 \dots \hat{\rho}_{2m-1}}^I, \quad \hat{G}_{\hat{\rho}_1 \dots \hat{\rho}_{2m-1}}^I = + \frac{1}{2} \hat{\epsilon}_{\hat{\rho}_1 \dots \hat{\rho}_{2m-1}}^{\hat{\mu} \hat{\nu}} \hat{F}_{\hat{\mu} \hat{\nu}}^I. \quad (\text{A14})$$

The confirmation of G - and H -BIDs is just parallel to the $D = 2m + 2$ case. The ansätze for DR is

$$\hat{G}_{\hat{\mu}_1 \dots \hat{\mu}_{2m-1}}^I = + \frac{2c'}{3} \hat{F}_{[\hat{\mu}_1 \hat{\mu}_2]}^I \hat{P}_{|\hat{\mu}_3 \hat{\mu}_4|}^{(1)} \dots \hat{P}_{|\hat{\mu}_{2m-5} \hat{\mu}_{2m-4}|}^{(m-3)} \hat{Q}_{|\hat{\mu}_{2m-3} \hat{\mu}_{2m-2} \hat{\mu}_{2m-1}|}, \quad (\text{A15a})$$

$$\hat{H}_{\hat{\mu}_1 \dots \hat{\mu}_{2m-2}}^I = + \frac{2c'g}{2m-1} \hat{F}_{[\hat{\mu}_1 \hat{\mu}_2]}^I \hat{P}_{|\hat{\mu}_3 \hat{\mu}_4|}^{(1)} \dots \hat{P}_{|\hat{\mu}_{2m-5} \hat{\mu}_{2m-4}|}^{(m-3)} \hat{Y}_{|\hat{\mu}_{2m-3} \hat{\mu}_{2m-2}|}, \quad (\text{A15b})$$

$$\hat{P}_{\hat{\mu} \hat{\nu}}^{(k)} \equiv \hat{\partial}_{\hat{\mu}} \hat{X}_{\hat{\nu}}^{(k)} - \hat{\partial}_{\hat{\nu}} \hat{X}_{\hat{\mu}}^{(k)}, \quad \hat{Q}_{\hat{\mu} \hat{\nu} \hat{\rho}} \equiv + \hat{\partial}_{\hat{\mu}} \hat{Y}_{\hat{\nu} \hat{\rho}} + \hat{\partial}_{\hat{\nu}} \hat{Y}_{\hat{\rho} \hat{\mu}} + \hat{\partial}_{\hat{\rho}} \hat{Y}_{\hat{\mu} \hat{\nu}}, \quad (\text{A15c})$$

$$\hat{P}_{\hat{\mu} \hat{\nu}}^{(k)} = \hat{P}_{2k+3, 2k+4}^{(k)} = -\hat{P}_{2k+4, 2k+3}^{(k)} = \epsilon_{2k+3, 2k+4}^{(k)} = -\epsilon_{2k+4, 2k+3}^{(k)} = +1, \quad (\text{A15d})$$

$$\hat{Q}_{\hat{\mu} \hat{\nu} \hat{\rho}} = \hat{Q}_{2m-3, 2m-2, 2m-1} = \epsilon_{2m-3, 2m-2, 2m-1} = +1 \quad (\text{for } [\hat{\mu} \hat{\nu} \hat{\rho}] = [2m-3, 2m-2, 2m-1]), \quad (\text{A15e})$$

$$\hat{F}_{\hat{\mu} \hat{\nu}}^I = F_{\mu\nu}^I \quad (\text{for } \hat{\mu} = \mu, \hat{\nu} = \nu), \quad (\text{A15f})$$

$$\hat{\epsilon}_{\hat{\mu}_1 \dots \hat{\mu}_{2m+1}} = \epsilon^{\mu\nu\rho\sigma} \epsilon^{\alpha_1 \dots \alpha_{2m-3}} = \epsilon^{\mu\nu\rho\sigma} \epsilon_{(1)}^{[\alpha_1 \alpha_2]} \dots \epsilon_{(m-3)}^{[\alpha_{2m-7} \alpha_{2m-6}]} \epsilon_{(m-3)}^{[\alpha_{2m-5} \alpha_{2m-4} \alpha_{2m-3}]}. \quad (\text{A15g})$$

The totally antisymmetric constant tensor $\epsilon^{\alpha\beta\gamma}$ is for the last three coordinates in $D = 2m + 1$. The satisfaction of the duality Eq. (A14) fixes the constant $c' = [(m-3)! \cdot (2m-7)!!]^2$.

- [1] M. Henneaux, V.E. Lemes, C.A. Sasaki, S.P. Sorella, O.S. Ventura, and L.C. Vilar, *Phys. Lett. B* **410**, 195 (1997).
- [2] B. de Wit and H. Samtleben, *Fortschr. Phys.* **53**, 442 (2005).
- [3] B. de Wit, H. Nicolai, and H. Samtleben, *J. High Energy Phys.* **02** (2008) 044.
- [4] C. Hull and P.K. Townsend, *Nucl. Phys.* **B438**, 109 (1995); E. Witten, *Nucl. Phys.* **B443**, 85 (1995); P.K. Townsend, in *Proceedings of ICTP Summer School on High Energy Physics and Cosmology* (Cambridge University, Trieste, 1996); P.K. Townsend, “*M-Theory from Its Superalgebra*,” Cargese Lectures 1997; T. Banks, W. Fischler, S.H. Shenker, and L. Susskind, *Phys. Rev. D* **55**, 5112 (1997); K. Becker, M. Becker, and J.H. Schwarz, *String Theory and M-Theory: A Modern Introduction* (Cambridge University Press, Cambridge, UK, 2007).
- [5] H. Samtleben, E. Sezgin, and R. Wimmer, *J. High Energy Phys.* **12** (2011) 062.
- [6] C.-S. Chu, arXiv:1108.5131.
- [7] E.C.G. Stueckelberg, *Helv. Phys. Acta* **11**, 225 (1938), <http://inspirehep.net/record/45337>; A. Proca, *J. Phys. Radium* **7**, 347 (1936); See, e.g., R. Delbourgo and G. Thompson, *Phys. Rev. Lett.* **57**, 2610 (1986); D. Feldman, Z. Liu, and P. Nath, *Phys. Rev. Lett.* **97**, 021801 (2006); For reviews see, e.g., H. Ruegg and M. Ruiz-Altaba, *Int. J. Mod. Phys. A* **19**, 3265 (2004).
- [8] S. Ferrara, D.Z. Freedman, and P. van Nieuwenhuizen, *Phys. Rev. D* **13**, 3214 (1976); S. Deser and B. Zumino, *Phys. Lett. B* **62**, 335 (1976); P. van Nieuwenhuizen, *Phys. Rep.* **68**, 189 (1981); J. Wess and J. Bagger, *Superspace and Supergravity* (Princeton University Press, Princeton, 1992).
- [9] J.M. Kunimasa and T. Goto, *Prog. Theor. Phys.* **37**, 452 (1967); A.A. Slavnov, *Theor. Math. Phys.* **10**, 99 (1972); M.J.G. Veltman, *Nucl. Phys.* **B7**, 637 (1968); A.A. Slavnov and L.D. Faddeev, *Theor. Math. Phys.* **3**, 312 (1970); A.I. Vainshtein and I.B. Khriplovich, *Yad. Fiz.* **13**, 198 (1971), <http://inspirehep.net/record/69554>; K.I. Shizuya, *Nucl. Phys.* **B121**, 125 (1977); Y.N. Kafiev, *Nucl. Phys.* **B201**, 341 (1982).